tion of boracic acid was kept constantly applied on boracic lint, and the odour rapidly ceased. For the lotion an ointment recommended by Dr Thin was substituted. This ointment is made by dissolving boracic acid in glycerine to saturation, and then adding as much olive oil and white wax as will make a consistent ointment, or at least a thick cream. The foot steadily healed, and when it had been entirely covered with new epidermis for some time, and seemed in a condition to stand it, a spirit lotion containing a little glycerine and a very little liq. carbonis detergens was gently pencilled over it. Instead of hardening the skin, the tar set up fresh irritation, and had speedily to be laid aside, and the ointment, which we thought had done its work, resumed for some time longer.

This case shows how an unheathy state of the secretions may, if uncorrected, maintain the disease indefinitely, for many forms of treatment had been tried ere I saw the patient; and also that even weak preparations of tar cannot be used with advantage in all cases, though apparently well suited for them.

With these remarks I must close. I have merely touched superficially on many topics connected with this subject, but what I have said may serve to indicate the general line of treatment which should be adopted.

ARTICLE II.—*Note on the Anatomy of the Pia Mater.* By Dr J. BATTY TUKE.

(Read before the Medico-Chirurgical Society of Edinburgh, 3rd May 1882.)

IN the most recently published standard work on anatomy, Turner's *Introduction to Human Anatomy*, we find the membranes of the brain described as threefold—the dura mater, the arachnoid, and the pia mater. I take the liberty of quoting largely from this admirable work, and of criticising its statements, as it is certain to contain the most recent and the most generally received opinion of anatomists.

The dura mater is described as a tough fibrous membrane, the inner surface of which has the appearance of a mucous membrane, and is covered by a layer of squamous cells. "The dura mater is well provided with lymph vessels, which in all probability open by stomata on the free inner surface." The description of the membrane concludes thus:—"Between the dura mater and the arachnoid membrane is a space of only a capillary lumen, containing a minute quantity of limpid serum, which moistens the smooth inner surface of the dura, and the corresponding smooth outer surface of the arachnoid. It is regarded as equivalent to the cavity of a serous membrane, and is named the *arachnoid cavity*, or, more appropriately, the *sub-dural space*."

The *arachnoid* is described as "a non-vascular connective tissue membrane of great delicacy and transparency, which envelops both the brain and spinal cord. *It is separated from these organs by the pia mater; but between it and the latter membrane is a distinct space, called sub-arachnoid.*¹ The free surface next the sub-dural space is smooth and covered by a layer of squamous endothelium. The opposite surface is connected to the pia by threads and membranes of delicate sub-arachnoid connective tissue which traverse the sub-arachnoid space and divide it into numerous loculi freely communicating with each other. The connexion is much closer opposite the summits of the convolutions than opposite the sulci between them, for the arachnoid does not dip between the convolutions: it is loose also opposite the interpeduncular space and around the pons, medulla, and spinal cord, in which localities, therefore, the sub-arachnoid space is most distinct. . . . *Endothelial cells line the inner surface of the arachnoid, cover the outer surface of the pia, and invest the various bands and membranes which traverse the sub-arachnoid space.*"¹

Of the *pia mater* it is said, "This tender *vascular* membrane consists of delicate connective tissue closely investing the whole outer surface of the brain and cord. It dips into the fissures between the convolutions. . . . *The arteries of the brain and spinal cord ramify and divide into small branches in the pia mater and sub-arachnoid tissue*¹ before they penetrate the nervous substance, and the veins conveying the blood from the nerve centres traverse these membranes before they open into the blood sinuses of the cranial dura mater and into the extra-dural venous plexus of the spinal canal. . . . The vessels which pass from the pia mater into the brain and spinal cord are invested by a loose funnel-shaped sheath, which has been described as forming the wall of a perivascular lymphatic vessel; but Axel Key and Retzius consider that the space between the bloodvessel and the sheath opens into the sub-arachnoid space, and contains sub-arachnoid fluid. A network of lymph vessels ramifies freely in the pia mater."

If we reduce these statements to a diagrammatic scheme (Figure 1), we have, *1st*, the vascular dura mater; *2nd*, a sub-dural space; *3rd*, a non-vascular fibrous membrane, the arachnoid; *4th*, a sub-arachnoid space; and *5th*, a vascular membrane dipping into all the sulci.

Reference to the drawings illustrating this paper will, I think, convince the observer that a modification of this view of the membranes must be adopted. Certain of the sections from which these drawings were made were taken from the brains of monkeys recently killed, others from human brains. The organs were removed from the calvaria after careful reflection of the dura mater, all the other investments being left *in situ*, hardened in

¹ The italics are my own.

chromic acid, and thin sections were cut perpendicularly to the plane of the surface.

Study of these specimens shows that over the curved surface of each convolution a membrane exists investing it closely. This membrane, the pia mater, consists of two layers of a distinctly fibrous character, for the most part intimately bound together by connective tissue, so intimately as to be inseparable, and to give us every right to consider it as one membrane. *Between these two layers, and in the connective tissue, the vessels permeate. No membrane external to this exists except the dura mater.* But when this investment approaches a sulcus, the two layers separate, the outer bridging the fissure, the inner dipping into it and closely investing its surface throughout all its ramifications, being reflected at the extremities of the sulcus, and becoming continuous on emerging from it with the bridging layer. In the angular space thus formed the arteries lie naked, or at most are supported by very fine fibrous trabeculæ; but as they enter the brain substance they receive from the investing layer fine hyaline sheaths (Diagram 2).

It is true that over the sulci, over the interpeduncular recess, and over the wide gap between the cerebellum and the medulla oblongata, a purely fibrous membrane exists; but when we refer to microscopic sections an explanation of its true character is obtained. It is found to be the outer layer of the vascular pia mater, which in these positions is separated widely from the inner layer and bridges the fissures and spaces.¹ Over the cerebellum these two layers are much less intimately connected by interstitial connective tissue than over the cerebrum. Over the spinal cord the two layers are completely separated, the inner very closely investing the organ, the outer becoming thick and coarse, loosely enveloping the cord, and being separated from the inner layer by vessels and bundles of nerves as they enter or emerge. This thick, definite membrane is, however, distinctly continuous with the outer layer of the cerebral pia mater.

From these observations I am forced to the conclusion that a separate arachnoid membrane does not exist—that the name should be dispensed with, and, as a corollary, that the theory of the existence of a sub-arachnoid space should be departed from; that we should speak of only two membranes of the brain and spinal cord, the *dura mater* and the *pia mater*, and of two spaces, the *sub-dural space* and the *intra-pia-matral space or spaces*.

Slight consideration will show that something much more important than mere accuracy of anatomical description is involved in this contention, although that is in itself a not inconsiderable matter. The practically important question is, What relation do the pia-matral spaces hold to the vessels and lymphatics of the brain?

¹ It has been suggested to me by Mr Symington that an analogue of this arrangement is found in the *falx cerebri*, *falx cerebelli*, and *tentorium cerebelli*.

Between each convolution a pia-matral space exists, and it is easily demonstrable by injections that they communicate freely with one another on the surface of the cerebrum by wide channels. Over the cerebellum and at the base of the cerebrum they become larger, but not actually continuous. Around the spinal cord the pia-matral space is continuous. In my opening statement I said that the inner layer of pia mater affords fine hyaline sheaths to the arteries as they enter the brain. This opinion was grounded on the writings of Robin, His, Fohmann and Arnold, Key and Retzius, Boll, Rindfleisch, and on my own observations. I present to you specimens which show the sheath as it enters the cerebral substance (Fig. 4), and as it loosely encapsulates the vessel in the interior of the brain. Although the existence of these perivascular hyaline sheaths is now an acknowledged fact in anatomy, the manner in which they debouch on the surface of the brain is not clearly stated, in consequence, I believe, of the prevalence of the theory of the presence of an arachnoid membrane apart from the pia mater, and of a sub-arachnoid space. Boll held that they empty themselves into a space which he called the great epi-cerebral lymph space, and which he believed to exist between the pia mater and the cerebral substance. This position is quite untenable, for the hyaline membrane is a prolongation of the inner layer of pia mater; how, then, can the channels they form debouch *below* that membrane? Key and Retzius ridicule this idea, but assert that the hyaline perivascular sheaths debouch into the sub-arachnoid space. It is needless to reiterate my arguments in proof of the non-existence of such a cavity; but admitting its existence for a moment for the sake of argument, how are Key and Retzius to overcome the difficulty of the intervening vascular pia mater, which must, if their theory be correct, be penetrated by the sheaths before they could reach it?

The simple and plain fact is that these channels debouch along with the vessels into the cavities of the pia mater, as shown in Diagram No. 2.

It is now pretty generally held that these conduits are lymphatics. Key and Retzius assert that by their means free communication is obtained between the ventricles of the brain and its outer surface. The anastomosis between the arteries supplying the tracts around the lateral ventricles and those entering the outer surface renders this statement easily appreciable. Obersteiner holds that by certain "spur-like" processes there is maintained a lymphatic communication between spaces surrounding each nerve-cell and the perivascular lymph spaces. He has proved this by injections. But whether we regard them as lymph conduits, or as channels for the transference of cerebro-spinal fluid from the brain to the spinal canal, or as serving both purposes, the position I have asserted renders the route taken by

fluid discernible. No patent route is open if we hold to the usual statement of a vascular pia mater encapsulating the cerebrum, on which is superimposed a fibrous arachnoid, for the simple reason that any perivascular conduit must needs penetrate the pia mater before it could reach a space between that investment and an arachnoid membrane. But if we hold to a vascular pia mater with broad interconvolutional spaces which communicate freely one with the other and with a wide intra-pia-matral space in the spinal canal, the transference of fluid is easily comprehensible.

To the pathologist the question is of great importance, as it involves the possible consequences to brain health which may be brought about by the interruption of the route by distention of vessels.

DESCRIPTION OF PLATES.

Diagrams.

FIG. 1. Projection of membranes and spaces according to the usually accepted anatomical description. V. D. M., Vascular dura mater; S. D., Sub-dural space or sac of arachnoid; F., Fibrous non-vascular arachnoid; S. A., Sub-arachnoid space; V. P. M., Vascular pia mater, dipping as a whole into a sulcus.

FIG. 2. Projection of membranes and spaces as indicated in the paper. V. D. M., Vascular dura mater; S. D., Sub-dural space; V. P. M., Vascular pia mater, outer layer bridging a sulcus, inner layer investing surface of sulcus, and affording fine hyaline sheaths to the vessels; $\times \times$, A Pia-matral space, bounded by the inner layer investing a sulcus, and superiorly by the bridging layer: in it the vessels lie naked or receive support from trabeculae; into it the hyaline sheaths debouch.

Lithographs.

No. 1, taken from a section of a monkey's brain, shows pia mater as a two-layered membrane at *a b*, and *a b*, and behaving as stated above. The cut ends of vessels are seen supported by trabeculae; *xx*, Brain substance. $\times 80$.

No. 2, from a human brain, shows a vessel in the pia mater, on the summit of a convolution, sending a branch into the brain substance, surrounded by a hyaline sheath. The sheath is represented somewhat too strongly. The communicating pia-matral space is shown between the fibrous tissue of the two layers. $\times 250$.

ARTICLE III.—*On the Diagnosis and Prognosis of Idiocy and Imbecility.* By WILLIAM W. IRELAND, M.D., Home and School for Imbeciles, Preston Lodge, Prestonpans.

THE object of this paper is to point out those features of idiocy and imbecility which it is most important that the physician in ordinary practice should know. I have, therefore, no intention of going deeper into the pathology of the subject than what seems useful in illustrating my remarks or removing the possibility of erroneous ideas arising in the mind of the reader.

FIG. 1.

V. D. M.
S. D. Space
V. P. M.
V. P. M.

Brain Substance

FIG. 2.

V. D. M.
S. D. Space
V. P. M.
X X Pia mater
Space

2