

Frequent moving of grazing dairy cows to new paddocks increases the variability of milk fatty acid composition

M. Coppa¹, A. Farruggia^{2,3}, P. Ravaglia^{1,2,3}, D. Pomiès^{2,3,4}, G. Borreani¹, A. Le Morvan^{2,3} and A. Ferlay^{2,3†}

¹Department of Agricultural Forest and Food Sciences, University of Turin, Via L. da Vinci 44, 10095, Grugliasco, Italy; ²INRA, UMR1213 Herbivores, F-63122 Saint-Genès-Champanelle, France; ³UMR1213 Herbivores, Clermont Université, VetAgro Sup, BP 10448, F-63000 Clermont-Ferrand, France; ⁴INRA, UE1296 Monts d'Auvergne, F-63820 Laqueuille, France

(Received 21 October 2013; Accepted 16 October 2014; First published online 8 December 2014)

The aim of this work was to investigate the variations of milk fatty acid (FA) composition because of changing paddocks in two different rotational grazing systems. A total of nine Holstein and nine Montbéliarde cows were divided into two equivalent groups according to milk yield, fat and protein contents and calving date, and were allocated to the following two grazing systems: a long duration (LD; 17 days) of paddock utilisation on a heterogeneous pasture and a medium duration (MD) of paddock utilisation (7 to 10 days) on a more intensively managed pasture. The MD cows were supplemented with 4 kg of concentrate/cow per day. Grazing selection was characterised through direct observations and simulated bites, collected at the beginning and at the end of the utilisation of two subsequent MD paddocks, and at the same dates for the LD system. Individual milks were sampled the first 3 days and the last 2 days of grazing on each MD paddock, and simultaneously also for the LD system. Changes in milk FA composition at the beginning of each paddock utilisation were highly affected by the herbage characteristics. Abrupt changes in MD milk FA composition were observed 1 day after the cows were moved to a new paddock. The MD cows grazed by layers from the bottom layers of the previous paddock to the top layers of the subsequent new paddock, resulting in bites with high organic matter digestibility (OMD) value and CP content and a low fibre content at the beginning of each paddock utilisation. These changes could induce significant day-to-day variations of the milk FA composition. The milk fat proportions of 16:0, saturated FA and branched-chain FA decreased, whereas proportions of de novo-synthesised FA, 18:0, c9-18:1 and 18:2n-6 increased at paddock change. During LD plot utilisation, the heterogeneity of the vegetation allowed the cows to select vegetative patches with higher proportion of leaves, CP content, OMD value and the lowest fibre content. These small changes in CP, NDF and ADF contents of LD herbage and in OMD values, from the beginning to the end of the experiment, could minimally modify the ruminal ecosystem, production of precursors of de novo-synthesised FA and ruminal biohydrogenation, and could induce only small day-to-day variations in the milk FA composition.

Keywords: grazing system, grazing selection, milk fatty acids, upland pasture, dairy cow

Implications

As cow milk fat proportions of fatty acids (FA) affect cheese characteristics (i.e. texture and appearance), considerable FA variations could make farmhouse cheese quality difficult to standardise. Therefore, it seems preferable to reduce variations of milk FA composition when milk fat proportions of specific FA are considered for milk payment. Thus, the identification of factors (e.g. moving of dairy cows to new paddocks, grazing selection) that influence variations of milk fat proportions of FA would be useful for farmers to control farmhouse cheese quality, and therefore milk payment.

Introduction

In upland dairy farming systems, considerable day-to-day variations of milk yield and quality are frequently reported by farmers and cheese makers, resulting in difficult management of dairy product quality for producers. These variations can be attributed to daily changes in milk fat and protein contents and fatty acid (FA) composition. Among these quality parameters, FA composition of milk fat plays an important role for human nutrition (Kratz *et al.*, 2013) and concurs to determine butter spreadability, cheese texture and sensory notes of dairy products (Martin *et al.*, 2005; Hurtaud *et al.*, 2010; Coppa *et al.*, 2011b). However, the origin of these day-to-day FA variations is not well understood,

† E-mail: anne.ferlay@clermont.inra.fr

resulting in difficult management of dairy product quality for producers.

The FA composition of pasture-derived milk is affected by several factors. Herbage allowance exerts an effect through the composition of herbage consumed (Dewhurst *et al.*, 2006). The total FA and 18:3n-3 contents of grass decreased with the advancement of herbage phenological stage (Chilliard *et al.*, 2007). Different phenological stages induce variations in the herbage chemical composition (i.e. CP and fibre contents), influencing the ruminal biohydrogenation of polyunsaturated fatty acid (PUFA; Gerson *et al.*, 1986). The effect of the botanical composition of upland pasture is due to specific secondary plant ingredients abundant in dicotyledons, inhibiting ruminal bacteria implied in the ruminal biohydrogenation of PUFA (Willems *et al.*, 2014), resulting in larger amounts of PUFA available in the duodenum and then in milk (Leiber *et al.*, 2005). These different factors could interact with cow grazing selection, which can be managed through the grazing system applied (Coppa *et al.*, 2011a).

In the upland dairy cow farming systems of central Europe, rotational grazing systems differed mainly in stocking density and duration of pasture utilisation (Sollemberg and Newman, 2007). Under low stocking density, cows have unlimited and interrupted access to an extensively managed plot for several weeks. By contrast, at higher stocking density, cows usually move to new paddocks quite frequently and have access to a more stable herbage allowance and chemical composition during the season (Sollemberg and Newman, 2007; Farruggia *et al.*, 2014). Vlaeminck *et al.* (2010) showed important variations of the milk FA composition from cows on intensively managed lowland grasslands with a short duration of paddock utilisation (4 days). Moreover, high variations of milk yield and milk FA composition have also been reported with continuous grazing systems, according to the grazing period over the year (Farruggia *et al.*, 2014). However, little information is still available with regard to the day-to-day variations of milk fat proportions of FA at paddock changes, according to the rotational grazing management.

This work aimed to investigate the variations of milk FA composition induced by moving grazing dairy cows to new paddocks with different frequencies. We explored that question within the grazing system and studied two experimental rotational grazing systems, differing by the duration of the rotation, the stocking rate and the type of vegetation.

Material and methods

Experimental design and animal management

The experiment was conducted at the INRA experimental farm of Marcenat in an upland area of Cantal, central France (latitude 45°15'N, longitude 2°55'E; altitude 1135 to 1215 m; annual rainfall of 1100 mm). The experiment presented in this paper represents a focus made within a main long-term 'system experiment' (still in progress), described by Pomiès *et al.* (2013), consisting of two different low-input pasture-based production systems with different input levels. Concerning the

'system experiment', a total of 24 Holstein and 24 Montbéliarde spring-calving cows were divided into two equivalent groups and were turned out to pasture at the end of April on two different grazing managements: LD characterised by a long duration of paddock utilisation (15 to 20 days) on a highly biodiversified pasture and not fertilised and MD characterised by a medium duration of paddock utilisation (7 to 10 days) on a moderately biodiversified and more intensively managed pasture (about 150 kg/ha of ammonium nitrate per year).

After calving, a sub-group of five Holstein and five Montbéliarde multiparous cows from each experimental system was selected according to individual milk yield, milk protein and fat contents, calving date, BW and parity to create two equivalent groups of 10 cows for the experiment presented in this paper. All the selected cows were at the beginning of the lactation (39 days in milk (DIM); standard deviation: 10.8). Two cows (one per each system) had a case of hypocalcaemia at pasture before the experiment, and thus they were removed. Before the experimental period, from 21 April to 21 May for the MD cows and from 27 April to 21 May for the LD cows, the animals grazed on similar grassland pastures according to the experimental grazing management. The experiment was carried out for 17 days (from 21 May to 6 June) in 2011, aiming to graze the pasture at the first seasonal grazing cycle. During the experimental period, the values for daily temperature and rainfall were on average 11.8°C (minimal and maximal values = 6.1°C and 17.6°C/day) and 5.8 mm/day (minimal and maximal values = 1.2 and 10.9 mm/day, with cumulate rainfall during on 17 days = 46.2 mm), respectively. During this period, the LD group grazed a 9.6-ha permanent mesotrophic pasture at a moderate stocking density (2.5 livestock unit (LU)/ha; 1 LU = 600 kg BW/ha), having a high botanical diversity (139 botanical species) and a marked structural heterogeneity of vegetation. The LD cows were fed only this herbage. The MD group grazed a 6.4-ha moderately biodiversified pasture at a higher stocking density (3.75 LU/ha), which was managed to offer cows leafy edible biomass throughout the grazing season. The MD pasture area was divided into two paddocks that were used consecutively. The change between Paddock 1 to Paddock 2 was managed on the basis of the daily milk yield of the group: cows were moved when the milk yield decreased by about 10% of the average of the three consecutive maximum daily milk yields achieved on the paddock. The MD cows were fed herbage and received 4 kg/cow per day of concentrate (ingredients expressed as % of the concentrate dry matter (DM): 38.8% wheat flour, 32.6% barley flour, 24.1% corn flour, 3% cane molasses and 1.5% carbonate-binding agent; FA composition expressed as g/100 g total FA: 12:0, 0.05; 14:0, 0.25; 16:0, 21.48; 18:0, 1.80; c9-18:1, 18.48; 18:2n-6, 54.39; 18:3n-3, 2.46; 20:0, 0.23; 22:0, 0.31; and 24:0, 0.54). The MD cows grazed for 9 days on Paddock 1 (from days 1 to 9) and 8 days on Paddock 2 (from days 10 to 17). The LD cows entered into the plot when the MD cows started grazing Paddock 1, and the LD cows grazed on the plot throughout the duration of the experiment (17 days).

Table 1 Botanical composition expressed as proportion of total ground cover on the different paddocks grazed during the experiment

LD Paddock ¹		MD Paddock 1 ¹		MD Paddock 2 ¹	
Botanical group	%	Botanical group	%	Botanical group	%
Grasses	41.2	Grasses	50.5	Grasses	73.7
Legumes	16.3	Legumes	18.3	Legumes	10.8
Non-legume forbs	42.5	Non-legume forbs	31.2	Non-legume forbs	15.5
Main species	%	Main species	%	Main species	%
<i>Agrostis capillaris</i>	17.2	<i>Trifolium repens</i>	17.5	<i>Agrostis capillaris</i>	15.0
<i>Trifolium repens</i>	12.5	<i>Lolium perenne</i>	14.2	<i>Dactylis glomerata</i>	15.0
<i>Festuca gr. rubra</i>	11.3	<i>Festuca gr. rubra</i>	10.4	<i>Festuca gr. rubra</i>	14.2
<i>Achillea millefolium</i>	5.2	<i>Agrostis capillaris</i>	8.3	<i>Trifolium repens</i>	10.8
<i>Plantago lanceolata</i>	4.6	<i>Taraxacum gr. officinale</i>	8.1	<i>Festuca pratensis</i>	10.0
<i>Veronica chamaedrys</i>	3.4	<i>Poa trivialis</i>	5.0	<i>Cynosurus cristatus</i>	3.3
<i>Carex caryophylla</i>	2.7	<i>Cynosurus cristatus</i>	4.2	<i>Holcus mollis</i>	3.3
<i>Poa pratensis</i>	2.6	<i>Dactylis glomerata</i>	3.8	<i>Phleum pratense</i>	3.3
<i>Helianthemum nummularium</i>	2.4	<i>Ranunculus acris</i>	2.9	<i>Poa pratensis</i>	3.3
<i>Dactylis glomerata</i>	2.1	<i>Hypochoeris radicata</i>	1.8	<i>Arrhenatherum elatius</i>	2.5

¹LD = long duration of heterogeneous pasture utilisation; MD = medium duration of paddock utilisation on intensively managed pastures.

The botanical composition of the vegetation (Table 1) was determined before the experiment by 40 and 33 surveys (Braun-Blanquet, 1932) on the LD and MD pastures, respectively. The herbage availability was estimated using the sward-cutting method (Smith *et al.*, 2005) the day before the cows started grazing on each paddock. The herbage availability was 0.93 t/DM per ha on LD and 1.45 and 1.29 t/DM per ha on MD Paddocks 1 and 2, respectively. The average maximum herbage height, measured by an electronic plate meter (30 × 30 cm², 4.5 kg/m²; Urban and Caudal, 1990), was 9.3 ± 0.3 cm on LD when cows entered into the paddock and 7.3 ± 0.4 cm at the end of the experiment, whereas it was 16.3 ± 0.2 and 12.4 ± 0.2 cm at the beginning and 10.8 ± 0.2 and 8.3 ± 0.2 cm at the end of Paddock 1 and Paddock 2 utilisation for the MD system, respectively.

Cow grazing selection measurements

Aiming to relate the cow diet composition to the milk FA composition, the grazing selection of each cow was characterised by direct observations and by simulated bites sampled at the beginning (days 1 and 10) and at the end (days 8 and 17) of each MD paddock utilisation, and at the same dates for LD.

Direct observations of grazing selection

All the grazing cows were observed continuously and individually during days 1, 8, 10 and 17 of each MD paddock utilisation, and at the same dates for the LD group, by an observer standing within <5 m at the side of the target animal (Farruggia *et al.*, 2008). The observer recorded the composition and structure of every fifth bite grazed by the target cow. The patches were classified as follows: short vegetative patches (SV), when patch height was <7 cm and the herbage was at vegetative stage; tall vegetative patches (TV), when patch height was >7 cm and the herbage was at vegetative stage; and tall mature patches (TM) when patch

height was >7 cm and the herbage was at mature stage (Farruggia *et al.*, 2008).

Simulated bites

Once having observed the composition and structure of each target cow bite, the observer collected a 'simulated bite': a sward patch similar to that of the bites grazed by the target cow, considering herbage height, phenological stage and botanical composition (Farruggia *et al.*, 2008; Verheyden-Tixier *et al.*, 2008). Simulated bites per cow were sampled several times during the day of observation (5 to 10 simulated bites in the morning and 5 to 10 simulated bites in the afternoon). The simulated bites were immediately stored at 4°C. The different bite samples of each cow collected at the same days of sampling were pooled to constitute one sample, representative of the herbage grazed daily by the cow. Each pooled sample was then divided into three homogeneous sub-samples. Two sub-samples were stored at -20°C until the analysis of chemical composition and FA composition, respectively. The third sub-sample was used to determine the botanical group composition and grass leaf-to-stem ratio (L/S). The botanical group composition was determined by hand separating the plants into groups of grasses, legumes and non-legume forbs. Each botanical group was weighed after oven drying at 60°C for 72 h to determine the respective proportions on DM basis. Before drying, 100 plants from the group of grasses were randomly selected and divided by hand into stems and leaves and then weighed after oven drying to determine the respective proportions (on DM basis) of the two fractions.

Herbage analyses

For the determination of the chemical composition, a sub-sample of simulated bites was dried at 60°C for 24 h and analysed for total nitrogen content by combustion (LECO method (Sweeney and Rexroad, 1987)), NDF and ADF

contents (Van Soest *et al.*, 1991) and pepsin-cellulase solubility (Aufrère and Michalet-Doreau, 1983) as a measure of the organic matter digestibility (OMD). The sub-samples of simulated bites were lyophilised (ThermovacTM-20, Froilabo, Ozoir-la-Ferriere, France), ground and analysed for FA composition as described by Ferlay *et al.* (2010).

Milk sampling and analyses

The individual milk yield was recorded at each milking (days 1 to 17). Individual milk samples were collected at the beginning of each MD paddock utilisation for 3 consecutive days (days 1, 2, 3 and 10, 11, 12 for Paddocks 1 and 2, respectively) and at the end of each MD paddock utilisation for the last 2 days (days 9, 10 and 16, 17 for Paddocks 1 and 2, respectively). Milk samples from the LD cows were collected on the same dates as the MD cows. A sub-sample was preserved with bronopol-B2, stored at 4°C and analysed for fat and protein contents by standard procedures (MilkoScan 4000, Foss System, Hillerød, Denmark, AOAC, 1997). Another sub-sample was stored at -20°C until the FA analysis. After having lyophilised (Thermovac TM-20; Froilabo S.A., Ozoir-la-Ferriere, France), the milk samples were methylated and analysed according to Ferlay *et al.* (2010). The FA methyl esters (FAME) were injected (0.6 µl) by an auto-sampler into a gas chromatograph equipped with a flame ionisation detector (Agilent Technologies 7890A, Wilmington, DE, USA). The FAMEs from all the samples were separated on a 100 m × 0.25 mm i.d. fused-silica capillary column (CP-Sil 88; Chrompack, Middelburg, The Netherlands). The injector temperature was maintained at 255°C and the detector temperature at 260°C. The initial oven temperature was held at 70°C for 1 min, increased to 100°C at a rate of 5°C/min (held for 2 min), then increased by 10°C/min to 175°C (held for 42 min) and 5°C/min to a final temperature of 225°C (held for 15 min). The carrier gas was hydrogen and the pressure was maintained constant (158.6 kPa) during analysis. Peaks were routinely identified by comparing retention times against commercial authentic standards. The FAME proportions were corrected to FA proportions according to their respective molecular weight.

Statistical analyses

Statistical analyses were performed using the SPSS for Windows software package (Version 16.0; SPSS Inc., Chicago, IL, USA). One cow from the LD system was removed from the statistical analyses because of severe digestive problems. This animal was fed hay just before the experiment. The milk FA composition was affected by the hay-based diet, resulting in milk samples not representative of the grazing system. For each grazing system, the data of grazing selection, simulated bites and milk yield and composition were analysed using the repeated measures model of ANOVA, with the day as the repeated factor. For data of the MD system, the paddock was used as a fixed factor and the paddock × day interaction was also tested. In the first step of data exploration, the statistical model included the day as the repeated factor, the breed effect as a fixed factor and the DIM were used as covariate. As the covariate and the breed effect were never

significant, a new repeated measures model of ANOVA was tested with only the day as the repeated factor. A principal component analysis (PCA) was performed on the main milk FA, parameters of grazing selection and composition of simulated bites, aiming to show the relationships between the milk fat proportions of FA and grazing selection of cows in the two grazing systems. As the literature data showed a 1-day delay in the response of milk fat proportions of FA to paddock changes (Khanal *et al.*, 2008; Vlaeminck *et al.*, 2010; Coppa *et al.*, 2012), to perform the PCA, data of cow grazing selection were associated with FA composition of milk sampled the day after the grazing selection measurements.

Results

Milk FA composition

The results of the proportions of the complete FA profiles for LD and MD milk samples are given in Supplementary Tables S1 and S2, respectively.

Rapid changes in milk FA composition were observed within the first 2 days after moving cows to a new paddock on both systems (Figure 1). A change from day 1 to day 2 was observed in the LD milk fat proportions of *de novo*-synthesised FA, saturated fatty acids (SFA) and monounsaturated fatty acids (MUFA), as well as the τ 10-18:1/ τ 11-18:1 ratio (-2.2, -2.7, +2.3/100 g and -0.02, respectively). The proportions of almost all FA in the LD milk were stable after day 2 (Figure 1), as in the case of *de novo*-synthesised FA and MUFA, whereas the *c*9, τ 11-CLA and *trans* FA proportions slightly decreased until day 17. The 16:0, *c*9-18:1, 18:2n-6 and SFA proportions and the 18:2n-6/18:3n-3 ratio slightly increased from day 3 to day 17 in the LD pasture (Figure 1).

The MD milk fat proportions of almost all FA showed significant changes at paddock moving (Figure 1). The MD milk fat proportions of *de novo*-synthesised FA, *c*9, τ 11-CLA and SFA decreased from day 3 to day 9 for both paddocks, but important changes were observed for day 9 of Paddock 1 and day 2 of Paddock 2 (Figure 1). The MD milk fat proportion of 16:0 followed the same trend. The MD milk fat proportions of *c*9-18:1 and MUFA and 18:2n-6/18:3n-3 and *c*9-18:1 *cis*9/16:0 ratios increased from day 2 to day 9 of each paddock utilisation, showing significant changes with an opposite trend when cows were moved from Paddock 1 to Paddock 2 (Figure 1).

Cow grazing selection

Direct observations of grazing selection. The LD cows selected a similar proportion of TV and SV patches all along the LD paddock utilisation, except the last days for which the proportion of SV patches decreased in favour of that of TV patches (Table 2). The selection of TM patches by LD cows was higher during the last week of utilisation than during the 1st week.

The MD cows selected more SV patches at the end than at the beginning of each paddock utilisation (Table 2). On the contrary, the selection of TV patches was lower at the end than at the beginning of Paddock 2 utilisation.




Figure 1 Changes in milk fatty acid composition¹ during utilisation of the medium duration (MD) and long duration (LD) paddocks. FA = fatty acid; SFA = sum of saturated fatty acids (from 4:0 to 24:0, including odd- and branched-chain fatty acids); OCFA = sum of odd-chain fatty acid (from 5:0 to 23:0); BCFA = sum of branched-chain fatty acids (from iso-13:0 to iso-18:0); MUFA = sum of monounsaturated fatty acid (from c9-10:1 to c11-20:1); PUFA = sum of polyunsaturated fatty acids (from t9-12-18:2 to 22:6n-3). In MD values with different lowercase letters differed at $P < 0.05$ for day effect; P = paddock effect; d = day effect; l = paddock x day interaction effect; † $P < 0.10$; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; when the significance is not given $P > 0.10$. In LD values with different lowercase letters differed at $P < 0.05$ for day effect; † $P < 0.10$; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; when the significance symbol is not given $P > 0.10$.

Simulated bites. Grasses were the dominant botanical group in the simulated bites of both the MD and LD cows (>85% of DM), whereas legumes were poorly represented (Table 3).

No significant differences in the botanical composition of simulated bites were observed among days during the experiment for both the grazing systems. The L/S ratio for LD-simulated

Table 2 Characteristics of grazed patches of the long (LD) or medium (MD) duration paddocks, obtained by direct observations of cow grazing selection

LD	d1	d8	d10	d16	s.e.m.	Significance ¹			
Short vegetative patches	48.3 ^b	47.7 ^b	41.6 ^b	75.5 ^a	1.36	***			
Tall vegetative patches	43.1 ^a	37.5 ^a	34.0 ^a	2.0 ^b	1.19	***			
Tall mature patches	8.6 ^b	14.8 ^b	24.4 ^a	22.5 ^a	0.80	*			
MD	Paddock 1		Paddock 2		s.e.m.	Significance ¹			
	d1	d8	d10	d16		P	d	Int	
Short vegetative patches	12.6 ^c	26.8 ^b	16.7 ^c	63.8 ^a	1.41	**	***	**	
Tall vegetative patches	51.7 ^b	48.6 ^b	74.3 ^a	14.9 ^c	1.66	ns	***	***	
Tall mature patches	35.8 ^a	24.6 ^{ab}	9.0 ^b	21.3 ^{ab}	0.86	**	ns	**	

^{a,b,c}Values within a row with different superscripts differ significantly at $P < 0.05$.

¹P = paddock; d = day; Int = paddock × day interaction.

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

Table 3 Botanical composition, chemical composition, oleic, linoleic and linolenic acid proportions of simulated bites of the long (LD) or medium (MD) paddocks

Simulated bites composition	LD						MD				s.e.m.	Effect and significance ¹			
	Paddock 1		Paddock 2						P	d		Int			
	d1	d8	d10	d16	d1	d8	d1	d8							
Botanical composition (% of DM)															
Grasses	90.6	86.8	93.7	94.0	0.92	†	92.3	94.0	88.8	96.9	1.00	ns	†	ns	
Legumes	2.1	1.7	0.8	0.9	0.26	ns	0.1	0.2	2.1	0.2	0.38	ns	ns	ns	
Non-legume forbs	7.3	11.5	5.4	5.1	0.77	†	7.7	5.8	9.1	2.9	0.87	ns	†	ns	
Nutritive value															
L/S	1.01 ^a	0.48 ^b	0.54 ^b	0.51 ^b	0.11	**	0.56	0.31	0.73	0.33	0.07	†	*	ns	
DM (% of fresh herbage)	28.9 ^b	29.6 ^b	32.3 ^a	32.2 ^a	0.47	*	22.3	22.7	22.8	24.7	0.49	ns	†	ns	
CP (% of DM)	15.5 ^a	14.6 ^{ab}	13.9 ^{ab}	12.7 ^b	0.26	***	13.5 ^{ab}	12.0 ^b	17.1 ^a	11.7 ^b	0.45	**	***	**	
OMD (%)	67.6 ^a	63.5 ^{ab}	63.3 ^{ab}	59.5 ^b	0.62	***	71.2 ^b	64.9 ^c	76.2 ^a	62.1 ^c	1.18	ns	***	**	
NDF (% of DM)	55.9 ^c	57.3 ^{bc}	58.8 ^{ab}	60.1 ^a	0.40	***	53.0 ^b	56.1 ^a	50.6 ^b	57.6 ^a	0.64	ns	***	*	
ADF (% of DM)	25.1 ^c	26.7 ^{bc}	27.4 ^{ab}	28.6 ^a	0.28	***	25.0	27.5	23.0	28.2	0.48	ns	***	†	
FA composition (g/100 g FA)															
c9-18:1	2.52	2.74	2.47	2.85	0.12	ns	3.95	4.69	2.27	2.69	0.27	**	†	ns	
18:2n-6	13.5	14.4	14.5	14.5	0.18	ns	15.3	15.1	14.1	15.3	0.18	†	ns	†	
18:3n-3	49.0 ^a	47.6 ^{ab}	45.6 ^b	45.0 ^b	0.44	**	47.0	41.1	51.5	43.7	0.82	**	***	ns	

DM = dry matter; OMD = organic matter digestibility; L/S = grass leaf-to-stem ratio.

^{a,b,c}Values within a row with different superscripts differ significantly at $P < 0.05$.

¹P = paddock; d = day; int = paddock × day interaction.

† $P < 0.10$; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

bites decreased after the 1st day, and then it remained constant. The L/S for MD-simulated bites decreased from the beginning to the end of each paddock utilisation. The CP content and OMD value of the LD-simulated bites decreased slightly from the beginning to the end of the experiment, while the NDF and ADF contents increased. The CP content and OMD value of the MD-simulated bites increased when moving cows from Paddock 1 to Paddock 2, whereas the opposite trend was observed for the NDF and ADF contents. The 18:3n-3 proportion of the LD- or MD-simulated bites decreased slightly from the beginning to the end of each

paddock utilisation, with an important increase at paddock change. The 18:2n-6 proportion of the LD- or MD-simulated bites did not change during the experimental period. Only the c9-18:1 proportion of MD-simulated bites increased during the utilisation of Paddock 1.

Dairy cow performance

From the beginning to the end of the experiment, the LD daily milk yield decreased regularly, whereas the MD value decreased more dramatically, but with a recovery of 2 kg/day between the two paddocks (Table 4). The milk protein

Table 4 Milk production and composition of cows offered the long (LD) or medium (MD) duration paddocks

Milk yield and composition	LD										s.e.m.	Significance ¹
	d1	d2	d3	d8	d9	d10	d11	d12	d16	d17		
Milk yield (kg/day)	28.2 ^a	28.7 ^a	28.1 ^a	26.0 ^{ab}	27.1 ^{ab}	25.5 ^b	25.6 ^b	25.6 ^b	23.5 ^c	24.2 ^{bc}	0.40	***
Fat (g/kg)	39.6	41.3	39.3	39.2	37.1	35.2	38.9	37.5	36.9	37.9	0.56	†
Protein (g/kg)	33.4 ^a	32.3 ^{ab}	32.2 ^{ab}	30.9 ^{ab}	30.1 ^b	30.4 ^{ab}	30.9 ^{ab}	30.5 ^{ab}	29.9 ^b	29.3 ^b	0.24	***

Milk yield and composition	MD										s.e.m.	Effect and significance		
	Paddock 1					Paddock 2						P	d	Int
	d1	d2	d3	d8	d9	d1	d2	d3	d7	d8				
Milk yield (kg/day)	34.1 ^a	33.5 ^a	32.2 ^{ab}	28.6 ^{bc}	29.1 ^b	28.4 ^{bc}	30.4 ^b	29.6 ^b	27.1 ^{bc}	26.6 ^c	0.66	ns	***	*
Fat content (g/kg)	34.8	35.7	34.7	35.8	34.9	35.7	36.9	36.6	35.8	33.6	0.53	ns	ns	ns
Protein content (g/kg)	31.6	30.4	30.4	29.4	29.0	30.0	30.0	29.9	28.5	28.0	0.22	ns	***	ns

^{a,b,c}Values within a row with different superscripts differ significantly at $P < 0.05$.

¹ P = paddock; d = day; Int = interaction day × paddock; ns = not significant. † $P < 0.10$; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

content decreased from the beginning to the end of the experiment in both systems, whereas the milk fat content tended to decrease ($P < 0.10$) for LD and was stable for the MD system (Table 4).

Relationships between grazing selection and milk FA composition

On the PCA plot of individual distribution, samples were separated according to the day of paddock utilisation for principal component 1 (PC1 scores increasing from the beginning to the end of the utilisation) and according to the grazing system for both PC1 and PC2 (Figure 2). On PCA plot of variable distribution, PC1 (27.4% of variance) was positively and very closely correlated to the DM and NDF contents for the simulated bites, SV patches and milk fat proportions of *cis* isomers of 18:1 and MUFA and *c9*-18:1/16:0 ratio (Figure 3; correlation coefficients with PC1 > 0.64). The PC1 was also negatively and closely correlated to the OMD value of simulated bites, total TV patches, milk fat proportions of SFA and *de novo*-synthesised FA (Figure 3; correlation coefficients with PC1 < -0.66). The PC2 (20.3% of variance) was correlated to the CP content, OMD value, the proportion of legumes and non-legume forbs, the L/S of simulated bites and milk fat proportions of *c9*, *t11*-CLA, *trans* isomers of 18:1 and PUFA (Figure 3; correlation coefficients with PC2 > 0.44), and negatively correlated to the NDF content and grasses proportion of simulated bites, TM patches and milk 18:2n-6/18:3n-3 and *t10*-18:1/*t11*-18:1 ratios (Figure 3; correlation coefficients with PC2 < -0.40).

Discussion

The original result of this study is the broad day-to-day changes in milk FA composition when dairy cows moved to new paddocks in relationship with the changes in herbage composition and grazing selection of the cows.




Figure 2 Principal component analysis (PCA) performed on the main milk fatty acid proportions, the grazing selection and composition of simulated bites. Plot of sample distribution of the medium (MD) and long (LD) grazing systems and day of paddock utilisation, projected on the two principal components (PC1 and PC2). LD = long duration of paddock utilisation on heterogeneous pasture; MD = medium duration of paddock utilisation on intensively managed pastures; b = beginning; m = middle; e = end; MD 1 and MD 2 indicated the Paddocks 1 and 2, respectively.

An abrupt change in milk FA composition in the MD milk was highlighted when cows were moved to Paddock 2, changing their selection from the bottom layers of the previous paddock (rich in short patches and with a high proportion of stems and fibre content) to the top vegetative layers of the new paddock rich in leaves. The PCA confirms this marked separation between the samples derived from the beginning to the end of each MD paddock utilisation. These findings are supported by our results on the simulated bites. Indeed, the MD-simulated bites at the beginning of each paddock utilisation were characterised by a higher CP content and OMD value than at the end of utilisation. The high L/S and the selection of TV patches support the hypothesis that cows grazed 'by layers' (Teague and




Figure 3 Principal component analysis (PCA) performed on the main milk fatty acid (FA) proportions, the grazing selection and composition of simulated bites³. Plot of variables distribution projected on the two principal components (PC1 and PC2). BCFA = branched-chain fatty acids; C18:1c = *cis*-18:1; C18:1t = *trans*-18:1; CLAc9t11 = *c9,t11*-18:2; *de novo* FA = *de novo*-synthesised FA; linoleic acid/ α -linolenic acid (LA/ALA) = 18:2n-6/18:3n-3 ratio; MUFA = monounsaturated fatty acids; OA/PA = *c9*-18:1/16:0 ratio; PUFA = polyunsaturated fatty acids; SFA = saturated fatty acids; t10/t11 = t10-18:1/t11-18:1 ratio. TM = tall mature patches; TV = tall vegetative patches; SV = short vegetative patches. Chemical composition and species proportion of stimulated bites: DM = dry matter; F = non-legume forbs proportion; G = grasses proportion; L = legumes proportion; L/S = grass leaf-to-stem ratio; OMD = organic matter digestibility; SB-ALA: 18:3n-3 proportion of stimulated bites; SB-LA: 18:2n-6 proportion of stimulated bites.

Dowhower, 2003; Coppa *et al.*, 2011a). The SV patches were selected as the paddock utilisation progressed. Consequently, the quality of consumed patches rapidly decreased, whereas the proportion of stems increased, mainly located in the lower sward layers, as shown by Abrahamse *et al.* (2008). Thus, a rapid change in the quality and composition of grazed patches is in agreement with a shift from the lower level of a paddock to the upper layer of the following one. The increase in the selection of short patches with high DM proportion during paddock utilisation was found much more important with increasing stocking density and utilisation rate (Abrahamse *et al.*, 2008).

This grazing 'by layers' causes a rapid increase in the nutritional quality of ingested patches from the end to the beginning of paddock utilisation at paddock change (Abrahamse *et al.*, 2008). As a consequence, the dietary readily available energy supply increased and could have favoured mammary lipogenesis, as suggested by the higher milk fat proportions of *de novo*-synthesised FA obtained in our study and in agreement with Dewhurst *et al.* (2006). The increase in the 18:3n-3 proportion of simulated bites could have increased the ruminal production of intermediate products, including t11-18:1, and thus their proportions in milk fat (Chilliard *et al.*, 2007). Moreover, due to a lower proportion of stems, the reduced herbage NDF and ADF contents at the beginning of MD paddock utilisation at paddock change could disfavour the development of cellulolytic bacteria in the rumen. As these bacteria are implicated in the production

of branched-chain fatty acid (BCFA) (Vlaeminck *et al.*, 2006), this can explain the lower milk fat proportion of BCFA at changing MD paddock. Furthermore, the *c9*-18:1/16:0 ratio varied rapidly about 1.0 to 0.7 at MD paddock change. Couvreur *et al.* (2006) showed that a decrease in *c9*-18:1/16:0 ratio below 0.8 was sufficient to change thermal characteristics of butter fat, and a decrease below 0.9 made perceptible in mouth a softer butter texture. In addition, the variations of SFA and MUFA proportions, linked to dairy product texture, were large (differences of about 6/100 g FA in SFA and MUFA proportions).

Such extent of FA variations of the MD milk at paddock change was quite unexpected, considering that each cow additionally received 4 kg of concentrate/day. Indeed, the marked effect of concentrate intake was a high milk fat proportion of 18:2n-6 for the MD cows (on average 1.03 v. 1.35/100 g FA for LD and MD groups, respectively), in agreement with Chilliard *et al.* (2007). The concentrate intake did not change the milk fat proportions of 18:3n-3 between the two groups (0.74 v. 0.77/100 g FA for LD and MD groups, respectively). Furthermore, the extent of our variations was similar to those of Vlaeminck *et al.* (2010), obtained with cows grazing on temporary lowland grassland, although in this experiment cows were supplemented only with 2.4 kg of concentrate/day. The range of variations at MD paddock change was also similar to those observed by Coppa *et al.* (2011c) from the beginning to the end of a grazing season. These authors showed significant effects of variations of FA proportions across the season on cheese texture and appearance.

Regarding the MD milk production, the cyclic and consistent decrease in milk yield during each MD paddock utilisation and the increase at paddock change were already observed by Hoden *et al.* (1986) and by Farruggia *et al.* (2014), and are consistent with the evolution of the quality of patches grazed by layers.

The FA proportions in LD milk were stable from day 3, after the adaptation of the cows to the characteristics of the herbage of the new paddock. The milk fat proportions of *de novo*-synthesised FA, BCFA and MUFA (including *c9*-18:1, *cis*-18:1 and *trans*-18:1) changed slightly between the beginning to the end of LD paddock utilisation. This slight differentiation was also confirmed by the poor separation of these milk samples on the PCA axes (Figure 2).

The free-grazing selection allowed the LD cows to ingest herbage with fibre and CP contents and OMD values that were quite constant throughout the experiment (Dumont *et al.*, 2007; Coppa *et al.*, 2011a). The overgrazing of preferred patches was also observed on continuous grazing systems at the beginning of the season (Coppa *et al.*, 2011a).

The biodiversity and heterogeneity in species distribution of the vegetation of LD pasture can result in patches with different palatabilities and nutritive value (Dumont *et al.*, 2007; Coppa *et al.*, 2011a; Farruggia *et al.*, 2014), allowing cows to select vegetative patches with high proportion of leaves, CP content and OMD value and low NDF and ADF contents, in agreement with our results. As the LD pasture

utilisation progressed, the cows selected less TV patches in favour of the SV and TM patches that became dominant at the end of utilisation. The decrease in CP content and OMD value of the simulated bites and the increase in the NDF and ADF contents appear to be contradictory to the increase in the number of short patches at vegetative stage. However, the overgrazing of the preferred and already partially consumed patches – so-called ‘patch grazing’ (Adler *et al.*, 2001), characteristic of extensive grazing management – could easily explain this apparent contradiction. The short patches, although at a vegetative stage, contained grasses with a high proportion of stems and with higher NDF and ADF contents (Coppa *et al.*, 2011a).

The low selection of legumes and non-legume forbs by grazing cows may be because of the small size of legumes and non-legume forbs, compared with the tall dominant grass, which could limit their prehension by grazing cows (Carpino *et al.*, 2003). This result could explain the small changes in milk fat proportions of n-3 FA when compared with other studies in upland pastures (Collomb *et al.*, 2002; Leiber *et al.*, 2005; Coppa *et al.*, 2011c), because legumes are rich in 18:3n-3 (Chilliard *et al.*, 2007). However, in those studies on upland pasture, the cow grazing selection and the chemical composition, phenology and leafiness of selected patches were poorly investigated.

The moderate decrease observed in milk yield for LD is in agreement with the high herbage allowance and the selection of high nutritive value patches by grazing cows through patch grazing (Adler *et al.*, 2001; Farruggia *et al.*, 2014). Further investigation would be required to determine whether this rapid variation in milk yield would also be observed at LD paddock changes. Indeed, the LD system could be sensitive in late season, when the regrowth of high-quality patches is limited (i.e. by the dryness), and the cows have been obliged to also graze the less-preferred patches, with a consequent decrease in milk yield (Coppa *et al.*, 2011a; Farruggia *et al.*, 2014).

In conclusion, our study highlighted important variations of milk FA composition at each paddock change. These variations could be related to the characteristics of the vegetation grazed by the cows. The extent of milk FA variations could be frequent in intensive grazing systems and sufficient to influence cheese sensory properties such as texture and appearance. Furthermore, these milk FA variations could be difficult to evaluate and manage for farmers who deliver milk to dairy plants in the case of milk payment based on FA proportions, which is occurring in some European countries (Borreani *et al.*, 2013). The variations observed in our experiment could be a criticism if milk payment based on proportions of FA will be extensively introduced. In this case, adaptation to rotational grazing systems should be tested, aiming to reduce these variations, such as moving cows to a new paddock before the end of herbage utilisation and introduce other livestock categories to finish the utilisation. Applying different grazing systems in several geographical areas, with various vegetation types, and studying day-to-day variations of milk FA composition all along the grazing

season would be needed to give farmers valuable tools to manage these variations.

Acknowledgements

This work was funded by the transversal programme on dairy systems of the INRA-UMR 1213 Herbivores. The authors also thank D. Bany and E. Tixier for the determination of the FA composition of the stimulated bites and milk, respectively.

Supplementary material

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/S1751731114003000>.

References

- Abrahamse PA, Dijkstra J, Vlaeminck B and Tamminga S 2008. Frequent allocation of rotationally grazed dairy cows changes grazing behavior and improves productivity. *Journal of Dairy Science* 91, 2033–2045.
- Adler PB, Raff DA and Lauenroth WK 2001. The effect of grazing on the spatial heterogeneity of vegetation. *Oecologia* 128, 465–479.
- AOAC 1997. Official methods of analysis, 16th edition. Association of Official Analytical Chemists, Gaithersburg, MD, USA.
- Aufrère J and Michalet-Doreau B 1983. In vivo digestibility and prediction of digestibility of some by-products. In Feeding value of by-products and their use by beef cattle – EEC seminar EUR 8918 EN (ed. CV Boucqué, LO Fimes and GB Cottyn), pp. 25–35. Commission of the European Union, Brussels, Luxembourg.
- Borreani G, Coppa M, Revello-Chion A, Comino L, Giaccone D, Ferlay A and Tabacco E 2013. Effect of different feeding strategies in intensive dairy farming systems on milk fatty acid profiles, and implications on feeding costs in Italy. *Journal of Dairy Science* 96, 6840–6855.
- Braun-Blanquet J 1932. Plant sociology: the study of plant communities (ed. GD Fuller, HS Conard), pp. 438. McGraw-Hill book company Inc., New York, NY, USA.
- Carpino S, Licita G and Van Soest PJ 2003. Selection of forage species by dairy cattle on complex Sicilian pasture. *Animal Feed Science and Technology* 105, 205–214.
- Chilliard Y, Glasser F, Ferlay A, Bernard L, Rouel J and Doreau M 2007. Diet, rumen biohydrogenation and nutritional quality of cow and goat milk fat. *European Journal of Lipid Science and Technology* 109, 828–855.
- Collomb M, Bütikofer U, Sieber R, Jeangros B and Bosset JO 2002. Correlations between fatty acids in cows’ milk fat produced in the lowland, mountain and highlands of Switzerland and botanical composition of the fodder. *International Dairy Journal* 12, 661–666.
- Coppa M, Farruggia A, Pradel P, Lombardi G and Martin B 2011a. An improved grazed class method to estimate species selection and dry matter intake by cows at pasture. *Italian Journal of Animal Science* 10, 58–65.
- Coppa M, Gorlier A, Lonati M, Martin B, Russo EM and Lombardi G 2012. The management of the transition from hay- to pasture-based diets affects milk fatty acid kinetics. *Dairy Science and Technology* 92, 279–295.
- Coppa M, Verdier-Metz I, Ferlay A, Pradel P, Didiene R, Farruggia A, Montel MC and Martin B 2011b. Effect of different grazing systems on upland pastures compared with hay diet on cheese sensory properties evaluated at different ripening times. *International Dairy Journal* 21, 815–822.
- Coppa M, Ferlay A, Monsallier F, Verdier-Metz I, Pradel P, Didiene R, Farruggia A, Montel MC and Martin B 2011c. Milk fatty acid composition and cheese texture and appearance from cows fed hay or different grazing systems on upland pastures. *Journal of Dairy Science* 94, 1132–1145.
- Couvreur S, Hurtaud C, Lopez C, Delaby L and Peyraud JL 2006. The linear relationship between the proportion of fresh grass in the cow diet, milk fatty acid composition, and butter properties. *Journal of Dairy Science* 89, 1956–1969.
- Dewhurst RJ, Shingfield KJ, Lee MRF and Scollan ND 2006. Increasing the concentrations of beneficial polyunsaturated fatty acids in milk produced by dairy cows in high-forage systems. *Animal Feed Science and Technology* 131, 168–206.

- Dumont B, Garel JP, Ginane C, Decuq F, Farruggia A, Pradel P, Rigolot C and Petit M 2007. Effect of cattle grazing a species-rich mountain pasture under different stocking rates on the dynamics of diet selection and sward structure. *Animal* 1, 1042–1052.
- Farruggia A, Dumont B, D'Hour P and Egal D 2008. How does protein supplementation affect the selectivity and performance of Charolais cows on extensively grazed pastures in late autumn? *Grass and Forage Science* 63, 314–323.
- Farruggia A, Pomiès D, Coppa M, Ferlay A, Verdier-Metz I, Le Morvan A, Bethier A, Pompanon F, Troquier O and Martin B 2014. Animal performances, pasture biodiversity and dairy product quality: how it works in contrasted mountain grazing systems. *Agriculture, Ecosystems and Environment* 185, 231–244.
- Ferlay A, Martin B, Lerch S, Gobet M, Pradel P and Chilliard Y 2010. Effect of supplementation of maize silage diets with extruded linseed, vitamin E and plant extracts rich in polyphenols, and morning v. evening milking on milk fatty acid profile in Holstein and Montbéliarde cows. *Animal* 4, 627–640.
- Gerson T, John A and King ASD 1986. Effects of feeding ryegrass of varying maturity on the metabolism and composition of lipids in the rumen of sheep. *Journal Agricultural Science* 106, 445–448.
- Hoden A, Muller A, Jourmet M and Faverdin P 1986. Pâturage pour vaches laitières 1. Comparaison des systèmes de pâturage 'rationné' et 'tournant simplifié' en zone normande. *Bulletin Technique du Centre de Recherches Zootechniques et Vétérinaires de Theix* 64, 25–35.
- Hurtaud C, Faucon F, Couvreur S and Peyraud JL 2010. Linear relationship between increasing amounts of extruded linseed in dairy cow diet and milk fatty acid composition and butter properties. *Journal of Dairy Science* 93, 1429–1443.
- Khanal RC, Dhiman TR and Boman RL 2008. Changes in fatty acids composition of milk from lactating dairy cows during transition to and from pasture. *Livestock Science* 114, 164–175.
- Kratz M, Baars T and Guyenet S 2013. The relationship between high-fat dairy consumption and obesity, cardiovascular, and metabolic disease. *European Journal of Nutrition* 52, 1–24.
- Leiber F, Kreuzer M, Nigg D, Wettstein HR and Scheeder MRL 2005. A study on the causes for the elevated n-3 fatty acids in cows' milk of Alpine origin. *Lipids* 40, 191–202.
- Martin B, Verdier-Metz I, Buchin S, Hurtaud C and Coulon JB 2005. How does the nature of forages and pastures diversity influence the sensory quality of dairy livestock products? *Animal Science* 81, 205–212.
- Pomiès D, Martin B, Pradel P, Verdier-Metz I, Constant I, Delbès-Paus C, Troquier O, Fournier F, Montel MC and Farruggia A 2013. Design of low-input dairy farming systems in mountain areas: animal performances and cheese sensory properties. In *Proceedings of the 17th Meeting of the FAO-CIHEAM Mountain Pasture Network – pastoralism and ecosystem conservation* (ed. G Lombardi, E Mosimann, A Gorlier, G Iussig, M Lonati, M Pittarello, and M Probo), pp. 22–26. Department of Agricultural, Forest and Food Sciences – University of Turin, Grugliasco, Italy.
- Smith HJ, Taweel HZ, Tas BM, Tamminga S and Elgersma A 2005. Comparison of techniques for estimating herbage intake of grazing dairy cows. *Journal of Dairy Science* 88, 1827–1836.
- Sollemberg LE and Newman YC 2007. Grazing management. In *Forages, the science of grassland agriculture* (vol. II 6th edition, ed. RF Barnes, CJ Nelson, KJ Moore and MC Collins), pp. 651–659. Blackwell Publishing Professional, Ames, IA, USA.
- Sweeney RA and Rexroad PR 1987. Comparison of LECO FP-228 "nitrogen determinator" with AOAC copper catalyst Kjeldahl method for crude protein. *Journal Association of Official Agricultural Chemists International* 70, 1028–1030.
- Teague WR and Dowhower SL 2003. Patch dynamics under rotational and continuous grazing management in large, heterogeneous paddocks. *Journal of Arid Environment* 53, 211–229.
- Urban B and Caudal JP 1990. Herbomètre automatisé. In *Les Journées de la mesure électronique, informatique, automatique* (ed. INRA Département Informatique), pp. 57–59. INRA Département Informatique, Port Leucate, France.
- Van Soest PJ, Robertson JB and Lewis BA 1991. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy Science* 74, 3583–3597.
- Verheyden-Tixier H, Renaud PC, Morellet N, Jamot NJ, Besle JM and Dumont B 2008. Selection for nutrients by red deer hinds feeding on a mixed forest edge. *Oecologia* 156, 715–726.
- Vlaeminck B, Fievez V, Cabrita ARJ, Fonseca AJM and Dewhurst RJ 2006. Factors affecting odd- and branched-chain fatty acids in milk: a review. *Animal Feed Science and Technology* 131, 389–417.
- Vlaeminck B, Abrahamse PA, Fievez V, Lourenco M, Dijkstra J and Tamminga S 2010. The effect of allocation frequency in rotational grazing systems on the fatty acid profile in milk fat of dairy cows. In *Proceeding of the 23rd General Meeting of the European Grassland Federation – grassland in a changing world* (ed. H Schnyder, J Isselstein, F, Taube, K Auerswald, J Schelleberg, M Wachendorf, A Hermann, M Gierus, N Grage, and A Hopkins), pp. 586–588. Universität Göttingen, Göttingen, Germany.
- Willems H, Kreuzer M and Leiber M 2014. Alpha-linolenic and linoleic acid in meat and adipose tissue of grazing lambs differ among alpine pasture types with contrasting plant species and phenolic compound composition. *Small Ruminant Research* 116, 153–164.