AMERICAN JOURNAL OF AGRICULTURAL & BIOLOGICAL SCIENCES

Volume 8, issue 1

ISSN: 1557-4989

Science
Publications

American Journal of Agricultural and Biological Sciences

Editors Board

Tanka P Kandel

Aarhus University
Denmark

Expertise: Agriculture, Crop production, Crop physiology, Bioenergy crop production, Greenhouse gas emissions, Biomass quality, Biogas production

Jean-paul Lachaud

(1 El Colegio de la Frontera Sur - 2) CNRS
Mexico

Expertise: Ethology of social insects (ants) Behavioral ecology Predatory behavior Biological control using ants Interactions ants-parasites

H C Ho

Mississippi State University
United States

Expertise: Geosciences

Paulo Canas Rodrigues

Nova University of Lisbon, Portugal
Portugal

Expertise: Principal component analysis; Low rank approximations; Time series and singular spectrum analysis; Quantitative genetics; Genotype by environment interaction; QTL by environment interaction.

Georgia Thomas Tanou

Aristotle University of Thessaloniki
Greece

Expertise: Agriculture, Plant physiology, Fruit tree science, Fruit postharvest physiology, proteomics,

Vasudeo Zambare

South Dakota School of Mines and Technol
United States

Expertise: Biotechnology

Cristina L.M. Sgherri

University of Pisa
Italy

Expertise: Biochemical and metabolic responses of plants to oxidative stress caused by biotic and abiotic stresses. Generation and detection of Reactive Oxygen Species (ROS) following stress conditions. Changes in the composition and fluidity of membranes following stress conditions. Antioxidative metabolism in resurrection plants subjected to dehydration and rehydration. Changes in nutraceutical properties of plants grown in high salt and boron conditions. Study and simulation of antioxidative potential of plant extracts with Electron Spin Resonance technique.

Fulu Tao

Chinese Academy of Sciences
China

Absalom Zamorano

Instituto Politecnico Nacional
Mexico

Expertise: Biophysics, Biomathematics, and Bioinformatics. Specifically Computational Simulation.

Neelesh Sharma

Faculty of Veterinary Medicine
India

Expertise: Veterinary medicine, Stem Cell Biology, Bovine mastitis

Ajai Kumar Srivastav

DDU Gorakhpur University
India

Expertise: Endocrinology, Toxicology and Fishery Biology.

Rajeshwar P. Sinha

Centre of Advanced Study in Botany
India

Expertise: Photobiology and Molecular Microbiology

Editors Board

Junzeng Xu

State key laboratory of hydrology water
China

Expertise: Hydrology-Water Resources and Hydraulic Engineering

Amr Amin

United Arab Emirates University
UAE

Expertise: Cancer biology, diagnostic and treatment, nanoparticles and drug delivery.

Srinivas Koduru

Yale University
United States

Expertise: Molecular cell signaling pathways in cancer biology, natural compounds and cancer prevention, microenvironment and immunobiology.

Wei Chen

Changchun Institute of Applied Chemistry
China

Expertise: Surface electrochemistry, Spectroelectrochemistry, Nano electrochemistry, Design and the application of electrocatalysts for fuel cells

Hua Lu

Department of Biological Sciences
United States

Expertise: biological science

Claudio Caprara

University of Bologna
Italy

Expertise: Agricultural process control and automation, precision agriculture, agricultural systems, logistics and production chains, biosystems engineering, renewable energy and GIS models.

Rajan Ghimire

Department of Ecosystem Science and Management
United States

Expertise: Soil fertility and plant nutrition Sustainable soil management and long-term farming system research
Soil organic carbon and nitrogen dynamics Soil microbial community structure and functions
Greenhouse gases emission and climate change

Saso B. Koceski

University Goce Delcev
Macedonia

Expertise: Artificial intelligence Robotics Human - computer interaction Web technologies Medical imaging and 3D medical visualization Image processing Automation and control Mobile applications development

Malek Bajbouj

Freie Universität Berlin
Germany

Expertise: Niederlandische Philologie

Barbara Manachini

Università degli Studi di Palermo
Italy

Expertise: Biologia ambientale e biodiversità (ex Biologia Animale)

De Yi Wang

Polymer Institute
Germany

Expertise: Environmental Friendly High Performance Polymer (Nano)composites
(Bio-based polymer, Petro-based Polymer and Rubber nanocomposites)

Dilip Shah

Massachusetts General Hospital
United States

Expertise: Immunology

Editors Board

Fei Gu

The Ohio State University
United States

Expertise: Biomedical Informatics

Nara Gavini

Mississippi State University,
United States

Expertise: Biological Sciences

Hongjin Zheng

Howard Hughes Medical Institute
United States

Expertise: Biochemistry, structural biology, electron microscopy

Zhang Jiu-quan

Tobacco Research Institute
China

Expertise: Agricultural Sciences

Kenneth Blum, Ph.d.

University of Florida
United States

Expertise: neuropsychiatry and genetics, nutritional genetics and pharmacogenetics

Kumar Venkitanarayanan

University of Connecticut
United States

Expertise: molecular mechanisms by which food-borne pathogens tolerate and survive adverse environmental stresses encountered in nature, foods, and in hosts

Laron Nelson

University of Toronto
Canada

Expertise: Medicine, development of integrated biomedical, behavioural, and social/structural interventions

Oliver Ebert

Technical University of Munich
Germany

Expertise: The main focus of our laboratory centers on the use of replication-competent viral vectors for the treatment of cancer including vesicular stomatitis virus (VSV) and Newcastle disease virus (NDV). The majority of efforts are directed at designing and testing oncolytic viral therapies in the treatment of gastrointestinal cancers

Paul M Rea

University of Glasgow
United Kingdom

Expertise: haematology, gastroenterology, paediatric surgery, cardiology and accident and emergency

C S Prakash

Tuskegee University
United States

Expertise: Food crops and importance to developing countries, training of scientists and students in plant biotechnology, global societal awareness of food biotechnology issues

Yanhong Shi

Beckman Research Institute
United States

Expertise: Cell Biology and Neuroscience

Contents

Title: TOLERANCE OF ACCESSIONS OF GLAGAH (*SACCHARUM SPONTANEUM*) TO DROUGHT STRESS AND THEIR ACCUMULATION OF PROLINE

Author: Aminatun Munawarti, Taryono, Endang Semiarti, Paul Holford and Sismindari

Source: American Journal of Agricultural and Biological Sciences 8(1): 1-11, 2013

Title: ENVIRONMENTAL EDUCATION AND BIODIVERSITY CONCERN: BEYOND THE ECOLOGICAL LITERACY

Author: Carlos Hiroo Saito

Source: American Journal of Agricultural and Biological Sciences 8(1): 12-27, 2013

Title: MANIPULATION OF PHOTOSYNTHESIS IN GRAPE (*VITIS VINIFERA*) CV. 'FLAME' BY THE APPLICATION OF TWO SUCROSE ANALOGS

Author: M.E. Tiznado-Hernandez, J.A. Orozco-Avitia, A.J. Ojeda-Contreras and A.A. Gardea-Bejar

Source: American Journal of Agricultural and Biological Sciences 8(1): 28-33, 2013

Title: G-2548A LEPTIN PROMOTER AND Q223R LEPTIN RECEPTOR POLYMORPHISMS IN OBESE MEXICAN SUBJECTS

Author: P. Carrillo-Vazquez Jonathan, Lopez-Alcantar Jeronimo, Chimal-Vega Brenda, Benitez-Cardoza Claudia, Zamorano-Carrillo Absalom, Reyes-Lopez Cesar, Lopez-Camarillo Cesar and A. Marchat Laurence

Source: American Journal of Agricultural and Biological Sciences 8(1): 34-43, 2013

Title: VARIABILITY AND GENETIC STRUCTURE IN A COMMERCIAL FIELD OF TEQUILA PLANTS, *AGAVE TEQUILANA* WEBER (AGAVACEAE)

Author: Martha Isabel Torres-Moran, Ana Paulina Velasco-Ramirez, Salvador Antonio Hurtado-de la Pena, Andres Rodriguez-Garcia and Salvador Mena-Munguia

Source: American Journal of Agricultural and Biological Sciences 8(1): 44-53, 2013

Title: ASSESSMENT OF SOIL COMPACTION UNDER CENTER PIVOT IRRIGATION SYSTEMS AND ITS EFFECT ON CROP PERFORMANCE

Author: Khalid Ali Al-Gaadi

Source: American Journal of Agricultural and Biological Sciences 8(1): 54-66, 2013

Title: ENHANCEMENT OF BIOGAS PRODUCTION POTENTIAL FOR ANAEROBIC CO-DIGESTION OF WASTEWATER USING DECANTER CAKE

Author: Thaniya Kaosol and Narumol Sohgrathok

Source: American Journal of Agricultural and Biological Sciences 8(1): 67-74, 2013

TOLERANCE OF ACCESSIONS OF GLAGAH (*SACCHARUM SPONTANEUM*) TO DROUGHT STRESS AND THEIR ACCUMULATION OF PROLINE

^{1,2}Aminatun Munawarti, ³Taryono, ⁴Endang Semiarti, ⁵Paul Holford and ⁶Sismindari

¹Postgraduate Program of Biotechnology, Universitas Gadjah Mada, Yogyakarta, Indonesia

²Department of Biology, Faculty of MIPA, Universitas Brawijaya, Malang, Indonesia

³Department of Agronomy, Faculty of Agriculture, Universitas Gadjah Mada, Yogyakarta, Indonesia

⁴Faculty of Biology, Universitas Gadjah Mada, Yogyakarta, Indonesia

⁵School of Science and Health, University of Western Sydney, New South Wales, Australia

⁶Faculty of Pharmacy, Universitas Gadjah Mada, Yogyakarta, Indonesia

Received 2012-06-15, Revised 2013-01-30; Accepted 2013-02-04

ABSTRACT

Glagah (wild sugarcane, *Saccharum spontaneum*) is a perennial grass that grows well in marginal soils. It has a high carbohydrate and fibre content, making it suitable for ethanol production and the study industry. Thus, glagah has potential as a crop species in its own right and may also be used in sugarcane breeding programs. However, glagah germplasm has not been extensively utilised in breeding programs, especially in relation to improving drought tolerance. This study was performed to evaluate the effect of drought stress over an eight week period on plant height, stalk diameter, green leaf number and leaf proline content of eight, two month-old accessions of glagah to identify their drought tolerance and to determine whether proline accumulation can be used as a metabolic marker of drought tolerance. Accessions, BOT-53, BOT-54 and BOT-62, were the most tolerant and productive. Two patterns of proline accumulation were shown in drought-stressed plants. In four accessions, proline increased after both four and eight weeks of drought stress. While, in the others, proline increased after four weeks and then declined. Significant, positive correlations were found between leaf proline contents (after both two and eight weeks of drought) and plant height at all assessment times. A significant, positive correlation was also discovered between proline content after eight weeks and green leaf number after four weeks of drought. Glagah accessions of BOT-53, BOT-54 and BOT-62 show drought-tolerance and have potential for use as a crop for arid regions or in breeding programs to improve production of sugarcane. Drought tolerance in glagah appears to be mediated by proline and accumulation of this amino acid has potential as a metabolic marker of drought tolerance.

Keywords: *Saccharum Spontaneum*, Proline, Metabolic Marker, Glagah, Drought Stress, Drought Tolerance, Proline Contents, Breeding Programs, Proline Accumulation, Ethanol Production

1. INTRODUCTION

Sugarcane (*Saccharum officinarum* L.) (Poaceae) is an important crop globally primarily for sugar production but is increasingly being used as a source of bioenergy due to its phenomenal capacity for dry matter production; however, a lack of water often limits sugarcane production (Naik, 2001; Silva *et al.*, 2008; Tammisola, 2010). Drought stress simultaneously affects a number of

morphological and physiological traits in plants thereby causing a loss of productivity (Bray, 1997; Jamaux *et al.*, 1997; Yordanov *et al.*, 2003). Therefore, traits associated with tolerance of drought stress are suitable criteria for screening germplasm collections for accessions that may contribute to breeding programs to reduce the influence of water deficit on crop yield (Silva *et al.*, 2008).

Maintenance of plant water potential during water deficits is essential to continued growth and can be achieved

Corresponding Author: Aminatun Munawarti, Postgraduate Program of Biotechnology, Universitas Gadjah Mada, Yogyakarta, Indonesia

by osmotic adjustment resulting from the accumulation of molecules, such as proline, in the cytoplasm (Ingram and Bartels, 1996). Proline in drought-stressed plants is not only used for osmotic adjustment but also for scavenging and detoxifying oxidants (Yamada *et al.*, 2005; Vallidoyan and Nguyen, 2006) and for preventing membrane damage and protein denaturation during severe drought stress (Bandurska, 1998; Ain-Lhout *et al.*, 2001). Thus, proline content has been proposed as a metabolic marker of drought stress (Dib *et al.*, 1994).

Glagah (wild sugarcane, *Saccharum spontaneum* L. (Poaceae) is a tall perennial grass with deep roots and rhizomes that can grow well in marginal soils where no other crop can be cultivated (Amalraj *et al.*, 2008) and can established itself in diverse habitats such as rocky regions, deserts and sandy flats (Kandasami *et al.*, 1983). Therefore, glagah can be considered as a drought-tolerant, perennial grass and a preliminary, unpublished study by the authors suggested there are considerable differences among accessions of glagah in their resistance to drought stress. Glagah has a high carbohydrate content, making the biomass from this species a suitable substrate for ethanol production (Chandel *et al.*, 2009). Glagah also has a high fibre content in its stalk and a dry matter content of 22%. This is higher than in the stalks of cultivated sugarcane, which only have ~14% dry matter, suggesting that this species has potential as a possible renewable fibre resource for the study industry (Amalraj *et al.*, 2008). Thus, glagah has not only potential value as a crop species in its own right but also may be used in breeding programs to improve the production of sugarcane.

Crop improvement has played a significant role in yield improvement (Swamy *et al.*, 2003). However, to date, the genetic resources in the germplasm of glagah, especially those related to drought stress, have not been extensively utilized in sugarcane breeding programs. To improve drought tolerance, it is necessary to understand the mechanisms by which plants respond to drought and studies on the physiological response of different glagah accessions to drought could be a useful to develop an understanding of these mechanisms. Therefore, in this study, we evaluated the effect of drought stress on plant height, stalk diameter and green leaf number of eight accessions of glagah (BOT-53, BOT-54, BOT-83, BOT-84, IM76-238, BOT-62, BOT-77 and BOT-91) to determine their drought tolerance. As outlined above, changes in proline content have an adaptive role in plants during stress. Therefore, leaf proline contents were also assayed to determine whether proline accumulation is involved with drought tolerance and whether concentrations of this amino acid can be used as a metabolic marker of drought tolerance.

2. MATERIALS AND METHODS

2.1. Plant Material and Experimental Conditions

Eight accessions of glagah, BOT-53, BOT-54, BOT-83, BOT-84, IM76-238, BOT-62, BOT-77 and BOT-91, were used in this study. The accessions were kindly provided by the Indonesian Sugar Research Institute (ISRI/P3GI), Pasuruan, East Java. The accessions were propagated from single node stalk segments (bud sets) in polybags (400×200×400 mm³) containing 10 kg of a mixture of soil and manure (2:1) for two months before being subjected to drought stress. The plants were watered every two days until the treatments were imposed and were fertilized three times (1 day, 1 week and 1 month after planting) with 2 g of ammonium sulphate and 1 g of triple superphosphate per polybag. After two months, the plants were transferred from the field to a glasshouse and were then subjected to drought stress by withholding watering for 8 weeks. Untreated, control plants were watered every two days.

2.2. Growth Measurement

Plant height and green leaf number were measured two, four and eight weeks after commencement of the treatment for both control and drought-stressed plants of each accession; stalk diameter was measured after four and eight weeks. Plant height was measured from the soil surface to the first fully-expanded leaf. Green leaf number was calculated from number of green leaves divided by total number of leaves; stalk diameter was measured at 50 mm above the soil surface. A Drought Susceptibility Index (DSI) was calculated as the plant height data according to the formula proposed by Fischer and Maurer (1978): $DSI = (1 - Y_d/Y_w)/(1 - X_d/X_w)$, where Y_d = height of droughted accession, Y_w = height of the watered accession, X_d = mean height of all droughted accessions and X_w = mean height of all watered accessions.

2.3. Estimation of Proline Accumulation

Assessment of proline content was performed in Laboratory of Biochemistry, Research Center for Biotechnology, Universitas Gadjah Mada, Yogyakarta. Proline was assayed in the youngest fully-expanded leaves from both stressed and control plants. Proline content was determined using the colorimetric method of Bates *et al.* (1973). DL-proline (Sigma) was used as a standard.

2.4. Experimental Design and Data Analysis

The experiment of was set out using a completely randomized design with two replicates. Data were checked for heteroscedastisity using Bartlett's test before being subjected to ANOVA using Statistica (Version 9.1., StatSoft, Inc. 2010) and means were separated using Fisher's LSD test at $P = 0.05$. Due to

large interactions between the effects of accession and drought treatment, data for plant height, green leaf number and proline content of the control plants were analysed separately from the droughted plants.

3. RESULTS

3.1. Plant Height

Plant height was significantly affected by accession ($F = 39.7-64.2$, d.f. = 7, $p < 0.0001$) and by treatment ($F = 64.2-187.9$, d.f. = 1, $p < 0.0001$) and there was a significant

accession*treatment interaction ($F = 3.8-4.5$, d.f. = 7, $P = 0.01-0.006$) at all assessment times (**Fig. 1**). Accessions, BOT-53, BOT-54, BOT-62 and IM76-238, appeared to show some degree of drought tolerance and, by the end of the eighth week of the drought stress, these accessions were significantly taller than the other accessions. After the eight weeks of drought stress, these accessions had reductions in height of 34.6, 9.6, 31.2 and 21.5%, respectively, compared to their control plants, whereas all of the other accessions had reductions in height of more than 41% compared to their controls.

Fig. 1. Plant height of eight glagah accessions subjected to drought stress for two (A), four (B) and eight weeks (C) and the height of well-watered, control plants. Each value is the mean of two independent samples. For each panel, means associated with the same letter are not statistically different from each other according to Fisher's LSD test at $P = 0.05$. Data for the control and droughted plants at each assessment period were analysed separately. The drought susceptibility index was calculated according to Fischer and Maurer (1978)

The DSI calculated from the plant height data from Week 8 suggests that the accessions can be split into three groups based on this criterion with BOT-54 and BOT-62 in the most drought tolerant group, BOT-83 and BOT-53 in an intermediate group with the remainder being the least tolerant.

3.2. Stalk Diameter

For stalk diameter, there were significant differences due to accession ($F = 34.7$ and 10.3 , d.f. = 7 , $p = <0.0001$) and to treatment ($F = 61.4$ and 413.7 ,

d.f. = 1 , $p = <0.0001$) at both the fourth and eighth weeks of drought stress (**Fig. 2**). There was no significant accession*treatment interaction. However, despite these statistical differences, the differences in diameter found among the control or drought-treated plants at each assessment time were small in real terms. Nor was there a large amount of variation in the percentage reduction in stalk diameter due to the drought treatment. After eight weeks, all accessions showed a 20-26% reduction in radial growth.

Fig. 2. Stalk diameter of eight glagah accessions subjected to drought stress for four (A) and eight weeks (B) and the diameter of well-watered, control plants. Each value is the mean of two independent samples. For each panel, means associated with the same letter are not statistically different from each other according to Fisher's LSD test at $P = 0.05$. Data at each assessment period were analysed separately

Fig. 3. Percentage of leaves remaining green of eight glagah accessions subjected to drought stress for two (A) and four (B) and the percentage for well-watered, control plants. Each value is the mean of two independent samples. For each panel, means associated with the same letter are not statistically different from each other according to Fisher's LSD test at $P = 0.05$. Data for the control and droughted plants at each assessment period were analysed separately

3.3. Green Leaf Number

Data concerning the percentage of leaves which remain green are shown in **Fig. 3**. After two weeks of drought, 50-90% of leaves on the droughted plants remained green and there were significant effects due to accession ($F = 38.8$, d.f. = 7, $p < 0.0001$), to treatment ($F = 68.4$, d.f. = 1, $p < 0.0001$) and a significant interaction ($F = 10.5$, d.f. = 7, $p < 0.0001$). After the four weeks of drought, there was a large reduction in the percentage of leaves that remained green and again the effects of accession, treatment and the accession*treatment interaction were all significant ($p < 0.0001$ in all cases). At this assessment point, green leaves only remained on accessions BOT-53, BOT-54, BOT-62 and BOT-77. After eight weeks of drought stress, green leaves only remained on one plant of BOT-53 and BOT-77 (**Fig. 4**).

3.4. Proline Content

The results of the proline analysis are presented in **Fig. 5**. Two weeks after drought treatment, there was little difference among the accessions in terms of proline content. Also, there was little difference between drought-stressed and control plants, with all plants containing between ~ 7 - $25 \mu\text{g g (FW)}$ **Fig. 5A**. The proline content of leaves of the control, watered plants did not increase during the assessment period and did not differ significantly among the accessions. However, differences emerged after four and eight weeks imposition of drought ($p < 0.0001$) (**Fig. 5B and C**) and the leaf proline content significantly increased in plants under drought stress. Two patterns of proline accumulation can be seen in the data from these plants.

(a)

(b)

Fig.4. Influence of drought stress on vegetative growth: (A) watered control plants, (B) potential drought tolerant accessions (BOT-53 and BOT-54) and an accession (BOT-77) with intermediate tolerance showing necrosis eight weeks after the imposition of drought stress

Table 1. Correlation coefficients and their statistical significance between leaf proline contents and plant height of eight droughted glagah accessions after two, four and eight weeks of imposed drought

Proline content	Plant height		
	Week 2	Week 4	Week 8
Week 2	$r = 0.5800$ $P = 0.0180$	$r = 0.5900$ $P = 0.0170$	$r = 0.610$ $P = 0.011$
Week 4	$r = 0.3500$ $P = 0.1900$	$r = 0.2500$ $P = 0.3400$	$r = 0.038$ $P = 0.890$
Week 8	$r = 0.8400$ $p < 0.0001$	$r = 0.8700$ $p < 0.0001$	$r = 0.910$ $p < 0.0001$

The leaf proline content of four accessions, BOT-53, BOT-54, IM76-238 and BOT-62 increased after both four and eight weeks of drought stress (**Fig. 5B and C**). The greatest increases in concentration were $339.25 \mu\text{g g}^{-1}$ FW (17-fold increase) and $353.25 \mu\text{g g}^{-1}$ FW (28-fold increase) in accessions, BOT-53 and BOT-54, respectively. In the other accessions (BOT-83, BOT-84, BOT-77 and BOT-91), proline concentrations increased significantly after four weeks of drought stress and then declined. However, at Week 8, the proline concentrations of accessions, BOT-84 and BOT-91, were still higher than in the watered control plants, whereas, the proline concentrations in accessions BOT-77 and BOT-83 were little different from their controls.

3.5. Correlation Between Proline Contents and Morphological Traits

Correlations were found between proline contents and morphological characters assessed in this study. Strong positive correlations were found between proline contents in plants after two weeks of drought and plant height after two, four and eight weeks (**Table 1**). Also, the proline contents after eight weeks showed strong positive correlations with plant heights after two, four and eight weeks. The correlation coefficients were stronger between the proline contents after eight weeks and plant height than between proline contents after two weeks and this character. The only other significant correlation was between proline content at eight weeks and green leaf number at four weeks ($r = 0.59$; $P = 0.016$). No significant correlations were found with stalk diameter.

Fig. 5. The proline content of eight glagah accessions subjected to drought stress for two (A), four (B) and eight weeks (C) and the proline content of well-watered, control plants. Each value is the mean of two independent samples. For each panel, means associated with the same letter are not statistically different from each other according to Fisher's LSD test at P = 0.05. Data for the control and droughted plants at each assessment period were analysed separately

4. DISCUSSION

Drought is one of the most important environmental stress factors that can restrict the potential yield of

sugarcane and tillering (beginning approximately on the fortieth day after planting and lasting up to 120 days) and the grand growth phase (taking up to 9 months after planting) are known as critical stages of drought-

sensitivity due to the high need for water at these times (Zingaretti *et al.*, 2012). Plants respond to drought stress at the molecular, cellular and physiological levels and this response depends on the species and genotype, the length and severity of water loss, the age and stage of development, the organ and cell type and the sub-cellular compartment (Barnabas *et al.*, 2008). In this study, the first obvious symptoms of stress were observed two weeks after the drought treatment and commenced with changes being leaf discoloration. Chlorosis is a common symptom associated with drought stress and is due to decreases in leaf photosynthetic pigments, as has been shown by Chaves *et al.* (2002) in their studies on *Quercus ilex* and *Q. suber* and by Manivannan *et al.* (2008) on sunflower. Studies have shown that the majority of chlorophyll loss in plants in response to water deficit occurs in the mesophyll cells with a lesser amount being lost from the bundle sheath cells (Anjum *et al.*, 2011). Low concentrations of photosynthetic pigments can directly limit photosynthetic potential and, hence, primary production. The decrease in chlorophyll content under drought stress has been considered as a typical symptom of oxidative stress and may be the result of pigment photo-oxidation and chlorophyll degradation and may be the cause of the reduction of green leaf number found in this study.

Besides green leaf number, drought stress also decreased plant height and stalk diameter in the eight accessions of glagah studied. However, stalk diameter appears to be a poor character for the assessment of changes in plant growth due to drought stress due to the low variation in this character. The reduction in plant height and stalk diameter may be due to a decline in the cell enlargement and greater leaf senescence under drought stress. Similar results were found in rice (Perez-Molphe-Balch *et al.*, 1996), avocado (Chartzoulakis *et al.*, 2002), sunflower cultivars (Manivannan *et al.*, 2008) and apple (Liu *et al.*, 2012).

The data for three morphological characteristics (plant height, stalk diameter and green leaf number) suggested that the accessions can be divided into three groups with respect to drought tolerance. Three accessions, BOT-53, BOT-54 and BOT-62, appear to be the most drought-tolerant and productive. Plants of these three accessions were the tallest at the end of the drought period and still had green leaves. Two accessions, BOT-77 and IM76-238, showed intermediate tolerance. IM76-238 grew to the same height as BOT-62, but all leaves became necrotic between four to eight weeks after the

drought treatment commenced. BOT-77 still had green leaves at the end of the experiment; however, it did not grow as tall as the other accessions. The remaining three accessions (BOT83, BOT-84 and BOT-91) showed the least tolerance to drought. They had no green leaves by the end of the experiment and were the shortest of the eight accessions. The ranking of plants according to their drought susceptibility index calculated from the plant height data from Week 8 supported the grouping of the accessions with the exception of BOT-83. This accession did not grow as high as the other accessions even when watered; however, its ranking showed that its DSI was the third smallest. It is necessary to further study these accessions to confirm their tolerance to drought and to establish their recovery after rewatering.

Proline is known to be involved in a plant's response to a various environmental stresses, including drought stress and proline accumulation was first reported in wilting perennial rye grass (*Lolium perenne*) (Kemble and Macpherson, 1954). An increase in proline content is a common response of plants to drought stress (Mostajeran and Rahimi-Eichi, 2009) and has been found in other species such as wheat (Johari-Pireivatlou, 2010; Keyvan, 2010; Akhkha *et al.*, 2011), chickpea (Mafakheri *et al.*, 2010), rice (Vajrabhaya *et al.*, 2001; Hien *et al.*, 2003; Mostajeran and Rahimi-Eichi, 2009), cotton (Parida *et al.*, 2008) and potato (Farhad *et al.*, 2011) and this study has shown a similar increase in *S. spontaneum*.

Studies on the use of proline content as a marker of drought tolerance have produced different conclusions. Hanson *et al.* (1979), working with *Hordeum vulgare* L., suggested that proline accumulation was of no practical use in breeding, even though they found a heritable component to this trait. Ceh *et al.* (2009) found no correlation between proline content and drought-resistance in hops (*Humulus lupulus* L.) and Ilahi and Dorffling (1982) found that drought-susceptible cultivars of *Zea mays* had higher proline contents than drought-resistant ones. In contrast, Singh *et al.* (1972; 1973) working with barley (*Hordeum vulgare* L.), Vajrabhaya *et al.* (2001) working on rice lines, Ma *et al.* (2004) working with *Brassica* spp., Sofo *et al.* (2004) working on olive trees, Bayoumi *et al.* (2008) studying *Triticum aestivum* L., Naser *et al.* (2010) working on the Persian walnut (*Juglans regia* L.) and Sharada and Naik (2011) working with groundnut (*Arachis hypogaea* L.) all found higher concentrations of proline in drought-tolerant types. Ma *et al.* (2004); Bayoumi *et al.* (2008) and Naser *et al.* (2010) all

suggested that proline accumulation could be used as a marker of drought tolerance. Differences in the role of proline with respect to drought tolerance in sugarcane have also been found. Zhao *et al.* (2010) suggested that proline was not a sensitive water stress indicator, whereas Rao and Asokan (1978) found that drought-resistant varieties of sugarcane accumulated more proline than susceptible ones and suggested that proline accumulation could be used as an index of drought tolerance. The significant positive correlations between plant height and proline content found in this study suggest that levels of this amino acid can be used as a marker of drought tolerance in glagah. However, this needs to be further confirmed with a wider range of genotypes. It is not unlikely that proline could act as a marker of drought tolerance in glagah as a mechanism for tolerance which has been suggested by Molinari *et al.* (2007) for sugarcane. These authors reported that transgenic sugarcane containing the P5CS gene from *Vigna aconitifolia* (Jacq.) Maréchal driven by a stress inducible promoter accumulated significantly higher amounts of proline and biomass under drought conditions. In these plants, there was no clear association between proline accumulation and osmotic adjustment. However, the increased biomass production was associated with lower levels of lipid peroxidation resulting in less damage to the plants' photosynthetic apparatus.

5. CONCLUSION

The results of this study indicated that glagah accessions, BOT-53, BOT-54 and BOT-62, are drought tolerant and have potential for use as a crop for arid regions and for use in breeding programs to improve production of sugarcane. Drought tolerance in glagah seems to be mediated by proline and concentrations of this amino acid have potential use as a marker of drought tolerance.

6. ACKNOWLEDGEMENT

The researchers are grateful to Prof. Dr. Ir. Mirzawan, P.D.N. and Ir. Eka Sugiyarta, MSc. from the Indonesian Sugar Research Institute (ISRI/P3GI), Pasuruan, East Java, Indonesia for kindly providing the glagah accessions. Thanks are also to Prof. Sutiman B. Sumitro, SU, DSc. and Dr. Wahyu Widoretno (Dept. of Biology, Universitas Brawijaya, Malang, Indonesia) for reading this manuscript and giving suggestions to it. Special thanks to

Chandra Kirana, MS, MAgrSc. PhD (School of Biological Sciences, Victoria University, Wellington, New Zealand) for her critical review. This research was financially supported by BPPS (DIKTI) and Universitas Brawijaya.

7. REFERENCES

- Ain-Lhout, F., M. Zunzunegui, M.C.D. Barradas, R. Tirado and A. Clavijo *et al.*, 2001. Comparison of proline accumulation in two Mediterranean shrubs subjected to natural and experimental water deficit. *Plant Soil*, 230: 175-183. DOI: 10.1023/A:1010387610098
- Akhkha, A., T. Boutraa and A. Alhejely, 2011. The rates of photosynthesis, chlorophyll content, dark respiration, proline and Absciscic Acid (ABA) in wheat (*Triticum durum*) under water deficit conditions. *Int. J. Agric. Biol.*, 13: 215-221.
- Amalraj, V.A., P. Rakkiyappan, D. Neelamathi, S. Chinnaraj and S. Subramanian, 2008. Wild cane as a renewable source for fuel and fiber in paper industry. *Curr. Sci.*, 95: 1599-1602.
- Anjum, S.A., X.Y. Xie, L.C. Wang, M.F. Saleem and C. Man *et al.*, 2011. Morphological, physiological and biochemical responses of plants to drought stress. *Afr. J. Agric. Res.*, 6: 2026-2032.
- Bandurska, H., 1998. Implication of ABA and proline on cell membrane injury of water deficit stressed barley seedlings. *Acta Physiol. Plant.*, 20: 375-381. DOI: 10.1007/s11738-998-0023-3
- Barnabas, B., K. Jager and A. Feher, 2008. The effect of drought and heat stress on reproductive processes in cereals. *Plant Cell Environ.*, 31: 11-38. DOI: 10.1111/j.1365-3040.2007.01727.x
- Bates, L.S., R.P. Waldren and I.D. Teare, 1973. Rapid determination of free proline for water-stress studies. *Plant Soil*, 39: 205-207. DOI: 10.1007/BF00018060
- Bayoumi, T.Y., M.H. Eid and E.M. Metwali, 2008. Application of physiological and biochemical indices as a screening technique for drought tolerance in wheat genotypes. *Afr. J. Biotechnol.*, 7: 2341-2352.
- Bray, E.A., 1997. Plant responses to water deficit. *Trends Plant Sci.*, 2: 48-54. DOI: 10.1016/S1360-1385(97)82562-9
- Ceh, B., A. Cerenak and D. Majer, 2009. Potassium and proline content in hop leaves as biochemical marker for drought stress tolerance. *Hop Bull.*, 16: 43-52.

- Chandel, A.K., M.L. Narasu, G. Chandrasekhar, A. Manikyam and L.V. Rao, 2009. Use of *Saccharum spontaneum* (wild sugarcane) as biomaterial for cell immobilization and modulated ethanol production by thermotolerant *Saccharomyces cerevisiae* VS3. *Bioresource Technol.* 100: 2404-2410. DOI: 10.1016/j.biortech.2008.11.014, PMID: 19114303
- Chartzoulakis, K., A. Patakas, G. Kofidis, A. Bosabalidis and A. Nastou, 2002. Water stress affects leaf anatomy, gas exchange, water relations and growth of two avocado cultivars. *Sci. Hortic.*, 95: 39-50. DOI: 10.1016/S0304-4238(02)00016-X
- Chaves, M.M., J.S. Pereira, J. Maroco, M.L. Rodriguez and C.P. Ricardo *et al.*, 2002. How plants cope with water stress in the field. *Photosynthesis and growth.* *Ann. Bot.*, 89: 907-916. PMID: 12102516
- Silva, M.D.A., J.A.G.D. Silva, J. Enciso, V. Sharma and J. Jifon, 2008. Yield components as indicators of drought tolerance of sugarcane. *Sci. Agric. (Piracicaba, Braz.)* 65: 620-627. DOI: 10.1590/S0103-90162008000600008
- Dib, T.A., P. Monneveux, E. Acevedo and M.M. Nachit, 1994. Evaluation of proline analysis and chlorophyll fluorescence quenching measurements as drought tolerance indicators in durum wheat (*Triticum turgidum* L. var. *durum*). *Euphytica*, 79: 65-73. DOI: 10.1007/BF00023577
- Farhad, M.S., A.M. Babak, Z.M. Reza, R.S.M. Hassan and T. Afshin, 2011. Response of proline, soluble sugars, photosynthetic pigments and antioxidant enzymes in potato (*Solanum tuberosum* L.) to different irrigation regimes in greenhouse condition. *Aust. J. Crop Sci.*, 5: 55-60.
- Fischer, R.A. and R. Maurer, 1978. Drought resistance in spring wheat cultivars. I. Grain yield responses. *Aust. J. Agric. Res.*, 29: 897-912. DOI: 10.1071/AR9780897
- Hanson, A.D., C.E. Nelsen, A.R. Pedersen and E.H. Everson, 1979. Capacity for proline accumulation during water stress in barley and its implications for breeding for drought resistance. *Crop Sci.*, 19: 489-493.
- Hien, D.T., M. Jacobs, G. Angenon, C. Hermans and T.T. Thu *et al.*, 2003. Proline accumulation and Δ^1 -pyrroline-5-carboxylate synthetase gene properties in three rice cultivars differing in salinity and drought tolerance. *Plant Sci.*, 165: 1059-1068. DOI: 10.1016/S0168-9452(03)00301-7
- Ilahi, I. and K. Dorffling, 1982. Changes in abscisic acid and proline levels in maize varieties of different drought resistance. *Physiol. Plant.*, 55: 129-135. DOI: 10.1111/j.1399-3054.1982.tb02275.x
- Ingram, J. and D. Bartels, 1996. The molecular basis of dehydration tolerance in plants. *Ann. Rev. Plant Physiol. Plant Mol. Biol.*, 47: 377-403. DOI: 10.1146/annurev.arplant.47.1.377
- Jamaux, I., A. Steinmetz and E. Belhassen, 1997. Looking for molecular and physiological markers of osmotic adjustment in sunflower. *New Phytol.*, 137: 117-127. DOI: 10.1046/j.1469-8137.1997.00817.x
- Johari-Pireivatlou, M., 2010. Effect of soil water stress on yield and proline content of four wheat lines. *Afr. J. Biotechnol.*, 9: 36-40.
- Kandasami, P.A., T.V. Sreenivasan, T.C.R. Rao, K. Palanichami and B.V. Natarajan *et al.*, 1983. Catalogue on sugarcane genetic resources-I: *Saccharum Spontaneum* L. Sugarcane Breeding Institute.
- Kemble, A.R. and H.T. Macpherson, 1954. Liberation of amino acids in perennial rye grass during wilting. *Biochem. J.*, 58: 46-49.
- Keyvan, S., 2010. The effects of drought stress on yield, relative water content, proline, soluble carbohydrates and chlorophyll of bread wheat cultivars. *J. Anim. Plant Sci.*, 3: 1051-1060.
- Liu, B., L. Cheng, M. Li, D. Liang and Y. Zou *et al.*, 2012. Interactive effects of water and nitrogen supply on growth, biomass partitioning and water-use efficiency of young apple trees. *Afr. J. Agric. Res.*, 7: 978-985. DOI: 10.5897/AJAR11.1212
- Ma, Q., D.W. Turner, D. Levy and W.A. Cowling, 2004. Solute accumulation and osmotic adjustment in leaves of Brassica oilseeds in response to soil water deficit. *Aust. J. Agric. Res.*, 55: 939-945.
- Mafakheri, A., A. Siosewarden, B. Bahramnejad, P.C. Struik and Y. Sohrabi, 2010. Effect of drought stress on yield, proline and chlorophyll contents in three chickpea cultivars. *Aust. J. Crop Sci.*, 4: 580-585.
- Manivannan, P., C.A. Jaleel, Z. Chang-Xing, R. Somasundaram and M.M. Azooz *et al.*, 2008. Variations in growth and pigment composition of sunflower varieties under early season drought stress. *Global J. Mol. Sci.*, 3: 50-56.

- Molinari, H.B.C., C.J. Marur, E. Daros, M.K.F.D. Campos and J.F.R.P.D. Carvalho *et al.*, 2007. Evaluation of the stress-inducible production of proline in transgenic sugarcane (*Saccharum* spp.): Osmotic adjustment, chlorophyll fluorescence and oxidative stress. *Physiol. Plantarum*, 130: 218-229. DOI: 10.1111/j.1399-3054.2007.00909.x
- Mostajeran, A. and V. Rahimi-Eichi, 2009. Effects of drought stress on growth and yield of rice (*Oryza sativa* L.) cultivars and accumulation of proline and soluble sugars in sheath and blades of their different ages leaves. *Am. Euras. J. Agric. Environ. Sci.*, 5: 264-272.
- Naik, G.R., 2001. Sugarcane Biotechnology. 1st Edn., Science Publishers, Enfield, ISBN-10: 1578081637, pp: 165.
- Naser, L., V. Kourosh, K. Bahman and A. Reza, 2010. Soluble sugars and proline accumulation play a role as effective indices for drought tolerance screening in Persian walnut (*Juglans regia* L.) during germination. *Fruits*, 65: 97-112. DOI: 10.1051/fruits/20010005
- Parida, A.K., V.S. Dagaonkar, M.S. Phalak and L.P. Aurangabadkar, 2008. Differential responses of the enzymes involved in proline biosynthesis and degradation in drought tolerant and sensitive cotton genotypes during drought stress and recovery. *Acta Physiol. Plant.*, 30: 619-627. DOI: 10.1007/s11738-008-0157-3
- Perez-Molphe-Balch, E., M. Gidekel, M. Segura-Nieto, L. Herrera-Estrella and N. Ochoa-Alejo, 1996. Effects of water stress on plant growth and root proteins in three cultivars of rice (*Oryza sativa*) with different levels of drought tolerance. *Physiol. Plantarum*, 96: 284-290. DOI: 10.1111/j.1399-3054.1996.tb00215.x
- Rao, K.C. and S. Asokan, 1978. Studies of free proline association to drought resistance in sugar cane. *Sugar J.*, 40: 23-24.
- Sharada, P. and G.R. Naik, 2011. Physiological and biochemical responses of groundnut genotypes to drought stress. *World J. Sci. Technol.*, 1: 60-66.
- Singh, T.N., D. Aspinall and L.G. Paleg, 1972. Proline accumulation and varietal adaptability to drought in barley: A potential metabolic measure of drought resistance. *Nature*, 236: 118-190. DOI: 10.1038/newbio236188a0
- Singh, T.N., L.G. Paleg and D. Aspinall, 1973. Stress metabolism III. Variations in Response to water deficit in the barley plant. *Aust. J. Biol. Sci.*, 26: 65-76. DOI: 10.1071/BI9730065
- Sofo, A., B. Dichio, C. Xiloyannis and A. Masia, 2004. Lipoxygenase activity and proline accumulation in leaves and roots of olive trees in response to drought stress. *Physiol. Plantarum*, 121: 58-65. DOI: 10.1111/j.0031-9317.2004.00294.x
- Swamy, B.P.M., H.D. Upadhyaya, P.V.K. Goudar, B.Y. Kullaiswamy and S. Singh, 2003. Phenotypic variation for agronomic characteristics in a groundnut core collection for Asia. *Field Crops Res.*, 84: 359-371. DOI: 10.1016/S0378-4290(03)00102-3
- Tammisola, J., 2010. Towards much more efficient biofuel crops-can sugarcane pave the way? *GM Crops*, 1: 181-198. DOI: 10.4161/gmcr.1.4.13173, PMID: 21844673
- Vajrabhaya, M., W. Kumpun and S. Chadchawan, 2001. The solute accumulation: The mechanism for drought tolerance in RD23 rice (*Oryza sativa* L) lines. *Sci. Asia* 27: 93-97.
- Vallidoyan, B. and H.T. Nguyen, 2006. Understanding regulatory networks and engineering for enhanced drought tolerance in plants. *Curr. Opin. Plant Biol.*, 9: 189-195. DOI: 10.1016/j.pbi.2006.01.019
- Yamada, M., H. Morishita, K. Urano, N. Shiozaki and K. Yamaguchi-Shinozaki *et al.*, 2005. Effects of free proline accumulation in petunias under drought stress. *J. Exp. Bot.*, 56: 1975-1981. DOI: 10.1093/jxb/eri195
- Yordanov, I., V. Velikova and T. Tsonev, 2003. Plant responses to drought and stress tolerance. *Bulg. J. Plant Physiol. Special*, 2003: 187-206.
- Zhao, D., B. Glaz and J.C. Comstock, 2010. Sugarcane response to water-deficit stress during early growth on organic and sand soils. *Am. J. Agric. Biol. Sci.*, 5: 403-414. DOI: 10.3844/ajabssp.2010.403.414
- Zingaretti, S.M., F.A. Rodrigues, J.P.D. Graca, L.D.M. Pereira and M.V. Lourenco, 2012. Sugarcane Responses at Water Deficit Conditions. In: *Water Stress*, Rahman, I.M.M. and H. Hasegawa (Eds.), InTech, ISBN: 9789533079639, pp: 13-13.

Simple Publication Process

Submission

Evaluation

Publication

CONTACT US

America
Science Publications
S-207, 244, 5th Avenue
New York, NY 10001

E: support@thescipub.com
F: +1 (866) 350 7082

Australia
Science Publications
S-71, 1a, 400 King William Street
Adelaide, SA 5000

E: support.au@thescipub.com
F: +61 8 7100 1123