test its efficacy, and he has arrived at the following conclusions:—

1. The direct contact of cold water with the normal spleen causes it to contract appreciably, and the degree of contraction varies according to the temperature of the water and the length of time for which it is employed.
2. Cold water also acts similarly on the normal spleen, if applied over the abdominal walls, but it is less effectual. In such cases the cold douche is to be preferred to cold compresses or bits of ice, but in all of them it accomplishes less than quinine.
3. In all forms of enlarged spleens, the result of disease, whether acute or chronic, the action of cold water causes contraction of the organ.
4. The paroxysms of intermittent fever can be arrested by Fleury's method of using the cold douche, but it has no advantage over quinine, either in new or old cases.
5. The cold douche does not accomplish a permanent cure of this fever. It does not always prevent recurrence, and leaves the spleen enlarged.
6. The spleen of typhoid fever is diminished in size by the same remedy.
7. In all forms of enlargement of the spleen, whether acute or chronic, the application of cold water to the spleen, either in the form of cold baths, ice-bags, or cold douches, with the internal use of quinine at the same time, is to be preferred to either of these methods alone.
8. As a substitute for quinine in chronic cases, quinoidine is recommended as being very much cheaper.—*Archiv. f. Fath. Anat.*, lvii. 1.

THE PULVIS GLYCYRRHIZÆ COMPOSITUS.—The compound liquorice powder is a preparation of the Prussian Pharmacopœia, especially useful in simple constipation resulting from atony of the bowel. It is composed of the following constituents, so prepared as to form when incorporated an almost impalpable powder: Senna leaves, ℥vj.; liquorice root, ℥vj.; fennel seeds, ℥ij.; sulphur, ℥ij.; refined sugar, ℥xvij. The usual dose is a small teaspoonful at bedtime, in water, forming an agreeable draught.—*New York Med. Rec.*

NEPHROTOMY.—W. W. Dawson, M.D., of Cincinnati, Ohio (*N. Y. Med. Jour.*), reports a case of extraction of a calculus from the kidney of a married woman, aged 50 years, who had borne nine children, and had always been a healthy and robust woman until eight years ago, when she had an attack of hæmaturia. The patient died on fifth day. The calculus was quite light, weighing only $20\frac{3}{4}$ grains, and measuring seven-eighths of an inch in the length by half an inch in breadth at its widest part, being uniform in shape. It was ammoniac magnesian phosphate, and situated on the internal wall at a point corresponding to the hilus of the kidney, and was embedded in partly disorganized fibrin. The kidney tissue which he incised was about one half-inch in thickness, and bled but little.

THE EARLY SYMPTOMS AND TREATMENT OF INSANITY.—In a paper read before the Bristol North District Medical Society, Dr W. W. Godding, Physician to Lunatic Asylum, Taunton, Mass.

(*Boston Med. and Surg. Jour.*), gives the following symptoms and management of insanity:—In nine cases out of ten, there will be marks of disorder about the clothing or person in some way, for, as a rule, the insane show untidiness of dress; the insane are often uneasy, in a constant motion, getting up, walking about the room, busy with nothing. Later in the disease, they will sit for hours in the same position; sometimes this is so from the commencement, not usually. Depression, as shown in the whole manner as well as in the countenance, is seen in the early stage of insanity, in a majority of cases. Dr Saukey goes so far as to claim that no attack of insanity occurs primarily without an incubating stage of depression. He doubts the truth of this, but it is a very common condition. Hallucinations of hearing are not very uncommon in insanity, and must be regarded as an unfavourable symptom. Hallucinations of sight are, in his experience, much rarer in insanity than in delirium. Want of sleep occurs in a great majority of the cases of insanity, and it is surprising how long they will sometimes survive apparently without sleep, and yet in the end recover. The return of sleep is often the commencement of convalescence. The digestion is impaired in the early stages. Sometimes the moral sentiment alone seems to be at fault. A not very uncommon form of this is seen in young persons recovering from typhoid fever. Paresis seems the very inflorescence of insanity. In regard to the early treatment, in the active forms of insanity, the two important points are to keep up the strength and to secure sleep. The first is best done by nourishment, and in many cases he thinks an early resort to the feeding tube is the humane and the only justifiable course to pursue. There is no doubt but it might be used to advantage much oftener than it is outside of hospitals. No year passes that he does not see lives saved by its use. A full meal is often a better sleep-compeller than an opiate. Milk, egg, and brandy, with a little nutmeg and sugar, is very palatable and anodyne in its action. As a hypnotic, chloral hydrate is the favourite reliance, in doses of 20–30 grains at bed-time; it will bear to be repeated, but not indefinitely. But in mania all drugs are uncertain in their effect. He holds that insane persons are not to be taken indiscriminately to a hospital simply because they are insane.

A RARE CASE OF RESTITUTION OF THE MEMBRANA TYMPANI AFTER FIFTEEN YEARS OF DISEASE. By C. H. Burnett, M.D., Aural Surgeon to Philadelphia Dispensary and the Presbyterian Hospital.—On the last day of July 1872, Christian L., 15 years old, a German, presented himself for treatment at the Philadelphia Dispensary. His complaint was, that since infancy he has had a discharge from the right ear, with progressive hardness of hearing. He says that there has never been any discharge from the left ear; all of which statements were corroborated by his father, who accompanied him.