Indian Medical Gazette.

FEBRUARY.

THE PATHOGENESIS OF
DIPHTHERITIC PARALYSIS.

In the *Quarterly Journal of Medicine*, Vol. 12, Nos. 45 and 46, F. M. R. Walshe publishes a most interesting and instructive paper on the above subject. His conclusions are based on the examination of a large number of cases of post-diphtheritic paralysis, following faucial, cutaneous and wound diphtheria.

The author states that most clinical and experimental observers have regarded the nervous complications of diphtheria as the result of selective hæmatogenous toxæmia of the nervous system. Other hypotheses have also been put forward, e.g., Mendel attributed the palatal paralysis to the fact that the nerves of this structure were bathed in toxin from the neighbouring infective focus; Babonneix and others hold that the process is of the nature of an ascending neuritis from the infective focus.

The author in 1918 advanced strong evidence of a close anatomical relationship between the site of the infective focus and the incidence of the initial paralytic symptoms—evidence which goes to prove that we are dealing with an ascending perineural lymphatic intoxication of the nervous system.

The author emphasises the fact that palatal paralysis occurs only in faucial diphtheria and is the initial symptom of involvement of the nervous system. The palatal musculature is innervated by fibres arising from the hinder end of the glosso-pharyngeal-vagus-accessorius nucleus (nucleus ambiguus) and carried in the pharyngeal branches of the vagus. The tensor palati supplied by the trigeminal nerve is an exception. The sensory innervation of the soft palate, uvula, tonsil and fauces comes from the tonsillar branches of the glosso-pharyngeal, the small and external palatine branches of the maxillary division of the trigeminal, and possibly from the facial *viâ* the large superficial petrosal nerve. Both the site of the infective focus and the structures paralysed therefore are largely within the innervation of the glosso-pharyngeal-vagus-accessorius nuclear system. The pharyngeal and laryngeal palsies and the weakness of the sterno-mastoids therefore constitute a true "local paralysis."

In extra-faucial diphtheria local paralyses are less evident, perhaps on account of the longer nervous path from the infective focus to the central nervous system. They were noted in eight cases and the condition was especially well marked in the case of a medical officer who developed a diphtheritic infection of the right thumb. This officer developed paresis of the hand and arm muscles, followed a few weeks later by multiple neuritis.

The ocular (paralysis of accommodation) and generalized (multiple neuritis) symptoms were common in both faucial and extra-faucial diphtheria. The ocular symptoms are due to a selective action of the toxin on the oculo-motor nerve nuclei.

The paper is summarized as follows:—

Examination of a series of cases of diphtheritic paralysis following faucial and extra-faucial (wound and cutaneous) infections reveals the following points:—

1. That palatal paralysis does not occur except after faucial diphtheria.
2. The musculature of the palate and the region of the infective focus derive their innervation from the same source, the glosso-pharyngeal-vagus-accessorius nuclear system and its peripheral fibres, and are thus closely related anatomically.
3. Similarly in extra-faucial diphtheria the paralysis often shows (27 per cent.) an onset anatomically related to the infective focus.
4. Polyneuritis follows both faucial and extra-faucial diphtheria equally, and, irrespective of the site of the infective focus, presents a remarkably constant symptom-complex.
5. Paralysis of accommodation also follows both forms of infection, though the faucial more frequently (86 per cent.) than the extra-faucial (33 per cent.). It is, therefore, not a "local" paralysis, and its more constant association with faucial infections may be explained by the close proximity of the local central nervous lesion in this case to the oculo-motor nuclei. To this extent a "local" factor may be admitted.
6. The grouping of the symptoms of tetanus, adopted by Meyer and Ransom, into local, specific, and generalized, may be applied equally well to the nervous phenomena of diphtheria. The palatal paralysis constitutes the "local," the ocular affection the "specific," and the polyneuritis the "generalized" forms of diphtheritic paralysis. In extra-faucial diphtheria, as in tetanus, the "local" paralysis varies with the site of the infective focus.

7. As for the pathological processes underlying this symptom-complex, it seems highly probable that as regards the local paralysis we are dealing with an "ascending lymphogenous toxi-infection" (by the perineural lymphatics) of the central nervous system from the infective focus. The essential pathological lesion here is central and not peripheral, and is situated in the nuclear complex already described.

8. The ocular and generalized symptoms are probably the result of the circulation of the toxin in the blood-stream, whence it gains access to the whole nervous system, central and peripheral. The essential nervous lesion here is probably both central and peripheral.

These conclusions, derived from clinical observations, clearly need confirmation by experimental and pathological investigations on lines suggested by the researches of Orr and Rows and various workers on tetanus.

Those that have been made are inconclusive, probably from a lack of close correlation with the clinical aspects of the disease, and in part from the absence of any clear conception of the pathological processes involved.

EPINEPHRIN IN TYPHOID.

BARREIRO, writing in the *Journal of American Medical Association*, August, 1919, says he has only twenty odd cases to report, but the benefit from epinephrin treatment was so great that he does not wait for further experience before proclaiming the advantages of suprarenal treatment in typhoid in the tropics. The benefit from epinephrin in the exhaustion of the soldiers in the trenches was mentioned early in the war, and he was impressed with the similarity between the conditions as described in this war—exhaustion from over-work and the exhaustion of patients early in typhoid. This suggested suprarenal treatment in typhoid, as the patients are often in an exhausted condition from the cares and work of business or from late hours and dissipation. He adds that the food in the tropics is too often defective and inappropriate, so that dyspepsia and liver disease are more prevalent in Porto Rico than among the soldiers in the trenches, as the diet of the latter was more rational than is the custom in the tropics. The perpetual summer climate promotes intestinal fermentations and auto-intoxication. The suprarenals soon become unequal to their task, as is evident from the intense weakness and low blood pressure early

in typhoid. The theoretical indication to supply the lacking suprarenal element was abundantly confirmed by the prompt improvement under suprarenal treatment. In a day or two the benefit was plainly evident, the patient feeling stronger, replying lucidly to questions, and the amount of urine increasing while it grew more normal in appearance. It was an actual resurrection in some cases by the second day after 3 drops of the 1:1,000 epinephrin had been given by the mouth in a glass of water twice a day. The improved muscle tone was particularly evident in the tongue which at first was not under control at all. The epinephrin was continued until defervescence in one case described in detail. After suspension for four days, the pulse ran up to 120. An immediate injection of 0.5 c.c. of the epinephrin solution corrected this, and convalescence continued without further mishap. In another case the fever kept up for thirty-three days, but the epinephrin tided the patient along into convalescence. There was no tendency to intestinal hæmorrhage in any of the patients given the epinephrin. Another feature of epinephrin treatment is that the stools lose their diarrhœic character; the epinephrin passing through the alimentary canal seems to induce vaso-constriction and thus checks the tendency to diarrhœa and averts danger of hæmorrhage. Since he began this routine use of epinephrin in tropical typhoid he has not lost a case.

Current Topics.

Glycosuria in Meningitis.

Journal of the American Medical Association,
August, 1919.

GLYCOSURIA seems to be rare in meningitis. De Massary and Tockmann discovered it in 3 of 45 cases; Cazamian in 3 of 9, and Loeper in 1 of 7. In the 3 cases reported in detail the meningitis had become partitioned off in the ventricles; the antiserum by lumbar puncture failed to reach this ventricle focus, and the men died. As these seemed to be the only ones with this involvement of the ventricles, and as these were the only ones with glycosuria, the inference seems justified that the glycosuria was a warning signal that the disease had invaded the ventricles. The irritation from this acted like the sugar puncture of the third ventricle. Trepine puncture of the ventricles thus seems the logical indication with glycosuria in meningitis, judging from the necropsy findings in these 3 cases. The glycosuria is not inevitably a sign of