ABSTRACTS FROM CURRENT MEDICAL LITERATURE.

EDITED BY T. K. MACLACHLAN AND R. G. LENDRUM.

MEDICINE.

Studies concerning Digitalis Therapy in Lobar Pneumonia.

—Niles and Wyckoff (*American Journal of the Medical Sciences*, vol. clxxx, No. 3, September, 1930) give a number of statistics to show the effect of digitalis therapy on the mortality-rate in lobar pneumonia. Cases are grouped under four headings:—(a) Received neither serum nor digitalis; (b) serum only; (c) digitalis only; and (d) both serum and digitalis. Thus groups (a) and (b) acted as controls to groups (c) and (d). The observations were carried out over a period of two years, and, for statistical purposes, the cases have been grouped and tabulated according to the factors which are known materially to affect the mortality-rate in lobar pneumonia, namely, sex, age, type of pneumococcus, severity of infection as indicated by septic complications and positive blood cultures. With only three exceptions the comparisons of mortality per cent in the digitalis treated and in the untreated control cases are in favour of the latter. The first exception is in the group classified according to the type of pneumococcus where in 108 Type II treated cases the mortality was 45·4 per cent against 47·7 per cent in 130 control cases. The second exception is in the series in which the pneumococcus type groups are further subdivided according to the age factor—i.e., patients aged 10 to 39 years, and patients aged 40 and over. In Type II, ages 10 to 39 group, 54 treated cases show a mortality of 31·5 per cent against 39·6 per cent in 53 untreated controls. In explanation of this it is pointed out that if all cases which were admitted to the series (not excluding cases discharged at their own risk), cases in (c) and (d) series which, for other reasons, such as systemic disease, &c., received no digitalis, and deaths occurring within twenty-four hours after admission) are analysed, it is found that in common with all other types the mortality of the digitalis-treated group in Type II infections is higher than in the control group.

In conclusion, the authors state that although they would prefer to continue the investigation and observe a larger number of cases over a series of years, they are of the opinion that the results obtained so far do *not* justify continuing the routine administration of digitalis to patients suffering from lobar pneumonia.—P. M. WOOD.

The Normal Filament and Non-filament Polymorphonuclear Neutrophil Count: Its Practical Value as a Diagnostic Aid. By Farley, St. Clair and Reisinger (*American Journal of the Medical Sciences*, vol. clxxx, No. 3, September, 1930).—Working from the results

obtained by Arneth, Schilling, Piney, Pons and Krumbhaar, and more directly from those achieved by Cooke and Ponder, the authors have evolved a simple method of classifying the different types of polymorphonuclear neutrophils for routine use by those without special training in hæmatology. They have divided the cells into two classes—(1) “non-filament” neutrophils (*i.e.*, cells with an undivided nucleus), and (2) “filament” neutrophils (*i.e.*, cells with a divided nucleus, whether into two, three, four or five lobes). For the basis of classification, Cooke’s criterion of division is taken, namely, “If there is any band of nuclear material except this chromatin filament connecting the different parts of the nucleus, that nucleus cannot for the purposes of the count be said to be divided.” Distinct myelocytes or myeloblasts are to be counted as such.

The authors have attempted to find the upper limit of the normal non-filament count, and an average normal filament and non-filament count, in two ways—(1) by mathematical reduction of the normal chart of Cooke and Ponder; (2) by an actual series of counts on 100 normal individuals. From their work they believe that 16 per cent can be used to practical advantage as the upper limit in health of normal young polymorphonuclear leucocytes (non-filament forms). While admitting that such a figure is somewhat arbitrary, they stress its great value in differentiating between a psycho-neurosis and symptoms due to a hidden focus of disease when there is no increase in the total number of leucocytes. As regards the average non-filament count, they found by their method in 190 presumably healthy subjects that the figure was 8 per cent.

In a brief *résumé*, they state that their method is to differentiate 100 consecutive leucocytes to determine thus the percentage of non-filament neutrophils, filament neutrophils, lymphocytes, monocytes, eosinophils, and basophils, using Cooke’s criterion to separate the polymorphonuclear neutrophils into old and young groups, and, in conclusion, emphasize that their method is not intended to take the place of the more elaborate methods of Cooke and Ponder, Arneth and others, but is designed especially for routine work.

—P. M. WOOD.

BOOKS, PAMPHLETS, &c., RECEIVED.

- Aids to Histology, by Alexander Goodall, M.D., F.R.C.P. Edin. Third edition. London: Baillière, Tindall & Cox. 1930. (3s. 6d. net.)
- Aids to Bacteriology, by William Partridge, F.I.C. Fifth edition. London: Baillière, Tindall & Cox. 1930. (5s. net.)
- An Introduction to Human Experimental Physiology, by F. W. Lamb, M.D. With a Foreword by A. V. Hill, Sc.D., F.R.S. With diagrams. London: Longmans, Green & Co. 1930. (12s. 6d.)
- Elementary Zoology for Medical Students, by L. A. Borradaile, Sc.D. London: Humphrey Milford. 1930. (10s. 6d. net.)