Biogeosciences Discussions is the access reviewed discussion forum of *Biogeosciences*

Microbiotic crusts on soil, rock and plants: neglected major players in the global cycles of carbon and nitrogen?

W. Elbert¹, B. Weber², B. Büdel², M. O. Andreae¹, and U. Pöschl¹

¹Max Planck Institute for Chemistry, Biogeochemistry Department, P. O. Box 3060,
55020, Mainz, Germany

²University of Kaiserslautern, Dept. Biology, Plant Ecology and Systematics, P. O. Box 3049,
67653 Kaiserslautern, Germany

Received: 17 June 2009 – Accepted: 6 July 2009 – Published: 14 July 2009

Correspondence to: W. Elbert (elbert@mpch-mainz.mpg.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

6983

Abstract

Microbiotic crusts consisting of bacteria, fungi, algae, lichens, and bryophytes colonize most terrestrial surfaces, and they are able to fix carbon and nitrogen from the atmosphere. Here we show that microbiotic crusts are likely to play major roles in the global biogeochemical cycles of carbon and nitrogen, and we suggest that they should be further characterized and taken into account in studies and models of the Earth system and climate.

For the global annual net uptake of carbon by microbiotic crusts we present a first estimate of $\sim 3.6 \text{ Pg a}^{-1}$. This uptake corresponds to $\sim 6\%$ of the estimated global net carbon uptake by terrestrial vegetation (net primary production, NPP: $\sim 60 \text{ Pg a}^{-1}$), and it is of the same magnitude as the global annual carbon turnover due to biomass burning. The estimated rate of nitrogen fixation by microbiotic crusts ($\sim 45 \text{ Tg a}^{-1}$) amounts to $\sim 40\%$ of the global estimate of biological nitrogen fixation (107 Tg a^{-1}). With regard to Earth system dynamics and global change, the large contribution of microbiotic crusts to nitrogen fixation is likely to be important also for the sequestration of CO_2 by terrestrial plants (CO_2 fertilization), because the latter is constrained by the availability of fixed nitrogen.

1 Introduction

Microbiotic crusts consisting of bacteria, fungi, algae, lichens, and bryophytes colonize most terrestrial surfaces (Figs. 1 and S1, see <http://www.biogeosciences-discuss.net/6/6983/2009/bgd-6-6983-2009-supplement.pdf>). In the form of biological soil crusts (BSC) and rock crusts (BRC), they inhabit soils and rocks in arid and semiarid regions as well as in other climatic zones (Friedmann, 1980; Friedmann et al., 1993; Matthes-Sears et al., 1997; Kappen et al., 1998; Büdel, 1999; Kurina and Vitousek, 1999; Zaady et al., 2000; Büdel, 2002; Belnap, 2003; Belnap and Lange, 2003; Rascher et al., 2003; Boisson et al., 2004; Hoppert et al., 2004; Walker et al., 2005; Bhatnagar et al., 2008;

6984

Büdel et al., 2008; Schmidt et al., 2008; Weber et al., 2008; Wilske et al., 2008). Microbiotic crusts also cover a large proportion of the surface of terrestrial plants, including stems and branches of trees (epiphytic crusts) as well as tree leaves (epiphyllic crusts) (Ruinen, 1956; Last and Deighton, 1965; Berrie and Eze, 1975; Bentley, 1987; Sheridan, 1991a; Peveling et al., 1992; Coley et al., 1993; Freiberg, 1999; Büdel et al., 2000; Freiberg and Freiberg, 2000; Sillett and Rambo, 2000; Campbell and Coxson, 2001; Lücking and Matzer, 2001; Anthony et al., 2002; Radies and Coxson, 2004; Wanek and Portl, 2005; Lakatos et al., 2006; Cáceres et al., 2007). For the sake of brevity, we shall include both types in the term “epiphytic crusts” (EPC).

It is well known that these communities are able to fix carbon and nitrogen from the atmosphere (Mayland and McIntosh, 1966; Jones, 1970; Bentley, 1987; Beymer and Klopatek, 1991; Lange et al., 1992; Freiberg, 1998; Evans and Johansen, 1999; Juhász et al., 2002; Belnap, 2003; Boison et al., 2004; Yeager et al., 2007). They produce biomass and they release C- and N-containing compounds like carbohydrates and amino acids to the surrounding ecosystem (Boucher and Nash, 1990; Beymer and Klopatek, 1991; Coxson et al., 1992; Belnap and Lange, 2003; Turetsky, 2003; Dojani et al., 2007; Schmidt et al., 2008). However, their contribution to the global cycles of carbon and nitrogen has not yet been quantified. In this study we calculate and present first estimates of the global annual net uptake of CO₂ and fixation of N₂ by microbiotic crusts.

2 Methods

From earlier studies of microbiotic crusts we have compiled as many estimates of annual uptake fluxes and surface coverage values as we could find in a thorough literature search. For global upscaling we used median values of the flux data in combination with global estimates of the surface areas of (semi)arid regions and plants, respectively. All used data and references are tabulated in the Appendix (Tables A1–A9).

Note that we used median rather than arithmetic mean values in order to obtain

6985

conservative estimates. The median values were generally in fair agreement with the corresponding arithmetic mean values, and the relative standard errors of the mean values (relative standard deviation divided by the square root of the number of data points) ranged up to ~80%. Thus we assume that the presented global estimates are uncertain by a factor of ~2.

3 Results and discussion

For the net uptake of carbon by soil crusts in arid and semi-arid regions, we obtained a median flux of $\sim 16 \text{ g m}^{-2} \text{ a}^{-1}$ (Table A1a). Multiplication with the global dry-land area (Table A2) yields an estimate of $\sim 1.0 \text{ Pg a}^{-1}$ for the net uptake of carbon by BSC in arid and semi-arid regions. With regard to the carbon balance, the reported net uptake values (photosynthesis minus respiration) should be considered as net primary production (NPP) (Chapin et al., 2006).

Soil crusts occur also in non-vegetated gaps of grasslands, tundra and steppe formations, sparsely vegetated grounds of temperate and boreal forests, burnt forest areas, formerly permafrosted soils, and previously ice-covered glacier grounds (Forman and Dowden, 1977; Eversman and Horton, 2004; Schmidt et al., 2008). The median net carbon uptake by soil crusts in these areas is $\sim 23 \text{ g m}^{-2} \text{ a}^{-1}$ (Table A1b), and that by rock crusts is $\sim 8 \text{ g m}^{-2} \text{ a}^{-1}$ (Table A1c). Because the global area colonized by non-arid BSC and by rock crusts are not known, we cannot provide a quantitative estimate for their carbon uptake, but obviously any non-zero flux will increase the total amount of carbon taken up by biological crusts to more than the value of $\sim 1 \text{ Pg a}^{-1}$ given above for arid and semiarid soil crusts alone.

For the net uptake of carbon by epiphytic crusts, we obtained a median flux of $\sim 28 \text{ g m}^{-2} \text{ a}^{-1}$ (Table A3). Multiplication with the corresponding global surface areas of evergreen leaves, branches and stems of trees, and tropical lianas (Tables A4, A5, and A7), assuming coverages of 30–50% (Tables A6 and A7), yields an estimate of $\sim 2.6 \text{ Pg a}^{-1}$ for the global net uptake of carbon by epiphytic crusts (Table A7).

6986

The total value of $\sim 3.6 \text{ Pg a}^{-1}$ estimated for the global net uptake of carbon by microbiotic crusts corresponds to $\sim 6\%$ of the estimated global net carbon uptake by terrestrial vegetation (net primary production, NPP: $\sim 60 \text{ Pg a}^{-1}$, Running et al., 2004). To put this flux into perspective: it is of the same magnitude as the global annual carbon turnover due to biomass burning, which has been estimated at 3.6 Pg a^{-1} (Andreae and Merlet, 2001).

The reported values of biomass in microbiotic crusts are in the range of $1\text{--}1200 \text{ g m}^{-2}$ (dry mass) with median values of 260 g m^{-2} for soil crusts and 130 g m^{-2} for epiphytic crusts (Table A8). By multiplication of these values with the global areas of dry-lands and forests/shrubs, respectively (Tables A2 and A7) we obtain global estimates of $\sim 17 \text{ Pg}$ for the biomass of BSC in arid and semi-arid regions and $\sim 6 \text{ Pg}$ for the biomass of EPC on evergreen leaves, on branches and stems of trees, and on tropical lianas. The total of $\sim 23 \text{ Pg}$ estimated for the global dry biomass of microbiotic crusts corresponds to $\sim 10 \text{ Pg}$ of carbon (conversion factor ~ 2) (Whitman et al., 1998), and thus to $\sim 2\%$ of the estimated global mass of carbon in terrestrial vegetation ($470\text{--}650 \text{ Pg}$) (Prentice et al., 2001).

Microbiotic crusts do not only fix carbon, but are also able to assimilate atmospheric nitrogen. The reported average fluxes of nitrogen fixation are in the range of $0.1\text{--}10 \text{ g m}^{-2} \text{ a}^{-1}$ (Table A9). With the median values of $\sim 0.4 \text{ g m}^{-2} \text{ a}^{-1}$ for BSC and $\sim 0.35 \text{ g m}^{-2} \text{ a}^{-1}$ for EPC (Table A9) and using the same surface area values as above, we obtain global estimates of $\sim 30 \text{ Tg a}^{-1}$ and $\sim 15 \text{ Tg a}^{-1}$ for nitrogen fixation by BSC and EPC, respectively. Thus, the total rate of nitrogen fixation by microbiotic crusts ($\sim 45 \text{ Tg a}^{-1}$) appears to be a major contribution to global biological nitrogen fixation, which is estimated to be about 107 Tg a^{-1} (Galloway, 2005). Note that nitrogen fixation by microbiotic crusts is likely to be important also for the sequestration of CO_2 by terrestrial plants (CO_2 fertilization), because the latter is constrained by the availability of fixed nitrogen (Reich et al., 2006).

6987

4 Conclusions

Overall, our calculations suggest that microbiotic crusts on soil, rock, and plants are major players in the global biogeochemical cycles of carbon and nitrogen and should thus be considered in climate and Earth system models. Regional and seasonal patterns as well as long-term trends regarding their diversity, abundance and gas exchange with the atmosphere need to be better characterized for a full mechanistic and quantitative understanding of the influence of the biosphere on climate.

Appendix A

See Tables A1–A9.

Acknowledgements. This work has been funded by the Max Planck Society (W. E., M. O. A., U. P.) and by the German Research Foundation (B. B., B. W.). Thanks to J. Steinkamp and E. Falge for support in the determination of geographical surface areas and leaf area indices, respectively.

The service charges for this open access publication have been covered by the Max Planck Society.

References

- Ajtay, G. L., Ketner, P., and Duvigneaud, P.: Terrestrial primary production and phytomass, in: The Global Carbon Cycle, SCOPE 13, edited by: Bolin, B., Degens, E. T., and Ketner, P., John Wiley & Sons, New York, 129–182, 1979.
- André, J., Gourbière, F., and Bardin, R.: Epiphytic lichens and microfungi associated to needles in a fir forest of Massif Central, *Oecologia Plantarum*, 10, 13–23, 1975.
- Andreae, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global Biogeochem. Cy.*, 15, 955–966, 2001.

6988

- Anthony, P. A., Holtum, J. A. M., and Jackes, B. R.: Shade acclimation of rainforest leaves to colonization by lichens, *Funct. Ecol.*, 16, 808–816, 2002.
- Antoine, M. E.: An ecophysiological approach to quantifying nitrogen fixation by *Lobaria oregana*, *Bryologist*, 107, 82–87, 2004.
- 5 Arora, V. K. and Boer, G. J.: A parameterization of leaf phenology for the terrestrial ecosystem component of climate models, *Glob. Change Biol.*, 11, 39–59, 2005.
- Asner, G. P., Archer, S., Hughes, R. F., Ansley, R. J., and Wessman, C. A.: Net changes in regional woody vegetation cover and carbon storage in Texas Drylands, 1937–1999, *Glob. Change Biol.*, 9, 316–335, 2003.
- 10 Becker, V. E.: Nitrogen fixing lichens in forests of the southern Appalachian Mountains of North Carolina, *Bryologist*, 83, 29–39, 1980.
- Bell, R. A. and Sommerfeld, M. R.: Algal biomass and primary production within a temperate zone sandstone, *Am. J. Bot.*, 74, 294–297, 1987.
- Belnap, J., Kaltenecker, J. H., Rosentreter, R., Williams, J., Leonard, S., and Eldridge, D.: Biological soil crusts: ecology and management, Technical reference 1730-2, US Department of the Interior, Denver, 1–118, 2001.
- 15 Belnap, J.: Nitrogen fixation in biological soil crusts from southeast Utah, USA, *Biol. Fert. Soils*, 35, 128–135, 2002.
- Belnap, J.: The world at your feet: desert biological soil crusts, *Frontiers in Ecology and the Environment*, 1, 181–189, 2003.
- 20 Belnap, J. and Lange, O. L.: Structure and functioning of biological soil crusts: synthesis, in: *Biological soil crusts: structure, function, and management*, 2nd ed., edited by: Belnap, J. and Lange, O. L., Ecological Studies, Springer-Verlag, Berlin Heidelberg, 471–479, 2003.
- Benson, S. and Coxson, D. S.: Lichen colonization and gap structure in wet-temperate rainforests of northern interior British Columbia, *Bryologist*, 105, 673–692, 2002.
- 25 Bentley, B. L.: Nitrogen fixation by epiphylls in a tropical rainforest, *Ann. Mo. Bot. Gard.*, 74, 234–241, 1987.
- Bernstein, M. E. and Carroll, G. C.: Microbial populations on Douglas fir needle surfaces, *Microbial Ecol.*, 4, 41–52, 1977.
- 30 Berrie, G. K. and Eze, J. M. O.: Relationship between an epiphyllous liverwort and host leaves, *Ann. Bot.-London*, 39, 955–963, 1975.
- Beymer, R. J. and Klopatek, J. M.: Potential contribution of carbon by microphytic crusts in pinyon-juniper woodlands, *Arid Soil Res. Rehab.*, 5, 187–198, 1991.

6989

- Bhatnagar, A., Makandar, M. B., Garg, M. K., and Bhatnagar, M.: Community structure and diversity of cyanobacteria and green algae in the soils of Thar Desert (India), *J. Arid Environ.*, 72, 73–83, 2008.
- Boison, G., Mergel, A., Jolkver, H., and Bothe, H.: Bacterial life and dinitrogen fixation at a gypsum rock, *Appl. Environ. Microb.*, 70, 7070–7077, 2004.
- 5 Boucher, V. L. and Nash, T. H.: Growth patterns in *Ramalina menziesii* in California: coastal vs. inland populations, *Bryologist*, 93, 295–302, 1990.
- Breda, N. J. J.: Ground-based measurements of leaf area index: a review of methods, instruments and current controversies, *J. Exp. Bot.*, 54, 2403–2417, doi:10.1093/jxb/erg263, 2003.
- 10 Brostoff, W. N., Sharifi, M. R., and Rundel, P. W.: Photosynthesis of cryptobiotic soil crusts in a seasonally inundated system of pans and dunes in the western Mojave Desert, CA: Field studies, *Flora*, 200, 592–600, 2005.
- Brown, P. J. and Dalton, D. A.: *In situ* physiological monitoring of *Lobaria oregana* transplants in an old-growth forest canopy, *Northwest Science*, 76, 230–239, 2002.
- 15 Büdel, B.: Ecology and diversity of rock-inhabiting cyanobacteria in tropical regions, *Eu. J. Phycol.*, 34, 361–370, 1999.
- Büdel, B., Meyer, A., Salazar, N., Zellner, H., Zotz, G., and Lange, O. L.: Macrolichens of montane rain forests in Panama, Province Chiriqui, *Lichenologist*, 32, 539–551, 2000.
- 20 Büdel, B.: Diversity and ecology of biological crusts, in: *Progress in Botany*, edited by: Lüttge, U. E., Beyschlag, W., Büdel, B., and Francis, D., Springer-Verlag, Berlin – Heidelberg, 386–404, 2002.
- Büdel, B., Weber, B., Köhl, M., Pfanz, H., Sültemeyer, D., and Wessels, D.: Reshaping of sandstone surfaces by cryptoendolithic cyanobacteria: bioalkalization causes chemical weathering in arid landscapes, *Geobiology*, 2, 261–268, 2004.
- 25 Büdel, B., Bendix, J., Bicker, F. R., and Green, T. G. A.: Dewfall as a water source frequently activates the endolithic cyanobacterial community in the granites of Taylor Valley, Antarctica, *J. Phycol.*, 44, 1415–1424, 2008.
- Buermann, W., Wang, Y. J., Dong, J. R., Zhou, L. M., Zeng, X. B., Dickinson, R. E., Potter, C. S., and Myneni, R. B.: Analysis of a multiyear global vegetation leaf area index data set, *J. Geophys. Res.-Atmos.*, 107, 4646, doi:10.1029/2001JD000975, 2002.
- 30 Cáceres, M. E. S., Lücking, R., and Rambold, G.: Phorophyte specificity and environmental parameters versus stochasticity as determinants for species composition of corticolous

6990

- crustose lichen communities in the Atlantic rain forest of northeastern Brazil, *Mycological Progress*, 6, 117–136, 2007.
- Caldwell, M. M., Meister, H.-P., Tenhunen, J. D., and Lange, O. L.: Canopy structure, light microclimate and leaf gas exchange of *Quercus coccifera* L. in a Portuguese macchia: measurements in different canopy layers and simulations with a canopy model, *Trees*, 1, 25–41, 1986.
- Campbell, J. and Coxson, D. S.: Canopy microclimate and arboreal lichen loading in subalpine spruce-fir forest, *Can. J. Bot.*, 79, 537–555, 2001.
- Carroll, G. C.: Needle microepiphytes in a Douglas-fir canopy – biomass and distribution patterns, *Can. J. Bot.-Revue Canadienne De Botanique*, 57, 1000–1007, 1979.
- Castellanos, V. A. E., Duran, R., Guzman, S., Briones, O., and Feria, M.: Three-dimensional space utilization of lianas: a methodology, *Biotropica*, 24, 396–401, 1992.
- Chapin, F. S., Woodwell, G. M., Randerson, J. T., Rastetter, E. B., Lovett, G. M., Baldocchi, D. D., Clark, D. A., Harmon, M. E., Schimel, D. S., Valentini, R., Wirth, C., Aber, J. D., Cole, J. J., Goulden, M. L., Harden, J. W., Heimann, M., Howarth, R. W., Matson, P. A., McGuire, A. D., Melillo, J. M., Mooney, H. A., Neff, J. C., Houghton, R. A., Pace, M. L., Ryan, M. G., Running, S. W., Sala, O. E., Schlesinger, W. H., and Schulze, E. D.: Reconciling carbon-cycle concepts, terminology, and methods, *Ecosystems*, 9, 1041–1050, 2006.
- Clark, D. B., Olivas, P. C., Oberbauer, S. F., Clark, D. A., and Ryan, M. G.: First direct landscape-scale measurement of tropical rain forest leaf area index, a key driver of global primary productivity, *Ecol. Lett.*, 11, 163–172, 2008.
- Clark, D. L., Nadkarni, N. M., and Gholz, H. L.: Growth, net production, litter decomposition, and net nitrogen accumulation by epiphytic bryophytes in a tropical montane forest, *Biotropica*, 30, 12–23, 1998.
- Coley, P. D., Kursar, T. A., and Machado, J.-L.: Colonization of tropical rain forest leaves by epiphylls: effects of site and host plant leaf lifetime, *Ecology*, 74, 619–623, 1993.
- Coxson, D. S., McIntyre, D. D., and Vogel, H. J.: Pulse release of sugars and polyols from canopy bryophytes in tropical montane rain-forest (Guadeloupe, French West-Indies), *Biotropica*, 24, 121–133, 1992.
- Currin, C. A. and Pearl, H. W.: Epiphytic nitrogen fixation associated with standing dead shoots of smooth cordgrass, *Spartina alterniflora*, *Estuaries*, 21, 108–117, 1998.
- DeLuca, T. H., Zackrisson, O., Nilsson, M. C., and Sellstedt, A.: Quantifying nitrogen-fixation in feather moss carpets of boreal forests, *Nature*, 419, 917–920, 2002.

6991

- Devakumar, A. S., Prakash, P. G., Sathik, M. B. M., and Jacob, J.: Drought alters the canopy architecture and micro-climate of *Hevea brasiliensis* trees, *Trees-Struct. Funct.*, 13, 161–167, 1999.
- Dojani, S., Lakatos, M., Rascher, U., Wanek, W., Lüttge, U., and Büdel, B.: Nitrogen input by cyanobacterial biofilms of an inselberg into a tropical rainforest in French Guiana, *Flora*, 202, 521–529, 2007.
- Dufrêne, E. and Bréda, N.: Estimation of deciduous forest leaf area index using direct and indirect methods, *Oecologia*, 104, 156–162, 1995.
- Ellyson, W. J. T. and Sillett, S. C.: Epiphyte communities on Sitka spruce in an old-growth redwood forest, *Bryologist*, 106, 197–211, 2003.
- Evans, R. D. and Johansen, J. R.: Microbiotic crusts and ecosystem processes, *Cr. Rev. Plant Sci.*, 18, 183–225, 1999.
- Evans, R. D. and Lange, O. L.: Biological soil crusts and ecosystem nitrogen and carbon dynamics, in: *Structure, function, and management*, 2nd ed., edited by: Belnap, J. and Lange, O. L., Ecological studies, Springer-Verlag, Berlin, 263–279, 2003.
- Eversman, S. and Horton, D.: Recolonization of burned substrates by lichens and mosses in Yellowstone National Park, *Northwest Science*, 78, 85–92, 2004.
- Falge, E., Ryel, R. J., Alsheimer, M., and Tenhunen, J. D.: Effects of stand structure and physiology on forest gas exchange: a simulation study for Norway spruce, *Trees-Struct. Funct.*, 11, 436–448, 1997.
- Forman, R. T. T.: Canopy lichens with blue-green algae: a nitrogen source in a Colombian rain forest, *Ecology*, 56, 1176–1184, 1975.
- Forman, R. T. T. and Dowden, D. L.: Nitrogen fixing lichen roles from desert to alpine in the Sangre de Cristo Mountains, New Mexico, *Bryologist*, 80, 561–570, 1977.
- Frahm, J.-P.: Die Zunahme von epiphytischen Hängemoosen in Europa am Beispiel einer Lokalität in den Vogesen, *Archive for Bryology*, 35, 1–10, 2008.
- Freiberg, E.: Microclimatic parameters influencing nitrogen fixation in the phyllosphere in a Costa Rican premontane rain forest, *Oecologia*, 17, 9–18, 1998.
- Freiberg, E.: Influence of microclimate on the occurrence of cyanobacteria in the phyllosphere in a premontane rain forest of Costa Rica, *Plant Biol.*, 1, 244–252, 1999.
- Freiberg, M. and Freiberg, E.: Epiphyte diversity and biomass in the canopy of lowland and montane forests in Ecuador, *J. Trop. Ecology*, 16, 673–688, 2000.
- Friedmann, E. I.: Endolithic microbial life in hot and cold deserts, *Orig. Life Evol. Biosph.*, 10,

6992

- 223–235, 1980.
- Friedmann, E. I., Kappen, L., Meyer, M. A., and Nienow, J. A.: Long-term productivity in the cryptoendolithic microbial community of the Ross Desert, Antarctica, *Microbial Ecol.*, 25, 51–69, 1993.
- 5 Fűrnkranz, M., Wanek, W., Richter, A., Abell, G., Rasche, F., and Sessitsch, A.: Nitrogen fixation by phyllosphere bacteria associated with higher plants and their colonizing epiphytes of a tropical lowland rainforest of Costa Rica, *The ISME Journal*, 2, 561–570, 2008.
- Galloway, J. N.: The global nitrogen cycle, in: *Biogeochemistry*, edited by: Schlesinger, W. H., *Treatise on Geochemistry*, Elsevier-Pergamon, Oxford, 557–583, 2005.
- 10 Garcia-Pichel, F. and Belnap, J.: Microenvironments and microscale productivity of cyanobacterial desert crusts, *J. Phycol.*, 32, 77–782, 1996.
- Garcia-Pichel, F., Belnap, J., Neuer, S., and Schanz, F.: Estimates of global cyanobacterial biomass and its distribution, in: *Algological Studies*, edited by: Chapman, R., Hoffmann, L., Komárková, J., Krienitz, L., and Kristiansen, J., E. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, 213–227, 2003.
- 15 Gaydarova, P. N.: Deciduous forest communities in the Black Sea coastal Strandzha region: temporal and spatial characteristics of leaf area index and density, *Trees-Struct. Funct.*, 17, 237–243, 2003.
- Gerwing, J. J. and Farias, D. L.: Integrating liana abundance and forest stature into an estimate of total aboveground biomass for an eastern Amazonian forest, *J. Trop. Ecology*, 16, 327–335, 2000.
- 20 Gerwing, J. J. and Vidal, E.: Changes in liana abundance and species diversity eight years after liana cutting and logging in an eastern Amazonian forest, *Conserv. Biol.*, 16, 544–548, 2002.
- 25 Goldewijk, K. K.: Estimating global land use change over the past 300 years: The HYDE Database, *Global Biogeochem. Cy.*, 15, 417–433, 2001.
- Gratani, L. and Varone, L.: Plant crown traits and carbon sequestration capability by *Platanus hybrida* Brot. in Rome, *Landscape Urban Plan.*, 81, 282–286, 2007.
- Hanson, R. B.: Nitrogen fixation (acetylene reduction) in a salt marsh amended with sewage sludge and organic carbon and nitrogen compounds, *Appl. Environ. Microb.*, 33, 846–852, 30 1977.
- Hoppert, M., Flies, C., Pohl, W., Gunzl, B., and Schneider, J.: Colonization strategies of lithobiontic microorganisms on carbonate rocks, *Environ. Geol.*, 46, 421–428, 2004.

6993

- Horne, A. J.: The ecology of nitrogen fixation on Signy Island South Orkney Islands, *Br. Antarct. Surv. Bull.*, 27, 1–18, 1972.
- Housman, D. C., Powers, H. H., Collins, A. D., and Belnap, J.: Carbon and nitrogen fixation differ between successional stages of biological soil crusts in the Colorado Plateau and Chihuahuan Desert, *J. Arid Environ.*, 66, 620–634, 2006.
- 5 Isichei, A. O.: Nitrogen fixation by blue-green algal soil crusts in Nigerian savanna, in: *Nitrogen cycling in West Africa ecosystems*, edited by: Rosswall, T., Royal Swedish Academy of Sciences, Stockholm, 191–198, 1980.
- Jeffries, D. L., Link, S. O., and Klopatek, J. M.: CO₂ fluxes of cryptogamic crusts I. Response to resaturation, *New Phytol.*, 125, 163–173, 1993.
- 10 Jones, K.: Nitrogen fixation in phyllosphere of Douglas-fir, *Pseudotsuga-Douglasii*, *Ann. Bot.-London*, 34, 239–244, 1970.
- Juhász, A., Balogh, J., Csintalan, Z., and Tuba, Z.: Carbon sequestration of the poikilohydric moss carpet vegetation in semidesert sandy grassland ecosystem, *Acta Biologica Szegediensis*, 46, 223–225, 2002.
- 15 Kappen, L., Lange, O. L., Schulze, E.-D., Buschbom, U., and Evenari, M.: Ecophysiological investigations on lichens of the Negev Desert. VII. The influence of the habitat exposure on dew imbibition and photosynthetic productivity., *Flora*, 169, 216–229, 1980.
- Kappen, L., Schroeter, B., Green, T. G. A., and Seppelt, R. D.: Chlorophyll *a* fluorescence and CO₂ exchange of *Umbilicaria aprina* under extreme light stress in the cold, *Oecologia*, 113, 325–331, 1998.
- 20 Kazda, M. and Salzer, J.: Leaves of lianas and self-supporting plants differ in mass per unit area and in nitrogen content, *Plant Biol.*, 2, 268–271, 2000.
- Klopatek, J. M.: The use of cryptogamic crusts as indicators of disturbance in semiarid landscapes, in: *Ecological indicators*, edited by: Kenzie, D. H. M., Hyatt, D. E., and MacDonald, V. J., Elsevier, London, 773–786, 1992.
- 25 Kucharik, C. J., Foley, J. A., Delire, C., Fisher, V. A., Coe, M. T., Lenters, J. D., Young-Molling, C., Ramankutty, N., Norman, J. M., and Gower, S. T.: Testing the performance of a Dynamic Global Ecosystem Model: Water balance, carbon balance, and vegetation structure, *Global Biogeochem. Cy.*, 14, 795–825, 2000.
- 30 Kurina, L. M. and Vitousek, P. M.: Controls over the accumulation and decline of a nitrogen-fixing lichen, *Stereocaulon vulcani*, on young Hawaiian lava flows, *J. Ecol.*, 87, 784–799, 1999.

6994

- Lakatos, M., Rascher, U., and Büdel, B.: Functional characteristics of corticolous lichens in the understory of a tropical lowland rain forest, *New Phytol.*, 172, 679–695, 2006.
- Lange, O. L., Schulze, E.-D., and Koch, W.: Experimentell-ökologische Untersuchungen an Flechten der Negev-Wüste. III. CO₂-Gaswechsel und Wasserhaushalt von Krusten- und Blattflechten am natürlichen Standort während der sommerlichen Trockenperiode, *Flora*, 159, 525–528, 1970.
- Lange, O. L., Kidron, G. J., Büdel, B., Meyer, A., Kilian, E., and Abeliovich, A.: Taxonomic composition and photosynthetic characteristics of the 'biological soil crusts' covering sand dunes in the wester Negev Desert, *Funct. Ecol.*, 6, 519–527, 1992.
- 10 Lange, O. L., Büdel, B., Meyer, A., and Kilian, E.: Further evidence that activation of net photosynthesis by dry cyanobacterial lichens requires liquid water, *Lichenologist*, 25, 175–189, 1993.
- Lange, O. L., Büdel, B., Zellner, H., Zotz, G., and Meyer, A.: Field measurements of water relations and CO₂ exchange of the tropical, cyanobacterial basidiolichen *Dictyonema glabratum* in a Panamanian rainforest, *Botanica Acta*, 107, 279–290, 1994a.
- 15 Lange, O. L., Meyer, A., Zellner, H., and Heber, U.: Photosynthesis and water relations of lichen soil crusts: field measurements in the coastal fog zone of the Namib Desert, *Funct. Ecol.*, 8, 253–264, 1994b.
- Lange, O. L., Belnap, J., Reichenberger, H., and Meyer, A.: Photosynthesis of green algal soil crust lichens from arid lands in southern Utah, USA: Role of water content on light and temperature responses of CO₂ exchange, *Flora*, 192, 1–15, 1997.
- 20 Lange, O. L., Belnap, J., and Reichenberger, H.: Photosynthesis of the cyanobacterial soil-crust lichen *Collema tenax* from arid lands in southern Utah, USA: Role of water content on light and temperature responses of CO₂ exchange, *Funct. Ecol.*, 12, 195–202, 1998a.
- 25 Lange, O. L., Hahn, S. C., Meyer, A. and Tenhunen, J. D.: Upland tundra in the foothills of the Brooks Range, Alaska, USA: lichen long-term photosynthetic CO₂ uptake and net carbon gain, *Arctic Alpine Res.*, 30, 252–261, 1998b.
- Lange, O. L.: Die Lebensbedingungen von Bodenkrusten-Organismen: Tagesverlauf der Photosynthese einheimischer Erdflechten, *Hoppea*, 61, 423–443, 2000a.
- 30 Lange, O. L.: Photosynthetic performance of a gelatinous lichen under temperate habitat conditions: long-term monitoring of CO₂ exchange of *Collema cristatum*, *Bibliotheca Lichenologica*, 75, 307–332, 2000b.
- Lange, O. L., Büdel, B., Meyer, A., Zellner, H., and Zotz, G.: Lichen carbon gain under trop-

6995

- ical conditions: water relations and CO₂ exchange of three *Leptogium* species of a lower montane rainforest in Panama, *Flora*, 195, 172–190, 2000.
- Lange, O. L.: Photosynthetic productivity of the epilithic lichen *Lecanora muralis*: long-term field monitoring of CO₂ exchange and its physiological interpretation – III. Diel, seasonal, and annual carbon budgets, *Flora*, 198, 277–292, 2003.
- 5 Lange, O. L., Büdel, B., Meyer, A., Zellner, H., and Zotz, G.: Lichen carbon gain under tropical conditions: water relations and CO₂ exchange of *Lobariaceae* species of a lower montane rainforest in Panama, *Lichenologist*, 36, 329–342, 2004.
- Lange, O. L. and Green, T. G. A.: Photosynthetic performance of the squamulose soil-crust lichen *Squamarina lentigera*: laboratory measurements and long-term monitoring of CO₂ exchange in the field, *Bibliotheca Lichenologica*, 88, 363–360, 2004.
- 10 Lange, O. L., Green, T. G. A., Melzer, B., Meyer, A., and Zellner, H.: Water relations and CO₂ exchange of the terrestrial lichen *Teloschistes capensis* in the Namib fog desert: Measurements during two seasons in the field and under controlled conditions, *Flora*, 201, 268–280, 2006.
- 15 Last, F. T. and Deighton, F. C.: Non-parasitic microflora on surfaces of living leaves, *British Mycological Society Transactions*, 48, 83–99, 1965.
- Lawrence, P. J. and Chase, T. N.: Representing a new MODIS consistent land surface in the Community Land Model (CLM 3.0), *J. Geophys. Res.-Biogeosciences*, 112, G01023, doi:10.1029/2006JG000168, 2007.
- 20 Lücking, R. and Matzer, M.: High foliicolous lichen alpha-diversity on individual leaves in Costa Rica and Amazonian Ecuador, *Biodivers. Conserv.*, 10, 2139–2152, 2001.
- Lüttge, U., Büdel, B., Ball, E., Strube, F., and Weber, P.: Photosynthesis of terrestrial cyanobacteria under light and desiccation stress as expressed by chlorophyll fluorescence and gas exchange, *J. Exp. Bot.*, 46, 309–319, 1995.
- 25 Matthes-Sears, U., Gerrath, J. A., and Larson, D. W.: Abundance, biomass, and productivity of endolithic and epilithic lower plants on the temperate-zone cliffs of the Niagara Escarpment, Canada, *Int. J. Plant Sci.*, 158, 451–460, 1997.
- Matthews, E.: Global vegetation and land use: new high-resolution data bases for climate studies, *J. Clim. Appl. Meteorol.*, 22, 474–487, 1983.
- 30 Mayland, H. F. and McIntosh, T. H.: Availability of biologically fixed atmospheric nitrogen-15 to higher plants, *Nature*, 209, 421–422, 1966.
- Mayland, H. F., McIntosh, T. H., and Fuller, W. H.: Fixation of isotopic nitrogen on a semiarid soil

6996

- by algal crust organisms, *Soil Science Society of America Proceedings*, 30, 56–60, 1966.
- McCune, B.: Gradients in epiphyte biomass in three *Pseudotsuga-Tsuga* forests of different ages in Western Oregon and Washington, *Bryologist*, 96, 405–411, 1993.
- McCune, B.: Using epiphyte litter to estimate epiphyte biomass, *Bryologist*, 97, 396–401, 1994.
- 5 McCune, B., Amsberry, K. A., Camacho, F. J., Clery, S., Cole, C., Emerson, C., Felder, G., French, P., Greene, D., Harris, B., Hutten, M., Larson, B., Lesko, M., Maiors, S., Markwell, T., Parker, G. G., Pendergrass, K., E. B. Peterson, Peterson, E. T., Platt, J., Proctor, J., Rambo, T., Rosso, A., Shaw, D., Turner, R., and Widmer, M.: Vertical profile of epiphytes in a Pacific northwest old-growth forest, *Northwest Science*, 71, 145–152, 1997.
- 10 Pankratova, E. M.: Functioning of cyanobacteria in soil ecosystems, *Eurasian Soil Science*, 39, S118–S127, 2006.
- Peveling, E., Burg, H., and Tenberge, K. B.: Epiphytic algae and fungi on spruce needles, *Symbiosis*, 12, 173–187, 1992.
- Philips, S. and Belnap, J.: Shifting carbon dynamics due to the effect of *Bromus tectorum* invasion on biological soil crust, *Bull ESA*, 205, 1998 Meeting Abstracts, 1998.
- 15 Pike, L. H., Rydell, R. A., and Denison, W. C.: A 400-year-old Douglas fir tree and its epiphytes: biomass, surface area, and their distributions, *Can. J. Forest Res.*, 7, 680–699, 1977.
- Pike, L. H.: The importance of epiphytic lichens in mineral cycling, *Bryologist*, 81, 247–257, 1978.
- 20 Prentice, C., Farquhar, G. D., Fasham, M. J. R., Goulden, M., Heimann, M., Jaramillo, V. J., Kheshgi, H. S., LeQuere, C., Scholes, R. J., and Wallace, D. W. R.: The carbon cycle and atmospheric carbon dioxide. The scientific basis (Contribution of Working Group 1 to the Third Assessment Report of the Intergovernmental Panel on Climate Change), in: *Climate Change 2001*, edited by: Houghton, J., Ding, Y., Griggs, D. J., Noguer, M., Linden, P. J. v. d., Dai, X., Maskell, K., and Johnson, C. A., Cambridge University Press, Cambridge, 183–237, 2001.
- 25 Prigent, C., Aires, F., Rossow, W., and Matthews, E.: Joint characterization of vegetation by satellite observations from visible to microwave wavelengths: A sensitivity analysis, *J. Geophys. Res.-Atmos.*, 106, 20665–20685, 2001.
- 30 Putz, F. E.: Liana biomass and leaf area of a “Tierra Firme” forest in the Rio Negro Basin, Venezuela, *Biotropica*, 15, 185–189, 1983.
- Radies, D. N. and Coxson, D. S.: Macrolichen colonization on 120-140 year old *Tsuga heterophylla* in wet temperate rainforests of central-interior British Columbia: a comparison of

6997

- lichen response to even-aged versus old-growth stand structures, *Lichenologist*, 36, 235–247, 2004.
- Ramankutty, N. and Foley, J. A.: Estimating historical changes in global land cover: Croplands from 1700 to 1992, *Global Biogeochem. Cy.*, 13, 997–1027, 1999.
- 5 Rascher, U., Lakatos, M., Büdel, B., and Lüttge, U.: Photosynthetic field capacity of cyanobacteria of a tropical inselberg of the Guiana Highlands, *Eur. J. Phycol.*, 38, 247–256, 2003.
- Reich, P. B., Hobbie, S. E., Lee, T., Ellsworth, D. S., West, J. B., Tilman, D., Knops, J. M. H., Naeem, S., and Trost, J.: Nitrogen limitation constrains sustainability of ecosystem response to CO₂, *Nature*, 440, 922–925, 2006.
- 10 Rhoades, F. M.: Biomass of epiphytic lichens and bryophytes on *Abies lasiocarpa* on a Mt. Baker lava flow, Washington, *Bryologist*, 84, 39–47, 1981.
- Roskoski, J. P.: Epiphyll dynamics of a tropical understory, *Oikos*, 37, 252–256, 1981.
- Rossi, F., Facini, O., Rotondi, A., Loreti, S., and Georgiadis, T.: Optical properties of juniper and lentisk canopies in a coastal Mediterranean macchia shrubland, *Trees-Struct. Funct.*, 15, 462–471, 2001.
- 15 Ruinen, J.: Occurrence of *Beijerinckia* species in the phyllosphere, *Nature*, 177, 220–221, 1956.
- Running, S. W., Nemani, R. R., Heinsch, F. A., Zhao, M. S., Reeves, M., and Hashimoto, H.: A continuous satellite-derived measure of global terrestrial primary production, *Bioscience*, 54, 547–560, 2004.
- 20 Sala, A., Sabate, S., Gracia, C., and Tenhunen, J. D.: Canopy structure within a *Quercus ilex* forested watershed – variations due to location, phenological development, and water availability, *Trees-Struct. Funct.*, 8, 254–261, 1994.
- San José, J. J. and Bravo, C. R.: CO₂ exchange in soil algal crusts occurring in the Trachypogon savannas of the Orinoco Llanos, Venezuela, *Plant Soil*, 135, 233–244, 1991.
- 25 Sánchez-Azofeifa, G. A., Kalácska, M., Espírito-Santo, M. M. D., Fernandes, G. W., and Schnitzer, S.: Tropical dry forest succession and the contribution of lianas to wood area index (WAI), *Forest Ecol. Manage.*, in press, 2009.
- Schlesinger, W. H., Phippen, J. S., Wallenstein, M. D., Hofmockel, K. S., Klepeis, D. M., and Mahall, B. E.: Community composition and photosynthesis by photoautotrophs under quartz pebbles, southern Mojave Desert, *Ecology*, 84, 3222–3231, 2003.
- 30 Schmidt, S. K., Reed, S. C., Nemergut, D. R., Grandy, A. S., Cleveland, C. C., Weintraub, M. N., Hill, A. W., Costello, E. K., Meyer, A. F., Neff, J. C., and Martin, A. M.: The earliest stages of

6998

- ecosystem succession in high-elevation (5000 metres above sea level), recently deglaciated soils, *Proceedings of the Royal Society B-Biological Sciences*, 275, 2793–2802, 2008.
- Scurlock, J. M. O., Asner, G. P., and Gower, S. T.: Global leaf area index data from field measurements, 1932–2000. Data set, online available at: <http://www.daac.ornl.gov> from the
- 5 Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, USA, 2001.
- Sheridan, R. P.: Epicaulous, nitrogen-fixing microepiphytes in a tropical mangal community, Guadeloupe, French West-Indies, *Biotropica*, 23, 530–541, 1991a.
- Sheridan, R. P.: Nitrogenase activity by *Hapalosiphon flexuosus* associated with *Sphagnum*
- 10 *erythrocalyx* mats in the cloud forest on the volcano La Soufriere, Guadeloupe, French West Indies, *Biotropica*, 23, 134–140, 1991b.
- Sherwood, M. and Carroll, G.: Fungal succession on needles and young twigs of old-growth Douglas fir, *Mycologia*, 66, 499–506, 1974.
- Sillett, S. C.: Branch epiphyte assemblages in the forest interior and on the clearcut edge of a
- 15 700-year-old Douglas fir canopy in Western Oregon, *Bryologist*, 98, 301–312, 1995.
- Sillett, S. C. and Rambo, T. R.: Vertical distribution of dominant epiphytes in Douglas-fir forests of the central Oregon Cascades, *Northwest Science*, 74, 44–49, 2000.
- Skarpe, C. and Henriksson, E.: Nitrogen fixation by cyanobacterial crusts and by associative-symbiotic bacteria in western Kalahari, Botswana, *Arid Soil Res. Rehabil.*, 1, 55–59, 1987.
- 20 Søchting, U.: Epiphyllic cover on spruce needles in Denmark, *Ann. Bot. Fennici*, 34, 157–164, 1997.
- Solhaug, K. A., Gauslaa, Y., and Haugen, J.: Adverse effects of epiphytic crustose lichens upon stem photosynthesis and chlorophyll of *Populus tremula* L, *Botanica Acta*, 108, 233–239, 1995.
- 25 Thiet, R. K., Boerner, R. E. J., Nagy, M., and Jardine, R.: The effect of biological soil crusts on throughput of rainwater and N into Lake Michigan sand dune soils, *Plant Soil*, 278, 235–251, 2009.
- Tretiach, M.: Ecophysiology of calcicolous lichens: progress and problems, *Giornale Botanico Italiano*, 129, 159–184, 1995.
- 30 Tretiach, M. and Geletti, A.: CO₂ exchange of the endolithic lichen *Verrucaria baldensis* from Karst habitats in northern Italy, *Oecologia*, 111, 515–522, 1997.
- Turetsky, M. R.: The role of bryophytes in carbon and nitrogen cycling, *Bryologist*, 106, 395–409, 2003.

6999

- Veluci, R. M., Neher, D. A., and Weicht, T. R.: Nitrogen fixation and leaching of biological soil crust communities in mesic temperate soils, *Microbial Ecol.*, 51, 189–196, 2006.
- Vincke, C., Granier, A., Breda, N., and Devillez, F.: Evapotranspiration of a declining *Quercus robur* (L.) stand from 1999 to 2001. II. Daily actual evapotranspiration and soil water reserve,
- 5 *Ann. For. Sci.*, 62, 615–623, 2005.
- Walker, J. J., Spear, J. R., and Pace, N. R.: Geobiology of a microbial endolithic community in the Yellowstone geothermal environment, *Nature*, 434, 1011–1014, 2005.
- Wanek, W., and Portl, K.: Phyllosphere nitrogen relations: reciprocal transfer of nitrogen between epiphyllous liverworts and host plants in the understorey of a lowland tropical wet
- 10 forest in Costa Rica, *New Phytol.*, 166, 577–588, 2005.
- Watson, R. T., Noble, I. R., Bolin, B., Ravindranath, N. H., Verardo, D. J., and Dokken, D. J.: Land use, land-use change and forestry, in: IPCC Special Reports on Climate Change, edited by: Watson, R. T., Noble, I. R., Bolin, B., Ravindranath, N. H., Verardo, D. J., and Dokken, D. J., Cambridge University Press, Cambridge, 377 pp., 2000.
- 15 Weaver, P. L., Medina, E., Pool, D., Dugger, K., Gonzales-Liboy, J., and Cuevas, E.: Ecological observations in the dwarf cloud forest of the Luquillo Mountains in Puerto Rico, *Biotropica*, 18, 79–85, 1986.
- Weaver, P. L. and Murphy, P. G.: Forest structure and productivity in Puerto Rico's Luquillo Mountains, *Biotropica*, 22, 69–82, 1990.
- 20 Weber, B.: Biologie und Ökologie gesteinsbewohnender Blaualgen/Cyanobakterien in der Dornbuschsavanne Nord-Transvaals, Südafrika, University of Würzburg, Germany, 1994.
- Weber, B., Scherr, C., Reichenberger, H., and Büdel, B.: Fast reactivation by high air humidity and photosynthetic performance of alpine lichens growing endolithically in limestone, *Arctic, Antarct. Alp. Res.*, 39, 309–317, 2007.
- 25 Weber, B., Olechowski, C., Knerr, T., Hill, J., Deutschewitz, K., Wessels, D. C. J., Eitel, B., and Büdel, B.: A new approach for mapping of biological soil crusts in semidesert areas with hyperspectral imagery, *Remote Sens. Environ.*, 112, 2187–2201, 2008.
- Weiskittel, A. R. and Maguire, D. A.: Branch surface area and its vertical distribution in coastal Douglas-fir, *Trees-Struct. Funct.*, 20, 657–667, 2006.
- 30 Wessels, D. C. A. and Kappen, L.: Photosynthetic performance of rock-colonising lichens in the Mount Zebra National Park, South Africa, *Koedoe*, 36, 27–48, 1993.
- Whitman, W. B., Coleman, D. C., and Wiebe, W. J.: Prokaryotes: The unseen majority, *Proceedings of the National Academy of Sciences of the United States of America*, 95, 6578–

7000

6583, 1998.

Whittaker, R. H. and Likens, G. E.: Primary production: Biosphere and man, Hum. Ecol., 1, 357–369, 1973.

Wilske, B., Burgheimer, J., Karnieli, A., Zaady, E., Andreae, M. O., Yakir, D., and Kesselmeier, J.: The CO₂ exchange of biological soil crusts in a semiarid grass-shrubland at the northern transition zone of the Negev desert, Israel, Biogeosciences, 5, 1411–1423, 2008, <http://www.biogeosciences.net/5/1411/2008/>.

Wiman, B. L. B. and Gaydarova, P. N.: Spectral composition of shade light in coastal-zone oak forests in SE Bulgaria, and relationships with leaf area index: a first overview, Trees-Struct. Funct., 22, 63–76, 2008.

Winkler, J. B. and Kappen, L.: Photosynthetic capacity of endolithic lichens from South-Africa, Bibliotheca Lichenologica, 67, 165–181, 1997.

Yeager, C. M., Kornosky, J. L., Morgan, R. E., Cain, E. C., Garcia-Pichel, F., Housman, D. C., Belnap, J., and Kuske, C. R.: Three distinct clades of cultured heterocystous cyanobacteria constitute the dominant N₂-fixing members of biological soil crusts of the Colorado Plateau, USA, FEMS Microbiol. Ecol., 60, 85–97, 2007.

Yoneda, T.: Surface area of woody organs of an evergreen broadleaf forest tree in Japan and Southeast Asia, J. Plant Res., 106, 229–237, 1993.

Zaady, E., Kuhn, U., Wilske, B., Sandoval-Soto, L., and Kesselmeier, J.: Patterns of CO₂ exchange in biological soil crusts of successional age, Soil Biol. Biochem., 32, 959–966, 2000.

Zotz, G., Büdel, B., Meyer, A., Zellner, H., and Lange, O. L.: *In situ* studies of water relations and CO₂ exchange of the tropical macrolichen, *Sticta tomentosa*, New Phytol., 139, 525–535, 1998.

Zotz, G.: Altitudinal changes in diversity and abundance of non-vascular epiphytes in the Tropics – an ecophysiological explanation, Selbyana, 20, 256–260, 1999.

7001

Table A1. Annual net carbon uptake fluxes (photosynthesis minus respiration) of biological soil crusts (BSC) in arid and semiarid regions, (b) BSC in other sparsely vegetated regions, and (c) biological crusts on rocks (BRC).

Carbon uptake flux (g m ⁻² a ⁻¹)	BSC components	Habitat	Location	Calculation background	Reference
(a)					
74.9	Early and later successional stage BSC (<i>Nostoc</i> / <i>Scytonema</i> or <i>Placidium</i> / <i>Collema</i> dominated)	1. Pinyon-juniper woodland; 2. Shrubland	1. Cold desert Canyonlands, SE Utah 2. Warm desert Jornada site, New Mexico	Mean value of balances minus respiration	(Housman et al., 2006)
49.1	Soilcrust cyanolichen (<i>Collema tenax</i>)	Pinyon-juniper woodland	Colorado Plateau, USA	Calculated from maximum values of net photosynthesis (NP _{max}) ⁽²⁾	(Lange et al., 1998a)
32.7	Lichen-dominated soilcrust (<i>Ramalina macrospora</i>)	Desert mesa	Highlands of the Central Negev Desert, Israel	Adopted balance	(Kappen et al., 1980)
29.9	Soilcrust lichens (<i>Diploschistes diacapsis</i> , <i>Peziza cerebriiformis</i> , <i>Squamaria lenigera</i>)	Pinyon-juniper woodland	Colorado Plateau, USA maximum values of net	Calculated from photosynthesis (NP _{max}) ⁽²⁾	(Lange et al., 1997)
28.8	Lichen-dominated soilcrust	Pinyon-juniper woodlands	Colorado Plateau, Utah	Adopted balance	(Kipatek, 1992)
24.6	Mixed crust, cyanobacteria and chlorophyta	Orinoco Llanos	Savannah, Venezuela maximum values of net	Calculated from photosynthesis (NP _{max}) ⁽²⁾	(San José and Bravo, 1991)
24.5	Lichen of BSC (<i>Teleoschistes capensis</i>)	Namib lichen fields	Namib desert, Namibia	Mean value of balances	(Lange et al., 2006)
21.0	BSC dominated by cyanolichen (<i>Gloeohoplia turgida</i>)	Slopes of the Dead Sea	Judean Desert, Israel maximum values of net	Calculated from photosynthesis (NP _{max}) ⁽²⁾	(Lange et al., 1993)
17.0	Lichen-dominated soilcrust	Blackbrush community	Colorado Plateau, Arizona	Adopted balance	(Kipatek, 1992)
16.0	Soilcrust lichens (<i>Acarospora schleicheri</i> , <i>Caloplaca volkii</i> , <i>Leccidella crystallina</i>)	Namib lichen fields	Namib desert, Namibia	Adopted balance	(Lange et al., 1994b)
15.4	Hypolithic community dominated by cyanobacteria	Desert pavement	Joshua Tree National Park, California	Calculated from maximum values of net photosynthesis (NP _{max}) ⁽²⁾	(Schlesinger et al., 2003)
14.0	Polkilohydric moss carpet (<i>Tortula ruralis</i>)	Calcareous semiarid grassland	Sandy grassland, <i>Festucetum vaginatae</i> <i>danubiale</i> , Hungary	Calculated from maximum values of net photosynthesis (NP _{max}) ⁽²⁾	(Juhász et al., 2002)
13.0	BSC, lichen dominated (<i>Peltula richardii</i> , <i>P. patellata</i> , <i>Placidium squamulosum</i> , <i>Peziza desipiens</i>)	Floodplains	Sonoran Desert, Baja California	Calculated from maximum values of net photosynthesis (NP _{max}) ⁽²⁾	Büdel and Lange, unpublished
10.5	Mixed BSC with cyanobacteria, algae, mosses, and lichens	Pinyon-juniper woodland	Canyonlands National Park, Utah	Calculated from maximum values of net photosynthesis (NP _{max}) ⁽²⁾	(Philips and Belnap, 1998)
7.8	Cyanobacteria-dominated BSC (<i>Microcoleus</i> and <i>Scytonema</i> -dominated)	Seasonally inundated system of pans and dunes	Western Mojave Desert	Mean value of balances minus respiration	(Brostoff et al., 2005)
6.6	Soilcrust lichens (<i>Squamaria cf.</i> <i>crassa</i> , <i>Diploschistes steppicus</i>)	Central Negev	Negev Desert, Israel	Calculated from maximum values of net photosynthesis (NP _{max}) ⁽²⁾	(Lange et al., 1970)
5.8	Cyanobacteria-dominated BSC (<i>Microcoleus</i> sp.)	Nizzana Dunes	Negev Desert, Israel	Calculated from maximum values of net photosynthesis (NP _{max}) ⁽²⁾	(Lange et al., 1992)
1.1	Cyanobacteria-dominated BSC (<i>Microcoleus vaginatus</i> , <i>Nostoc</i> sp., <i>Scytonema</i> sp.)	Pinyon-juniper woodland	Colorado Plateau, southeastern Utah	Mean value of balances	(Garcia-Pichel and Belnap, 1996)
0.8	Cyanobacteria-dominated soilcrust (<i>Scytonema</i> sp., <i>Microcoleus</i> sp.)	Blackbrush community	Kaiparowits Basin, southern Utah	Mean value of balances	(Jeffries et al., 1993)
16.0 (median), 20.7 (arithmetic mean), 17.8 (standard deviation)					

7002

Table A1. Continued.

Carbon uptake flux (g m ⁻² a ⁻¹)	BSC components	Habitat	Location	Calculation background	Reference
(b)					
80.7	BSC dominated by green algae (Klebsormidium spp.)	Forest clear cut area	Summer green deciduous forest zone, Czech Republic	Calculated from maximum values of net photosynthesis (NP _{max}) ^{a)}	Büdel and Lukesová, unpublished
29.9	Sclerotium chlorococcum (Diploschistes muscorum, Squamaria fenigera, Fulgensia fulgens)	Xerothermic steppe formation	Summer green deciduous forest zone, Germany	Calculated from maximum values of net photosynthesis (NP _{max}) ^{a)}	(Lange, 2000a)
28.2 (Squamaria fenigera)	Chlorolichen	Xerothermic steppe formation forest zone, Germany	Summer green deciduous forest zone, Germany	Adopted balance	(Lange and Green, 2004)
29.8	Cyanolichen (Collema cristatum)	Xerothermic steppe formation	Summer green deciduous forest zone, Germany	Adopted balance	(Lange, 2000b)
20.4	Lichen community in dry dwarf shrub heath	Upland tundra	Footfills of the Brooks Range, Alaska	Adopted balance	(Lange et al., 1998b)
15.7	Lichen heath	Upland tundra	Footfills of the Brooks Range, Alaska	Adopted balance	(Lange et al., 1998b)
9.6	Lichen community in Dryas heath	Upland tundra	Footfills of the Brooks Range, Alaska	Adopted balance	(Lange et al., 1998b)
4.7	Lichen community in moist dwarf shrub heath	Upland tundra	Footfills of the Brooks Range, Alaska	Adopted balance	(Lange et al., 1998b)
23.1 (median), 26.9 (arithmetic mean), 23.5 (standard deviation)					
(c)					
21.7	Chlorolichen (Lecanora muralis)	Sandstone wall	Spessart midlands, Germany	Adopted balance	(Lange, 2003)
20.4	Epi- and endolithic lichens	Sandstones	Mount Zebra National Park, Cape Province and Savannah, Mutamba, northern Transvaal	Calculated from maximum values of net photosynthesis (NP _{max}) ^{a)}	(Wessels and Kappen, 1993)
16.8	Cyanobacterial mat 2. Limestone rock outcrop, 3. vertical loamy surface	1. epilithic on granite rock, Endolithic habitat	1. Inselberg, Ivory coast, 2. Paraguaná Peninsula, Venezuela, 3. Northern Coastal Range, Venezuela	Calculated from maximum values of net photosynthesis (NP _{max}) ^{a)}	(Lüttge et al., 1995)
15.8	Endolithic lichens (Hymenelia preceps, H. concolor)	Endolithic habitat	Calcareous rock outcrops, Untersberg, northern Alps	Calculated from maximum values of net photosynthesis (NP _{max}) ^{a)}	(Weber et al., 2007)
12.7	Endolithic lichens (Lecidea aff. sarcogynoides, Sarcogyna cf. austroafricana, Lecidea confluenta, Lithoglypha aggregata)	Endolithic habitat	1. Sandstone Inselberg on Mutamba Ranch, South Africa; 2. Golden Gate Highlands National Park, South Africa	Calculated from maximum values of net photosynthesis (NP _{max}) ^{a)}	(Winkler and Kappen, 1997)
8.7	Epi- and endolithic lichens	Limestone	Karst Plateau, Trieste, Italy	Calculated from maximum values of net photosynthesis (NP _{max}) ^{a)}	(Trellich and Gelelli, 1997)
8.6	Cyanobacterial mat	1. epilithic on granite rock, 2. Limestone rock outcrop, 3. vertical loamy surface	1. Inselberg, Ivory coast, 2. Paraguaná Peninsula, Venezuela, 3. Northern Coastal Range, Venezuela	Calculated from maximum values of net photosynthesis (NP _{max}) ^{a)}	(Lüttge et al., 1995)
8.0	Endolithic cyanobacteria	Cryptendolithic habitat	Sandstone plateau along Brakrivier, Northern Transvaal, South Africa	Calculated from maximum values of net photosynthesis (NP _{max}) ^{a)}	(Weber, 1994)
7.4	Cyanobacteria/ cyanolichen crusts	Inselberg	Dry savanna, South Africa	Adopted balance	(Büdel, 1999)
7.0	Epilithic lichen (Acaropora sp.)	Sandstones	Mount Zebra National Park, South Africa	Calculated from maximum values of net photosynthesis (NP _{max}) ^{a)}	(Wessels and Kappen, 1993)
6.9	Epi- and endolithic lichens	Epilithic and endolithic on limestone; Calypso ehrenbergii on chert	Central Negev Desert, Israel	(Lange et al., 1970) maximum values of net photosynthesis (NP _{max}) ^{a)}	
3.9	Cryptendolithic cyanobacteria together with green algae	Temperate zone sandstone	Colorado Plateau, Arizona	Calculated from maximum values of net photosynthesis (NP _{max}) ^{a)}	(Bell and Sommerfeld, 1987)
3.5	Endolithic lichen (Verrucaria baldensis)	Limestone	Karst Plateau, Trieste, Italy	Calculated from maximum values of net photosynthesis (NP _{max}) ^{a)}	(Trellich and Gelelli, 1997)
3.3	Cyanobacteria/ cyanolichen crusts	Inselberg	Humid savanna, Venezuela	Adopted balance	(Büdel, 1999)
2.3	Cyanobacteria/ cyanolichen crusts	Inselberg	Thorn bush savannah, South Africa	Adopted balance	(Büdel, 1999)
0.6	Lichen-dominated endolithic community with green algae and cyanobacteria	Cryptendolithic habitat	Beacon sandstone of the Ross Desert, Antarctica	Adopted balance	(Friedmann et al., 1993)
7.7 (median), 9.2 (arithmetic mean), 6.5 (standard deviation)					

^{a)} NP_{max} values were scaled to estimated average ambient conditions.

7003

Table A2. Global surface area of terrestrial arid and semiarid regions (dry-lands).

Area (10 ¹³ m ²)	References
7.8	(Watson et al., 2000)
6.8	(Asner et al., 2003)
6.6	(Ramankutty and Foley, 1999)
6.5	(Whittaker and Likens, 1973)
6.4	(Garcia-Pichel et al., 2003)
6.0	(Kucharik et al., 2000)
5.5	(Goldewijk, 2001)
5.0	(Ajtay et al., 1979)
5.0	(Lawrence and Chase, 2007)
6.4 (median), 6.2 (arithmetic mean), 0.9 (standard deviation)	

Table A3. Annual net carbon uptake flux (photosynthesis minus respiration) of epiphytic and epiphyllic crusts (EPC).

Carbon uptake flux (g m ⁻² a ⁻¹)	EPC components	Habitat	Location	Calculation background	Reference
167.7	Homoiomerous cyanolichen (<i>Leptogium cyanescens</i>)	Lower montane rainforest	Cordillera Central, Republic of Panama	Adopted balance	(Lange et al., 2000)
116.3	Homoiomerous cyanolichen (<i>Leptogium phyllocarpum</i>)	Lower montane rainforest	Cordillera Central, Republic of Panama	Adopted balance	(Lange et al., 2000)
86.0	Epiphytic Lobariaceae (<i>Sticta weigeli</i>)	Lower montane rainforest	Cordillera Central, Republic of Panama	Adopted balance	(Lange et al., 2004)
78.9	Cyanobacterial basidiolichen (<i>Dictyonema glabratum</i>)	Lower montane rainforest	Cordillera Central, Republic of Panama	Adopted balance	(Lange et al., 1994a)
74.2	Epiphytic Lobariaceae (<i>Sticta sublimbata</i>)	Lower montane rainforest	Cordillera Central, Republic of Panama	Adopted balance	(Lange et al., 2004)
60.2	Epiphytic Lobariaceae (<i>Sticta tomentosa</i>)	Lower montane rainforest	Cordillera Central, Republic of Panama	Adopted balance	(Zotz et al., 1998)
29.5	Epiphytic Lobariaceae (<i>Pseudocyphellaria intricata</i>)	Lower montane rainforest	Cordillera Central, Republic of Panama	Adopted balance	(Lange et al., 2004)
26.8	Homoiomerous cyanolichen (<i>Leptogium azureum</i>)	Lower montane rainforest	Cordillera Central, Republic of Panama	Adopted balance	(Lange et al., 2000)
25.4	Epiphytic Lobariaceae (<i>Pseudocyphellaria aurata</i>)	Lower montane rainforest	Cordillera Central, Republic of Panama	Adopted balance	(Lange et al., 2004)
9.3	Corticolous lichen (<i>Coenogonium linkii</i>)	Tropical evergreen lowland rainforest	Les Nouragues National Park, French Guiana	Calculated from maximum values of net photosynthesis (NP _{max}) ^{a)}	(Lakatos et al., 2006)
5.6	Epiphytic Lobariaceae (<i>Lobaria crenulata</i>)	Lower montane rainforest	Cordillera Central, Republic of Panama	Calculated from maximum values of net photosynthesis (NP _{max}) ^{a)}	(Lange et al., 2004)
2.7	Corticolous lichen (<i>Cryptothecia rubrocincta</i>)	Tropical evergreen lowland rainforest	Les Nouragues National Park, French Guiana	Calculated from maximum values of net photosynthesis (NP _{max}) ^{a)}	(Lakatos et al., 2006)
1.9	Corticolous lichen (<i>Thelotrema albobolivaceum</i>)	Tropical evergreen lowland rainforest	Les Nouragues National Park, French Guiana	Calculated from maximum values of net photosynthesis (NP _{max}) ^{a)}	(Lakatos et al., 2006)
1.1	Corticolous lichen (<i>Phyllopsora corallina</i>)	Tropical evergreen lowland rainforest	Les Nouragues National Park, French Guiana	Calculated from maximum values of net photosynthesis (NP _{max}) ^{a)}	(Lakatos et al., 2006)
(37–64)	Epiphytic bryophytes lower montane wet forest	Tropical	Costa Rica	Adopted balance	(Clark et al., 1998)
28.2 (median), 49.0 (arithmetic mean), 50.5 (standard deviation)					

^{a)} NP_{max} values were scaled to estimated average ambient conditions.

7005

Table A4. Global tropical forest area.

Forest area (10 ¹³ m ²)	References
1.9	(Kucharik et al., 2000)
1.8	(Watson et al., 2000)
1.7	(Matthews, 1983)
1.7	(Whittaker and Likens, 1973)
1.7	(Ramankutty and Foley, 1999)
1.6	(Prigent et al., 2001)
1.5	(Matthews, 1983)
1.5	(Ajtay et al., 1979)
1.4	(Lawrence and Chase, 2007)
1.3	(Matthews, 1983)
1.7 (median), 1.6 (arithmetic mean), 0.2 (standard deviation)	

7006

Table A5. Plant surface area of (a) leaves of tropical evergreen trees (leaf area index, LAI), (b) tropical lianas (LAI), (c) leaves of evergreen trees and shrubs (LAI; temperate and boreal), and (d) stems and branches of trees and shrubs (wood area index, WAI; stem area index, SAI).

Area index (m ² m ⁻²)	Forest type	Location	References
(a)			
12.3	Tropical lowland rainforest	Thailand	(Arora and Boer, 2005)
8.0	Tropical rainforest	Global average	(Whittaker and Likens, 1973)
7.4	Evergreen tropical forest	Thailand	(Putz, 1983)
7.3	Subtropical wet forest	Puerto Rico	(Weaver et al., 1986)
7.3	Tropical lowland rainforest	Malaysia	(Putz, 1983)
6.0 (up to 13)	Tropical upland wet forest	Costa Rica	(Clark et al., 2008)
5.5	Rubber tree forest	Maharashtra, India	(Devakumar et al., 1999)
5.2	Tropical rainforest	Rio Negro Basin, Venezuela	(Putz, 1983)
5.0	Lower montane rainforest	Puerto Rico	(Weaver and Murphy, 1990)
4.8	Tropical evergreen broadleaf forest	Global average	(Scurlock et al., 2001)
4.7	Tropical deciduous forest	Mexico	(Arora and Boer, 2005)
4.0	Broadleaf forests	Amazon	(Buermann et al., 2002)
3.3	Lower montane rainforest	Puerto Rico	(Weaver and Murphy, 1990)
2.0	Lower montane rainforest	Puerto Rico	(Weaver et al., 1986)
(5–8)	Tropical forest	Amazon	(Arora and Boer, 2005)
(6–7)	Subtropical wet forest	Puerto Rico	(Weaver and Murphy, 1990)
(3–5)	Lower montane rainforest	Puerto Rico	(Weaver and Murphy, 1990)
(3–3.5)	Lower montane rainforest	Puerto Rico	(Weaver and Murphy, 1990)
5.4 (median), 5.9 (arithmetic mean), 2.5 (standard deviation)			

7007

Table A5. Continued.

Area index (m ² m ⁻²)	Forest type	Location	References
(b)			
3.3	Lianas in tropical forest	Brazilian Amazon	(Gerwing and Farias, 2000)
3.3	Lianas in tropical forest	Brazilian Amazon	(Gerwing and Vidal, 2002)
1.7	Lianas and vines in tropical deciduous forest	Mexico	(Castellanos et al., 1992)
1.2	Lianas in evergreen tropical rain forest	Rio Negro Basin, Venezuela	(Putz, 1983)
0.7	Lianas in tropical upland wet forest	Costa Rica	(Clark et al., 2008)
(2–7)	Lianas in lowland tropical rain forest	Gabon	(Kazda and Salzer, 2000)
1.7 (median), 2.0 (arithmetic mean), 1.2 (standard deviation)			
(c)			
5.3	Mediterranean oak forest	Spain	(Sala et al., 1994)
4.6	Mediterranean oak forest	Spain	(Sala et al., 1994)
4.3	Mediterranean evergreen maquis	Portugal	(Caldwell et al., 1986)
2.9	Coastal oak forest	Bulgaria	(Wiman and Gaydarova, 2008)
2.5	Mediterranean evergreen maquis	Italy	(Gratani and Varone, 2007)
2.5	Mediterranean evergreen maquis	Sardinia, Italy	(Rossi et al., 2001)
(1.4–3.2)	Coastal oak forest	Bulgaria	(Gaydarova, 2003)
3.6 (median), 3.7 (arithmetic mean), 1.2 (standard deviation)			
(d)			
1.9 ^{a)}	Douglas fir forest	Oregon coast range, USA	(Weiskittel and Maguire, 2006)
1.3 ^{a)}	Mixed forest	Belgium	(Vincke et al., 2005)
1.3 ^{b)}	Mediterranean oak forest	Spain	(Sala et al., 1994)
0.7 ^{a)}	Tropical dry forest	Mexico, Costa Rica, Brazil	(Sánchez-Azofeifa et al., in press)
0.5 ^{a)}	Tropical dry forest	Mexico, Costa Rica, Brazil	(Sánchez-Azofeifa et al., in press)
(2.0–2.6) ^{c)}	Rainforests	Japan and Southeast Asia	(Yoneda, 1993)
(0.4–2.5) ^{a)}	Mixed forest	Alsace, France	(Breda, 2003)
(0.5–1.9) ^{c)}	Norway spruce forest	Germany	(Falge et al., 1997)
(0.3–1.5) ^{a)}	Mixed forest	France	(Dufrêne and Bréda, 1995)
1.3 (median), 1.1 (arithmetic mean), 0.6 (standard deviation)			

a) WAI

b) SAI, but main trunks excluded

c) SAI

7008

Table A6. Plant surface coverage by EPC on (a) leaves, (b) needles, and (c) stems and branches.

Coverage (%)	EPC components	Location, host plant	Reference
(a)			
~100	Liverworts	Humid tropical forests	(Berrie and Eze, 1975)
> 90	Bryophytes	Tropical lowland rainforest in Costa Rica	(Furnkranz et al., 2008)
(50–100)	Lichens	Tropical rainforests	(Anthony et al., 2002)
(25–31)	Epiphyll cover	Coffee plants, semi-hot humid climate	(Roskoski, 1981)
(20–30) (up to 70)	Bryophytes	Rainforest in Costa Rica, understory plants	(Freiberg, 1998, 1999)
(22–35)	Lichens	Moist forests of Panama	(Coley et al., 1993)
(2–20)	Liverworts	Moist forests of Panama	(Coley et al., 1993)
(2–10)	Cyanobacteria	Rainforest in Costa Rica, understory plants	(Freiberg, 1998, 1999)
(b)			
58	Alectoroid lichens	Old-growth Pseudotsuga-Tsuga forests, Pacific Northwest, USA	(McCune et al., 1997)
55	Epiphytic lichen (<i>Hypogymnia physodes</i>)	Conifers, temperate European sites	(Sechting, 1997)
27	Bryophytes	Old-growth Pseudotsuga-Tsuga forests, Pacific Northwest, USA	(McCune et al., 1997)
25	Various lichens	Old-growth Pseudotsuga-Tsuga forests, Pacific Northwest, USA	(McCune et al., 1997)
17	Cyanolichens	Old-growth Pseudotsuga-Tsuga forests, Pacific Northwest, USA	(McCune et al., 1997)
8	Actinomycetes, fungi, and algae	Douglas fir, Oregon, USA	(Bernstein and Carroll, 1977)
7	Microepiphyte cover	Douglas fir, Oregon, USA	(Pike et al., 1977)
5	Actinomycetes, fungi, and algae	Douglas fir, Oregon, USA	(Carroll, 1979)
(45–80)	Green algae, fungi, and bacteria	Spruce trees, Germany	(Pevling et al., 1992)
(32–77)	Epiphytes (chlorolichens, cyanolichens, and bryophytes)	Sitka spruce, California, USA	(Ellyson and Sillett, 2003)
(c)			
~100	Crustose lichens	Lowland forest, Panama	(Zotz, 1999)
38	Crustose lichens	Lowland rainforest, Costa Rica	(Forman, 1975)
(40–100)	Bryophytes (2–14 cm in thickness)	Montane rainforests, Ecuador	(Freiberg and Freiberg, 2000)
(3–83) (most between 20–50)	Corticolous and crustose microlichens	Atlantic rainforests, Brazil	(Cáceres et al., 2007)
(40–80)	Foliose and crustose lichens	Forest in Norway	(Solhaug et al., 1995)
(25–60) (up to 80)	Mosses, liverworts, cyanolichens and other lichens	Douglas fir, Oregon, USA	(Sherwood and Carroll, 1974; Sillett, 1995; Sillett and Rambo, 2000)
(2–50)	Bryophytes (2–14 cm in thickness)	Lowland rainforests, Ecuador	(Freiberg and Freiberg, 2000)

7009

Table A7. Annual net carbon uptake by EPC in forest and shrub areas.

Substrate	Area index ^{a)} (m ² m ⁻²)	Forest/ shrub area (10 ¹³ m ²)	Plant surface area (10 ¹³ m ²)	EPC coverage ^{d)} (%)	EPC covered surface area (10 ¹³ m ²)	Annual net carbon uptake ^{e)} (Pg a ⁻¹)
Leaves of tropical evergreen trees	5.4	1.7 ^{b)}	9.2	50	4.7	1.3
Tropical lianas	1.7	1.7 ^{b)}	2.9	50	1.6	0.4
Leaves of evergreen trees and shrubs; temperate and boreal	3.6	1.0 ^{c)}	3.6	30	1.0	0.3
Stems and branches of trees and shrubs	1.3	4.4 ^{c)}	5.7	35	2.3	0.6
Total		4.4 ^{c)}			9.6	2.6

a) see Table A5;

b) see Table A4;

c) from “Community Land Model” (CLM 3.0) (Lawrence and Chase, 2007);

d) estimated from values given in Table A6;

e) calculated assuming a median carbon net fixation rate of 28 g m⁻² (Table A3).

7010

Table A8. Estimated global biomass (dry weight) of (a) BSC, (b) BRC, (c, d) EPC. Values in (a–c) in g m^{-2} of ground surface area, (d) in g m^{-2} of thallus or branch surface area.

Biomass (g m^{-2})	Component	Location	Reference
(a)			
1187	Soilcrust cyanolichen (<i>Collema tenax</i>)	Pinyon-juniper woodland, Colorado Plateau, USA	(Lange et al., 1998a)
684	Chlorolichen (<i>Squamaria lentigera</i>)	Xerothermic steppe formation, Germany	(Lange and Green, 2004)
447	Lichen of BSC (<i>Teloschistes capensis</i>)	Namib Desert, Namibia	(Lange et al., 2006)
306	Cyanolichen (<i>Collema cristatum</i>)	Xerothermic steppe formation, Germany	(Lange, 2000b)
258	Lichen community in dry dwarf shrub heath	Upland Tundra, Brooks Range, Alaska	(Lange et al., 1998b)
219	Lichen heath	Upland Tundra, Brooks Range, Alaska	(Lange et al., 1998b)
155	Lichen community in Dryas heath	Upland Tundra, Brooks Range, Alaska	(Lange et al., 1998b)
65	Lichen-dominated soilcrust (<i>Ramalina maciformis</i>)	Desert mesa, Central Negev Desert, Israel	(Kappen et al., 1980)
57	Lichen community in moist dwarf shrub heath	Upland Tundra, Brooks Range, Alaska	(Lange et al., 1998b)
(1–16)	Edaphic cyanobacteria	Various locations on Earth	(Garcia-Pichel et al., 2003)
(1–12)	Soil crust cyanobacteria	Various locations, Russia	(Pankratova, 2006)
258 (median), 375.3 (arithmetic mean), 362.0 (standard deviation)			
(b)			
594	Epilithic chlorolichen (<i>Aspicillia radiosa</i>)	Limestone karst plateau, Trieste, Italy	(Tretiach, 1995)
243	Cyanobacterial biofilm	Inselberg, tropical rainforest, French Guiana	(Dojani et al., 2007)
336	Endolithic cyanobacteria community	Mt. Falconer, Antarctica	(Büdel et al., 2008)
117	Endolithic chlorolichen (<i>Verrucaria</i> sp.)	Limestone karst plateau, Trieste, Italy	(Tretiach, 1995)
50	Nearly endolithic chlorolichen (<i>Acrocordia conidea</i>)	Limestone karst plateau, Trieste, Italy	(Tretiach, 1995)
50	Endolithic chlorolichen (<i>Verrucaria marmorea</i>)	Limestone karst plateau, Trieste, Italy	(Tretiach, 1995)
30	Nearly endolithic chlorolichen (<i>Petractis clausa</i>)	Limestone karst plateau, Trieste, Italy	(Tretiach, 1995)
30	Endolithic chlorolichen (<i>Rinodina immersa</i>)	Limestone karst plateau, Trieste, Italy	(Tretiach, 1995)
(38–185)	Endolithic microbial communities in sandstone and granite	Mojave Desert, California; Sonoran Desert, Mexico; Negev Desert, Israel	(Friedmann, 1980)
(32–177)	Endolithic microbial communities	Beacon sandstone, Dry Valley, Antarctica	(Friedmann, 1980)
(4–107)	Endolithic cyanobacteria community	Tshipise sandstone formation, South Africa	(Büdel et al., 2004)
(2–73)	Endo- and epilithic algae, cyanobacteria, and lichens	Niagara Escarpment, Ontario, Canada	(Matthes-Sears et al., 1997)
84 (median), 181.3 (arithmetic mean), 200.9 (standard deviation)			

7011

Table A8. Continued.

Biomass (g m^{-2})	Component	Location	Reference
(c)			
1013	Epiphytic bryophytes (<i>Frullania atrata</i> , <i>Phyllogonium fulgens</i>) and epiphyllous algae (<i>Trentepohlia</i>)	Tropical rainforest (at greater depth within cloud forest canopy), Guadeloupe, French West Indies	(Coxson et al., 1992)
422	Epiphytic non-crustose lichens and bryophytes	Mt. Baker, Washington, USA	(Rhoades, 1981)
332	Epiphytic lichens and bryophytes	Old-growth conifer forest, Pacific northwest, USA	(McCune et al., 1997)
221	Epiphytic bryophytes (<i>Frullania atrata</i> , <i>Phyllogonium fulgens</i>) and epiphyllous algae (<i>Trentepohlia</i>)	Tropical rainforest (upper 1.5 m of cloud forest canopy), Guadeloupe, French West Indies	(Coxson et al., 1992)
200	Epiphytic mosses (mainly <i>Hypnum andoi</i>)	Vosges Mountains, France	(Frahm, 2008)
155	Epiphytic cyanolichens, alectorioid and other lichens, and bryophytes	Pseudotsuga, Thuja and Tsuga stands in Oregon and Washington, USA	(McCune, 1993)
125	Epiphytic cyanolichens, alectorioid and other lichens, and bryophytes	Pseudotsuga, Thuja and Tsuga stands in Oregon and Washington, USA	(McCune, 1994)
104	Epiphytic lichens (dominated by <i>Pseudevernia</i> spp.)	Fir plantation, Massif du Pilat, Loire, France	(André et al., 1975)
84	Epiphytic lichens (<i>Lobaria</i> , <i>Usnea</i> , <i>Hypogymnia</i> , <i>Alectoria</i> , and <i>Parmelia</i> spp.)	Conifer, oak and northern hardwood forests, Canada and USA	(Pike, 1978)
55	Cyanolichen (<i>Lobaria oregana</i>)	Douglas fir and western hemlock dominated forest, Pacific Northwest, USA	(Brown and Dalton, 2002)
41	Epiphytic lichens (<i>Alectoria</i> spp., <i>Bryoria</i> spp., and foliose lichens)	Entire needle trees, British Columbia, Canada	(Campbell and Coxson, 2001)
1	Epiphytic noncrustose cyanolichens	Very moist submontane forest, (2700 m a.s.l.), Colombia	(Forman, 1975)
(63–350)	Non-crustose lichens and bryophytes	Varies forests in USA and Canada	(Rhoades, 1981)
(90–260)	Epiphytic lichens (<i>Alectoria</i> spp., <i>Bryoria</i> spp., <i>Usnea</i> spp., fruticose lichens, and cyanolichens) and mosses	Old-growth Douglas fir forest in Oregon, USA	(Sillett and Rambo, 2000)
(108–133)	Canopy lichen community	Wet-temperate rainforest, British Columbia, Canada	(Benson and Coxson, 2002)
(39–50)	Lichen spp.	Douglas fir forest, Oregon, USA	(Becker, 1980)
(<1)	Epiphytic cyanolichens	Gray beech forests, North Carolina, USA	(Becker, 1980)
140 (median), 204.5 (arithmetic mean), 278.7 (standard deviation)			

7012

Table A8. Continued.

Biomass (g m ⁻²)	Component	Location	Reference
(d)			
192 ^{a)}	Lichens (<i>Sticta</i> sp.)	Rainforest, Colombia	(Forman, 1975)
167 ^{a)}	Lichens with green algae	Rainforest, Colombia	(Forman, 1975)
162 ^{a)}	Epiphytic cyanolichens	Appalachian Mountains, North Carolina, USA	(Becker, 1980)
153 ^{a)}	Lichens with non-heterocystic cyanobacteria	Rainforest, Colombia	(Forman, 1975)
151 ^{a)}	Homoiomerous cyanolichen (<i>Leptogium phyllocarpum</i>)	Lower montane rainforest, Cordillera Central, Republic of Panama	(Lange et al., 2000)
136 ^{a)}	Lichens (<i>Leptogium</i> sp.)	Rainforest, Colombia	(Forman, 1975)
134 ^{a)}	Epiphytic <i>Lobariaceae</i> (<i>Sticta tomentosa</i>)	Lower montane rainforest, Cordillera Central, Republic of Panama	(Zotz et al., 1998)
129 ^{a)}	Homoiomerous cyanolichen (<i>Leptogium cyanescens</i>)	Lower montane rainforest, Cordillera Central, Republic of Panama	(Lange et al., 2000)
105 ^{a)}	Homoiomerous cyanolichen (<i>Leptogium azureum</i>)	Lower montane rainforest, Cordillera Central, Republic of Panama	(Lange et al., 2000)
70 ^{a)}	Cyanobacterial basidiolichen (<i>Dictyonema glabratum</i>)	Lower montane rainforest, Cordillera Central, Republic of Panama	(Lange et al., 1994a)
53 ^{a)}	Corticolous lichen (<i>Coenogonium linkii</i>)	Tropical evergreen lowland forest, Les Nouragues National Park, French Guiana	(Lakatos et al., 2006)
(10–720) ^{b)}	Epiphytic lichens and bryophytes	Montane and lowland rainforests, Ecuador	(Freiberg and Freiberg, 2000)
136 (median), 132 (arithmetic mean), 41.7 (standard deviation)			

a) per m² of thallus surface area;b) per m² of branch surface area.

7013

Table A9. Estimated annual fluxes of nitrogen fixation by (a) BSC and (b) EPC.

Nitrogen fixation flux (g m ⁻² a ⁻¹)	Crust components	Location	Reference
(a)			
5.7	BSC, cyanobacteria	Western Kalahari, Botswana	(Skarpe and Henriksson, 1987)
1.3	BSC, undisturbed	Cold desert, Utah, USA	(Belnap, 2002)
1.0 ^{a)}	BSC, cyanobacteria	Semiarid grassland, Arizona, USA	(Mayland et al., 1966)
0.6	BSC, cyanobacteria	Savanna, Nigeria	(Isichei, 1980)
0.4	BSC, cyanobacterium	Tropical volcano, Guadeloupe	(Sheridan, 1991b)
0.4	BSC	Lake Michigan sand dunes, USA	(Thiet et al., 2009)
0.2	Cyanobacteria on soil	Signy Island, South Orkney Islands	(Horne, 1972)
0.14	BSC, disturbed	Cold desert, Utah, USA	(Belnap, 2002)
0.13	BSC	Southern Utah, USA	(Veluci et al., 2006)
(0.2–37)	BSC, cyanobacteria and cyanolichens	Desert ecosystems	(Belnap et al., 2001)
(2.5–10.0)	BSC, cyanobacteria	Great Basin desert, Utah, USA	(Veluci et al., 2006)
(0.1–10.0)	BSC	Various (semi-)arid locations	(Evans and Lange, 2003)
(0.7–1.8)	BSC	Sonoran Desert, Arizona, USA	(Veluci et al., 2006)
(0.9–1.3)	BSC, undisturbed	Cold desert, Utah, USA	(Belnap, 2002)
(0.1–0.2)	Bryophyte-cyanobacteria	Antarctica	(Turetsky, 2003)
0.4 (median), 1.1 (arithmetic mean), 1.8 (standard deviation)			
(b)			
42.3	EPC, cyanobacteria (<i>Nostoc</i> and <i>Stigonema</i> spp.)	Mangrove boles and branches, Guadeloupe, French West Indies	(Sheridan, 1991a)
3.8	EPC, cyanobacteria (<i>Calothrix</i> and <i>Lyngbya</i> spp.)	<i>Spartina alterniflora</i> stems, transplanted salt marsh, North Carolina, USA	(Currin and Pearl, 1998)
2.6	EPC, cyanobacteria (<i>Calothrix</i> and <i>Lyngbya</i> spp.)	<i>Spartina alterniflora</i> stems, natural salt marsh, North Carolina, USA	(Currin and Pearl, 1998)
0.5 ^{b)}	EPC, cyanobacteria (<i>Calothrix</i> and <i>Anabaena</i> spp.)	<i>Spartina alterniflora</i> leaves and stems, salt marsh, Georgia, USA	(Hanson, 1977)
0.2 ^{c)}	EPC, cyanobacteria (<i>Calothrix</i> and <i>Anabaena</i> spp.)	<i>Spartina alterniflora</i> leaves and stems, salt marsh, Georgia, USA	(Hanson, 1977)
0.2	EPC, symbiotic cyanobacterium (<i>Nostoc</i> sp.) within feather moss (<i>Pleurozium schreberi</i>)	Boreal forests	(DeLuca et al., 2002)
0.2	Epiphytic lichens	Old-growth Douglas fir forest of Pacific Northwest, USA	(Brown and Dalton, 2002)
0.15	Epiphytic cyanolichens (<i>Lobaria oregana</i>)	Pseudotsuga, Thuja and Tsuga stands (WRCC site), Washington, USA	(Antoine, 2004)
(0–6)	EPC	Temperate and tropical regions	(Freiberg, 1998)
(0.3–1.7)	Epiphytic cyanolichens (<i>Lobaria oregana</i>)	Pseudotsuga, Thuja and Tsuga stands (HJA site), Washington, USA	(Antoine, 2004)
(0.2–0.8)	Epiphytic lichens	Rainforest in Colombia (2700 m a.s.l.)	(Forman, 1975)
(0.2–0.5)	Understory EPC, cyanobacteria	Premontane rainforest in Costa Rica	(Freiberg, 1998)
(0.1–0.5)	Epiphytic lichens	Scotland and Sweden	(Forman, 1975)
0.35 (median), 6.2 (arithmetic mean), 14.6 (standard deviation)			

a) maximum; b) average from control and experimental plot – 10 h fixation period estimate; c) average from control and experimental plot – 24 hour fixation period estimate.

7014

Fig. 1. Exemplary photographs of microbiotic crusts. **(A)** Biological soil crust (BSC) showing cyanobacteria and chlorolichen *Psora decipiens*, Nama Karoo semi desert, Western Cape, South Africa. **(B)** Biological rock crust (BRC) showing chlorolichen (*Rhizocarpon geographicum* aggr.), Sadnig, Eastern Alps, Austria. **(C)** Epiphytic crust (EPC) showing lichen and bryophytes on *Acer pseudoplatanus* tree in a city park, Trier, Germany.