

# Injurious tail biting in pigs: how can it be controlled in existing systems without tail docking?

R. B. D'Eath<sup>1†</sup>, G. Arnott<sup>1,2</sup>, S. P. Turner<sup>1</sup>, T. Jensen<sup>3</sup>, H. P. Lahrmann<sup>3</sup>, M. E. Busch<sup>3</sup>, J. K. Niemi<sup>4</sup>, A. B. Lawrence<sup>1</sup> and P. Sandøe<sup>5</sup>

<sup>1</sup>SRUC, West Mains Road, Edinburgh EH9 3JG, UK; <sup>2</sup>Queen's University Belfast, School of Biological Sciences, 97 Lisburn Road, Belfast BT9 7BL, UK; <sup>3</sup>Danish Agriculture & Food Council, Pig Research Centre, Axeltovej 3, 1609 Copenhagen V, Denmark; <sup>4</sup>MTT Agrifood Research Finland, Economic Research, Kampusranta 9, FI-60320 Seinäjoki, Finland; <sup>5</sup>Department of Large Animal Sciences and Department of Food and Resource Economics, University of Copenhagen, Grønnegårdsvej 8, 1870 Frederiksberg C, Copenhagen, Denmark

(Received 28 October 2013; Accepted 9 April 2014; First published online 6 June 2014)

---

*Tail biting is a serious animal welfare and economic problem in pig production. Tail docking, which reduces but does not eliminate tail biting, remains widespread. However, in the EU tail docking may not be used routinely, and some 'alternative' forms of pig production and certain countries do not allow tail docking at all. Against this background, using a novel approach focusing on research where tail injuries were quantified, we review the measures that can be used to control tail biting in pigs without tail docking. Using this strict criterion, there was good evidence that manipulable substrates and feeder space affect damaging tail biting. Only epidemiological evidence was available for effects of temperature and season, and the effect of stocking density was unclear. Studies suggest that group size has little effect, and the effects of nutrition, disease and breed require further investigation. The review identifies a number of knowledge gaps and promising avenues for future research into prevention and mitigation. We illustrate the diversity of hypotheses concerning how different proposed risk factors might increase tail biting through their effect on each other or on the proposed underlying processes of tail biting. A quantitative comparison of the efficacy of different methods of provision of manipulable materials, and a review of current practices in countries and assurance schemes where tail docking is banned, both suggest that daily provision of small quantities of destructible, manipulable natural materials can be of considerable benefit. Further comparative research is needed into materials, such as ropes, which are compatible with slatted floors. Also, materials which double as fuel for anaerobic digesters could be utilised. As well as optimising housing and management to reduce risk, it is important to detect and treat tail biting as soon as it occurs. Early warning signs before the first bloody tails appear, such as pigs holding their tails tucked under, could in future be automatically detected using precision livestock farming methods enabling earlier reaction and prevention of tail damage. However, there is a lack of scientific studies on how best to respond to outbreaks: the effectiveness of, for example, removing biters and/or bitten pigs, increasing enrichment, or applying substances to tails should be investigated. Finally, some breeding companies are exploring options for reducing the genetic propensity to tail bite. If these various approaches to reduce tail biting are implemented we propose that the need for tail docking will be reduced.*

---

**Keywords:** pigs, housing, enrichment, tail biting, behaviour

## Implications

Tail biting in growing pigs is a serious welfare and economic problem, and there is pressure to avoid tail docking. For the first time relying only on studies where tail damage was recorded, we review the evidence on controlling tail biting in pigs that are not tail docked. Adequate feeder space and manipulable substrate provision are important, but more work is needed on the type and quantity of substrate needed.

Vigilance for behavioural signs which occur before the first damaging biting would enable rapid detection and prevention/early response to outbreaks. Genetic selection could play a role in reducing tail biting.

## Introduction

Tail biting in domestic pigs occurs when pigs bite and chew the tails of pen-mates. It is a considerable animal welfare (Munsterhjelm *et al.*, 2013) and economic problem, causing painful injuries which are a site for further infection

---

† E-mail: rick.death@sruc.ac.uk

(Sihvo *et al.*, 2012), resulting in carcass losses for producers (Valros *et al.*, 2004; Kritas and Morrison, 2007) and reducing weight gain (Wallgren and Lindahl, 1996; Sinisalo *et al.*, 2012). Several risk factors have been proposed, suggesting multi-factorial causation (Schröder-Petersen and Simonsen, 2001; EFSA, 2007) and three different aetiologies have been proposed (Taylor *et al.*, 2010). Removal of part of the tail (tail docking) a few days after birth usually reduces the likelihood and severity of tail biting (Sutherland and Tucker, 2011). Where tail docking is banned, tail biting incidence usually increases, even when the housing environment and management are improved (D'Eath *et al.*, 2014).

However, even though tail docking reduces tail biting, it does not eliminate it and has significant drawbacks: it is an acutely painful mutilation, and it may 'mask' the real underlying problems in housing and management that result in tail biting (Sutherland and Tucker, 2011). For these reasons, the EU Council Directive (2001/93/EC amending Directive 91/630/EEC, The Council of The European Union, 2001b) came into force from January 2003 banning the 'routine' tail docking of pigs, unless 'there is evidence that injuries... to other pigs' ears or tails have occurred' and insisting that before resorting to tail docking 'other measures shall be taken to prevent tail biting... taking into account environment and stocking densities'. It goes on to state that '...pigs must have permanent access to a sufficient quantity of material to enable proper investigation and manipulation activities, such as straw, hay, wood, sawdust, mushroom compost, peat or a mixture of such, which does not compromise the health of the animals'. Despite this clear legal signal, tail docking continues for 95% or more of pigs in European pig producing countries such as Germany, Denmark, Belgium, France, Ireland, the Netherlands and Spain, and for over 80% in the United Kingdom (EFSA, 2007; Harley *et al.*, 2012).

Perhaps in response to this gap between policy and reality, the European Commission (Directorate-General for Health and Consumers, DG Sanco; Bergersen, 2013) is currently engaging in a process to agree and clarify the guidance to farmers associated with the mentioned Directive and its later versions (the latest being 2008/120/EC on the protection of pigs). Some countries already go further than the EU directives in restricting tail docking (Mul *et al.*, 2010). In Denmark, no more than half of the tail may be docked, and in the Netherlands, a voluntary agreement exists between farmers and government to phase out tail docking entirely by 2023 (Spoolder *et al.*, 2011). A few countries already have either a complete ban on tail docking (Sweden, Finland, Switzerland; EFSA, 2007; Swiss Federal Council, 2008) or a ban on docking without anaesthesia (Norway; EFSA, 2007) so that tail docking is rare. At the same time animal welfare protection organisations in many European countries focus on tail docking as a sign of welfare problems in intensive pig production; and in some countries political pressure is building up in favour of an effective ban on tail docking.

In another article, we consider the decisions facing farmers under current EU rules as to whether to tail dock, and the economic, legal and pig welfare consequences of this decision

(D'Eath *et al.*, 2014). In the present article, we ask how farmers can become better at controlling tail biting without the use of tail docking. Our review focuses on changes that would be possible in existing systems, rather than considering radical system re-design (De Greef *et al.*, 2011). Various knowledge gaps are identified and promising areas for future innovation are proposed. We begin by introducing the nature of tail biting, and then review risk factors relating to the pigs' environment. For the first time, we rely only on studies which reported effects on tail injuries, rather than those which describe pigs' non-injurious interactions with tails (tail in mouth). The relationships between these risk factors and the underlying process(es) that govern the expression of tail biting are poorly understood, and we present a new illustration of the diversity of hypotheses. A novel illustrated meta-analysis quantifies the effectiveness of enrichment on tail biting in undocked pigs, and the practical experiences of countries and production systems in which tail docking is banned are considered. The next section of the review then focuses on risk factors that relate to characteristics of the pigs themselves, including the possibility of genetic selection to reduce tail biting. Finally we consider the prospects for early detection of tail biting outbreaks, possibly by automated means, which could facilitate targeted prevention measures. Farmers react to tail biting in various ways but little is known about the efficacy of these measures in preventing the further escalation of an outbreak.

### **Tail biting: why it remains an intractable problem**

#### *Tail biting occurs in outbreaks*

Damaging tail biting occurs in a sporadic way, in unpredictable 'outbreaks', rather like an infectious disease (Blackshaw, 1981). For example, in one study using abattoir data, 'high incidence farms' were identified at one point in time, but when a similar 'high incidence' list was made a few months later, most of the farms were different – although there were a few farms with a persistent problem (Busch *et al.*, 2004). In general, while some of the risk factors that affect the overall incidence of tail biting are known, for any given outbreak, the specific triggering factor(s) are usually difficult to identify. Sometimes a change (weather, season, food, or disease outbreak) can be identified, but often no obvious change has occurred, and the cause may be down to variability in individual pigs' threshold of response to risk factors.

#### *Tail biting can spread quickly within the group*

Tail biting begins with one pig in the pen starting to bite. Tail damage can increase rapidly, with one study reporting that progress from bite marks to a clearly visible tail wound took on average 7 days (Zonderland *et al.*, 2010b), although practical experience suggests that it can occur even more quickly. Over time, biting pigs may continue or escalate their biting of existing victims, but also begin biting other pigs in the group (Niemi *et al.*, 2011). Additionally, other pigs in an affected pen begin tail biting too, perhaps because they copy the behaviour (social facilitation; Blackshaw, 1981) or the

bitten tails might stimulate investigation and biting (stimulus enhancement; Fraser, 1987a). Although never formally studied, there appears to be considerable variation in the rate at which a pig increases its tail biting behaviour, and in the rate of spread to new biters. In one study, a batch of pigs already showing tail biting, moved to an environment with considerable space and access to rooting substrates, subsequently showed healing and improvement over time (De Greef *et al.*, 2011), suggesting that escalation is not inevitable.

#### *Scientific investigation of tail biting is difficult*

Tail biting is challenging to study. Its apparently sudden, unpredictable appearance and rapid spread can make it hard to investigate the events immediately before and after an outbreak begins. Its sporadic occurrence also means that a number of experimental studies have failed to observe any damaging tail biting at all. Such studies often report the effects of experimental treatments on tail investigation behaviour ('tail in mouth' Petersen *et al.*, 1995; Schroder-Petersen *et al.*, 2004), which is at best an indirect indicator of tail biting, because 'tail in mouth' behaviour may or may not be a precursor to damaging tail biting (EFSA, 2007). Other studies include all pig-directed oral behaviours including ear and flank biting, and sometimes belly nosing, together in a single category (e.g. Jensen *et al.*, 2010; Zwicker *et al.*, 2013). This lack of precision makes interpretation difficult if the focus is on tail biting alone (Taylor *et al.*, 2010). In order to avoid these problems with indirect or imprecise indicators, this review focuses on studies where tail damage (with evidence of partial tail loss or of injury severe enough that blood was drawn) was the end point.

The sporadic occurrence of tail biting, and difficulties with experimental studies mean that multi-farm epidemiological studies (Moinard *et al.*, 2003; Goossens *et al.*, 2008) or abattoir data (Valros *et al.*, 2004; Harley *et al.*, 2012), sometimes combined with farm surveys (Hunter *et al.*, 2001), are often used to study tail biting. These usually record tail damage, and can find risk factors associated with it, but unlike experiments, are unable to determine cause and effect, so must be interpreted with caution. We begin by looking at risk factors in the pigs' environment, and then explore risk factors intrinsic to the pig. Some of these risk factors and causes of tail biting may also affect the related problems of ear- and flank biting (Brunberg *et al.*, 2011), but this is beyond the scope of this paper.

#### **Risk factors for tail biting in the pigs' environment and how to manage them**

Tail biting does not have a single cause. It is a multi-factorial problem, and a variety of risk factors have been identified which are associated with it. Various efforts have been made to review all the currently known risk factors to weight their importance in order to influence policy makers (Bracke *et al.*, 2006; EFSA, 2007; Spooler *et al.*, 2011) and to provide practical advice to farmers (Bracke *et al.*, 2004; Jensen *et al.*, 2004; Taylor *et al.*, 2012).

Taylor *et al.* (2010) in a recent review made a convincing case that there were at least two and possibly three different types of tail biting: two-stage, sudden-forceful and obsessive. 'Two stage' tail biting results from re-directed foraging due to a lack of suitable substrates. There is a progression from investigation and gentle manipulation of tails (stage 1) to damaging biting (stage 2). The second type, 'sudden forceful' tail biting is an aggressive behaviour (Van Putten, 1969; Moinard *et al.*, 2003) apparently resulting from frustration over a lack of access to food, water or lying space. Pigs approaching a fully occupied resource such as a feeder may resort to biting at tails as the most readily available target for aggression. For example, in a recent study, 60% of the tail biting by pigs, which had limited feeder access (Palander *et al.*, 2013), occurred within 1 m of the feeder (A. Valros, personal communication). The third type, 'obsessive tail biting' is characterised by certain individual pigs which appear to be fixated on tails and go from one tail to another, inflicting damaging bites (Beattie *et al.*, 2005; Van de Weerd *et al.*, 2005). Here, we consider 'obsessive' biters to be individuals which are more likely than other pigs either to begin or to continue tail biting through the mechanisms explained above (and illustrated in Figure 1) for two-stage or sudden forceful tail biting.


The mechanism of action of each possible risk factor on the underlying processes controlling the expression of tail biting is in many cases unknown. Figure 1 illustrates many of the possible connections between proposed environmental risk factors and the underlying processes of 'two stage' and 'sudden-forceful' tail biting (Taylor *et al.*, 2010). The nature of each possible connection is described in Supplementary Material S1. Evidence for the effect of risk factors on damaging tail biting is discussed further below.

#### **Availability of manipulable materials**

Manipulable materials which are attractive to pigs as measured by their motivation to access them (Holm *et al.*, 2008; Jensen *et al.*, 2008) or by the time pigs spend interacting with them over a sustained period have the characteristics 'ingestible', 'odorous', 'chewable', 'deformable' and 'destructible' (Van de Weerd *et al.*, 2003; Studnitz *et al.*, 2007; Van de Weerd and Day, 2009). The opportunity to perform investigation and manipulation behaviours are in themselves important for pig welfare (Studnitz *et al.*, 2007; Van de Weerd and Day, 2009), but here we focus on whether manipulable materials can reduce damaging tail biting apparently by providing an alternative outlet for investigatory behaviour.

#### *Difficulties with the provision of loose manipulable materials on the floor*

Systems making use of full or part-slatted floors, enabling automatic collection of pig faeces and urine (slurry) are common in indoor pig production. In comparison with straw-bedded systems, the labour (cleaning and waste handling) and input costs (e.g. of straw, peat and other substrates) are lower (Bornett *et al.*, 2003), some environmental impacts



**Figure 1** Postulated relationships between the underlying processes of tail biting (text in bold, connected in order by solid arrows) and various known or suspected risk factors (text in plain type) connected with dashed numbered arrows to show how some of the risk factors might influence each other or the underlying process of tail biting. Some proposed risk factors for which the evidence is currently weak (e.g. disease and parasitism, draughts) are included where a plausible hypothesis exists. The meaning of the numbered arrows is explained in Supplementary Material S1.

may be lower (Stern *et al.*, 2005), and liquid slurry is more valuable as a fertiliser than solid manure (Sanchez and Gonzalez, 2005). The requirement to provide manipulable materials to occupy pigs, presents a difficulty for farmers with systems which rely on slatted-floors and liquid slurry handling (via pumps). Materials such as long (unchopped) straw do not easily pass through slats leading to pen fouling. Additionally, too much straw can separate from the liquid slurry and build up in the slurry pit, or if it does flow, it can block parts of slurry-handling systems, such as holes, pipes or vacuum-based slurry pumps (Tuytens, 2005; Day *et al.*, 2008). Ways to reduce the problem of straw blockage may include use of chopped straw, in combination with engineering solutions, such as larger diameter pipes (Evira, 2013), slurry pumps fitted with chopper blades, the use of smaller, faster flowing slurry systems (PRC, 2011), or progressive cavity pumps which are suitable for viscous liquids. Depending on the quantity and type of substrate, these measures may not be 100% effective but are more likely to be successful if considered at the building design stage.

Non-destructible materials such as metal chains, or rubber or hard plastic objects have been tried. Although pigs may initially interact with these due to their novelty, interest in them usually declines rapidly over a few days (Van de Weerd *et al.*, 2003; Van de Perre *et al.*, 2011). Even a repeating cycle

of different objects may not be enough, as re-introduction of the same object after an interval of several weeks is usually not as effective at sustaining interest as a novel object would be (Van de Perre *et al.*, 2011). The European Commission have made it clear that chains and other non-destructible materials are not sufficient to comply with the EU Council directive (EC, 2009). Gradually destructible materials, which take days or weeks to be chewed through such as wooden poles (often mounted vertically in a tube at the side of the pen, or suspended from a chain) are popular with farmers in some countries as they require less regular replenishment than other more readily destructible substrates and appear to comply with the EU Council directive (2001/93/EC) which lists wood as a suitable material. Wooden poles were found to be an effective enrichment for reducing tail damage in a recent unpublished Finnish study using freshly felled tree trunks 5 to 10 cm in diameter suspended on chains horizontally below snout level (Telkänranta *et al.*, 2014a). However, since some of the features required to make a material attractive to pigs are lacking (ingestible, odorous) or weak (chewable, deformable, destructible) in hard wood poles, the need to use soft, fresh woods which do have these features could be important.

In the face of these difficulties, an important question is whether it is possible to provide sufficient manipulable

materials to pigs within existing intensive housing systems in order to reduce tail biting to a level which is acceptably low from a management, production and welfare perspective without the need to tail dock (D'Eath *et al.*, 2014).

#### *Alternative ways of providing manipulable materials*

In part-slatted floored pens, it is possible to provide loose material such as chopped straw, peat or sawdust, which in small quantities may be used with slurry pumps (Munsterhjelm *et al.*, 2009). Substrate can be provided on the solid floor, while pigs defecate and urinate in the slatted part. To limit the passage of substrate from the solid to the slatted part of the floor, pen designs incorporating barriers (e.g. 50 mm high wooden strip, Zwicker *et al.*, 2013) or where the slatted area is raised (BPEX, 2010) may be used. Practical experience suggests that such designs are usually not entirely successful, especially in high temperatures where pigs may choose to defecate in the lying area, wallowing in the wet faeces to keep cool, and at higher stocking densities where functional separation of lying and dunging areas becomes more difficult to achieve, particularly in older pigs (Jensen *et al.*, 2012).

Faecal contamination of manipulable substrates is a common problem which reduces their attractiveness to pigs (Scott *et al.*, 2009), and this contamination can be reduced by hanging objects in the pen. Hanging of substrates limits the form of interaction, for example chewing may be possible but not rooting (Day *et al.*, 2008). This might be important, or different forms of investigatory behaviour may substitute for one another in preventing tail biting, as long as the pigs are occupied. Hanging objects thus may have potential: in a meta-analysis of the time spent by pigs interacting with enrichment, properties promoting this interaction included enrichments which were suspended and/or deformable (Averós *et al.*, 2010). For example pigs show sustained interest in interacting with destructible ropes (Trickett *et al.*, 2009), or hanging objects with an edible component (Van de Weerd *et al.*, 2003), and 'flavoured rope' devices for pigs are being sold commercially in Finland. However, the effects of these forms of enrichment on tail biting have not been investigated.

Another approach is to deliver loose manipulable materials by means of an elevated rack, so that pigs can gradually obtain the material for themselves over a period (Beattie *et al.*, 2001; Van de Weerd *et al.*, 2006; Zwicker *et al.*, 2012 and 2013). This has the potential advantage of 'double' interaction (in the rack, and beneath it: on the floor, or in a box or feeder; Zwicker *et al.*, 2012) which might mean less material can be used for the same total amount of interest from the pigs. A related approach is to use a low-level rooting box which can contain loose materials and keeps them separate from slats (Van de Weerd *et al.*, 2003; De Greef *et al.*, 2011).

#### *Quantifying the effects of different enrichment methods on tail damage*

Studies published in refereed journals which compare the effect of different types and quantities of manipulable substrates on tail damage are summarised in Table 1. Most of

the studies had pigs with intact tails, but some were docked, as indicated in the table legend. The studies all focus on grower-finisher pigs, except for Zonderland *et al.* (2008) which used weaners. Different indices of tail damage were used by different authors: most studies report either the percentage of pigs removed from the study with severe tail injury, or the percentage of pigs or of pens having tail wounds. One study (Munsterhjelm *et al.*, 2009) used a tail lesion index (scoring from 0 to 2). To compare studies that used different measures, we calculated the fold-change in tail damage for each pair of treatments in these studies (i.e. the reduction in tail injury following the provision of one type of manipulable substrate compared with another). Where one of a pair of substrates had zero damage, it was not possible to calculate a fold-change, so 'max' was reported in Table 1, and this value did not contribute to the mean fold change, probably resulting in an underestimate of the effect size. Most studies compared deep straw with either no enrichment, or with minimal enrichment with chains or hanging toys, considered to represent commercial practice. In Figure 2, the information from the studies in Table 1 is summarised graphically, giving a quantification of the relative value of different materials as it is drawn to scale using the mean log 'fold' difference between observed levels of biting damage as the distance between the materials. A log scale was used so that 'fold' differences could be added together on the same scale and shown relative to each other in a single diagram (since e.g.  $\log 2 + \log 3 = \log 6$ ).

In terms of manipulable substrate treatments that are compatible with fully or part-slatted floors, straw racks and light straw ( $\leq 20$  g/pig per day) are probably the most promising treatments for which data are available. Provision of straw in racks reduces tail damage compared with a rubber hose, chain or hanging toy, with two studies finding a small but consistent reduction in the percentage of pens with tail wounds (fold-improvement of 1.9 or 1.7, Van de Weerd *et al.*, 2006; Zonderland *et al.*, 2008). In one study, the straw rack affected minor tail injuries, but was more effective at reducing severe tail damage (Van de Weerd *et al.*, 2006), which suggests that the straw rack might have reduced the rate of escalation of biting.

Light straw (10 g twice a day per pig, Zonderland *et al.*, 2008), or light chopped straw and wood shavings (12.5 g a day per pig, Munsterhjelm *et al.*, 2009) were both highly effective at reducing tail damage compared with minimally enriched treatments in two studies, with fold-differences almost as high as for deep straw studies (see Figure 2). Unfortunately, neither of these studies included plentiful loose material as a positive control. In a producer survey combined with an abattoir study, Hunter *et al.* (2001) found that 'light straw' use reduced tail biting damage risk compared with no straw. Despite these positive findings, chopped straw may not be as attractive to pigs as long straw: A behaviour study comparing chopped with long straw (each 400 g/pig per day) suggested that chopped straw offers fewer possibilities for interaction and observed tail biting behaviour increased (Day *et al.*, 2008). However, this study included non-injurious

**Table 1** Summary of comparative manipulable material studies in which tail injuries were reported for growing pigs

Manipulable material A (g/pig per day)	Outcome variable	Value of outcome variable for material A	Manipulable material B (g/pig per day)	Value of outcome variable for material B	Number of pigs (pens) per treatment	P-value of difference <sup>1</sup>	Fold-improvement in outcome of material B over A <sup>2</sup>	Mean fold-improvement of similar studies	Log <sub>10</sub> mean fold change	References
None <sup>3</sup>	% Removed for tail injury	17.5	Compost rack (500)	<1	108 (6)	<0.05	17.5+	17.5+	1.24	Beattie <i>et al.</i> (2001)*
None	% Pigs with tail wounds	2.3	Straw bedding	0.5	~4550 (40 farms)	<0.001	4.6	8.6	0.93	Courboulay <i>et al.</i> (2009) <sup>†</sup>
None	% Removed for tail injury	2.5	Straw (490)	0.2	512 (16)	ns	12.5			Scott <i>et al.</i> (2007) <sup>†</sup>
Hanging toy	% Pigs with tail wounds	42.2	Straw (400)	0	181 (12)	na	Max			Van de Weerd <i>et al.</i> (2005)*
Hanging toy	% Died or were removed for tail injury	11.7	Straw (400)	1.4	2048 (64)	<0.001	8.4	6.7+	0.83	Scott <i>et al.</i> (2006) <sup>†</sup>
Hanging toy	% Pens with tail wounds	83	Straw (5 cm deep)	17	72 (6)	<0.05	4.9			Van de Weerd <i>et al.</i> (2006)*
	% Removed for tail injury	11.1		0		na	Max			
None	Tail lesion index	0.7	Light chopped straw/wood shavings (12.5)	0.1	126 (31)	<0.05	7	7	0.85	Munsterhjelm <i>et al.</i> (2009)*
Rubber hose or chain	% Pens with tail wounds	56	Light straw (20)	8	240 (24)	<0.05	7	7	0.85	Zonderland <i>et al.</i> (2008)*
Rubber hose or chain	% Pens with tail wounds	56	Straw rack (5)	29	240 (24)	ns/<0.05 <sup>4</sup>	1.9			Zonderland <i>et al.</i> (2008)*
Chain and rubber-covered chain	% Prevalence of tail lesions	10.6	Straw rack	12.9	336 (12)	ns	0.8			Scollo <i>et al.</i> (2013) <sup>‡</sup>
Hanging toy	% Pens with tail wounds	83	Straw rack <sup>5</sup>	50	72 (6)	ns	1.7	1.3	0.11	Van de Weerd <i>et al.</i> (2006)*
	% Removed for tail injury	11.1		1.4		na	(7.9)			
Rootable feed dispenser	% Pens with tail wounds	33	Straw rack <sup>5</sup>	50	72 (6)	ns	0.6			Van de Weerd <i>et al.</i> (2006)*
	% Removed for tail injury	1.4		1.4		na	(1)			
Straw rack (5)	% Pens with tail wounds	29	Light straw (20)	8	240 (24)	ns	3.6	3.6	0.55	Zonderland <i>et al.</i> (2008)*
Straw rack <sup>5</sup>	% Pens with tail wounds	50	Straw (5 cm deep)	17	72 (6)	ns	2.9	2.9	0.46	Van de Weerd <i>et al.</i> (2006)*
	% Removed for tail injury	1.4		0		na	Max			

An asterisk (\*) by the reference indicates that pigs had intact (undocked) tails, a dagger (†) indicates they were tail docked, and a double dagger (‡) indicates that half the pigs were docked in a 2×2 experimental design. All of the studies were controlled experiments with the exception of Courboulay *et al.* (2009) which scored tails in part- or fully slatted and straw systems in an on-farm observational study of 82 farms. Fold-improvement is a result of the tail biting value for the first manipulable material divided by the value for the second. Where two similar materials are named, the average was taken.


<sup>1</sup>P-values are reported where these are available in the source paper; ns means that the difference was not significant, but numerical values have still been used to contribute to estimate the mean fold-change; na means that the P-value is not available as it was not reported in the source paper.

<sup>2</sup>'Max' indicates that the improvement was such that tail biting reduced to zero in the second treatment, and this information was not used for the calculation of average fold improvement. Values in parentheses in this column were not used for calculation of the 'Mean fold improvement' – where two different outcome variables were reported for the same study, one of them had to be chosen for use with other studies – the most comparable outcome variables were used where possible.

<sup>3</sup>For the Beattie *et al.* (2001) study, 'None' includes the average of pens with nothing and pens with the non-manipulable empty overhead racks.

<sup>4</sup>In this study, straw rack and metal chain had significantly different % tail wounds, but straw rack and rubber hose did not. Metal chain and rubber hose had very similar levels of tail wounds so were combined for simplicity.

<sup>5</sup>The straw rack was described as a metal tube with a chain mail base which was filled with long straw (and with a tray on the floor underneath) but the quantity provided/used was not reported.



**Figure 2** Enrichment materials' relative effect at reducing tail biting based on Log<sub>10</sub> fold reductions in tail damage, using studies from Table 1. Line thickness indicates the number of studies used; thinnest lines = 1 study, intermediate lines = 2 studies, thickest lines = 3 or more studies. Shading of the box indicates the amount of material that is used up. Compost and Straw (shown in black), at least 500 g/pig per day, light straw (in dark grey, 12.5 to 20 g/pig per day), straw rack (5 g/pig per day). None and Straw used as reference, as these are the most common materials used across studies. This means that none and straw each have only one horizontal line. Light straw and Straw rack have multiple lines, which show the range of positions they could occupy relative to other substrates based on a number of studies.

chewing and biting, and tail damage was not reported. In contrast, a Danish study suggested that chopped and long straw (each at 100 g/pig per day) occupied pigs for a similar amount of time (Lahrmann and Steinmetz, 2011).

Our comparative survey has identified a number of data gaps: only two studies included comparisons of more than one pair of treatments, allowing the substrates to be placed into an overall ranking (Van de Weerd *et al.*, 2006; Zonderland *et al.*, 2008), and there was a paucity of studies investigating how different quantities of straw or other materials affect tail damage. Time spent exploring and manipulating straw rather than other pigs increases with straw quantity until above 300 g/pig per day (Olsson, 2011) or at around 500 g/pig per day (Pedersen *et al.*, 2013). However, tail biting occurred at very low levels in these studies, even in treatments with only 20 g/pig per day (undocked pigs, Olsson, 2011) or 10 g/pig per day (docked pigs, Pedersen *et al.*, 2013). Also, no studies have compared hanging toys with no enrichment, and none have looked at the effect of hanging destructible enrichments such as ropes on tail damage, except for one recent report in suckling piglets (Telkänranta *et al.*, 2014b), so there is considerable scope for further research.

#### *Manipulable materials as fuel for anaerobic digesters (AD)*

Materials which act as foraging enrichment for pigs could double as fuel for AD. This idea is being tested in the 'Starplus' system at Wageningen (Verdoes, 2014). ADs enable farmers to deal with farm wastes, producing energy (methane) and digestates which can be used as fertiliser. Pig slurry provides

micronutrients and trace elements needed for bacterial growth, but its energy content is low, so (non-wood) biological materials are added, some which could provide rooting/foraging (and eating) opportunities for pigs: chopped grass, maize or grass silage, sugar beet and kitchen waste (if concerns over biosecurity could be addressed). For example, pigs prefer chopped straw mixed with Maize silage over straw (Jensen and Pedersen, 2007; Jensen *et al.*, 2010) possibly because it may include edible components. Many questions remain, however: materials must be compatible with floor slats/slurry systems, a method to deliver substrate to the pens is required, fungal growth in wet fermenting materials can be a problem (T. Jensen, personal communication), and there are hygiene issues if pigs are eating material from the floor. Finally, fuel source costs, energy prices and government policies affect the economic feasibility of AD.

#### **Social factors: space allowance (stocking density), group size, mixing**

At space allowances lower than those currently recommended in the EU, reduced space allowance increased tail damage in one experiment (Kruider *et al.*, 1975). At space allowances closer to or within the recommended range, one multi-farm study found an association between reduced space allowance and tail injuries (Goossens *et al.*, 2008) but another similar study did not (Smulders *et al.*, 2008), and no effect was found in an experimental study (Street and Gonyou, 2008). Group size (Schmolke *et al.*, 2003; Smulders *et al.*, 2008; Street and Gonyou, 2008) and mixing of groups (Smulders *et al.*, 2008; Zonderland *et al.*, 2008) had no effect in studies where tail damage was reported.

#### **Feeding: feeder space, feed restriction, feed type, nutrients, minerals**

Restricted feeder space increased tail biting in one experimental study (damaged tails, Hansen *et al.*, 1982) and is a risk factor in epidemiological studies (Hunter *et al.*, 2001; Moinard *et al.*, 2003). Other experimental studies, in which low levels of tail biting occurred, found no effect of feeder space (Georgsson and Svendsen, 2001 and 2002). The form and presentation of feed may be important: pigs fed pelleted diets showed higher levels of tail injury than meal or liquid fed pigs in one study (Hunter *et al.*, 2001) while Temple *et al.* (2012) found liquid feed in a trough increased tail injury compared with wet feed in a hopper.

Nutritional qualities of the diet: protein, specific amino acids, minerals or high energy density have all been suggested to affect tail biting (Edwards, 2011), but there is little direct evidence of nutritional manipulations affecting tail damage. Experiments using 'model' tails suggest that attraction to blood may be increased if the diet is nutritionally inadequate in terms of protein (Fraser *et al.*, 1991) or minerals (Fraser, 1987b). Tail biting pigs were more attracted to cords soaked with pig blood than their non-biting pen mates (McIntyre and Edwards, 2002b) and this preference

can be reduced by the addition of the amino acid tryptophan to their diets (McIntyre and Edwards, 2002a). Differences in serotonin metabolism in the prefrontal cortex and an altered pattern of tryptophan uptake have been reported in tail biting pigs in contrast to bitten and unaffected group-mates or unaffected pigs from another group (Valros *et al.*, 2013). Additional salt in the diet or on the floor of the pen can increase foraging and drinking behaviour (Brooks, 2005), which may reduce biting, but it is not clear whether this was effectively a foraging enrichment or addressing a nutritional deficiency. Jaeger *et al.* (2013) proposed a novel causal pathway for tail biting: high energy density diets for weaner pigs (as well as exposure to various pathogens) result in a build-up of endotoxins which cause ear or tail necrosis, which then attracts biting. If the necrotic tissue is itchy, this could increase the tolerance of tail investigation and biting in victim pigs.

### **Climate: temperature, draughts, seasonal effects**

Either low (Temple *et al.*, 2012), or both low and high temperatures (Geers *et al.*, 1989) have been identified by epidemiological studies as risk factors for tail damage, and providing access to a water misting system can reduce tail injury in hot climates (Couboulay *et al.*, 2008). Seasonal effects on tail damage have been identified (Schrøder-Petersen and Simonsen, 2001, Busch, personal communication), the exact nature of which varies between different studies. It seems plausible that rapid changes in temperature (either up or down), an increase in draughts at certain times of year (known to affect activity, Scheepens *et al.*, 1991), or heat stress are likely to be the underlying cause of seasonal effects (Figure 1), as there is a limit to the capacity of ventilation/heating/cooling systems in most pig buildings.

### **Disease, including parasitism**

Disease has been proposed to be a risk factor (Edwards, 2011). Levels of tail damage are higher in herds with higher levels of respiratory illness (Elst *et al.*, 1988 cited by Moinard *et al.*, 2003; Edwards, 2011), and in a study where health records from individual pigs were examined, leg disorders and tail damage were highly correlated (Niemi *et al.*, 2012). Caution is required with the interpretation of epidemiological data, as disease may result from infections that follow tail biting (Moinard *et al.*, 2003; Kritas and Morrison, 2007), or poor health status may be an indirect indicator of less technically efficient farms.

Controlled studies in which measures to improve health result in a reduction in tail damage provide better evidence. Currently there is only an anecdotal report of tail biting being reduced following anthelmintic treatment (Barnikol, 1978) and an as-yet unpublished study concerning PCV2 vaccination (Parker *et al.* in preparation, cited by Edwards, 2011). So at this stage, the evidence for disease as a cause of tail biting is weak.

### **The experience of countries and assurance schemes where tail docking is banned**

Pig producers in some countries and assurance schemes have already had to adapt their systems to cope with greater restrictions on tail docking, and the changes they have made are instructive. The tail docking and housing system rules for grower-finisher pigs applied by selected non-docking European countries and selected assurance schemes are summarised in Table 2, with some systems which permit tail docking included for comparison.

The 'tail docking restricted or banned' farms have a number of features in common, many of which may reduce tail biting risk. The space allowance is usually more generous, with up to 50% more space per pig being provided. Fully slatted pens are not allowed, enabling manipulable materials to be provided on the solid-floored part of the pen (although part slatted, part drained floors are permitted in Finland). Compared with the EU minimum provision, there are more specific rules on the quantity of materials, usually by specifying the frequency of replenishment, or the behaviour that pigs must be able to perform: in Finland pigs must be able to make small piles of material, Freedom Food requires sufficient quantities of material for rooting, pawing and chewing behaviour. The type of material provided is often also restricted, for example Sweden and the Danish assurance scheme Antonius require straw, and Norway repeats the EU list, but stipulates wood chips rather than 'wood' which rules out the use of wooden posts.

The smaller scale of farms in Finland, Norway and Switzerland, compared with the United Kingdom and Denmark (Table 2) might enable a greater supervision of the animals (assuming more staff per pig), making detection and prevention of tail biting easier, and smaller farms often have lower disease risk (Goldberg *et al.*, 2000). For example, Finland is free from Porcine Reproductive and Respiratory disease, and mycoplasma and Salmonella are at low levels, although a lower density of farms and fewer pig movements including imports may also be important here.

Tail docking is not completely outlawed in all of the assurance schemes in Table 2. The assurance schemes Antonius, Outdoor (including Organic) in Denmark and Freedom Food allow farmers to apply for a dispensation to use tail docking for a limited time if a tail biting outbreak occurs. For example, Freedom Food farmers must annually seek written permission to dock, and if tail biting in the previous year was low, they are encouraged to trial a cessation of docking for some pigs, with the aim of stopping docking altogether. At each application, farmers must document the other measures they have taken to prevent tail biting and quantify their success. The standards give a detailed list of a number of environmental improvements that should be tried including providing straw and increasing feeder space. In 2010, 30% of Freedom Food breeding farms supplying indoor wean to finish herds requested permission to dock (Kate Parkes RSPCA, personal communication). This suggests that the majority of scheme members are managing to rear intact-tailed pigs, while those with tail biting problems

**Table 2** Comparison of minimum standards for housing grower-finisher pigs across countries and selected assurance schemes (from the UK and Denmark) that restrict or completely ban tail docking, with housing standards where docking is widespread (EU, Denmark and UK standard indoor housing)

Country	EU Directives	Denmark	Denmark	Denmark	UK	UK	UK	Sweden	Finland	Norway	Switzerland
System	–	Standard indoor <sup>1</sup>	Antonius	Outdoor (includes organic) <sup>2</sup>	Standard indoor	Freedom food	Organic <sup>3</sup>	–	–	–	–
Farm size (finish pigs)	234 <sup>4</sup>	2538 <sup>4</sup>	–	–	1038 <sup>4</sup>	–	–	1046 <sup>4</sup>	485 <sup>4</sup>	264 <sup>4</sup>	420 <sup>4</sup>
Space allowance 41 kg pigs (m <sup>2</sup> /pig)	0.4 <sup>5</sup>	0.4 <sup>1</sup>	0.5 <sup>6</sup>	1.4 (includes outdoor area) <sup>7</sup>	0.4 <sup>8</sup>	0.4 (1.17 for straw yard, mucked out monthly) <sup>9</sup>	555 (0.8 indoor only in extreme weather) <sup>3</sup>	0.48 <sup>10</sup>	0.6 <sup>11</sup> 0.4 <sup>12</sup>	0.5 <sup>13</sup>	0.6 <sup>14</sup>
Space allowance 101 kg pigs (m <sup>2</sup> /pig)	0.65 <sup>5</sup>	0.65 <sup>1</sup>	0.85 <sup>6</sup>	2.3 (includes outdoor area) <sup>7</sup>	0.65 <sup>8</sup>	0.75 (1.54 for straw yard, mucked out monthly) <sup>9</sup>	625 (1.3 indoor only in extreme weather) <sup>3</sup>	0.94 <sup>10</sup>	0.9 <sup>11</sup> 0.65 <sup>12</sup>	0.8 <sup>13</sup>	0.9 <sup>14</sup>
Floor – minimum solid area (% pen)	0 <sup>5</sup>	33 (grower), 50 (weaner) solid or drained by July 2015. Most already comply <sup>1</sup>	33 to 50 <sup>6</sup>	50 (of indoor area) <sup>7</sup>	0 <sup>8</sup>	66 <sup>9</sup>	50 <sup>3</sup>	70 to 75 <sup>10</sup>	67 can include drained floor where perforations are up to 10% of the area <sup>11</sup>	Solid-floored area large enough for all pigs to lie. <sup>13</sup>	67 lying area, permitted to have 'low degree of perforation for the drainage of liquids' must be solid by 2018 <sup>14</sup>
Tail docking	Not allowed routinely, only if evidence of injuries to ears or tails. 'Before (tail docking) other measures shall be taken to prevent tail biting and other vices taking into account environment and stocking densities. For this reason inadequate environmental conditions or management systems must be changed'. (vet or competent person can dock <7-day-old piglets) <sup>5</sup>	As EU, but no more than half the tail, and only 2 to 4-day-old piglets <sup>1</sup> , Docking is widespread	No, but vet can give a time-limited dispensation for tail biting problems <sup>6</sup>	No, but possible to get a dispensation for 60 days for tail biting problems <sup>7</sup>	As EU. Docking is widespread. <sup>8</sup>	Outdoor no, indoor no but can apply for permission to dock to 6cm for 1 yr if they have tail biting (in 2010, 30% did). Must take other steps to reduce tail biting to prove docking is a last resort <sup>9</sup>	No <sup>3</sup>	No <sup>10</sup>	No <sup>15</sup>	Only by a vet using anaesthetic and longlasting analgesic <sup>13</sup>	No <sup>14</sup>

Table 2 (Continued)

Country	EU Directives	Denmark	Denmark	Denmark	UK	UK	UK	Sweden	Finland	Norway	Switzerland
Manipulable materials	'Pigs must have permanent access to a sufficient quantity of material to enable proper investigation and manipulation activities, such as straw, hay, wood, sawdust, mushroom compost, peat or a mixture of such, which does not compromise the health of the animals'. <sup>5</sup>	Denmark wide: Material must be of 'natural origin' and be used 'for rooting' and provided 'on the floor' <sup>1</sup>	Straw bedding – all pigs able to lie on straw. <sup>6</sup>	Straw-bedded indoor lying area, outdoor run can be concrete <sup>7</sup>	As EU: Chains alone not enough, tyres not allowed, objects not fouled and within reach of pigs <sup>8</sup>	Lying area and must have comfortable absorbent bedding (straw, sawdust, shredded paper). Permanent access to materials (straw, peat, silages, mushroom compost) in sufficient quantities to allow and encourage proper expression of rooting, pawing and chewing behaviours. <sup>9</sup>	Mainly outdoor, with soil, stones, green plants. If indoor, 'ample' bedding (straw, sawdust, sand, paper or natural materials such as bracken or rushes, not peat) <sup>3</sup>	Straw must be provided for all pigs <sup>10</sup>	Permanent and enough to make into small piles, or if not permanent, materials that can be re-shaped, replenished twice daily (typically straw, sawdust, wood shavings or peat are used), plus additional materials – ball, chain or sticks. <sup>16</sup>	As EU, but 'wood (chips)' in the list of materials rather than wood <sup>13</sup>	Solid floor bedded with sawdust, straw rack provided <sup>14</sup>
Abattoir scoring to estimate tail biting prevalence		0.5% to 1.5% <sup>17</sup>	–	1.0% to 4.0% <sup>17</sup>	1.0% 'severe tail lesions' 2.4% 'evidence of tail biting' <sup>8</sup>	–	–	<2.0% tail damage <sup>18</sup>	1.8% tail damage, 5.1% partial condemnations <sup>19</sup>	4.0% tail damage <sup>20</sup>	

<sup>1</sup>Ban on fully slatted floors applies from July 2000 for new buildings, and for all housing by July 2015. Drained floor defined as maximum 10% openings (Danish Government, 2000, 2003a and 2003b; DVFA, 2013).

<sup>2</sup>Outdoor access (can be concrete): 0.6 m<sup>2</sup>/pig at 40 kg and 1.0 m<sup>2</sup>/pig at 100 kg.

<sup>3</sup>Soil Association (2012), outdoor-based system, giving permanent access to soil and growing plant foods. Must provide summer wallows and/or shade. Rotational grazing required. Indoors only under exceptional circumstances and must have outside run allowing rooting and dunging.

<sup>4</sup>Farm sizes were calculated from Eurostat (2013) figures for 2010 for 'other pigs' which includes grower/finisher pigs (available only at country level). Low EU average is due to the inclusion of many member states which do not have a major pig industry.

<sup>5</sup>EU Directives 2001/88/EC and 2001/93/EC (The Council of The European Union, 2001a and 2001b).

<sup>6</sup>Antonius: Danish Crown (2007).

<sup>7</sup>Outdoor: Friland (2012), Ministeriet for Fødevarer, Landbrug og Fiskeri (MFLF, 2012).

<sup>8</sup>Defra (2003), BPEX (2010) 1% figure from Northern Ireland and Republic of Ireland (Harley *et al.*, 2012), 2.4% for six abattoirs in England (Hunter *et al.*, 2001).

<sup>9</sup>RSPCA (2012) and Kate Parkes (RSPCA, personal communication).

<sup>10</sup>Jordbruksverket (2010), Mul *et al.* (2010), SLU and LRF (2009).

<sup>11</sup>Council of State (2012). The regulations came into effect in the first of January 2013. If a facility was already operating at that time, the space allowance regulations come into effect on the first of January 2018, and the minimum solid floor area on the first of January 2028, or both come into effect upon renovation if that is sooner.

<sup>12</sup>MAF (1997). Applies to all facilities until the 31 December 2012 and to old facilities not renovated before 2018.

<sup>13</sup>LMD (2003).

<sup>14</sup>Swiss Federal Council (2008), Wechsler (2013), CIWF (2009).

<sup>15</sup>Tail docking prohibited since January 2003 (Council of State, 2002).

<sup>16</sup>Evira (2013).

<sup>17</sup>Taken from figure j, p. 86 in Forkman *et al.* (2010) 2008–2010 figures from one abattoir so are directly comparable between systems.

<sup>18</sup>Holmgren and Lundeheim (2004), Keeling *et al.* (2012).

<sup>19</sup>Partanen *et al.* (2012).

<sup>20</sup>Fjetland and Kjastad (2002).

are allowed to use tail docking to protect the welfare of potential victim pigs.

Finally, Table 2 gives figures on tail biting prevalence in the different countries and schemes, estimated from abattoirs. It is very difficult to compare the figures, since they are not collected in a standardised way, for example pigs with missing tails are usually not counted as injured (EFSA, 2007; Keeling *et al.*, 2012). It is also difficult to compare between docked and undocked pigs; if the number of lesions are counted, long tails provide a greater area for biting than docked or part-docked tails (Webster and Day, 1998). To avoid this problem, it might be best to use data on partial carcass condemnations (PCC), which can be used as indicators of the most severe cases of tail biting (Valros *et al.*, 2004; Kritas and Morrison, 2007), but these are not always available. Occasionally, different housing systems are assessed in the same way during the same period at one abattoir. Data from a Danish study (Forkman *et al.*, 2010) show that tail biting damage at slaughter was higher in intact-tailed pigs in organic (or outdoor) systems (range 1.0% to 4.0%), in comparison to docked pigs in conventional indoor housing (range 0.5% to 1.5%; Table 2). These data suggest that tail docking is more effective at reducing tail biting than the combined effect of various improvements to the environment (as reported by Hunter *et al.*, 2001). It also highlights how challenging it is for producers to rear intact-tailed pigs, even in improved environments. However, it would be better to have data comparing countries and systems which used a standard scoring method and included PCC.

### **Risk factors for tail biting which are characteristics of the pig and how to manage them**

Characteristics of the pigs themselves can affect their propensity to tail bite or to be bitten. This could involve any step of the tail biting process (Figure 1). To take the 'two-stage' form of tail biting for example pigs could vary in: (i) how much they perform exploratory behaviour (e.g. Zonderland *et al.*, 2011), (ii) whether they explore tails rather than other things, (iii) whether tail manipulation becomes biting, (iv) whether one bite becomes many, and (v) in the likelihood of learning tail biting from a pen mate. Also, differences between pigs in their propensity to tail bite or be bitten could occur due to a greater sensitivity to any environmental risk factor (Figure 1).

#### *Characteristics of victim pigs: sex and breed*

Certain pigs may be more likely to become victims of tail biting. In several studies, castrated males were more likely to be the victims of tail biting than females (Kritas and Morrison, 2004 and 2007; tail damage, Wallgren and Lindahl, 1996), with the risk to males rising with the proportion of females in the pen (Kritas and Morrison, 2004). Tail biting was higher in pigs grouped by sex (intact males and females) in one abattoir study (Hunter *et al.*, 2001), but no relationship was found in an on-farm study of castrates and females (Steinmetz and Pedersen, 2009). All-female

pens have been reported as having more (Zonderland *et al.*, 2010a) tail damage than pens of entire males. In another study, all female pens had less (Steinmetz and Pedersen, 2009) tail damage than pens of all-castrated males. It is not clear what causes these different findings in relation to sex.

Breed may affect the likelihood of being bitten. In a Swedish pedigree population of male pigs with low levels of tail biting, bitten pigs were: Yorkshire (3.5%), Landrace (1.8%) and Hampshire (0.1%, Westin, 2003). However this was based on small numbers (63 victims out of 3049) in mixed breed groups which were not composed in a systematic way. Thus the evidence for breed differences in the risk of being bitten is weak.

#### *Characteristics of tail biters: growth retardation, early experience, breed*

It has been suggested that tail biters are often the smaller pigs in a group (Sambraus, 1985) which is in agreement with data from some studies (Zonderland *et al.*, 2011) although others have found that tail biters were no more likely to be smaller than average (Breuer *et al.*, 2005). Once tail biting begins, certain individual pigs show much higher levels of biting than others (Beattie *et al.*, 2005; Van de Weerd *et al.*, 2005), and have been characterised as 'fanatical' or 'obsessive' biters (Taylor *et al.*, 2010). In one study with small numbers of 'obsessive' biters, these pigs were smaller than average (Van de Weerd *et al.*, 2005). Beattie *et al.* (2005) found that 'tail in mouth' behaviour was higher in pigs which grew poorly in the first 3 weeks after weaning. Although poorly supported by evidence from studies which include tail damage, it remains a popular theory with farmers, that smaller pigs in a pen begin biting perhaps because they resort to this biting as an aggressive tactic when excluded from food (Schröder-Petersen and Simonsen, 2001) and/or because of a problem with nutrition or metabolism (Edwards, 2006; EFSA, 2007).

EFSA (2007) concluded that the rearing environment is not as important as the current environment for tail biting risk. Where pigs receive manipulable materials during the grower stage, past housing experience makes little difference to tail manipulation behaviours (Day *et al.*, 2002; no tail damage reported, Simonsen, 1995; Statham *et al.*, 2011). However, a greater risk of tail lesions caused by tail biting occurs in pigs which have experienced manipulable materials in the farrowing pen early in life, but which are then absent during later stages (Ruiterkamp, 1985; Munsterhjelm *et al.*, 2009). In contrast to the small effects seen in experimental studies, epidemiological studies have found associations between early life factors and tail biting. These factors include slatted floors (Smulders *et al.*, 2008) or absence of substrates (Moinard *et al.*, 2003) in the farrowing pen, and limited feeder space or high temperatures in the nursery (Smulders *et al.*, 2008). Epidemiological studies of course do not prove a causal link and further research is required.

As well as breed differences in the propensity to become victims of tail biting (described above), some have reported breed differences in the propensity to perform tail biting.

In the Swedish pedigree population with mixed breed pens of male pigs described earlier, the biters were: Landrace (1.7%), Yorkshire (0.64%) and Hampshires (0.1%), but only 27 biters out of 3049 animals were observed (Westin, 2003). In a UK study, there was no effect of breed on tail biting, although there were breed differences in ear biting (Duroc > Large White > Landrace; Breuer *et al.*, 2003). Two other studies reported finding no breed differences in the performance of tail biting (Lund and Simonsen, 2000; Guy *et al.*, 2002). Thus, as with breed differences in being tail bitten, breed differences in biting might occur, but the evidence is fairly weak.

#### *Genetics of tail biting: biters, victims and unaffected pigs*

A single published quantitative genetic study exists which found that biting other pigs' tails was a heritable trait (Breuer *et al.*, 2005), at least in Landrace (but not in Large White) pigs. Heritability was low at  $0.05 \pm 0.02$ , although tail biters were rare (295 tail biters in a population of 9018 pigs) and tail biting was treated as a binary trait, which reduces the power of genetic analysis.

Commercial pig breeding mainly focuses on economically important traits of lean growth rate, food conversion and reproductive traits such as litter size. Some pig breeding companies are considering broadening their breeding goals, and traits relating to behaviour and welfare issues such as tail biting are of interest (Merks *et al.*, 2012; Canario *et al.*, 2013). The inclusion of additional traits in a breeding index inevitably leads to a reduced rate of genetic progress in other traits (Falconer and Mackay, 1996). However, breeding companies normally use economic weightings in breeding indices, and the considerable costs of tail biting could make it economically optimal to include a trait linked to lowered levels of tail biting in a multi-trait index (Lawrence *et al.*, 2004).

A number of other factors stand in the way of conventional genetic selection against tail biting (see discussion in Turner, 2011). The positive genetic correlation relationship between tail biting and lean tissue growth rate found by Breuer *et al.* (2005) could slow genetic progress if found in other populations. Phenotyping is also a challenge: identification of 'biting' pigs is considerably more difficult than identification of victims. Direct observation of biting may be the most accurate method, but is time consuming, especially since tail biting, as described above, often occurs in sporadic, unpredictable outbreaks. Also, it may be important to identify the individual pig which starts the outbreak (first biter), as once bloody tails appear in the pen, other pigs are more likely to begin biting. Because of these difficulties, there would be enormous value in identification of a proxy trait, associated with tail biting. Unfortunately, tests based on artificial tails have proved largely disappointing in terms of their predictive value for real tail biting (Breuer *et al.*, 2003; Beattie *et al.*, 2005; Statham, 2008), although in one study the time spent manipulating an enrichment device before tail biting began was higher in biters than victims (Zonderland *et al.*, 2011). Automated detection of tail biting might be possible using similar methods to those proposed for detection of early-warning signs (see next section).

There is some prospect of identifying molecular genetic markers of pigs at lower risk of tail biting, an approach which reduces the amount of phenotyping required. Single nucleotide polymorphism markers of biting and victim pigs (in contrast to non-biting controls from the same pen) have been identified (Wilson *et al.*, 2012). Brain gene expression studies also suggest that biters and victims have more in common than unaffected pigs from the same group (Brunberg *et al.*, 2013a) or a different group (Brunberg *et al.*, 2013b). These authors suggest that unaffected pigs may show a 'tail biting resistant' phenotype. If confirmed in other populations, this idea suggests that selection against both biters and victims and for unaffected pigs might be possible.

Another approach which side-steps the problem of phenotyping tail biters is to use 'associative genetic effects' (Turner, 2011; Camerlink *et al.*, 2012). Quantitative genetic models for pig growth can be modified allowing pigs to have heritable influences on the growth of their pen-mates (Bijma *et al.*, 2007a and 2007b; Rodenburg *et al.*, 2010). Depending on the context, these 'social breeding values' might reflect differences in positive social behaviours such as social nosing (Camerlink *et al.*, 2012) or in negative behaviours such as aggression, food competition, disease transmission and ear, flank or tail biting. To have a more direct effect on tail biting, modelling of associative genetic effects could be used in combination with phenotyping for tail damage. With sufficient representation of different sires across pens, pigs with a high genetic propensity to cause tail damage to pen mates could be identified, without the need to observe biting behaviour.

Selection to reduce tail biting behaviour could raise ethical concerns, particularly concerning 'naturalness' (D'Eath *et al.*, 2010). Because 'two-stage' tail biting results from frustrated foraging behaviour, we could speculate that selection for lower tail biting might also reduce foraging. Also, selecting animals to function well in poor environments, rather than improving the animals' environment to satisfy their needs might seem distasteful to some, and could lead to a decline in housing standards (Kanis *et al.*, 2004). However, given that pigs are already undergoing constant genetic change to alter production traits, alongside improvements to the housing environment we should perhaps consider whether genetic selection to reduce tail biting could be part of a solution which makes an end to tail docking possible (D'Eath *et al.*, 2014).

Although this is speculative, breeding to reduce tail length might be possible as tail length is heritable in various mammalian species (rodents, Barnett, 1965; cats, Howell and Siegel, 1966; sheep, Branford Oltenacu and Boylan, 1974), and naturally short-tailed pigs might be less prone to becoming victims of tail biting. There are probably difficulties though as tail length is likely to be genetically correlated with back length (a desired trait in bacon pigs) and tail-less mutations may have undesirable side-effects such as those seen in Manx cats (Howell and Siegel, 1966). Even if breeding to reduce tail length were successful, other concerns remain. The curly tail could be seen by consumers as an

essential pig characteristic (although it is absent in wild pig species) and may have a function in communication (Kiley-Worthington, 1976). Finally, breeding rather than docking to shorten tails still side steps the problem that pigs need an outlet for their foraging behaviour.

### Early detection and targeted prevention

An alternative approach to the problem of tail biting is to detect outbreaks before or as soon as they begin, and to carry out targeted intervention (such as those discussed in the next section) to ameliorate or even prevent an outbreak (FAWC, 2011). Regardless of the system, if pig producers could identify certain 'at risk' individuals, groups, or batches, and target them for preventive intervention, this would be cheaper and more practical compared with making changes for every pig.

Early detection of tail biting might be possible by identifying changes in pig behaviour that precede an outbreak. Four main types of early warning sign have been described, which appear in the days or weeks before an outbreak begins (first bloody tails): (1) General activity (restlessness) increases (Statham *et al.*, 2009; Zonderland *et al.*, 2011), particularly in biters (Svendsen *et al.*, 2006). (2) Non-damaging 'tail in mouth' behaviour increases (Fraser, 1987a; Feddes and Fraser, 1994; Schröder-Petersen and Simonsen, 2001). (3) Tails are held down or 'tucked under' (Statham *et al.*, 2009; Zonderland *et al.*, 2009 and 2010). (4) Feeding patterns might change. In one study, feeder visits tended ( $P < 0.1$ ) to be lower in groups which went on to tail bite 6 to 9 weeks pre-outbreak, and tended ( $P < 0.1$ ) to increase during weeks 2 to 5 pre-outbreak in pigs which would become tail biting victims (Wallenbeck and Keeling, 2013). More research on feeding patterns is needed.

Increased observation of pigs by staff might identify these early warning signs, but staff time has a cost, so automatic detection (precision livestock farming) would be attractive (reviewed by Rushen *et al.*, 2012). The detection of specific behaviours such as tail posture and tail in mouth behaviour may be possible (Sonoda *et al.*, 2013), but increases in activity (and perhaps feeding patterns) are perhaps the easiest of the 'early warning signs' to detect automatically (Costa *et al.*, 2013). One promising approach is 'optical flow', which estimates animal activity by quantifying overall pixel changes from moment to moment in a video image. 'Optical flow' has been used to detect the reduced activity of lame broiler chickens (Dawkins *et al.*, 2009), and the disturbance of behaviour in laying hen flocks when feather pecking is occurring (Lee *et al.*, 2011).

The use of on-board animal devices (such as electronic ID ear tags), combined with detectors in the pen to record pig location also has potential to detect changes in activity (or feeding patterns). Currently, the infrastructure and consumable costs associated with either video or electronic ID approaches to monitoring pig behaviour may be prohibitive. But with falling costs and various other benefits of electronic ID (easier record keeping for medicines and at weighing, or

even for detecting when a pig has not visited the feeder in a long while) and video (estimation of pig size average and variability) the use of these technologies could become more widespread in future. Detection of changes in pig vocalisations is also a plausible approach (Manteuffel *et al.*, 2004) but would require considerable further research and validation.

### Reacting to outbreaks

Once a tail biting outbreak occurs, pig producers react in various ways (Arey, 1991). Hunter *et al.* (2001) surveyed British pig producers and found that 67% removed the bitten pig(s), 51% added enrichment objects, 25% applied sprays or tar to injured tails, 16% added straw, 6% reduced stocking density and 6% gave antibiotics. In one study, moving pigs that were already biting to pens with substrates and more space resulted in reduced biting behaviour (De Greef *et al.*, 2011), and Edwards (2011) has suggested that the effectiveness of salt or other nutritional supplements should be investigated. There is considerable scope for more research in this area: only one scientific study of the effectiveness of interventions has been reported. Zonderland *et al.* (2008) compared the interventions of removing the biter or adding straw as soon as tail damage was detected and found them to be equally effective (the 'no intervention' control was considered unethical).

Where pigs are removed from a tail biting group, bitten pigs are easier to identify (Hunter *et al.*, 2001), but removing tail biters might have a greater impact. Since biting spreads rapidly to other pigs, the time window is small for removing the first biter. Even if biters are removed, leaving bitten pigs in the pen might encourage new biters (because they are attracted to the bloody or scabbed tails), so it has been suggested that removal of *both* biters *and* bitten pigs might be optimal (Zonderland *et al.*, 2008; Boyle and Lemos Teixeira, 2010).

The other difficulty with removing pigs is the question of how to manage the pigs that are removed (Boyle and Lemos Teixeira, 2010). In an outbreak where multiple pigs are removed, the farmer may be constrained by space to group house them. They must also decide whether to use a lower stocking density and/or substrates or other forms of enrichment for those removed pigs. There is obviously a concern that removing biters and putting them into new groups could result in further tail biting, although anecdotally, Zonderland *et al.* (2008) reported that they did not experience this problem. Removing pigs to different groups and/or returning them needs to be carefully managed as it can result in social aggression including fighting and bullying (Marchant-Forde and Marchant-Forde, 2005).

Another intervention worthy of further research is the application of aversive substances to tails. Bracke (2009) found that when pigs were offered untreated ropes, or ropes treated with Dippel's oil or Stockholm tar to chew on, they avoided the treated ropes, suggesting that these treatments might be aversive when applied to tails. On the other hand, a concern over adding substances (including antibiotic sprays)

to tails is that it might make them more novel, stimulating investigation and perhaps biting.

As suggested by Edwards (2011), there is clearly an urgent need for systematic research into the effectiveness of different methods for reacting to outbreaks. This research should: (i) investigate the different methods separately or in combination and develop others, (ii) investigate the quantity/frequency of enrichment that is necessary to reduce tail biting (iii) investigate the optimal timing of interventions (there may be a point after which certain methods cease to be effective), (iv) investigate whether it is most effective and efficient to target individuals, pens, or a whole room of pigs.

## Conclusions

The risk factors affecting tail damage caused by tail biting were reviewed. A number of risk factors that have been proposed and reviewed elsewhere (Schröder-Petersen and Simonsen, 2001; EFSA, 2007) are not currently well supported by experimental studies where damaging tail biting was the end point. These include group size, nutrition, disease incidence and pig breed. Surprisingly, the evidence for an effect of stocking density was also quite weak. Epidemiological evidence alone suggests that temperature and season might be important. The evidence was strongest for the provision of manipulable substrates, and an effect of feeder space was also found.

Housing systems using slatted floors and liquid slurry handling are in widespread use due to their economic advantages, but limit the amount of loose manipulable substrates that can be used. A crucial question for this review was whether, at commercial stocking densities, in part solid-, part slatted-floored pens, small quantities of straw or similar manipulable substrate (perhaps delivered via a rack), can reduce tail biting to the point where tail docking is no longer necessary. Very few studies have looked at this, but the few that have were promising. Damaging tail biting was greatly reduced in two studies with undocked pigs using light straw (10 g twice a day per pig, Zonderland *et al.*, 2008) or light chopped straw and wood shavings (12.5 g a day per pig, Munsterhjelm *et al.*, 2009), and the experience of Finland which uses small quantities of enrichment material is also positive. Further studies investigating the effect of quantity and type of enrichment material on tail biting risk are necessary, and such studies are especially valuable if treatments are compared to a negative control of very little enrichment and a positive control of a plentiful loose material. In particular, further studies of destructible hanging materials such as ropes and destructible fresh wood would be useful. As well as controlled scientific studies, investigations into the experiences of producers in assurance schemes which are working to phase out tail docking would also be worthwhile. As a way of reducing the cost of using enrichment materials, the possibility of using materials which could combine with pig slurry as fuel for AD is interesting but faces a number of technical hurdles.

The mechanisms by which the various proposed environmental risk factors might affect the underlying process(es) of

tail biting are largely unknown. Possible mechanisms are shown in Figure 1, but much of this is speculative (see Supplementary Material S1) and there is considerable scope for further research into whether and how these risk factors might cause or affect tail biting. Alongside optimising the environment, it may be possible to use genetic selection to reduce tail biting. The challenge of phenotyping by identifying biters (Breuer *et al.*, 2005) and especially the 'first biter' could potentially be made easier (using automatic detection or proxy measures), made into a smaller task (by identification of genetic markers) or side-stepped altogether (by the use of 'associative genetic effects' for growth, or possibly, for tail lesions). The possibility that a 'tail biting resistant' phenotype might exist is interesting (Brunberg *et al.*, 2013b), but identifying these pigs would still be challenging; they are the pigs in an affected pen which are neither biters nor victims.

Another potential area for innovation is the use of precision livestock farming methods to automatically detect the early warning signs of a tail biting outbreak at the pre-damaging stage. Various behavioural signs have been identified, including tail position, 'tail in mouth' behaviour, and increased activity, some of which might be detectable by automatic methods based on electronic tags (see <http://pigit.ku.dk> and [www.pigwise.eu](http://www.pigwise.eu)) or on video (Sonoda *et al.*, 2013). If farmers could identify when and where an outbreak of tail biting was about to begin, they could target preventative measures, which would be more economic in terms of time and materials than making changes for all pigs. A final potential area for innovation is to test the efficacy of measures to stop tail biting once it begins (or just before it begins), which has been the subject of only one scientific study (Zonderland *et al.*, 2008).

Spooler *et al.* (2011) suggested that 'an intact curly tail' can be regarded as the single most important welfare indicator in finishing pigs, since to achieve this requires a high standard of housing and management over a pig's lifetime, so it serves as an 'iceberg indicator' of welfare (FAWC, 2009) and demonstrates respect for the 'animal integrity' of the pig. Within a system type, it also indicates good management to prevent (or quickly deal with) tail biting. A current difficulty is that alternative systems with intact-tailed pigs usually suffer from higher levels of tail biting than conventional systems that tail dock (Hunter *et al.*, 2001; Table 2). This means there is an ethical question as to how we should weigh a welfare impact on many (all pigs being docked as a precaution) with a worse welfare impact for a few (victims of tail biting) (D'Eath *et al.*, 2014). Would we consider that a ban on tail docking had led to improved welfare if it increased tail damage at slaughter from 1% of pigs to 4% of pigs? The threshold for what constitutes an 'acceptably low level' of tail biting must be decided in a wider ethical debate which considers the pigs' perspective. The experience of countries with complete bans on tail docking is that farmers do learn to reduce tail biting in other ways, although the resulting economic costs of this adaptation may reduce competitiveness and participation in export markets.

In most EU countries where docking is permitted, the letter of the EU Directive (2001/93/EC) that requires provision of manipulable materials is being followed, although the pressure group Compassion in World Farming found that on the farms they visited in a number of EU countries it was not (CIWF, 2008). However, the Directive states that docking should be used as a last resort only when there is evidence of a tail biting problem and other environmental measures have been tried, and this is only being enacted in reality by a minority of producers, for example some assurance schemes (e.g. Danish Antonius and Organic, UK Freedom Food). This approach seems a logical middle road (which appears to be the 'spirit' of the EU directive), allowing the majority of pigs in these schemes to benefit by avoiding docking and having their behavioural needs met, while still allowing docking to protect the welfare of potential tail biting victims on farms with a problem. However, it would be more difficult to enforce than a complete ban on tail docking. Thus it would require considerably more detail in terms of the measures producers should take before resorting to tail docking, and these measures would most likely involve substantial changes from current practice, imposing considerable costs on producers.

### Acknowledgements

Funding from the Danish Pig Research Centre (Danish Agriculture & Food Council) supported this review. This funding is mainly from pig industry sources, and three of the authors of the paper also work for this organisation, resulting in a potential or perceived conflict of interest. However, the contract for this research with the independent academic partners (SRUC and University of Copenhagen) explicitly endorses that the work is at 'arms length' from commercial interests, and the first and senior author have led the process, and are satisfied that the result is an independent review of the available evidence. SRUC receives additional funding from the Rural and Environment Science and Analytical Services (RESAS) Division of the Scottish Government. Bouda Ahmadi, Emma Baxter, Sarah Ison, Karsten Klint Jensen and Kenny Rutherford contributed to valuable discussions and provided comments on drafts. Roland Weber, Linda Keeling, Kate Parkes and Lene Juul Pedersen also provided valuable information.

### Supplementary material

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/S1751731114001359>

### References

Arey DS 1991. Tail-biting in pigs. *Farm Building Progress* 20–23.

Averós X, Brossard L, Dourmad JY, De Greef KH, Edge HL, Edwards SA and Meunier-Salaün MC 2010. A meta-analysis of the combined effect of housing and environmental enrichment characteristics on the behaviour and performance of pigs. *Applied Animal Behaviour Science* 127, 73–85.

Barnett SA 1965. Genotype and environment in tail length in mice. *Experimental Physiology* 50, 417–429.

Barnikol M 1978. Observations of tail biting (cannibalism) amongst fattening hogs and breeding pigs in connection with protein and mineral nourishment. *Tieraerztliche Umschau* 33, 540.

Beattie VE, Sneddon IA, Walker N and Weatherup RN 2001. Environmental enrichment of intensive pig housing using spent mushroom compost. *Animal Science* 72, 35–42.

Beattie VE, Breuer K, O'Connell NE, Sneddon IA, Mercer JT, Rance KA, Sutcliffe MEM and Edwards SA 2005. Factors identifying pigs predisposed to tail biting. *Animal Science* 80, 307–312.

Bergersen B 2013. Extended working group meeting on the development of guidelines concerning Council Directive 2008/120/EC on the protection of pigs. European Commission Directorate – General for Health and Consumers (DG-Sanco), Brussels, Belgium.

Bijma P, Muir WA and van Arendonk JAM 2007a. Multilevel selection 1: quantitative genetics of inheritance and response to selection. *Genetics* 175, 277–288.

Bijma P, Muir WM, Ellen ED, Wolf JB and van Arendonk JAM 2007b. Multilevel selection 2: estimating the genetic parameters determining inheritance and response to selection. *Genetics* 175, 289–299.

Blackshaw JK 1981. Some behavioural deviations in weaned domestic pigs: persistent inguinal nose thrusting, and tail and ear biting. *Animal Production* 33, 325–332.

Bornett HLI, Guy JH and Cain PJ 2003. Impact of animal welfare on costs and viability of pig production in the UK. *Journal of Agricultural & Environmental Ethics* 16, 163–186.

Boyle L and Lemos Teixeira D 2010. Improving pig welfare will reduce carcass losses. *Teagasc Pig Farmers Conferences*. Moorepark, Ireland, pp. 47–56. Retrieved October 21, 2013, from [http://www.teagasc.ie/publications/view\\_publication.aspx?PublicationID=48](http://www.teagasc.ie/publications/view_publication.aspx?PublicationID=48)

BPEX 2010. Environmental enrichment for pigs. Retrieved October 21, 2013, from <http://www.bpex.org.uk/articles/301028>

Bracke MBM 2009. Rope test may indicate efficacy of tail-biting treatments in growing pigs. *Animal Welfare* 18, 263–266.

Bracke MBM, Hulsege B, Keeling L and Blokhuis HJ 2004. Decision support system with semantic model to assess the risk of tail biting in pigs. 2. 'Validation'. *Applied Animal Behaviour Science* 87, 45–54.

Bracke MBM, Zonderland JJ, Lenskens P, Schouten WGP, Vermeer H, Spooler HAM, Hendriks HJM and Hopster H 2006. Formalised review of environmental enrichment for pigs in relation to political decision making. *Applied Animal Behaviour Science* 98, 165–182.

Branford Oltenacu EA and Boylan WJ 1974. Inheritance of tail length in crossbred Finnsheep. *Journal of Heredity* 65, 331–334.

Breuer K, Sutcliffe MEM, Mercer JT, Rance KA, Beattie VE, Sneddon IA and Edwards SA 2003. The effect of breed on the development of adverse social behaviours in pigs. *Applied Animal Behaviour Science* 84, 59–74.

Breuer K, Sutcliffe MEM, Mercer JT, Rance KA, O'Connell NE, Sneddon IA and Edwards SA 2005. Heritability of clinical tail-biting and its relation to performance traits. *Livestock Production Science* 93, 87–94.

Brooks PH 2005. Effect of diet on the behaviour and welfare of pigs. *Proceedings of the Manitoba Swine Seminar*. Retrieved October 21, 2013, from <http://www.prairieswine.com/effect-of-diet-on-the-behaviour-and-welfare-of-pigs/>

Brunberg E, Wallenbeck A and Keeling LJ 2011. Tail biting in fattening pigs: associations between frequency of tail biting and other abnormal behaviours. *Applied Animal Behaviour Science* 133, 18–25.

Brunberg E, Jensen P, Isaksson A and Keeling LJ 2013a. Brain gene expression differences are associated with abnormal tail biting behavior in pigs. *Genes, Brain, and Behavior* 12, 275–281.

Brunberg E, Jensen P, Isaksson A and Keeling LJ 2013b. Behavioural and brain gene expression profiling in pigs during tail biting outbreaks – evidence of a tail biting resistant phenotype. *PLoS One* 8, e66513.

Busch ME, Wachmann H, Nielsen EO and Petersen HH 2004. Halebid – Kan slagtedata anvendes til at identificere besætninger med høj forekomst? (Tail biting – can slaughter data be used to identify herds with a high prevalence?). Danish Pig Research Centre (PRC), Copenhagen, Denmark. Report: Meddelelse 643. Retrieved October 21, 2013, from [http://vsp.lf.dk/Publikationer/Kilder/lu\\_medd/2004/643.aspx?full=1](http://vsp.lf.dk/Publikationer/Kilder/lu_medd/2004/643.aspx?full=1)

Camerlink I, Bijma P, Kemp B and Bolhuis JE 2012. Relationship between growth rate and oral manipulation, social nosing, and aggression in finishing pigs. *Applied Animal Behaviour Science* 142, 11–17.

Canario L, Mignon-Grasteau S, Dupont-Nivet M and Phocas F 2013. Genetics of behavioural adaptation of livestock to farming conditions. *Animal* 7, 357–377.

- CIWF 2008. The state of Europe's pigs: An exposé. Compassion in world farming, investigative report. Compassion in World Farming, Godalming, Surrey, UK.
- CIWF 2009. Welfare of pigs in the European Union – the urgent need for reform of existing legislation and effective enforcement. Compassion in World Farming, Godalming, Surrey, UK.
- Costa A, Ismayilova G, Borgonovo F, Leroy T, Berckmans D and Guarino M 2013. The use of image analysis as a new approach to assess behaviour classification in a pig barn. *Acta Veterinaria Brno* 82, 25–30.
- Council of State 2002. Valtioneuvoston asetus eläinsuojeluasetuksen muuttamisesta. VNa 910/2002 (Government decree on the regulation of the protection of animals). Retrieved October 21, 2013, from <http://www.finlex.fi/fi/laki/alkup/2002/20020910>
- Council of State 2012. Valtioneuvoston asetus sikojen suojelusta. VNa 629/2012 (Government decree on the protection of pigs). Retrieved October 21, 2013, from <http://www.finlex.fi/fi/laki/alkup/2012/20120629>
- Courboulay V, Eugene A and Delarue E 2009. Welfare assessment in 82 pig farms: effect of animal age and floor type on behaviour and injuries in fattening pigs. *Animal Welfare* 18, 515–521.
- Courboulay V, Souffran H, Guingand N, Massabie P and Dubois A 2008. Intérêt de la brumisation en engraissement pour améliorer le confort des porcs en croissance (Fogging system: a way to improve the welfare of growing-finishing pigs during summer periods). *Journées Recherche Porcine* 40, 243–246.
- D'Eath RB, Conington J, Lawrence AB, Olsson IAS and Sandøe P 2010. Breeding for behavioural change in farm animals: practical, economic and ethical considerations. *Animal Welfare* 19, 17–27.
- D'Eath RB, Niemi J, Ahmadi B, Arnott G, Turner SP, Baxter EM, Rutherford KMD, Jensen T, Lahrmann HP, Busch ME, Jensen KK, Lawrence AB and Sandøe P 2014. Tail docking vs. tail biting – a scientific, legal, socio-economic and ethical analysis. *Animal Welfare* (submitted).
- Danish Crown 2007. Kvalitetshåndbog for Antonius grisproduktion (Quality handbook for Antonius pig production), Copenhagen, Denmark. Retrieved October 21, 2013, from <http://www.danishcrown.dk/lib/file.aspx?fileID=286>
- Danish Government 2000. Lov no. 104 af 14 Februar 2000 om indendørs hold af smågrise, avls- og slagtesvin (Act on indoor keeping of pigs, breeding and fattening), Copenhagen, Denmark.
- Danish Government 2003a. BEK nr 323 af 06/05/2003. Bekendtgørelse om beskyttelse af svin (Order on the protection of pigs) Fødevareministeriet (Danish Ministry of Food), Copenhagen, Denmark. Retrieved October 21, 2013, from <https://www.retsinformation.dk/Forms/R0710.aspx?id=1571>
- Danish Government 2003b. BEK nr 324 af 06/05/2003, Bekendtgørelse om halekupering og kastration af dyr (Order on tail docking and castration of animals) Fødevareministeriet (Ministry of Food), Copenhagen, Denmark. Retrieved October 21, 2013, from <https://www.retsinformation.dk/Forms/R0710.aspx?id=1572>
- Dawkins MS, Lee HJ, Waitt CD and Roberts SJ 2009. Optical flow patterns in broiler chicken flocks as automated measures of behaviour and gait. *Applied Animal Behaviour Science* 119, 203–209.
- Day JEL, De Weerd HAV and Edwards SA 2008. The effect of varying lengths of straw bedding on the behaviour of growing pigs. *Applied Animal Behaviour Science* 109, 249–260.
- Day JEL, Burfoot A, Docking CM, Whittaker X, Spoolder HAM and Edwards SA 2002. The effects of prior experience of straw and the level of straw provision on the behaviour of growing pigs. *Applied Animal Behaviour Science* 76, 189–202.
- De Greef KH, Vermeer HM, Houwers HWJ and Bos AP 2011. Proof of principle of the comfort class concept in pigs. Experimenting in the midst of a stakeholder process on pig welfare. *Livestock Science* 139, 172–185.
- Defra 2003. Code of recommendations for the welfare of livestock: pigs. Defra Publications, London.
- DVFA 2013. Fødevarestyrelsens vejledning om beskæftigelses- og rodematerialer (DVFA guidance on employment and enrichment materials) Danish Veterinary and Food Administration (Fødevarestyrelsen), Copenhagen, Denmark. Retrieved October 21, 2013, from <http://www.foedevarestyrelsen.dk/Leksikon/Sider/Hold-af-svin.aspx>
- EC 2009. Answer from the European Commission to a written question (E-5360/2009) from Member of the European Parliament, Caroline Lucas, Strasbourg. Retrieved October 21, 2013, from <http://www.europarl.europa.eu/sides/getAllAnswers.do?reference=E-2009-5360&language=EN>
- Edwards S 2011. What do we know about tail biting today? *The Pig Journal* 66, 81–86.
- Edwards SA 2006. Tail biting in pigs: understanding the intractable problem. *Veterinary Journal* 171, 198–199.
- EFSA 2007. The risks associated with tail biting in pigs and possible means to reduce the need for tail docking considering the different housing and husbandry systems (question no. EFSA-Q-2006-013). Annex to the EFSA Journal 611, 1–13.
- Elst WET, Vaessen MA, Vos HJMP, Binnendijk GP, Huime REM and Backus GBC 1988. Proefverslag- Varkensproefbedrijf- Zuid-enwest-Nederland 200, 1–54.
- Eurostat 2013. Livestock: number of farms and heads of animals by type of farming in 2010 European Commission. Retrieved October 21, 2013, from <http://epp.eurostat.ec.europa.eu/portal/page/portal/agriculture/data/database>
- Evira 2013. Sikojen tonkimismateriaali (Materials for pigs to root) Evira (Finnish Food Safety Authority), Helsinki, Finland. Retrieved October 21, 2013, from [http://www.evira.fi/files/attachments/fi/elaimet/elainsuojelu\\_ja\\_elainten\\_pito/sikojen\\_tonkimismateriaali.pdf](http://www.evira.fi/files/attachments/fi/elaimet/elainsuojelu_ja_elainten_pito/sikojen_tonkimismateriaali.pdf)
- Falconer DS and Mackay TFC 1996. Introduction to quantitative genetics, 4th edition. Longman, New York.
- FAWC 2009. Farm animal welfare in Great Britain: past, present and future. Farm Animal Welfare Council, London.
- FAWC 2011. Opinion on mutilations and environmental enrichment in piglets and growing pigs. Farm Animal Welfare Council, London.
- Feddes JJR and Fraser D 1994. Non-nutritive chewing by pigs: implications for tail-biting and behavioural enrichment. *Transactions of the ASAE* 37, 947–950.
- Fjetland O and Kjastad HP 2002. Tail biting in pigs. *Norsk Veterinartidsskrift* 114, 249–253.
- Forkman B, Olsen P, Nørgaard N, Laursen PB, Pedersen LJ, Holm L, Viekilde K and Søndergaard JT 2010. Arbejdsgrupperapport om hold af svin (Working group report on the keeping of pigs) Justitsministeriet Departementet, Copenhagen, Denmark. Retrieved October 21, 2013, from <https://bdkv2.borger.dk/Lovgivning/Hoeringsportalen/Sider/Fakta.aspx?hpId=2146002290>
- Fraser D 1987a. Attraction to blood as a factor in tail-biting by pigs. *Applied Animal Behaviour Science* 17, 61–86.
- Fraser D 1987b. Mineral-deficient diets and the pig's attraction to blood: implications for tail-biting. *Canadian Journal of Animal Science* 67, 909–918.
- Fraser D, Berron DE and Ball RO 1991. Enhanced attraction to blood by pigs with inadequate dietary-protein supplementation. *Canadian Journal of Animal Science* 71, 611–619.
- Friland 2012. Frilandgris Produktionskoncept – en god begyndelse (Outdoor pigs production concept – a good start), Randers, Denmark. Retrieved October 21, 2013, from <http://www.friland.dk/lib/file.aspx?fileID=636>
- Geers R, Dellaert B, Goedseels V, Hoogerbrugge A, Vranken E, Maes F and Berckmans D 1989. An assessment of optimal air temperatures in pig houses by the quantification of behavioral and health-related problems. *Animal Production* 48, 571–578.
- Georgsson L and Svendsen J 2001. One or two feeders for groups of 16 growing-finishing pigs: effects on health and production. *Acta Agriculturae Scandinavica Section A – Animal Science* 51, 257–264.
- Georgsson L and Svendsen J 2002. Degree of competition at feeding differentially affects behavior and performance of group-housed growing-finishing pigs of different relative weights. *Journal of Animal Science* 80, 376–383.
- Goldberg TL, Weigel RM, Hahn EC and Scherba G 2000. Associations between genetics, farm characteristics and clinical disease in field outbreaks of porcine reproductive and respiratory syndrome virus. *Preventive Veterinary Medicine* 43, 293–302.
- Goossens X, Sobry L, Odberg F, Tuytens F, Maes D, De Smet S, Nevens F, Opsomer G, Lommelen F and Geers R 2008. A population-based on-farm evaluation protocol for comparing the welfare of pigs between farms. *Animal Welfare* 17, 35–41.
- Guy JH, Rowlinson P, Chadwick JP and Ellis M 2002. Behaviour of two genotypes of growing/finishing pig in three different housing systems. *Applied Animal Behaviour Science* 75, 193–206.
- Hansen LL, Hagelso AM and Madsen A 1982. Behavioural results and performance of bacon pigs fed ad libitum from one or several self-feeders. *Applied Animal Ethology* 8, 307–333.
- Harley S, More S, O'Connell N, Hanlon A, Teixeira D and Boyle L 2012. Evaluating the prevalence of tail biting and carcass condemnations in slaughter pigs in the Republic and Northern Ireland, and the potential of abattoir meat inspection as a welfare surveillance tool. *Veterinary Record* 171, 621–662.

- Holm L, Jensen MB, Pedersen LJ and Ladewig J 2008. The importance of a food feedback in rooting materials for pigs measured by double demand curves with and without a common scaling factor. *Applied Animal Behaviour Science* 111, 68–84.
- Holmgren N and Lundeheim N 2004. Risk factors for tail biting. *Proceedings of the 18th IPVS Congress, Hamburg, Germany*, p. 786.
- Howell JM and Siegel PB 1966. Morphological effects of the Manx factor in cats. *Journal of Heredity* 57, 100–104.
- Hunter EJ, Jones TA, Guise HJ, Penny RHC and Hoste S 2001. The relationship between tail biting in pigs, docking procedure and other management practices. *Veterinary Journal* 161, 72–79.
- Jaeger F 2013. Das Projekt "intakter Ringelschwanz" beim Schwein – stehen wir vor dem Durchbruch? (The project "intact curly tails" in pigs – are we just before a breakthrough?). *Tieraerztliche Umschau* 68, 3–11.
- Jensen MB and Pedersen LJ 2007. The value assigned to six different rooting materials by growing pigs. *Applied Animal Behaviour Science* 108, 31–44.
- Jensen MB, Studnitz M and Pedersen LJ 2010. The effect of type of rooting material and space allowance on exploration and abnormal behaviour in growing pigs. *Applied Animal Behaviour Science* 123, 87–92.
- Jensen MB, Studnitz M, Halekoh U, Pedersen LJ and Jorgensen E 2008. Pigs' preferences for rooting materials measured in a three-choice maze-test. *Applied Animal Behaviour Science* 112, 270–283.
- Jensen T, Kold Nielsen C, Vinther J and D'Eath RB 2012. The effect of space allowance for finishing pigs on productivity and pen hygiene. *Livestock Science* 149, 33–40.
- Jensen T, Riis AL, Maribo H, Vernersen A and Busch ME 2004. Halebid manual (tail-biting manual). Videncenter for Svineproduktion og danske slagterier, Copenhagen, Denmark.
- Jordbruksverket 2010. Statens jordbruksverks föreskrifter och allmänna råd om djurhållning inom lanbruket m.m. (Swedish State Board of Agriculture Regulations and Guidelines on Animal Husbandry in Agriculture, etc.). Statens jordbruksverk, Jönköping, Sweden.
- Kanis E, van den Belt H, Groen AF, Schakel J and De Greef KH 2004. Breeding for improved welfare in pigs: a conceptual framework and its use in practice. *Animal Science* 78, 315–329.
- Keeling LJ, Wallenbeck A, Larsen A and Holmgren N 2012. Scoring tail damage in pigs: an evaluation based on recordings at Swedish slaughterhouses. *Acta Veterinaria Scandinavica* 54, 32.
- Kiley-Worthington M 1976. Tail movements of ungulates, canids and felids with particular reference to their causation and function as displays. *Behaviour* 56, 69–115.
- Krider JL, Albright JL, Plumlee MP, Conrad JH, Sinclair CL, Underwood L, Jones RG and Harrington RB 1975. Magnesium supplementation, space and docking effects on swine performance and behavior. *Journal of Animal Science* 40, 1027–1033.
- Kritas SK and Morrison RB 2004. An observational study on tail biting in commercial grower-finisher barns. *Journal of Swine Health and Production* 12, 17–22.
- Kritas SK and Morrison RB 2007. Relationships between tail biting in pigs and disease lesions and condemnations at slaughter. *Veterinary Record* 160, 149–152.
- Lahrmann HP and Steinmetz H 2011. Effekt af halm længde på forekomsten af unormal adfærd og halmrettet adfærd hos slagtesvin (Effect of straw length on the incidence of abnormal behavior and straw directed behavior of pigs). Danish Pig Research Centre (VSP), Copenhagen, Denmark.
- Lawrence AB, Conington J and Simm G 2004. Breeding and animal welfare: practical and theoretical advantages of multi-trait selection. *Animal Welfare* 13, S191–S196.
- Lee HJ, Roberts SJ, Drake KA and Dawkins MS 2011. Prediction of feather damage in laying hens using optical flows and Markov models. *Journal of the Royal Society Interface* 8, 489–499.
- LMD 2003. FOR 2003-02-18 nr 175: Forskrift om hold av svin (Regulations on the keeping of pigs) Landbruks- og matdepartementet (Norwegian Ministry of Agriculture and Food). Retrieved October 21, 2013, from <http://www.lovdata.no/cgi-wifit/ldles?doc=/sf/sf-20030218-0175.html>
- Lund A and Simonsen HB 2000. Aggression and stimulus-directed activities in two breeds of finishing pig. *The Pig Journal* 45, 123–130.
- MAF 1997. Sikojen pidolle asetettavat eläinsuojeluväimukset (Animal protection requirements on concerning the keeping of pigs), Decision 13/EE0/1997. Ministry of Agriculture and Forestry, Helsinki, Finland.
- Manteuffel G, Puppe B and Schon PC 2004. Vocalization of farm animals as a measure of welfare. *Applied Animal Behaviour Science* 88, 163–182.
- Marchant-Forde JN and Marchant-Forde RM 2005. Minimizing inter-pig aggression during mixing. *Pig News and Information* 26, 63N–73N.
- McIntyre J and Edwards SA 2002a. An investigation into the effect of tryptophan on tail chewing behaviour of growing pigs. *Proceedings of the British Society of Animal Science*, p. 34.
- McIntyre J and Edwards SA 2002b. Preference for blood and behavioural measurements of known tailbiting pigs compared to control penmates. *Proceedings of the 36th International Congress of the International Society for Applied Ethology, Egmond an Zee, the Netherlands*, p. 93.
- Merks JWM, Mathur PK and Knol EF 2012. New phenotypes for new breeding goals in pigs. *Animal* 6, 535–543.
- MFLF 2012. Vejledning om økologisk jordbrugsproduktion (Guidance on organic agricultural production) Ministeriet for Fødevarer, Landbrug og Fiskeri (Danish Ministry of Food, Agriculture and Fisheries), Copenhagen, Denmark. Retrieved October 21, 2013, from [http://1.naturerhverv.fvm.dk/oekologi\\_vejledning\\_bedrifter.aspx?ID=2137](http://1.naturerhverv.fvm.dk/oekologi_vejledning_bedrifter.aspx?ID=2137)
- Moinard C, Mendl M, Nicol CJ and Green LE 2003. A case control study of on-farm risk factors for tail biting in pigs. *Applied Animal Behaviour Science* 81, 333–355.
- Mul M, Vermeij I, Hindle V and Spooler H 2010. EU Welfare Legislation on Pigs, Report 273 Livestock Research, Wageningen UR.
- Munsterhjelm C, Brunberg E, Heinonen M, Keeling L and Valros A 2013. Stress measures in tail biters and bitten pigs in a matched case-control study. *Animal Welfare* 22, 331–338.
- Munsterhjelm C, Peltoniemi OAT, Heinonen M, Halli O, Karhapää M and Valros A 2009. Experience of moderate bedding affects behaviour of growing pigs. *Applied Animal Behaviour Science* 118, 42–53.
- Niemi JK, Sinisalo A, Valros A and Heinonen M 2011. The timing and treatment of tail biting in fattening pigs. In 24th NJF Congress "Food, Feed, Fuel and Fun – Nordic Light on Future Land Use and Rural Development", NJF Report 7 (ed. J. Hulgren, P. Persson, E. Nadeau and F. Fogelberg), pp. 50–55. NJF, Uppsala, Sweden.
- Niemi JK, Sinisalo A, Valros A and Heinonen M 2012. Hännänpurenta – syy vai seuraus? (Tail biting – cause or effect?). Schulman, N. and Kauppinen, H. Maataloustieteen Päivät 2012 [Verkkojulkaisu] (Agricultural Science Days 2012 [Online]). Suomen Maataloustieteellisen Seuran julkaisu no. 28 (The Scientific Agricultural Society of Finland publications no. 28), Helsinki, Finland (<http://www.smts.fi/mtpjul2012.html>, accessed 30th May 2014).
- Olsson L 2011. Optimizing amount of straw for growing-finishing pigs – considering time spent in manipulative behaviour. SLU, Uppsala, Sweden.
- Palander PA, Heinonen M, Simpura I, Edwards SA and Valros AE 2013. Jejunal morphology and blood metabolites in tail biting, victim and control pigs. *Animal* 7, 1523–1531.
- Partanen K, Niemi J, Voutila L and Jukola E 2012. Vastuullisen tuotannon vaatimusten täyttäminen näkyy lihasikalan tuloksessa (Responsible pig meat production). In Maataloustieteen Päivät 2012 [Verkkojulkaisu] (Agricultural Science Days 2012 [Online]) Suomen Maataloustieteellisen Seuran julkaisu no. 29 (The Scientific Agricultural Society of Finland publications no. 29), p. 275 (ed. N. Schulman and H. Kauppinen). Helsinki, Finland. <http://www.smts.fi/mtpjul2012.html>, accessed 30th May 2014.
- Pedersen LJ, Herskin M and Forkman B 2013. How much straw do slaughter pigs need to meet the need for exploration. ISAE 47th Congress, Florianopolis, Brazil, p. 46.
- Petersen V, Simonsen HB and Lawson LG 1995. The effect of environmental stimulation on the development of behaviour in pigs. *Applied Animal Behaviour Science* 45, 215–224.
- PRC 2011. Annual report, Pig Research Centre, Danish Agriculture and Food Council, Copenhagen, Denmark.
- Rodenburg TB, Bijma P, Ellen ED, Bergsma R, de Vries S, Bolhuis JE, Kemp B and van Arendonk JAM 2010. Breeding amiable animals? Improving farm animal welfare by including social effects into the genetic model. *Animal Welfare* 19, 77–82.
- RSPCA 2012. Welfare standards for pigs. Freedom Food. Retrieved 30 May 2014, from <http://www.rspca.org.uk/sciencegroup/farmanimals/standards/pigs>
- Ruiterkamp WA 1985. Het Gedrag Van Mestvarkens in Relatie Tot Huisvesting (The Behaviour of Fattening Pigs in Relation to Housing). PhD Thesis, University of Utrecht, The Netherlands.

- Rushen J, Chapinal N and de Passille A 2012. Automated monitoring of behavioural-based animal welfare indicators. *Animal Welfare* 21, 339–350.
- Sambrasus HH 1985. Mouth-based anomalous syndromes. In *World animal science, A5, ethology of farm animals. A comprehensive study of the behavioural features of common farm animals* (ed. AF Fraser), pp. 391–422. Elsevier, Amsterdam.
- Sanchez M and Gonzalez JL 2005. The fertilizer value of pig slurry. I. Values depending on the type of operation. *Bioresource Technology* 96, 1117–1123.
- Scheepens CJM, Hessing MJC, Laarakker E, Schouten WGP and Tielen MJM 1991. Influences of intermittent daily draft on the behaviour of weaned pigs. *Applied Animal Behaviour Science* 31, 69–82.
- Schmolke SA, Li YZ and Gonyou HW 2003. Effect of group size on performance of growing-finishing pigs. *Journal of Animal Science* 81, 874–878.
- Schröder-Petersen DL and Simonsen HB 2001. Tail biting in pigs. *Veterinary Journal* 162, 196–210.
- Schröder-Petersen DL, Heiskanen T and Ersboll AK 2004. Tail-in-mouth behaviour in slaughter pigs, in relation to internal factors such as: age, size, gender, and motivational background. *Acta Agriculturae Scandinavica Section A – Animal Science* 54, 159–166.
- Scollo A, Di Martino G, Bonfanti L, Stefani AL, Schiavon E, Marangon S and Gottardo F 2013. Tail docking and the rearing of heavy pigs: the role played by gender and the presence of straw in the control of tail biting. Blood parameters, behaviour and skin lesions. *Research in Veterinary Science* 95, 825–830.
- Scott K, Taylor L, Gill BP and Edwards SA 2009. Influence of different types of environmental enrichment on the behaviour of finishing pigs in two different housing systems 3. Hanging toy versus rootable toy of the same material. *Applied Animal Behaviour Science* 116, 186–190.
- Scott K, Chennells DJ, Armstrong D, Taylor L, Gill BP and Edwards SA 2007. The welfare of finishing pigs under different housing and feeding systems: liquid versus dry feeding in fully-slatted and straw-based housing. *Animal Welfare* 16, 53–62.
- Scott K, Chennells DJ, Campbell FM, Hunt B, Armstrong D, Taylor L, Gill BP and Edwards SA 2006. The welfare of finishing pigs in two contrasting housing systems: fully-slatted versus straw-bedded accommodation. *Livestock Science* 103, 104–115.
- Sihvo H, Simola O, Munsterhjelm C and Syrjä P 2012. Systemic spread of infection in tail-bitten pigs. *Journal of Comparative Pathology* 146, 73.
- Simonsen HB 1995. Effect of early rearing environment and tail docking on later behaviour and production in fattening pigs. *Acta Agriculturae Scandinavica Section A – Animal Science* 45, 139–144.
- Sinialo A, Niemi JK, Heinonen M and Valros A 2012. Tail biting and production performance in fattening pigs. *Livestock Science* 143, 220–225.
- SLU and LRF 2009. Swedish pig production (pamphlet). Swedish University of Agricultural Sciences and Swedish Pig Producers Association, Uppsala, Sweden.
- Smulders D, Hautekiet V, Verbeke G and Geerst R 2008. Tail and ear biting lesions in pigs: an epidemiological study. *Animal Welfare* 17, 61–69.
- Soil Association 2012. Organic standards: farming and growing (revision 16.7), UK. Retrieved October 21, 2013, from <http://www.soilassociation.org/organicstandards>
- Sonoda LT, Fels M, Oczak M, Vranken E, Ismayilova G, Guarino M, Viazzi S, Bahr C, Berckmans D and Hartung J 2013. Tail Biting in pigs – causes and management intervention strategies to reduce the behavioural disorder. A review. *Berliner und Münchener Tierärztliche Wochenschrift* 126, 104–112.
- Spoolder H, Bracke M, Mueller-Graf C and Edwards S 2011. Preparatory work for the future development of animal based measures for assessing the welfare of pig – Report 2. Preparatory work for the future development of animal based measures for assessing the welfare of weaned, growing and fattening pigs including aspects related to space allowance, floor types, tail biting and need for tail docking. Technical report submitted to EFSA.
- Statham P 2008. Effects of experience and individuality on tail biting in pigs. PhD thesis, University of Bristol.
- Statham P, Green L and Mendl M 2011. A longitudinal study of the effects of providing straw at different stages of life on tail-biting and other behaviour in commercially housed pigs. *Applied Animal Behaviour Science* 134, 100–108.
- Statham P, Green L, Bichard M and Mendl M 2009. Predicting tail-biting from behaviour of pigs prior to outbreaks. *Applied Animal Behaviour Science* 121, 157–164.
- Steinmetz HV and Pedersen ML 2009. Kønsvis opstaldnings betydning for forekomsten af halebid (Single sex housing – its importance for tail biting). Report Meddelelse nr. 845 Videncenter for Svineproduktion, Danish Pig Research Centre.
- Stern S, Sonesson U, Gunnarsson S, Öborn I, Kumm KI and Nybrant T 2005. Sustainable development of food production: a case study on scenarios for pig production. *AMBIO: A Journal of the Human Environment* 34, 402–407.
- Street BR and Gonyou HW 2008. Effects of housing finishing pigs in two group sizes and at two floor space allocations on production, health, behavior, and physiological variables. *Journal of Animal Science* 86, 982–991.
- Studnitz M, Jensen MB and Pedersen LJ 2007. Why do pigs root and in what will they root? A review on the exploratory behaviour of pigs in relation to environmental enrichment. *Applied Animal Behaviour Science* 107, 183–197.
- Sutherland MA and Tucker CB 2011. The long and short of it: a review of tail docking in farm animals. *Applied Animal Behaviour Science* 135, 179–191.
- Svendsen J, Olsson AC and Botermans J 2006. Data on tail biting in pigs. Proceedings of the 19th IPVS Congress, Copenhagen, Denmark, p. 613.
- Swiss Federal Council 2008. 455.1 Animal Welfare Ordinance (TSchV). Retrieved October 21, 2013, from <http://www.bvet.admin.ch/themen/tierschutz/02443/index.html?lang=de>
- Taylor NR, Main DCJ, Mendl M and Edwards SA 2010. Tail-biting: a new perspective. *The Veterinary Journal* 186, 137–147.
- Taylor NR, Parker RMA, Mendl M, Edwards SA and Main DCJ 2012. Prevalence of risk factors for tail biting on commercial farms and intervention strategies. *The Veterinary Journal* 194, 77–83.
- Telkänranta H, Bracke M and Valros A 2014a. Developing chewable and rootable low-cost objects for commercial pig farms. Proceedings of the 25th ISAE Nordic Regional Meeting, Drobak, Norway, p. 10.
- Telkänranta H, Swan K, Hirvonen H and Valros A 2014b. Chewable materials before weaning reduce tail biting in growing pigs. *Applied Animal Behaviour Science* (in press <http://dx.doi.org/10.1016/j.applanim.2014.01.004>).
- Temple D, Courboulay V, Velarde A, Dalmau A and Manteca X 2012. The welfare of growing pigs in five different production systems in France and Spain: assessment of health. *Animal Welfare* 21, 257–271.
- The Council of The European Union 2001a. Commission Directive 2001/93/EC of 9 November 2001 amending Directive 91/630/EEC laying down minimum standards for the protection of pigs. *Official Journal L316*, 01/12/2001.
- The Council of The European Union 2001b. Council Directive 2001/88/EC of 23rd October 2001 amending Directive 91/630/EEC laying down minimum standards for the protection of pigs. *Official Journal L 316*, 01/12/2001.
- Trickett SL, Guy JH and Edwards SA 2009. The role of novelty in environmental enrichment for the weaned pig. *Applied Animal Behaviour Science* 116, 45–51.
- Turner SP 2011. Breeding against harmful social behaviours in pigs and chickens: state of the art and the way forward. *Applied Animal Behaviour Science* 134, 1–9.
- Tuytens FAM 2005. The importance of straw for pig and cattle welfare: a review. *Applied Animal Behaviour Science* 92, 261–282.
- Valros A, Ahlstrom S, Rintala H, Hakkinen T and Saloniemä H 2004. The prevalence of tail damage in slaughter pigs in Finland and associations to carcass condemnations. *Acta Agriculturae Scandinavica Section A – Animal Science* 54, 213–219.
- Valros A, Palander P, Heinonen M, Brunberg E, Keeling L and Piepponen P 2013. Preliminary evidence of an altered serotonin metabolism in the prefrontal cortex of tail biting pigs. ISAE 47th Congress, Florianopolis, Brazil, p. 105.
- Van de Perre V, Driessen B, Van Thielen J, Verbeke G and Geers R 2011. Comparison of pig behaviour when given a sequence of enrichment objects or a chain continuously. *Animal Welfare* 20, 641–649.
- Van de Weerd HA and Day JEL 2009. A review of environmental enrichment for pigs housed in intensive housing systems. *Applied Animal Behaviour Science* 116, 1–20.
- Van de Weerd HA, Docking CM, Day JEL and Edwards SA 2005. The development of harmful social behaviour in pigs with intact tails and different enrichment backgrounds in two housing systems. *Animal Science* 80, 289–298.
- Van de Weerd HA, Docking CM, Day JEL, Avery PJ and Edwards SA 2003. A systematic approach towards developing environmental enrichment for pigs. *Applied Animal Behaviour Science* 84, 101–118.

- Van de Weerd HA, Docking CM, Day JEL, Breuer K and Edwards SA 2006. Effects of species-relevant environmental enrichment on the behaviour and productivity of finishing pigs. *Applied Animal Behaviour Science* 99, 230–247.
- Van Putten G 1969. An investigation into tail-biting among fattening pigs. *British Veterinary Journal* 125, 511–517.
- Verdoes N 2014. Starplus stalconcept voor varkens (Starplus Stable concept for pigs). Retrieved January 29, 2014, from <http://www.wageningenur.nl/nl/show/Starplus-stalconcept-voor-varkens.htm>
- Wallenbeck A and Keeling LJ 2013. Using data from electronic feeders on visit frequency and feed consumption to indicate tail biting outbreaks in commercial pig production. *Journal of Animal Science* 91, 2879–2884.
- Wallgren P and Lindahl E 1996. The influence of tail biting on performance of fattening pigs. *Acta Veterinaria Scandinavica* 37, 453–460.
- Webster SD and Day JEL 1998. Tail-biting and tail-docking in pigs. *The Veterinary Record* 142, 375.
- Wechsler B 2013. Experience from Switzerland on ensuring the welfare of pigs. Extended working group meeting on the development of guidelines concerning Council Directive 2008/120/EC on the protection of pigs (DG SANCO), Brussels, EU.
- Westin R 2003. Svansbitning hos gris relaterat till individuell tillväxt och ras (Tail biting in pigs related to individual growth and breed), Department of Animal Environment and Health, Sveriges Lantbruksuniversitet (SLU), Veterinärmedicinska fakulteten, Veterinärprogrammet.
- Wilson K, Zanella R, Ventura C, Johansen HL, Framstad T, Janczak A, Zanella AJ and Neibergs HL 2012. Identification of chromosomal locations associated with tail biting and being a victim of tail-biting behaviour in the domestic pig (*Sus scrofa domestica*). *Journal of Applied Genetics* 53, 449–456.
- Zonderland JJ 2010. Talking tails: quantifying the development of tail biting in pigs. PhD, Wageningen UR.
- Zonderland JJ, Bracke MBM, den Hartog LA, Kemp B and Spoolder HAM 2010a. Gender effects on tail damage development in single- or mixed-sex groups of weaned piglets. *Livestock Science* 129, 151–158.
- Zonderland JJ, Kemp B, Bracke MBM, den Hartog LA and Spoolder HAM 2010b. Individual piglets' contribution to the development of tail biting. *Animal* 5, 601–607.
- Zonderland JJ, van Riel JW, Bracke MBM, Kemp B, den Hartog LA and Spoolder HAM 2009. Tail posture predicts tail damage among weaned piglets. *Applied Animal Behaviour Science* 121, 165–170.
- Zonderland JJ, Schepers F, Bracke MBM, den Hartog LA, Kemp B and Spoolder HAM 2011. Characteristics of biter and victim piglets apparent before a tail-biting outbreak. *Animal* 5, 767–775.
- Zonderland JJ, Wolthuis-Fillerup M, Van Reenen CG, Bracke MBM, Kemp B, den Hartog LA and Spoolder HAM 2008. Prevention and treatment of tail biting in weaned piglets. *Applied Animal Behaviour Science* 110, 269–281.
- Zwicker B, Gygax L, Wechsler B and Weber R 2012. Influence of the accessibility of straw in racks on exploratory behaviour in finishing pigs. *Livestock Science* 148, 67–73.
- Zwicker B, Gygax L, Wechsler B and Weber R 2013. Short- and long-term effects of eight enrichment materials on the behaviour of finishing pigs fed ad libitum or restrictively. *Applied Animal Behaviour Science* 144, 31–38.