Periscope

DIPLOPIA WITHOUT EXTRA-OCULAR PALSIES, CAUSED BY HETERONYMOUS DEFECTS IN THE VISUAL FIELDS ASSOCIATED WITH DEFECTIVE MACULAR VISION.

True diplopia may occur without any history or physical sign of strabismus, ptosis, or pupillary disturbances. Six cases of this type are described here in which a true diplopia was present without any weakness of the extra-ocular muscles, and in which the lesion was an adenoma of the hypophysis; and two cases where the condition was found in tumour of the hypophyseal stalk.

Such a diplopia is often most troublesome when the patient attempts to fix both eyes on an object for precise observation, and it is not influenced by conjugate deviation of the eyes in any direction, whereas diplopia which results from weakness of any of the muscles of the eye is always altered in some way by conjugate deviation of the eyes. This type of diplopia has not been found in any patients with homonymous defects in the visual fields, but only in the presence of bitemporal hemianopia and in association with periods of amblyopia. Where the amblyopia was transient, the diplopia was transient; and where the amblyopia was permanent, the diplopia persisted. This diplopia depends on the fact that in the presence of impaired macular vision the patient attempts to use the adjacent areas of the retina in which visual acuity is greater. In the presence of a coexistent bitemporal hemianopia the only available fields are the two nasal fields. To secure binocular vision of good acuity he must employ these. Since, however, they are not homologous parts of the retina, diplopia inevitably results. In order to bring into focus his two nasal fields he must, to some extent, diverge his visual axes. Whether this divergence is visible or not will depend upon how far into the perimacular field he must go to find greater visual acuity. Eight cases are recorded here, and in certain of them there was a slight divergent strabismus without any accompanying weakness of convergence or of internal rotation of either eye on conjugate lateral movements.

L. S. KUBIE and J. W. BECKMANN (*Brain*, 1929,
lii., pp. 317 to 333).

SUBACUTE COMBINED DEGENERATION OF THE CORD: SYMPTOMATOLOGY AND EFFECTS OF LIVER THERAPY.

While the beneficial effects of liver diet in pernicious anæmia are well recognised, there seems to be some doubt as to the value of this form of treatment in subacute combined degeneration of the cord. A series of 61 cases are here reviewed, of which 30 had been treated

Periscope

by liver. In 12 cases the first symptoms were attributable to anæmia. Collier in 1926 stated that no case had ever been reported in the literature in which a patient, already under observation for pernicious anæmia, has developed signs and symptoms of combined degeneration of the spinal cord. This was confirmed in 1927 by Piney, but in the present series it occurred definitely in two instances. In the one case the course of events was (1) a stage of pernicious anæmia without any symptoms or signs of spinal cord involvement; (2) rapid remission of the anæmia with liver treatment; (3) relapse of the disease following cessation of the liver diet, with the development of subacute combined degeneration of the cord; (4) gradual improvement of both blood and nervous phenomena after a further course of liver treatment. Prior to the introduction of liver therapy subacute combined degeneration of the cord was always regarded as a progressive and fatal disease. In the 31 cases not treated by liver diet, 28 died within an average period of 14.6 months from the onset of the disease. Of the 3 still alive, the disease had lasted 2.1, 3.7 and six years respectively. On the other hand, of the 30 liver-treated cases, 5 died, and the other 25 are still living after an average period of 31 months. The improvement which occurs on dietetic measures suggests a similarity to certain deficiency diseases, such as pellagra and lathyrism, in which involvement of the white matter also occurs. In addition, it is noteworthy that the histological changes in the cord in these diseases bear a marked similarity to those found in subacute combined degeneration. In view of the constancy of achlorhydria and the presence of a hereditary factor, it is possible that we are dealing with a deficiency disease occurring in persons with a certain constitutional diathesis.

As with pernicious anæmia, so with the nervous phenomena, the remission brought about by liver therapy is only maintained as long as the treatment is continued. The best results are obtained by an intensive course ($3\frac{1}{2}$ to 4 lb. a week) for 6 to 8 weeks, followed by $1\frac{1}{2}$ to 2 lb. of liver per week indefinitely. The extract is less efficient than whole liver. The amount of liver required in subacute combined degeneration is greater than that necessary in "pure" pernicious anæmia. Active sepsis retards the improvement and necessitates larger amounts.

The authors conclude that as all symptoms and signs of cord involvement may disappear, the degenerative changes in the white matter cannot be regarded as necessarily of a permanent character.

C. C. UNGLEY and M. M. SURMAN (*Brain*, 1929, lii., pp. 271 to 294).

A. N. B.