EQUATOR: reporting guidelines for health research

Douglas G. Altman, Iveta Simera, John Hoey, David Moher, Ken Schulz

Affiliations: Centre for Statistics in Medicine, University of Oxford, Oxford, UK (DGA, IS); Queen's University, Kingston, Ont., Canada (JH). Chalmers Research Group, University of Ottawa, Ottawa, Ont., Canada (DM); Quantitative Sciences, Family Health International, Durham, NC, USA (KS).

Competing interests: We are all involved in health-care initiatives and research that should benefit from wide uptake of reporting guidelines.

Correspondence: iveta.simera@cancer.org.uk

This article was first published in *The Lancet* on 5 April 2008 and is reproduced here with permission.

editors and peer reviewers, the quality of health-research reporting in journal articles is unsatisfactory. Guidelines that specify a minimum set of items for reporting can improve the accuracy and transparency of publications, thus facilitating easier and more reliable appraisal of quality and relevance. During the past 10 years several internationally respected guidelines for the reporting of health research have been developed. However, those guidelines are still not widely supported by medical journals are or adhered to by researchers, and thus their potential impact is lessened.

To remedy this situation the UK National Knowledge Service provided funding to start the EQUATOR project (Enhancing the QUAlity and Transparency Of health Research). This initiative seeks to improve the reliability of medical publications by promoting transparent and accurate reporting of health research. This movement grew out of the work of CONSORT⁹ and other groups.

EQUATOR is an umbrella for all areas of healthresearch reporting. The network aims to become a global centre that provides resources and training, and which assists in the development, dissemination, and implementation of robust reporting guidelines. EQUATOR's strategic plan reflects the needs of its major stakeholders: developers of reporting guidelines, researchers, journal editors, peer-reviewers, and research-funding bodies.

One of the first activities was to identify existing reporting guidelines to see how they were developed. The development methods of most guidelines were broadly similar, but with wide variation in important details. Development usually took a long time and only half the groups had strategies for dissemination and implementation of their guidelines. The difficulty of securing sufficient funding to develop, assess, and disseminate guidelines was widely acknowledged as a major problem.

The initial survey and discussions with the main stakeholders helped us to prioritise future activities. First, we developed an internet-based resource centre, which can be freely accessed on our website.¹³ At present, it provides a collection of available reporting guidelines. In the future, the website will also host other resources for authors of research articles, editors, peer-reviewers, and developers of guidelines, including a comprehensive digital library for health-research reporting, guidance for the development of robust reporting guidelines, tools to facilitate their use, and educational materials.

The availability of good reporting guidelines is not sufficient for the improvement of the quality of reporting. Our second priority will be active promotion of such guidelines and their use, by developing and running training for editors, peer-reviewers, and authors. The courses will concentrate on the important factors of research reporting and the efficient use of reporting guidelines.

Poor reporting reflects a collective failure of those involved. Collaboration with and the support of influential medical journals are vital for the success of this project. Benefits will be equally split between both communities—users will benefit from improved reliability of scientific information and journals will benefit from increased loyalty of their readers attracted by improved accuracy and reliability of reports. The EQUATOR network will regularly monitor how journals



implement reporting guidelines. We will annually audit the quality of reporting across the health-research literature and hope to document gradual improvements.

Sufficient funding is a necessary requirement for the development and implementation of robust reporting guidelines and widespread promotion of good reporting of research. In view of how much money funding agencies spend on health research, their lack of interest in ensuring that this research is reported accurately is deeply disappointing. Good reporting is not an optional extra; it is an essential component of research. Funding bodies should recognise this and support initiatives such as EQUATOR that aim to improve the current situation.

The EQUATOR Network will hold its official launch meeting on June 26, 2008, in London, UK. The meeting will focus on better understanding of problems associated with health-research reporting and use of reporting guidelines, and on finding potential solutions to improve the health-research literature.

REFERENCES

- Chan A, Altman DG. Epidemiology and reporting of randomised trials published in PubMed journals. *Lancet* 2005;365(9465): 1159-62. [CrossRef] [PubMed]
- Mallett S, Deeks JJ, Halligan S, Hopewell S, Cornelius V, Altman DG. Systematic reviews of diagnostic tests in cancer: review of methods and reporting. BMJ 2006;333(7565):413. [CrossRef] [PubMed] [Full Text]
- 3. Mills E, Loke YK, Wu P, Montori VM, Perri D, Moher D, et al. Determining the reporting quality of RCTs in clinical pharmacology. *Br J Clin Pharmacol* 2004;58(1):61–5. [CrossRef] [PubMed] [Full Text]
- Pocock SJ, Collier TJ, Dandreo KJ, de Stavola BL, Goldman MB, Kalish LA, et al. Issues in the reporting of epidemiological studies: a survey of recent practice. BMJ 2004;329(7471):883. [CrossRef] [PubMed] [Full Text]

- Riley RD, Abrams KR, Sutton AJ, Lambert PC, Jones DR, Heney D, et al. Reporting of prognostic markers: current problems and development of guidelines for evidence-based practice in the future. *Br J Cancer* 2003;88(8):1191–8. [CrossRef] [PubMed]
- Smidt N, Rutjes AW, van der Windt DA, Ostelo RW, Reitsma JB, Bossuyt PM, et al. Quality of reporting of diagnostic accuracy studies. Radiology 2005;235(2):347–53. [CrossRef] [PubMed] [Full Text]
- Bossuyt PM, Reitsma JB, Bruns DE, Gatsonis CA, Glasziou PP, Irwig LM, et al. Towards complete and accurate reporting of studies of diagnostic accuracy: the STARD initiative. Standards for Reporting of Diagnostic Accuracy. Clin Chem 2003;49(1):1–6. [PubMed] [Full Text]
- 8. Moher D, Cook DJ, Eastwood S, Olkin I, Rennie D, Stroup DF. Improving the quality of reports of meta-analyses of randomised controlled trials: the QUOROM statement. Quality of Reporting of Meta-analyses. *Lancet* 1999;354(9193):1896–900. [PubMed]
- 9. Moher D, Schulz KF, Altman DG. The CONSORT statement: revised recommendations for improving the quality of reports of parallel-group randomised trials. *Lancet* 2001;357(9263):1191–4. [PubMed]
- 10. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, STROBE Initiative. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *PLoS Med* 2007;4(10):e296. [CrossRef] [PubMed] [Full Text]
- Altman DG. Endorsement of the CONSORT statement by high impact medical journals: survey of instructions for authors. *BMJ* 2005;330(7499):1056-7. [CrossRef] [PubMed] [Full Text]
- Smidt N, Overbeke J, de Vet H, Bossuyt P. Endorsement of the STARD Statement by biomedical journals: survey of instructions for authors. Clin Chem 2007;53(11):1983-5. [CrossRef] [PubMed]
- EQUATOR Network. Resource centre. 2007. Accessed 2007 Nov 29. [Full Text]

Citation: Altman DG, Simera I, Hoey J, Moher D, Schulz K. EQUATOR: reporting guidelines for health research. *Open Med* 2008;2(2):e24-25.

Published: 28 April 2008

Copyright: This article is licenced under the Creative Commons Attibution-ShareAlike 2.5 Canada License, which means that anyone is able to freely copy, download, reprint, reuse, distribute, display or perform this work and that the authors retain copyright of their work. Any derivative use of this work must be distributed only under a license identical to this one and must be attributed to the authors. Any of these conditions can be waived with permission from the copyright holder. These conditions do not negate or supersede Fair Use laws in any country. For further information see http://creativecommons.org/licenses/by-sa/2.5/ca/.