

Title: Effect of Postactivation Potentiation on Short Sprint Performance: A Systematic Review and Meta-Analysis

Jose Jonas de Oliveira,^{1,2,*} Alex Harley Crisp,¹ Carolina Gabriela Reis Barbosa,¹ Alexandre de Souza e Silva,² Ronaldo Julio Baganha,^{1,2} and Rozangela Verlengia¹

¹Post-graduate Program in Human Movement Sciences, University of Methodist of Piracicaba (UNIMEP), Piracicaba, Sao Paulo, Brazil

²Physical Education Department, University Centre of Itajuba (FEPI), Itajuba, Minas Gerais, Brazil

*Corresponding author: Jose Jonas de Oliveira, Rodovia do Acucar, km 156, Zip code 13.400-911, Piracicaba Sao Paulo, Brazil. Tel: +55-1931241515, Fax: +55-1931241500, E-mail: joliveira63@gmail.com

Received 2017 February 17; Revised 2017 June 06; Accepted 2017 July 31.

Abstract

Background: Short sprint is an important physical ability that determines the success in various sports modalities and may influence different conditioning activities. Postactivation potentiation (PAP) is a conditioning method used in practicing sports, which can result in acute improvements in muscle power and performance by interaction of physiological and neural mechanisms.

Objectives: The current study aimed at investigating the effect of PAP on a sprint of less than 40 meters.

Methods: A systematic review and meta-analysis was conducted on the randomized, controlled studies. The search was conducted in several databases (PubMed, Web of Science, Cochrane, and Science Direct) using the following keywords: "Postactivation Potentiation", "Sprint", and "Complex Training and Sprint". The sprint time data (mean and standard deviation) of the selected studies were analyzed using the OpenMeta (Analyst)[®] program. Data were expressed as weighted mean differences (WMD) between the groups with 95% confidence interval (95% CI). The level of significance was $P < 0.05$.

Results: A total of 1859 records were initially identified out of which 15 studies were selected according to the inclusion criteria and included in the current study. The meta-analysis results showed a positive effect of PAP on sprint time in the distance 0 to 10 meter (WMD = -0.031 seconds, 95% CI: -0.050, -0.012, $P = 0.001$), 11 to 20 meter (WMD = -0.048 seconds, 95% CI: -0.089, -0.007, $P = 0.021$), 21 to 30 meter (WMD = -0.060 seconds, 95% CI: -0.094, -0.026; $P < 0.001$), and 31 to 40 meter (WMD = -0.109 seconds, 95% CI = -0.141, -0.077, $P < 0.001$).

Conclusion: In summary, PAP induced positive effects on short sprint performances.

Keywords: High-Intensity Exercise, Warm-Up, Postactivation Potentiation, Short Sprint

1. Introduction

Postactivation potentiation (PAP) refers to an initial muscular activation with moderate/high load intensity, which results in acute improvements in muscle power and performance in subsequent explosive activities. Thus, PAP is used to elaborate physical training sessions or prior competitions due to their potential positive effects on athletic performance (1, 2).

The main mechanisms suggested responsible to improve physical performance are the phosphorylation of the myosin regulatory light chain (3, 4), and this effect occurs in greater magnitude in type II muscle fibers, which consequently favors the performance in high intensity and short duration activities (1).

However, some controversial results are observed, with some studies indicating no significant improvements on sprint performance (5-9). Many factors such as muscle fiber type (10), training experience (11), gender (12) and muscle strength level (13) can influence the individual re-

sponsiveness to PAP. In addition, evidence indicates that manipulation of the PAP protocol (intensity load, volume, rest interval) exerts a significant influence on the subsequent performance response (14).

On the other hand, an important factor is the transfer of the PAP stimulus to specific sports muscle actions. In this context, most studies investigate the influence of PAP on lower limbs and evaluate the performance improvement through vertical jumps (12, 15-27). However, a smaller number of studies investigated the effects of PAP on sprint performance, with results expressed at different distances (5-7, 13, 28-34).

Short sprint is considered as an important physical ability that determines the success of various sports modalities. Therefore, the current study aimed at conducting a systematic review on the evidence and performing a meta-analysis to obtain information about the PAP effect on the performance of short sprints.

2. Methodology

2.1. Search Strategy

A systematic review and meta-analysis of randomized, controlled studies were conducted and evaluated the PAP effect on short sprint performance. The search was performed in the following databases: PubMed, Web of Science, Cochrane, and Science Direct. The following keywords were searched: “postactivation potentiation”, “sprint”, “complex training and sprint”. The search was conducted in November 2016. The systematic review was not limited to specific years.

2.2. Inclusion and Exclusion Criteria

The studies considered eligible based on the following inclusion criteria: (i) randomized, controlled studies; (ii) peer-reviewed and published in English; (iii) used the running sprint as a performance test.

The adopted exclusion criteria were (i) sprint test greater than 40 meters; (ii) PAP performed in upper limbs; (iii) PAP performed in isometric exercises; (iv) PAP performed in jumping exercises; (v) PAP performed using isokinetic equipment; (vi) PAP performed using a vibratory platform; (vii) load intensity less than 70% of 1RM (1-repetition maximum); (viii) rest interval of less than 4 minutes between the PAP and the sprint test.

2.3. Studies Selection

One of the researchers searched the databases and excluded the duplicate articles. The evaluation of the titles and abstracts was done by 2 independent researchers in which they selected the articles pertinent to the theme of the study. The same researchers performed a complete reading of the articles and selected the eligible ones based on the inclusion and exclusion criteria. In addition, the cited references from eligible studies were also analyzed to identify relevant studies not found in the search. Disagreements in the inclusion and exclusion of studies among the researchers were resolved by consulting a 3rd reviewer.

2.4. Data Extraction

Two independent researchers did the data extraction. The extracted data included (i) study design; (ii) number of subjects; (iii) age; (iv) sets and repetition in PAP protocol; (v) intensity load in PAP protocol; (vi) rest interval between PAP protocol and sprint test; (vii) distance from the sprint test; (viii) mean \pm standard deviation (SD) of sprint time. Disagreements among the researchers on data extraction were resolved by consensus.

2.5. Statistical Analysis

The sprint time data (mean \pm SD) of the selected studies were analyzed using the OpenMeta (Analyst)[®] program. The heterogeneity between the studies was assessed by the Cochrane Q-test and the I² inconsistency test. Due to the low heterogeneity, a fixed effect model was used for the analysis. Four meta-analyses were performed based on the final sprint distance: (i) up to 10 meter; (ii) 11 to 20 meter; (iii) 21 to 30 meter; (iv) 31 to 40 meter. Data were expressed as weighted mean differences (WMD) between the groups with 95% confidence interval (95% CI). Level of significance was $P < 0.05$.

3. Results

Figure 1 illustrates the flowchart of the excluded and included studies in the current systematic review. The search strategy identified 1859 records in the searched databases (PubMed, Web of Science, Cochrane, and Science Direct). After an initial reading of titles and abstracts, 35 relevant studies were selected. They were thoroughly read and according to the inclusion/exclusion criteria, 15 studies were selected. In the current meta-analysis, 11 studies were included, which presented mean \pm SD data for sprint time.

3.1. Study Characteristics

Table 1 presents the characteristics and results of each study included in the current systematic review. The studies were published from 2005 to 2016. The included studies ($n = 15$) constitute of totally 203 subjects (PAP condition ($n = 203$) and control ($n = 203$). All of the studies evaluated males aged 16 to 25 years. The studies presented data of athletes from sports of collective modality (5-7, 9, 13, 22, 28-34), athleticism (21, 35), and resistance trained subjects (8, 9).

The PAP protocol characteristics of the studies were constituted of 1 to 5 sets and 1 to 10 repetitions. The studies used the load intensity from 70% to 91% in 1RM (6, 8, 13, 28-35). Two studies (5, 22) used the load intensity of 3RM, and one study (7) used 5RM.

Okuno et al., (33) and Yetter et al., (31) used the progression of the load intensity (30% to 90% of 1RM) between the sets. Vanderka et al., (9) used the load intensity at the individual maximum power output. The rest interval between the PAP and the sprint test ranged from 4 to 16 minutes. The total distance covered in the sprint test ranged from 5 to 40 meters in a straight line. Only 1 study (33) evaluated the best sprint time of 30 meters with a change of direction (15 + 15 meters) in a repeated sprint protocol.




Figure 1. Flowchart of the included Studies

3.2. Meta-Analysis

Six studies evaluated the sprint time in the final distance of up to 10 meter (7, 8, 22, 28, 30, 35). In the study by Bevan et al. (32), sprint time data were included at distances of 5 and 10 meters. Till and Cooke (7) presented the sprint time in individuals with greater ($n = 6$) and lower muscular strength level ($n = 6$), according to the 5RM test. The study by Wyland et al., (35) included data from tradi-

tional squat exercises ($n = 20$) and squat exercises with an elastic band ($n = 20$). The PAP had a positive effect ($P = 0.001$) on the reduction of sprint time, compared with the control condition, with a WMD value of -0.031 seconds (95% CI: $-0.050, -0.012$). No heterogeneity ($I^2 = 0, P = 0.944$) was observed between the studies (Figure 2A).

Three studies evaluated the sprint time in the final distance of 11 to 20 meter (7, 8, 13). Till and Cooke (7) presented

Table 1. Results of the Studies on Short Sprint Performance of Postactivation Potentiation

| No. | Author, y | Subjects (Sports Modality) | Age | PAP Protocol | | | | Results |
|-----|----------------------------|---|-------------|---------------------|------------------------|--------------------|-----------------------|---|
| | | | | Set × Repetitions | Load Intensity, % | Rest Interval, min | Distance (Sprint), m | |
| 01 | Bevan et al. (2010) | 16 males (rugby) | - | 1 × 3 | 91 1RM | 4, 8, 12, 16 | 5 and 10 | Decrease in the sprint time -5 and 10 m |
| 02 | Chatzopoulos et al. (2007) | 15 males (amateur team game players) | 22.0 ± 2.0 | 1 × 10 | 90 1RM | 5 | 30 | Decrease in the sprint time -10 and 30 m |
| 03 | Crewther et al. (2011) | 9 males (rugby) | 20.1 ± 0.9 | 1 × 3 | 3RM | 4, 8, 12, 16 | 10 | ND in sprint time |
| 04 | Duncan et al. (2014) | 10 males (rugby) | 25.2 ± 5.02 | 1 × 3 | 90 1RM | 4 | 30 | ND in sprint time |
| 05 | Evetovich et al. (2015) | 7 males (soccer) | 20.4 ± 1.6 | 1 × 3 | 3RM | 8 | 36.6 | Decrease in the sprint time |
| 06 | Lim et al. (2013) | 12 males (sprinters) | 22.4 ± 3.2 | 1 × 3 | 90 1RM | 4 | 30 | ND in sprint time |
| 07 | Low et al. (2015) | 16 males (soccer) | 17.0 ± 0.6 | 1 × 3 | 91 1RM | 4 | 35 | Decrease in the sprint time |
| 08 | Mcbride et al. (2005) | 15 males (soccer) | 20.8 ± 1.0 | 1 × 3 | 90 1RM | 4 | 40 | Decrease in the sprint time -40 m, ND in the sprint time -10 and 30 m |
| 09 | Okuno et al. (2013) | 12 males (handball) | 18.7 ± 1.7 | 1 × 5, 1 × 3, 5 × 1 | 50 1RM, 70 1RM, 90 1RM | 5 | RSA 6 × 30, (15 + 15) | Improved best sprint time, Improved mean sprint time, ND fatigue |
| 10 | Rahimi (2007) | 11 males (soccer) | 22.4 ± 1.0 | 2 × 4 | 70 1RM, 85 1RM | 4 | 40 | Decrease in the sprint time |
| 11 | Seitz et al. (2014) | 13 males (rugby) | 18.3 ± 0.9 | 1 × 3 | 90 1RM | 7 | 20 | Decrease in the sprint time |
| 12 | Till et al. (2009) | 12 males (soccer) | 18.3 ± 0.72 | 1 × 5 | 5RM | 4 | 20 | ND in the sprint time |
| 13 | Vanderka et al. (2016) | 12 males (track and field) | 16 - 18 | 2 × 6 | Maximum Power | 6 | 40 | ND in the sprint time |
| | | 13 males (soccer) | 17 - 19 | | | | | |
| 14 | Wyland et al. (2015) | 20 males (resistance trained) | 23.3 ± 4.4 | 5 × 3 | 85 1RM | 4 | 5 × 9.1 | Decrease in sprint time - PAP with elastic band |
| 15 | Yetter et al. (2008) | 10 males (football, weightlifting, and track and field) | 22.3 ± 0.8 | 1 × 5, 1 × 4, 1 × 3 | 30, 50, 70 | 4 | 40 | Decrease in sprint time |

Abbreviations: ND, No Significant Difference; RM, Repetition Maximum.

the sprint time in subjects who had greater ($n = 6$) and lower strength levels ($n = 6$), according to the 5RM test. The study by Seitz et al., (13) included data from squats ($n = 13$) and power clean ($n = 13$) exercises. The PAP had a positive

effect ($P = 0.021$) on the reduction of sprint time compared with the control condition, with a WMD value of -0.048 seconds (95% CI: $-0.089, -0.007$). No heterogeneity was observed ($I^2 = 24.35\%$, $P = 0.252$) between the studies (Figure

2B).

Four studies evaluated the sprint time in the distance of 21 to 30 meter (8, 28, 30, 33). The PAP had a positive effect ($P < 0.001$) on the reduction of sprint time compared with the control condition, with a WMD value of -0.060 seconds (95% CI: -0.094, -0.026). No heterogeneity was observed ($I^2 = 0\%$, $P = 0.553$) between the studies (Figure 3A).

Four studies evaluated the sprint time in the final distance of 31 to 40 meter (22, 28, 29, 34). The study by Rahimi et al. (29), included the data of PAP with load intensities of 70% of 1RM ($n = 11$) and 85% of 1RM ($n = 11$). The PAP had a positive effect ($P < 0.001$) on the reduction of sprint time compared with the control condition, with a WMD value of -0.109 seconds (95% CI: -0.141, -0.077). No heterogeneity was observed ($I^2 = 0\%$, $P = 0.195$) between the studies (Figure 3B).

4. Discussion

The current study aimed at performing a systematic review and meta-analysis to determine the effectiveness of PAP on short sprints performances. The main finding was that PAP significantly reduced sprint time compared with the control condition.

The results of the current study corroborated with another meta-analysis (36) that indicated an overall moderate effect in sprint activities after PAP. However, the current study expanded these findings by showing that PAP increased short sprint performance considered essential in several sports modalities.

Sprint performance can be represented in 3 phases: (a) acceleration, (b) maximum speed, and (c) deceleration (37). The acceleration phase can be subdivided into initial (0 to 12 meter) and main (12 to 35 meter) (38); thus, the ability to produce high levels of muscular strength/power at the beginning of the sprint is important to increase acceleration and short sprint performance (39).

Evidence indicates a moderate to high correlation between the maximum muscle strength and power with short sprint performance (40-43). Wisloff et al., (41) observed a high and moderate correlation between muscle 1RM strength in back squat exercises with sprint time at distances of 10 meters ($r = 0.94$) and 30 meters ($r = 0.71$). In this sense, strategies that acutely enhance muscle strength and power capacity can affect the performance of short sprints.

The current study data indicated that PAP was an efficient preconditioning method to increase acceleration capacity and reduce sprint time at distances 0 to 10 meter (WMP = -0.031 seconds), 11 to 20 meter (WMP = -0.048 seconds), 21 to 30 meter (WMD = -0.060 seconds), and 31 to 40 meter (WMD = -0.109 seconds). The improvement of short sprint performances after PAP may be related to an

interaction of physiological and neural mechanisms, such as phosphorylation of the regulatory myosin light chain (4) and the increase of high-threshold motor units recruitment (11, 44, 45).

The manipulation of PAP protocols such as volume (sets and repetitions), intensity, load, and rest interval exerts on the subsequent athletic performance (14). According to the studies that observed positive effect of PAP on the short sprint performance (13, 22, 28-32, 34, 35), the exponential effect in the majority of the studies was a single set of 3 repetitions, performed at high load intensity (~90% 1RM) and rest interval between 4 and 8 minutes in back squat exercise. This finding is the important information for strength and conditioning coaches to prescribe PAP protocol prior to short sprints activities.

The current meta-analysis had some limitations that should be addressed. A small number of studies included the analysis of different sprint distances. Additionally, some of the included studies had small samples sizes. The major strength of the study was that the majority of studies included evaluated athletes.

Conclusively, the current systematic review and meta-analysis provide evidence that PAP induced positive effects on short sprint performance. Based on the practical applications, PAP can be an interesting, intense strategy to potentiate athletic performance if prescribed by strength and conditioning coaches before activities that involve short sprints.

References

1. Robbins DW. Postactivation potentiation and its practical applicability: a brief review. *J Strength Cond Res.* 2005;19(2):453-8. doi: 10.1519/R-14653.1. [PubMed: 15903390].
2. Kilduff LP, Bevan HR, Kingsley MI, Owen NJ, Bennett MA, Bunce PJ, et al. Postactivation potentiation in professional rugby players: optimal recovery. *J Strength Cond Res.* 2007;21(4):1134-8. doi: 10.1519/R-20996.1. [PubMed: 18076243].
3. Rassier DE, Macintosh BR. Coexistence of potentiation and fatigue in skeletal muscle. *Braz J Med Biol Res.* 2000;33(5):499-508. doi: 10.1590/S0100-879X2000000500003. [PubMed: 10775880].
4. Esformes JI, Bampouras TM. Effect of back squat depth on lower-body postactivation potentiation. *J Strength Cond Res.* 2013;27(11):2997-3000. doi: 10.1519/JSC.0b013e31828d4465. [PubMed: 23442291].
5. Crewther BT, Kilduff LP, Cook CJ, Middleton MK, Bunce PJ, Yang GZ. The acute potentiating effects of back squats on athlete performance. *J Strength Cond Res.* 2011;25(12):3319-25. doi: 10.1519/JSC.0b013e318215f560. [PubMed: 22076086].
6. Duncan MJ, Thurgood G, Oxford SW. Effect of heavy back squats on repeated sprint performance in trained men. *J Sports Med Phys Fitness.* 2014;54(2):238-43. [PubMed: 24509997].
7. Till KA, Cooke C. The effects of postactivation potentiation on sprint and jump performance of male academy soccer players. *J Strength Cond Res.* 2009;23(7):1960-7. doi: 10.1519/JSC.0b013e3181b8666e. [PubMed: 19855319].
8. Lim JJ, Kong PW. Effects of isometric and dynamic postactivation potentiation protocols on maximal sprint performance. *J Strength Cond*

- Res. 2013;**27**(10):2730–6. doi: [10.1519/JSC.0b013e3182815995](https://doi.org/10.1519/JSC.0b013e3182815995). [PubMed: 23302751].
9. Vanderka M, Krcmar M, Longova K, Walker S. Acute Effects of Loaded Half-Squat Jumps on Sprint Running Speed in Track and Field Athletes and Soccer Players. *J Strength Cond Res.* 2016;**30**(6):1540–6. doi: [10.1519/JSC.000000000001259](https://doi.org/10.1519/JSC.000000000001259). [PubMed: 26562707].
 10. Hamada T, Sale DG, MacDougall JD, Tarnopolsky MA. Postactivation potentiation, fiber type, and twitch contraction time in human knee extensor muscles. *J Appl Physiol (1985).* 2000;**88**(6):2131–7. [PubMed: 10846027].
 11. Chiu LZ, Fry AC, Weiss LW, Schilling BK, Brown LE, Smith SL. Postactivation potentiation response in athletic and recreationally trained individuals. *J Strength Cond Res.* 2003;**17**(4):671–7. doi: [10.1519/1533-4287\(2003\)017<0671:PPRIAA>2.0.CO;2](https://doi.org/10.1519/1533-4287(2003)017<0671:PPRIAA>2.0.CO;2). [PubMed: 14636093].
 12. Rixon KP, Lamont HS, Bemben MG. Influence of type of muscle contraction, gender, and lifting experience on postactivation potentiation performance. *J Strength Cond Res.* 2007;**21**(2):500–5. doi: [10.1519/R-18855.1](https://doi.org/10.1519/R-18855.1). [PubMed: 17530946].
 13. Seitz LB, de Villarreal ES, Haff GG. The temporal profile of postactivation potentiation is related to strength level. *J Strength Cond Res.* 2014;**28**(3):706–15. doi: [10.1519/JSC.0b013e3182a73ea3](https://doi.org/10.1519/JSC.0b013e3182a73ea3). [PubMed: 23965945].
 14. Wilson JM, Duncan NM, Marin PJ, Brown LE, Loenneke JP, Wilson SM, et al. Meta-analysis of postactivation potentiation and power: effects of conditioning activity, volume, gender, rest periods, and training status. *J Strength Cond Res.* 2013;**27**(3):854–9. doi: [10.1519/JSC.0b013e31825c2bdb](https://doi.org/10.1519/JSC.0b013e31825c2bdb). [PubMed: 22580978].
 15. Young WB, Jenner A, Griffiths K. Acute enhancement of power performance from heavy load squats. *J Strength Cond Res.* 1998;**12**(2):82–4. doi: [10.1519/1533-4287](https://doi.org/10.1519/1533-4287).
 16. Scott SL, Docherty D. Acute effects of heavy preloading on vertical and horizontal jump performance. *J Strength Cond Res.* 2004;**18**(2):201–5. doi: [10.1519/R-13123.1](https://doi.org/10.1519/R-13123.1). [PubMed: 15142025].
 17. Hilfiker R, Hubner K, Lorenz T, Marti B. Effects of drop jumps added to the warm-up of elite sport athletes with a high capacity for explosive force development. *J Strength Cond Res.* 2007;**21**(2):550–5. doi: [10.1519/R-20215.1](https://doi.org/10.1519/R-20215.1). [PubMed: 17530974].
 18. Weber KR, Brown LE, Coburn JW, Zinder SM. Acute effects of heavy-load squats on consecutive squat jump performance. *J Strength Cond Res.* 2008;**22**(3):726–30. doi: [10.1519/JSC.0b013e3181660899](https://doi.org/10.1519/JSC.0b013e3181660899). [PubMed: 18438248].
 19. Requena B, Gapeyeva H, Garcia I, Erelina J, Paasuke M. Twitch potentiation after voluntary versus electrically induced isometric contractions in human knee extensor muscles. *Eur J Appl Physiol.* 2008;**104**(3):463–72. doi: [10.1007/s00421-008-0793-8](https://doi.org/10.1007/s00421-008-0793-8). [PubMed: 18563434].
 20. Needham RA, Morse CI, Degens H. The acute effect of different warm-up protocols on anaerobic performance in elite youth soccer players. *J Strength Cond Res.* 2009;**23**(9):2614–20. doi: [10.1519/JSC.0b013e3181bf3ef](https://doi.org/10.1519/JSC.0b013e3181bf3ef). [PubMed: 19910812].
 21. Witmer CA, Davis SE, Moir GL. The acute effects of back squats on vertical jump performance in men and women. *J Sports Sci Med.* 2010;**9**(2):206–13. [PubMed: 24149687].
 22. Evetovich TK, Conley DS, McCawley PF. Postactivation potentiation enhances upper- and lower-body athletic performance in collegiate male and female athletes. *J Strength Cond Res.* 2015;**29**(2):336–42. doi: [10.1519/JSC.0000000000000728](https://doi.org/10.1519/JSC.0000000000000728). [PubMed: 25330084].
 23. West D, Cunningham D, Bevan H, Crewther B, Cook C, Kilduff L. Influence of active recovery on professional rugby union player's ability to harness postactivation potentiation. *J Sports Med Phys Fitness.* 2013;**53**(2):203–8. [PubMed: 23584329].
 24. Bogdanis GC, Tsoukas A, Veligekas P, Tsolakis C, Terzis G. Effects of muscle action type with equal impulse of conditioning activity on postactivation potentiation. *J Strength Cond Res.* 2014;**28**(9):2521–8. doi: [10.1519/JSC.0000000000000444](https://doi.org/10.1519/JSC.0000000000000444). [PubMed: 24584048].
 25. Arabatzis F, Patikas D, Zafeiridis A, Giavroudis K, Kannas T, Gourgoulis V, et al. The post-activation potentiation effect on squat jump performance: age and sex effect. *Pediatr Exerc Sci.* 2014;**26**(2):187–94. doi: [10.1123/pes.2013-0052](https://doi.org/10.1123/pes.2013-0052). [PubMed: 24225048].
 26. Fukutani A, Takei S, Hirata K, Miyamoto N, Kanehisa H, Kawakami Y. Influence of the intensity of squat exercises on the subsequent jump performance. *J Strength Cond Res.* 2014;**28**(8):2236–43. doi: [10.1519/JSC.0000000000000409](https://doi.org/10.1519/JSC.0000000000000409). [PubMed: 24513618].
 27. Naclerio F, Chapman M, Larumbe-Zabala E, Massey B, Neil A, Triplett TN. Effects of Three Different Conditioning Activity Volumes on the Optimal Recovery Time for Potentiation in College Athletes. *J Strength Cond Res.* 2015;**29**(9):2579–85. doi: [10.1519/JSC.0000000000000915](https://doi.org/10.1519/JSC.0000000000000915). [PubMed: 25719916].
 28. McBride JM, Nimphius S, Erickson TM. The acute effects of heavy-load squats and loaded counter-movement jumps on sprint performance. *J Strength Cond Res.* 2005;**19**(4):893–7. doi: [10.1519/R-16304.1](https://doi.org/10.1519/R-16304.1). [PubMed: 16287357].
 29. Rahimi R. The acute effects of heavy versus light-load squats on sprint performance. *Sci J Facta Univ.* 2007;**5**(2):163–9.
 30. Chatzopoulos DE, Michailidis CJ, Giannakos AK, Alexiou KC, Patikas DA, Antonopoulos CB, et al. Postactivation potentiation effects after heavy resistance exercise on running speed. *J Strength Cond Res.* 2007;**21**(4):1278–81. doi: [10.1519/R-21276.1](https://doi.org/10.1519/R-21276.1). [PubMed: 18076255].
 31. Yetter M, Moir GL. The acute effects of heavy back and front squats on speed during forty-meter sprint trials. *J Strength Cond Res.* 2008;**22**(1):159–65. doi: [10.1519/JSC.0b013e31815f958d](https://doi.org/10.1519/JSC.0b013e31815f958d). [PubMed: 18296970].
 32. Bevan HR, Cunningham DJ, Tooley EP, Owen NJ, Cook CJ, Kilduff LP. Influence of postactivation potentiation on sprinting performance in professional rugby players. *J Strength Cond Res.* 2010;**24**(3):701–5. doi: [10.1519/JSC.0b013e3181c7b68a](https://doi.org/10.1519/JSC.0b013e3181c7b68a). [PubMed: 20145565].
 33. Okuno NM, Tricoli V, Silva SB, Bertuzzi R, Moreira A, Kiss MA. Postactivation potentiation on repeated-sprint ability in elite handball players. *J Strength Cond Res.* 2013;**27**(3):662–8. doi: [10.1519/JSC.0b013e31825bb582](https://doi.org/10.1519/JSC.0b013e31825bb582). [PubMed: 22561976].
 34. Low D, Harsley P, Shaw M, Peart D. The effect of heavy resistance exercise on repeated sprint performance in youth athletes. *J Sports Sci.* 2015;**33**(10):1028–34. doi: [10.1080/02640414.2014.979857](https://doi.org/10.1080/02640414.2014.979857). [PubMed: 25554921].
 35. Wyland TP, Van Dorin JD, Reyes GF. Postactivation Potentiation Effects From Accommodating Resistance Combined With Heavy Back Squats on Short Sprint Performance. *J Strength Cond Res.* 2015;**29**(11):3115–23. doi: [10.1519/JSC.0000000000000991](https://doi.org/10.1519/JSC.0000000000000991). [PubMed: 25968229].
 36. Seitz LB, Haff GG. Factors Modulating Post-Activation Potentiation of Jump, Sprint, Throw, and Upper-Body Ballistic Performances: A Systematic Review with Meta-Analysis. *Sports Med.* 2016;**46**(2):231–40. doi: [10.1007/s40279-015-0415-7](https://doi.org/10.1007/s40279-015-0415-7). [PubMed: 26508319].
 37. Bishop D, Girard O, Mendez-Villanueva A. Repeated-sprint ability - part II: recommendations for training. *Sports Med.* 2011;**41**(9):741–56. doi: [10.2165/11590560-000000000-00000](https://doi.org/10.2165/11590560-000000000-00000). [PubMed: 21846163].
 38. Mackala K, Fostiak M, Kowalski K. Selected determinants of acceleration in the 100m sprint. *J Hum Kinet.* 2015;**45**:135–48. doi: [10.1515/hukin-2015-0014](https://doi.org/10.1515/hukin-2015-0014). [PubMed: 25964817].
 39. Bissas AI, Havenetidis K. The use of various strength-power tests as predictors of sprint running performance. *J Sports Med Phys Fitness.* 2008;**48**(1):49–54. [PubMed: 18212710].
 40. Newman MA, Tarpenning KM, Marino FE. Relationships between isokinetic knee strength, single-sprint performance, and repeated-sprint ability in football players. *J Strength Cond Res.* 2004;**18**(4):867–72. doi: [10.1519/13843.1](https://doi.org/10.1519/13843.1). [PubMed: 15574095].
 41. Wisloff U, Castagna C, Helgerud J, Jones R, Hoff J. Strong correlation of maximal squat strength with sprint performance and vertical jump height in elite soccer players. *Br J Sports Med.* 2004;**38**(3):285–8. doi: [10.1136/bjism.2002.002071](https://doi.org/10.1136/bjism.2002.002071). [PubMed: 15155427].
 42. Lopez-Segovia M, Marques MC, van den Tillaar R, Gonzalez-Badillo JJ. Relationships between vertical jump and full squat power outputs

- with sprint times in u21 soccer players. *J Hum Kinet.* 2011;**30**:135-44. doi: [10.2478/v10078-011-0081-2](https://doi.org/10.2478/v10078-011-0081-2). [PubMed: [23487438](https://pubmed.ncbi.nlm.nih.gov/23487438/)].
43. Wang R, Hoffman JR, Tanigawa S, Miramonti AA, La Monica MB, Beyer KS, et al. Isometric Mid-Thigh Pull Correlates With Strength, Sprint, and Agility Performance in Collegiate Rugby Union Players. *J Strength Cond Res.* 2016;**30**(11):3051-6. doi: [10.1519/JSC.0000000000001416](https://doi.org/10.1519/JSC.0000000000001416). [PubMed: [26982977](https://pubmed.ncbi.nlm.nih.gov/26982977/)].
44. Baker D. Acute effect of alternating heavy and light resistances on power output during upper-body complex power training. *J Strength Cond Res.* 2003;**17**(3):493-7. doi: [10.1519/00124278-200308000-00011](https://doi.org/10.1519/00124278-200308000-00011). [PubMed: [12930175](https://pubmed.ncbi.nlm.nih.gov/12930175/)].
45. Hodgson M, Docherty D, Robbins D. Post-activation potentiation: underlying physiology and implications for motor performance. *Sports Med.* 2005;**35**(7):585-95. doi: [10.2165/00007256-200535070-00004](https://doi.org/10.2165/00007256-200535070-00004). [PubMed: [16026172](https://pubmed.ncbi.nlm.nih.gov/16026172/)].

Uncorrected Proof