

Palaeartic species of *Rhamphomyia* (*Pararhamphomyia*) *anfractuosa* group (Diptera, Empididae)

Miroslav Barták¹, Štěpán Kubík¹

¹ Department of Zoology and Fisheries, Faculty of Agrobiolgy, Food and Natural Resources, Czech University of Life Sciences Prague, CZ-16521 Praha 6-Suchbát, Czech Republic

Corresponding author: Miroslav Barták (bartak@af.czu.cz)

Academic editor: M. Ivković | Received 10 February 2015 | Accepted 11 July 2015 | Published 27 July 2015

<http://zoobank.org/9F5794FC-3FF0-49FE-A7C3-4A410BBB44E5>

Citation: Barták M, Kubík S (2015) Palaeartic species of *Rhamphomyia* (*Pararhamphomyia*) *anfractuosa* group (Diptera, Empididae). ZooKeys 514: 111–127. doi: 10.3897/zookeys.514.9379

Abstract

Palaeartic species of the *Rhamphomyia* (*Pararhamphomyia*) *anfractuosa* group are revised. *Rhamphomyia* (*Pararhamphomyia*) *biflexata* **sp. n.**, *R.* (*Pararhamphomyia*) *lineodorsata* **sp. n.**, *R.* (*Pararhamphomyia*) *nudiscutellata* **sp. n.**, and *R.* (*Pararhamphomyia*) *shatalkini* **sp. n.** (all from Russian Far East) are described and illustrated. A key to Palaeartic species of the *Rhamphomyia* (*Pararhamphomyia*) *anfractuosa* group is provided.

Keywords

Empidoidea, *Rhamphomyia*, taxonomy, key, new species, Palaeartic

Introduction

Rhamphomyia Meigen is one of the three megadiverse groups of Empididae, alongside *Empis* Linnaeus and *Hilara* Meigen. Almost 600 species, distributed mostly in the Northern Hemisphere have been described worldwide (Yang et al. 2007; Barták 2007; Barták et al. 2007; Barták and Kubík 2008a, b, c, 2009, 2010, 2012; Saigusa 2012; Barták et al. 2014), but many more await description.

Rhamphomyia (*Pararhamphomyia*) *anfractuosa* group is delimited here as probably a natural group of *Pararhamphomyia* (as fixed by Barták and Sinclair 2003 and diagnosed by Barták 1982). The most important characters shared by members of this group are as follows (* for characters considered synapomorphic):

- acrostichals present, biserial
- male head holoptic
- legs brown to black
- proepisternal depression without setae
- phallus forms several tightly coiled loops*
- cercus simple (“subcercal process” or “posterior cercus” absent)*
- hypandrium membranose posteriorly, anterior part desclerotized medially*
- tip of epandrial lamellae with expanding membranose pouch (see Fig. 1 and 9)*
- tergite and sternite 8 fused*
- axillary angle acute to right angled
- anal vein complete or depigmented in middle
- halter yellow
- female leg parts broadly pennate

As usual in *Rhamphomyia*, it is not always easy to arrange single females into groups or even subgenera (compare species group approach in Barták 2002, 2003; Barták and Kubík 2009), so, separate key to females of this group would be meaningless.

The most allied species to this group are species of *Pararhamphomyia* sharing most characters with this group mentioned above, except the tightly coiled phallus and/or yellow halter (e.g., *R. slovaki* Barták, *R. plumifera* Zetterstedt, *R. deformata* Frey, *R. deformatella* Barták). The subgenus *Calorhamphomyia* Saigusa with similarly modified last abdominal segments and phallus may be allied to this group of species, however its members may be easily separated by at least partly yellow legs.

Material and methods

The material studied is deposited in the following collections: CULSP (Czech University of Life Sciences Prague), ZMMU (Zoological Museum of Moscow State University). Acronyms are used further in the text.

Terminal abdominal segments (Figs 1–8) and hind legs (Figs 13–16) were photographed by means of Nikon Digital sight DS Fi-1. Each image resulted usually from combining 7–10 layers by means of Nish element. Legs were illustrated from these images; details were added by direct observation. Genitalia (Figs 9–12) together with 2–3 pregenital segments were removed and macerated in potassium hydroxide solution (approx. 10 %) in small vials submerged in hot water for 1–2 hours. After neutralizing with 8 % acetic acid, the genitalia were dissected in glycerine and their parts photographed by means of an Olympus E-41 digital camera mounted on an Olympus BX51

compound microscope. Images were edited with the computer software Quick Foto micro 2.3 provided with Deep focus 3.1. Each image resulted usually from combining 7–15 layers. Images were improved by means of Adobe Photoshop.

The morphological terms used here follow Merz and Haenni (2000), Sinclair (2000) and Sinclair and Cumming (2006). All body measurements (including body and setae length) were taken from dry specimens (therefore the actual length may differ) by means of an ocular micrometer on a Nikon SMZ 1500 binocular microscope.

Length of antennal segments = length of scape: pedicel: postpedicel: style (in 0.01 mm). Male body length was measured from antennal base to the tip of last abdominal segment (without the genitalia) and female body length from the base of antennae to the tip of the cerci. Wing measurements: M_2/d = length of vein M_2 ; greatest length of discal medial cell (discal cell); CuA_1 ratio = length of apical: preapical sections of vein CuA_1 ; lw/ww = greatest length of wing (from basicosta to apex); greatest width of wing. Length of frons is measured from front margin of front ocellus to antennal base.

Taxonomic account

Rhamphomyia (*Pararhamphomyia*) *biflexata* sp. n.

<http://zoobank.org/2715DE4F-1E0A-4636-825F-751DC030B876>

Figs 2, 9, 14


Type material. HOLOTYPE male, Russia, Amurskaja oblast, g. (= city) Zeja, 8.vii.1981, A. Shatalkin, deposited in ZMMU; PARATYPE: 1 male, same data as holotype (CULSP).

Diagnosis. Species of the subgenus *Pararhamphomyia* with pair of long thumb-like dorsomarginal processes on tergite 6, lustrous abdomen, uniserial dorsocentrals and two outgoing loops of the phallus.

Etymology. The species epithet stresses two outgoing loops on the phallus.

Description. Male. Head holoptic, facets in dorsal third of eye enlarged. Frons blackish brown, light grey microtrichose, without setae. Ocellar setae less than half as long as frons, black, ocellar triangle without additional setae. Face blackish brown, light grey microtrichose dorsally and broadly lustrous in ventral third, 0.28 mm broad ventrally and 0.32 mm long, bare. Occiput blackish brown, microtrichose, sparsely black setose, setae rather thick and short, ventrally longer and finer. Antenna black, basal segment brown, length of antennal segments = 11: 12: 40: 19, setae on basal two segments up to 0.10 mm long. Labrum brown, lustrous, slightly longer than head is high. Palpus brown, short, covered with several short setae and one strong preapical seta (0.25 mm long). Gena narrow and lustrous, clypeus lustrous.

Thorax brownish-black, light grey microtrichose (without brownish-yellow tinge), darker stripes below rows of acrostichals and dorsocentrals scarcely visible. All setae black. Chaetotaxy: proepisternum with about 10 setae, both propleural depression and prosternum bare; acrostichals damaged but probably irregularly biserial and short; 8



Figures 1–4. Male terminalia, lateral view: **1** *Rhamphomyia* (*Pararhamphomyia*) *anfractuosa* Bezzi **2** *Rhamphomyia* (*Pararhamphomyia*) *biflexata* sp. n. **3** *Rhamphomyia* (*Pararhamphomyia*) *lineodorsata* sp. n. **4** *Rhamphomyia* (*Pararhamphomyia*) *multisinuosa* Frey.

uniserial dorsocentrals, 0.20 mm long in middle of rows, ending in 2 prescutellars, 1 short intrahumeral, 1 strong posthumeral; postpronotum with 1 strong seta and about 8–10 smaller setae; 3–4 notopleurals (1–2 setae on anterior part of notopleuron); 1 supraalar and 1 prealar; 1 long and 1 shorter postalar; 3 pairs of scutellars (outer pair short); laterotergite with black setae.

Legs brown, lustrous, black setose. Coxae blackish-brown, microtrichose (only hind coxa with lustrous spot anteriorly near base), black setose. One long seta in posteroapical comb of hind tibia. Fore femur with fine anteroventral setae $1/3$ as long as long as depth of femur, dorsal setae shorter, posteroventrals almost absent. Fore tibia with 4 posterodorsal setae slightly longer than width of tibia, ventral and anterodorsal setae short. Mid femur with two rows of spiny setae ventrally, anteroventrals half as long as depth of femur, posteroventrals longer than depth of femur, other setae short. Mid tibia with two rows of setae dorsally nearly $2\times$ longer than width of tibia (each row consists

of 3–4 setae including preapicals), two rows of ventral setae about as long as width of tibia. Hind femur (Fig. 14) with row of fine anteroventrals shorter than depth of femur, posteroventrals equally fine and irregularly arranged. Hind tibia with 5–6 pairs of antero- and posterodorsal setae 1.5× longer than width of tibia, ventral setae short. Basal tarsomeres of fore and mid legs thin and short setose, with several short ventral spines. Basal tarsomere of hind leg thin, with several setae dorsally 2× longer than diameter of tarsomere and with several ventral spines slightly longer than diameter of tarsomere.

Wing clear to light brownish, stigma slightly darker, veins brown, anal vein almost complete. Costal seta strong and long (several other setae between costal seta and wing base relatively strong and long), axillary angle acute. Measurements: $M_2/d = 1.3$, CuA_1 ratio = 1.6–1.7, $lw/ww = 2.5–3.1$. Halter yellow, calypter yellow with dark fringes.

Abdomen brown, lustrous (only segment 1 and part of segment 7 microtrichose as well as tips of thumb-like processes). All setae dark. Hind marginal setae on sides of tergites 2–4 nearly as long as their corresponding segments (discal setae shorter), on segments 5–6 shorter (but still slightly longer than discal setae), segment 7 very short setose. Dorsum of abdomen with short setae. Tergite 6 (Fig. 2) with two thumb-like processes on dorsal hind side; tergite 8 fused with sternite. Phallus (Fig. 9) with two outgoing loops; hypandrium membranose on posterior part (this part covers only part of ventral “ciliation” of phallus); phallus with hair-like “ciliation” ventrobasally and not much produced basal bulge.

Length: Body 5.5 mm, wing 5.5 mm.

Female. Unknown.

Remarks. *Rhamphomyia* (*Pararhamphomyia*) *biflexata* sp. n. may be easily identified according to the key. It is the only species with long dorsomarginal processes on tergite 6 (similarly as *R. shatalkini*) and simultaneously with two outgoing loops on phallus. Female is unknown with certainty, see also Remarks under *R. shatalkini* sp. n.

Distribution. Russia (Far East).

Dates of occurrence. July.

***Rhamphomyia* (*Pararhamphomyia*) *lineodorsata* sp. n.**


<http://zoobank.org/945770C7-B15A-40B5-B524-5B6548EEC453>

Figs 3, 10, 13

Type material. HOLOTYPE male: Russia, Amurskaja oblast, g. (= city) Zeja, 31.viii.1982, A. Shatalkin, deposited in ZMMU; PARATYPES: 2 males, same data as holotype; 1 male, same locality as holotype, 29.viii.1981, A. Shatalkin; paratypes depositories: ZMMU, CULSP.

Additional material examined (excluded from type series): 2 females, same locality as holotype, 4.ix.1981; 1 female, same locality as holotype, 13.ix.1981 – all A. Shatalkin (ZMMU and CULSP).

Diagnosis. Species of the subgenus *Pararhamphomyia* with phallus forming loops in space, dark brownish black mesoscutum with darker stripes below rows of setae and



Figures 5–8. Male terminalia, lateral view: **5** *Rhamphomyia* (*Pararhamphomyia*) *nudiscutellata* sp. n. **6** *Rhamphomyia* (*Pararhamphomyia*) *robustior* Frey **7** *Rhamphomyia* (*Pararhamphomyia*) *shatalkini* sp. n. **8** *Rhamphomyia* (*Pararhamphomyia*) sp. 2.

uniserial dorsocentrals. Female tergites 6 and 7 lustrous and hind tibia without penate ciliation dorsally.

Etymology. The name of the species is derived from dark stripes on the mesoscutum, differing from closely allied species, *R. robustior* Frey.

Description. Male. Head holoptic, facets in dorsal half of eye enlarged. Frons blackish brown, light grey microtrichose, without setae. Ocellar setae one third as long as frons, black, ocellar triangle with 1–2 pairs of additional setae. Face blackish brown, light grey microtrichose, 0.30 mm broad ventrally and equally long, without setae. Occiput blackish brown, microtrichose, sparsely black setose, setae rather thick and short, longer and finer ventrally. Antenna black, scape and pedicel brown, length of antennal segments = 16–18: 12: 48–50: 16, setae on basal two segments up to 0.12 mm long. Labrum brown, lustrous, slightly longer than height of head. Palpus brown, short, covered with several setae and one strong preapical seta (0.20 mm long). Gena narrow and lustrous, clypeus microtrichose.

Thorax brownish-black, rather dark brownish grey microtrichose, scutum with distinct darker stripes below rows of acrostichals and dorsocentrals (best visible in

posterior view). All setae black. Chaetotaxy: proepisternum with about 10 setae, both propleural depression and prosternum bare; 10–16 narrowly biserial (anteriorly almost uniserial), fine acrostichals about 0.20 mm long; 8–11 almost regularly uniserial dorsocentrals slightly longer (about 0.30 mm in middle of rows), ending in 2–3 long prescutellars, 1–2 strong intrahumerals, 1 strong posthumeral; postpronotum with 2–3 long and 10–15 gradually shorter setae; 4 notopleurals (2–3 long setae on anterior part of notopleuron); 1 long supraalar and 2–3 smaller prealar; 1 long and 1 shorter postalar; 3 pairs of scutellars; laterotergite with black setae.

Legs brown, microtrichose, black setose. Coxae blackish-brown, microtrichose (only hind coxa with lustrous spot anteriorly near base), black setose. One long seta present in comb at tip of hind tibia. Fore femur with fine anteroventral setae up to half as long as depth of femur, dorsal setae shorter, posteroventrals up to half as long as depth of femur, present mostly only on proximal half. Fore tibia with 4–6 posterodorsal setae about as long as width of tibia, ventral and anterodorsal setae short. Mid femur with two rows of spiny setae ventrally, anteroventrals one third as long as depth of femur, posteroventrals slightly longer than depth of femur, other setation short. Mid tibia with only two anterodorsal setae (one subbasal and one preapical – but holotype on one leg atypically with three such setae), and 4–6 posterodorsals slightly longer than depth of tibia, two rows of ventral setae somewhat shorter than width of tibia (several posteroventrals may be longer than remaining ones). Hind femur (Fig. 13) with ventral microtrichosity up to 0.04 mm long, with rather fine anteroventral setae about as long as depth of femur in basal half and sometimes apically, very short to absent on third quarter of femur, posteroventrals present only on extreme base of femur. Hind tibia with 3–5 antero- and 6–8 posterodorsal setae slightly longer than width of tibia, ventral setae short. Basal tarsomeres of fore and mid legs thin and short setose, mid one with several short ventral spines, basal tarsomere of hind leg slightly swollen, with several setae dorsally and spine like setae ventrally up to 2× longer than diameter of tarsomere.

Wing light brown, stigma slightly darker, veins brown (yellowish in basal part of wing), anal vein almost complete or indistinct about middle. Costal seta present, axillary angle acute. Measurements: $M_2/d = 1.4\text{--}1.6$, CuA_1 ratio = $1.4\text{--}2.1$, $lw/ww = 2.7\text{--}3.1$. Halter yellow, calypter brownish-yellow with dark fringes.

Abdomen brown, dark brown microtrichose (dark brown in both lateral and dorsal views), setae all dark. Hind marginal setae almost as long as corresponding segments, discal setae shorter. Dorsum of abdomen with short setae. Phallus (Figs 3, 10) with three twists in space.

Length: Body 4.5–5.3 mm, wing 5.1–5.8 mm.

Female. Head dichoptic, frons approximately 0.35 mm long and 0.25 mm wide, subparallel, with two rows of 5–7 relatively long setae on sides. Face approximately 0.25 mm long and subequally wide in middle (slightly broadening ventrally), without setae. Palpus lighter than in male, yellowish red. Thorax as in male, only setae shorter. Fore femur with anterodorsal row of almost pennate setae slightly shorter than depth of femur, with anteroventral row of thin setae as long as depth of femur and with posteroventral



Figures 9–12. Male genitalia (macerated), lateral view: **9** *Rhamphomyia* (*Pararhamphomyia*) *biflexata* sp. n. **10** *Rhamphomyia* (*Pararhamphomyia*) *lineodorsata* sp. n. **11** *Rhamphomyia* (*Pararhamphomyia*) *nudiscutellata* sp. n. **12** *Rhamphomyia* (*Pararhamphomyia*) *shatalkini* sp. n.

row of pennate setae slightly longer than depth of femur. Fore tibia as in male, only posterodorsal setae less differentiated. Mid femur with both (antero)dorsal and (postero)ventral pennation about as long as depth of femur. Mid tibia with short subpennate ciliation both dorsally and ventrally in addition to several slightly longer setae on both sides. Hind femur with long pennate ciliation ventrally in addition to several setae and shorter dorsal subpennate ciliation. Hind tibia with two rows of dorsal setae slightly longer than width of tibia, ventral setae short and slightly subpennate. Wing light brown as in male, measurements: $M_2/d = 1.4\text{--}1.6$, CuA_1 ratio = $1.5\text{--}1.9$, $lw/ww = 2.8\text{--}3.0$. Abdomen microtrichose, with tergites 6 to 8 and sternite 8 lustrous and sternite 7 sublustrous. Hind marginal setae on segments 2–6 $2/3$ as long as corresponding segments.

Length: Body 5.3–5.8 mm, wing 5.5–6.0 mm.

Remarks. *Rhamphomyia* (*Pararhamphomyia*) *lineodorsata* sp. n. is closely allied to remaining three Palaearctic species of *R. anfractuosa* group of species with phallus twisted in space, viz *R. anfractuosa* Bezzi, *R. nudiscutellata* sp. n. and *R. robustior*. Most

specimens of this complex examined differ from specimens of *R. multisinuosa* complex (with phallus twisted in a single plain) by absence of submedian anterodorsal setae on mid tibia in addition to characters given in the key. However, as mentioned above, holotype of the above described species has atypically one such seta present on one leg. We excluded females from type series because of problems with exact identification of females in this group of species and because we had no pairs taken in copula. But we believe we identified them properly. See under *R. nudiscutellata* for discussions of the differences between females of this complex.

Distribution. Russia (Far East).

Dates of occurrence. August–September.

***Rhamphomyia* (*Pararhamphomyia*) *nudiscutellata* sp. n.**

<http://zoobank.org/BB33F445-E025-4126-8C47-007857B89D56>

Figs 5, 11, 15


Type material. HOLOTYPE male: Russia, Amurskaja oblast, g. (gorod = city) Zeja, 14.ix.1981, A. Shatalkin, deposited in ZMMU; PARATYPES: 2 males, same data as holotype; 4 males, Tuva, okr. (=region) Baj-Chaaka, Berezovka, 5.ix.1973, V. Kovalev; 2 males, same locality, 3.ix.1973, V. Kovalev; 2 males, same locality, 7.ix.1973, V. Kovalev; 2 males, Tuva, okr. Saryg Sep, listvennicznik na granice lesa (= larch on forest boundary), 28.viii.1973, V. Kovalev; 1 male Chitin, r. (=river) Kuenga, vyche (= above) Chernyshevskaja, 24.viii.1977, V. Kovalev. Paratypes depositories: ZMMU, CULSP.

Additional material examined (excluded from type series): 2 females, Chitin, r. (reka = river) Kuenga, vyche Chernyshevskaja, 26.viii.1977, V. Kovalev; 1 female, Tuva, okr. (= region) Baj-Chaaka, Berezovka, 5.ix.1973, V. Kovalev; 1 female, same locality, 2.ix.1973, V. Kovalev; 2 females, Tuva, okr. (= region) Shagonar, Ishtii-Khem, 21.viii.1973, V. Kovalev; 1 female, same locality, 24.viii.1973, V. Kovalev; (ZMMU and CULSP).

Diagnosis. Species of the subgenus *Pararhamphomyia* with phallus forming loops in space, light grey mesoscutum and uniserial dorsocentrals. Female front femur with two rows of ventral setae longer than depth of femur, hind tibia not pennate, wing light brownish and tergites 6 and 7 lustrous.

Etymology. The name of the species is derived from the relatively naked scutellum bearing only four setae (*nudus*, Latin = naked).

Description. Male. Head holoptic, facets in dorsal half of eye enlarged. Frons blackish brown, light grey microtrichose, without setae. Ocellar setae one third as long as frons, black, ocellar triangle with 1–2 pairs of additional setae. Face blackish brown, light grey microtrichose, 0.20 mm broad ventrally and 0.25 mm long, without setae. Occiput brownish black, rather light grey microtrichose, black setose. Antenna black, scape and pedicel brown, length of antennal segments = 15–16: 11: 45–50: 12–14, setae on basal two segments about 0.10 mm long. Labrum brown, lustrous, slightly shorter than head is high. Palpus brown, short, covered with several rather long setae



Figures 13–16. Male hind legs (femur, tibia and basitarsus): **13** *Rhamphomyia* (*Pararhamphomyia*) *lineodorsata* sp. n. **14** *Rhamphomyia* (*Pararhamphomyia*) *biflexata* sp. n. **15** *Rhamphomyia* (*Pararhamphomyia*) *nudiscutellata* sp. n. **16** *Rhamphomyia* (*Pararhamphomyia*) *shatalkini* sp. n.

(0.20 mm long), preapical seta poorly differentiated. Gena narrow and lustrous, clypeus microtrichose.

Thorax brownish-black, rather light grey microtrichose, scutum without stripes, only in immature specimens with poorly visible darker stripes below rows of acros-tichals and dorsocentrals. All setae black. Chaetotaxy: proepisternum with 5–8 setae, both propleural depression and prosternum bare; 6–10 biserial, short and fine acros-tichals (about 0.15 mm long); 7–10 regularly uniserial dorsocentrals (about 0.25 mm in middle of rows), ending in 2 long prescutellars, 1–2 fine and long intrahumerals, 1 strong posthumeral; postpronotum with 2–3 long and 6–10 gradually shorter setae; 3 notopleurals (0–2 long setae on anterior part of notopleuron); 1 long supraalar, prealar absent; 1 long and 1 shorter postalar; 2 pairs of scutellars; laterotergite with black setae.

Legs brown, microtrichose, black setose. Coxae blackish-brown, microtrichose (only hind coxa with lustrous spot anteriorly near base), black setose. One long seta present in comb at tip of hind tibia. Fore femur with complete rows of fine antero- and posteroventral setae up to 2/3 as long as long as depth of femur, dorsal setae shorter. Fore tibia short setose, without differentiated setae except preapical. Mid femur with two rows of spiny setae ventrally, anteroventrals up to one-third as long as depth of femur, posteroventrals slightly longer than depth of femur, other setation short. Mid

tibia with only two anterodorsal setae (one short subbasal and one long preapical), and 3–4 posterodorsals slightly longer than depth of tibia, two rows of ventral setae somewhat shorter than width of tibia (several posteroventrals may be longer than remaining ones). Hind femur (Fig. 15) with ventral microtrichosity up to 0.05 mm long, with rather fine anteroventral setae about as long as depth of femur in basal half and in some specimens apically, very short to absent on third quarter of femur, posteroventrals present only on extreme base of femur. Hind tibia with 3–4 antero- and 5–6 posterodorsal setae slightly longer than width of tibia, ventral setae short. Basal tarsomeres of fore and mid legs thin and short setose, mid basal tarsomere with several short ventral spines; basal tarsomere of hind leg slightly swollen, with several setae dorsally and spine like setae ventrally up to 2× longer than diameter of tarsomere.

Wing clear, stigma scarcely darker, veins brown and yellowish in basal part of wing, anal vein indistinct about middle. Costal seta present, axillary angle right. Measurements: $M_2/d = 1.2\text{--}1.7$, CuA_1 ratio = 1.6–1.9, $lw/ww = 2.7\text{--}3.3$. Halter yellow, calypter yellow with dark fringes.

Abdomen brown, entirely light grey microtrichose (light grey from both lateral and dorsal views), setae all dark. Hind marginal setae on tergites 2–3 somewhat longer and on tergites 4–6 as long as corresponding segments, discal setae shorter. Dorsum of abdomen with short setae. Phallus (Figs 5, 11) with three twists in space.

Length: Body 3.8–4.0 mm, wing 4.5–5.8 mm.

Female. Head dichoptic, frons approximately 0.25 mm long and 0.20 mm wide, subparallel, with two rows of 4–6 relatively long setae on sides. Face approximately 0.20 mm long and subequally wide in middle (strongly divergent ventrally), without setae. Palpus brown as in male. Thorax as in male. Fore femur with both antero- and posteroventral rows of setae as long as depth of femur, posteroventrals mostly thin but in some specimens on one or both legs thickened – almost pennate, dorsal ciliation short and thin. Fore tibia as in male, only posteroventral setae slightly differentiated. Mid femur with both (antero)dorsal and (postero)ventral pennation about as long as depth of femur. Mid tibia short setose, most specimens with several antero- and posteroventral setae and/or several setae dorsally shorter than depth of tibia. Hind femur with long pennate ciliation ventrally in addition to several setae and shorter dorsal subpennate ciliation. Hind tibia slightly flattened, with two rows of dorsal setae slightly longer than width of tibia, ventral setae short, short ciliation slightly subpennate. Wing clear with only indistinct brownish tinge, measurements: $M_2/d = 1.3\text{--}1.6$, CuA_1 ratio = 1.6–1.8, $lw/ww = 2.8\text{--}3.0$. Abdomen microtrichose, with tergites 6 to 8 and sternites 7 and 8 lustrous. Hind marginal setae on segments 2–4 as long as corresponding segments, on segments 5–7 2/3 as long as corresponding segments.

Length: Body 4.0–4.4 mm, wing 4.6–5.2 mm.

Remarks. *Rhamphomyia* (*Pararhamphomyia*) *nudiscutellata* sp. n. is closely allied to the remaining three Palaeartic species of the *R. anfractuosa* group of species with phallus twisted in space, viz *R. anfractuosa*, *R. lineodorsata* sp. n. and *R. robustior*. All four species may be identified according to the key. We excluded females from the type series because of problems with exact identification of females in this group of species

and because we had no pairs taken in copula. But we believe we identified them properly. Female differs from *R. robustior* and *R. anfractuosa* in lustrous abdominal tergites 6 and 7 (microtrichose in both *R. anfractuosa* and *R. robustior*), from *R. lineodorsata* by brown palpus (yellowish red in *R. lineodorsata*) and from *R. anfractuosa* also by front femur with two rows of ventral setae longer than depth of femur, hind tibia without broad pennate ciliation and light brownish wing (in *R. anfractuosa*, front femur has almost no ventral setae, hind tibia is broadly pennate both dorsally and ventrally and wing is deep brown).

Distribution. Russia (Far East).

Dates of occurrence. August–September.

***Rhamphomyia (Pararhamphomyia) shatalkini* sp. n.**

<http://zoobank.org/11ADE0FE-A4E1-48FB-BAF6-1923EBE2BC2A>

Figs 7, 12, 16

Type material. HOLOTYPE male: Russia, Amurskaja oblast, g. (= city) Zeja, 22.vi.1978, leg. A. Shatalkin, deposited in ZMMU; PARATYPES: 2 males, same data as holotype; 2 males, same locality, 23.vi.1978; 2 males, same locality, 24.vi.1978; 1 male, same locality, 21.vi.1979 – all A. Shatalkin; 1 male, same locality, 25.vi.1982, A. Ozerov; 3 males, Russia, Juzhnoje Primorije, Kamenushka, 9.vi.1984, A. Shatalkin; 1 male, Russia, Irkutskaja o. (= oblast, = region), Listvjanka, 21.vi.1965, O.P. Negrobov; paratypes depositories: ZMMU, CULSP.

Diagnosis. Species of the subgenus *Pararhamphomyia* with uniserial dorsocentrals, lustrous abdomen, tergite 6 with two thumb-like dorsomarginal processes, mesoscutum without lustrous stripes and phallus with four outgoing loops.

Etymology. The species is named after Anatole Shatalkin, dipterist from Moscow Museum and collector of part of type series.

Description. Male. Head holoptic, facets in dorsal third of eye enlarged. Frons blackish brown, light grey microtrichose, without setae. Ocellar setae fine, half as long as frons, black, ocellar triangle without additional setae. Face blackish brown, light grey microtrichose dorsally and broadly lustrous along ventral margin, 0.25 mm broad ventrally and 0.30 mm long, without setae. Occiput blackish brown, grey microtrichose, sparsely black setose, setae rather thick and short, ventrally longer and finer. Antenna black, scape brown, pedicel and extreme base of postpedicel brownish-yellow, length of antennal segments = 13: 10: 40: 17, setae on basal two segments nearly 0.12 mm long. Labrum brown, lustrous, about as long as or slightly longer than height of head. Palpus brown, short, with several setae almost 0.30 mm long. Gena narrow and lustrous, clypeus lustrous.

Thorax brownish-black, light grey microtrichose (with slight brownish-yellow tinge), scutum with somewhat darker stripes below rows of acrostichals and dorsocentrals. All setae black. Chaetotaxy: proepisternum with 10–15 setae, both propleural depression and prosternum bare; 14–20 irregularly biserial, fine acrostichals about 0.20

mm long; almost regularly uniserial slightly longer dorsocentrals (about 0.25 mm in middle of rows), ending in 2–3 strong and long prescutellars, 1 small intrahumeral, 1 strong posthumeral; postpronotum with 1 strong seta and about 15 additional finer setae; 4 notopleurals (1–2 long setae on anterior part of notopleuron); 1 supraalar and 1 equally strong prealar; 1 long and 1 shorter postalar; 3 pairs of scutellars (rarely two pairs); laterotergite with black setae.

Legs brown, lustrous, black setose. Coxae blackish-brown, microtrichose (only hind coxa with two lustrous spots anteriorly near base and at apex), black setose. One long seta in posteroapical comb of hind tibia. Fore femur with fine anteroventral setae $1/3$ as long as depth of femur, posteroventral and dorsal setae shorter. Fore tibia with 4–5 strong posterodorsal setae $2\times$ longer than width of tibia, ventral and anterodorsal setae short. Mid femur with two rows of spiny setae ventrally, anteroventrals half as long as depth of femur and densely arranged, posteroventrals sparse and longer than depth of femur, other setation short. Mid tibia with two rows of setae dorsally nearly $2\times$ longer than width of tibia (each row consists of 4–5 setae), row of short posteroventral setae, anteroventrals more irregularly arranged and somewhat longer than posteroventrals. Hind femur (Fig. 16) with anteroventral row of rather fine setae nearly as long as depth of femur (1 or 2 of them may be stronger than remaining), other setae including posteroventrals short and fine. Hind tibia about as thick as hind femur, with 6–8 pairs of antero- and posterodorsal setae $1.5\times$ longer than width of tibia, ventral setae short. Basal tarsomeres of fore and mid legs thin and short setose, with several short ventral spines. Basal tarsomere of hind leg thin, with several setae dorsally $2\times$ longer than width of tarsomere and with several ventral spines slightly longer than width of tarsomere.

Wing light brownish, stigma slightly darker, veins brown, anal vein almost complete. Costal seta strong and long (several other setae between costal seta and wing base relatively strong and long), axillary angle acute. Measurements $M_2/d = 1.3\text{--}1.4$, CuA_1 ratio = $2.2\text{--}2.5$, $lw/ww = 2.6\text{--}3.0$. Halter yellow, calypter brownish-yellow with dark fringes. Abdomen brown, lustrous (only segment 1 and small spots on 3 pregenital segments microtrichose). All setae dark. Hind marginal setae on sides of tergite 2 nearly as long as segment, on segments 3–5 gradually shorter (discal setae shorter), marginals on tergite 6 short, tergite 7 bare. Dorsum of abdomen with short setae. Abdominal tergite 6 with two thumb-like processes dorsally (Fig. 7). Phallus (Fig. 12) with four outgoing loops in a single plain; phallus with hair-like “ciliation” ventrobasally and produced basal bulge; hypandrium membranose on posterior part (this part covers whole ventrobasal “ciliation” of phallus).

Length: Body 5.3–6.4 mm, wing 5.7–6.4 mm.

Female. Unknown.

Remarks. *Rhamphomyia* (*Pararhamphomyia*) *shatalkini* sp. n. may be easily distinguished from all other Palaeartic species of *Rhamphomyia* (except unnamed species *R.* sp. 1) by peculiar shape of phallus forming four outgoing loops in a single plain and simultaneously tergite 6 bearing two long thumb-like dorsomarginal processes. However, the mesoscutum in the new species is entirely microtrichose but with three lustrous stripes below lines of setae in *R.* sp. 1. Other species with similarly formed

phallus are *R. multisinuosa* Frey and *R. spectabilis* Frey, both without long thumb-like processes on tergite 6. Female of *R. shatalkini* remains unknown with certainty. We have at our disposal several females which may belong to either *R. shatalkini*, *R. spectabilis* or *R. biflexata*, but we are unable to associate them with particular males. Males of all three species have very similar microtrichosity pattern of mesoscutum which otherwise helps to associate males with females even if not taken in copula. These females differ from all other Palearctic *Pararhamphomyia* by the following combination of characters: body entirely dark setose, dorsocentrals almost regularly uniserial, halter yellow, clypeus lustrous, both mid and hind femora broadly pennate, tibiae without pennation and abdomen lustrous except the first segment.

Distribution. Russia (Far East).

Dates of occurrence. June.

Unnamed species

Rhamphomyia (*Pararhamphomyia*) sp. 1

Material examined. 1 male, Russia, Amurskaja oblast, g. (= city) Zeja, 26.vi.1982, M. Krivosheina, deposited in ZMMU.

Remarks. Species very similar to *R. shatalkini* sp. n. with the exception of characters given in the key. We hesitate to describe new species from only a single specimen.

Rhamphomyia (*Pararhamphomyia*) sp. 2

Material examined. 1 male, Russia, Amurskaja oblast, g. (= city) Zeja, 26.vii.1978, A. Shatalkin, deposited in ZMMU.

Remarks. Species very similar to *R. spectabilis* Frey with the exception of characters given in the key. We hesitate to describe new species from only a single specimen; moreover, we do not have *R. spectabilis* at our disposal.

Key to males of Palearctic species of the *Rhamphomyia anfractuosa* group

- | | | |
|-------|---|----------|
| 1 | Phallus forms loops in space. Abdomen microtrichose. Hind femur with ventral microtrichosity. No modifications of tergites 6 and 7. No lustrous stripes below rows of acrostichals and dorsocentrals..... | 2 |
| – | Phallus forms loops in flat plain. Abdomen lustrous. Hind femur without ventral microtrichosity. Tergites 6 and/or 7 modified. Lustrous stripes below rows of acrostichals and dorsocentrals present or absent..... | 5 |
| 2 (1) | Mesoscutum dark brownish black. Abdomen dark brown. Usually three (rarely four) pairs of scutellar setae. Axillary angle acute..... | 3 |

- Mesoscutum light grey microtrichose. Abdomen light grey. Usually two (rarely one or three) pairs of scutellar setae. Axillary right angled **4**
- 3 (2) Mesoscutum with distinct darker stripes below acrostichals and dorsocentrals. Dorsocentrals regularly uniserial (Additional characters opposite of *R. nudiscutellata*: fore tibia with differentiated posterodorsal setae, wing light brown). (Additional characters: female tergites 6 and 7 lustrous and hind tibia without pennate ciliation dorsally, palpus yellowish red). Phallus as in Fig. 3 ***R. lineodorsata* sp. n.**
- Mesoscutum subpolished, without distinct stripes. Dorsocentrals irregularly biserial. (Additional characters: female tergites 6 and 7 microtrichose and hind tibia with broad pennate ciliation dorsally). Phallus as in Fig. 6 ***R. robustior* Frey**
- 4 (2) Hind marginal setae on tergites 5–6 nearly absent. Wings brown. Antennal stylus slightly shorter than postpedicel. Phallus with five twists (Fig. 1). (Additional characters: female front femur nearly without setae, hind tibia broadly pennate both dorsally and ventrally, wing deep brown and tergites 6 and 7 microtrichose) ***R. anfractuosa* Bezzi**
- Hind marginal setae on tergites 5–6 about as long as these segments. Wings clear. Antennal stylus 1/4 as long as postpedicel. Phallus with three twists (Fig. 5). (Additional characters opposite of *R. lineodorsata*: fore tibia without differentiated posterodorsal setae, wing clear) (Additional characters: female fore femur with two rows of ventral setae longer than depth of femur, hind tibia without pinnate setae, wing light brownish and tergites 6 and 7 lustrous) ***R. nudiscutellata* sp. n.**
- 5 (1) Phallus with two outgoing loops (Fig. 2) ***R. biflexata* sp. n.**
- Phallus with 3 or 4 outgoing loops **6**
- 6 (5) Tergite 6 with two long thumb-like dorsomarginal processes **7**
- Tergite 6 without processes or with only small triangular shaped dorsomarginal projections **8**
- 7 (6) Mesoscutum microtrichose, without lustrous stripes. Dorsocentrals uniserial. Syntergosternite 8 without dorsomarginal projections. Phallus as in Fig. 7 ... ***R. shatalkini* sp. n.**
- Mesoscutum with lustrous stripes below rows of acrostichals and dorsocentrals. Dorsocentrals irregularly biserial. Syntergosternite 8 with two dorsomarginal triangular projections ***R. sp. 1***
- 8 (6) Tergite 6 with two small triangular shaped dorsomarginal projections. Mesoscutum with lustrous stripes below rows of acrostichals and dorsocentrals. Phallus as in Fig. 4 ***R. multisinuosa* Frey**
- Tergite 6 without projections. Mesoscutum microtrichose, without lustrous stripes **9**
- 9 (8) Dorsocentrals irregularly biserial. Phallus with four outgoing loops ***R. spectabilis* Frey**
- Dorsocentrals uniserial. Phallus with three outgoing loops (Fig. 8) ... ***R. sp. 2***

Acknowledgements

This paper was supported by S grant of MSMT (Ministry of Education, Sports and Youth). Our special thanks are due to anonymous reviewers for improving the manuscript.

References

- Barták M (1982) The Czechoslovak species of *Rhamphomyia* (Diptera, Empididae), with description of a new species from Central Europe. Acta Universitatis Carolinae - Biologica 1980 (1982): 381–461.
- Barták M (2002) Nearctic species of *Rhamphomyia* subgenus *Megacyttarus* (Diptera: Empididae). Acta Universitatis Carolinae Biologica 46(1-2): 3–215.
- Barták M (2003) Revision of Palaearctic species of *Rhamphomyia* (*Megacyttarus*) *argentea* group (Diptera: Empididae). Acta Universitatis Carolinae Biologica 47: 197–245.
- Barták M (2007) Five new European species of the *Rhamphomyia* (s.str.) *albosegmentata* group (Diptera: Empididae). Revue Suisse de Zoologie 114(2): 417–435.
- Barták M, Çiftçi MC, Hasbenli A (2007) A new species of *Rhamphomyia* (s. str.) Meigen (Diptera, Empididae) from southern Anatolia. Entomological News 118(2): 143–147. doi: 10.3157/0013-872X(2007)118[143:ANSORS]2.0.CO;2
- Barták M, Kubík Š (2008a) New peculiar Eastern Palaearctic *Rhamphomyia* (Diptera: Empididae). Entomological News 119(4): 338–344. doi: 10.3157/0013-872X-119.4.338
- Barták M, Kubík Š (2008b) A new species of *Rhamphomyia* (*Pararhamphomyia*) (Diptera: Empididae) from Thailand. Oriental Insects 42: 285–289. doi: 10.1080/00305316.2008.10417552
- Barták M, Kubík Š (2008c): Four new West Palaearctic species of the *Rhamphomyia* (s.str.) (Diptera: Empididae). Revue Suisse Zool. 115(1): 25–36.
- Barták M, Kubík Š (2009) Two new East Palaearctic *Rhamphomyia* (*Pararhamphomyia*) (Diptera: Empididae). Entomological News 120(1): 76–86. doi: 10.3157/021.120.0114
- Barták M, Kubík Š (2010) Three new European species of the *Rhamphomyia* (s.str.) *melania* group (Diptera: Empididae). Revue Suisse de Zoologie 117(1): 89–100.
- Barták M, Kubík Š (2012) A review of the Palaearctic species of *Rhamphomyia* subgenus *Holoclera* (Diptera: Empididae) with description of 5 new species. Revue Suisse de Zoologie, roč. 119(3): 385–407.
- Barták M, Kubík Š, Civelek H, Dursun O (2014) New species of *Rhamphomyia* (Diptera: Empididae) from Turkey with a key to species of the Middle East and adjacent territories. Zootaxa 3815(1): 68–78. doi: 10.11646/zootaxa.3815.1.4
- Barták M, Sinclair B (2003) Case 3269. *Rhamphomyia* (*Rhamphomyia*) Meigen, 1822 and *Rhamphomyia* (*Pararhamphomyia*) Frey, 1922 (Insecta: Diptera): proposed conservation of usage of the subgeneric names by designation of *Empis sulcata* Meigen, 1804 as the type species of *Rhamphomyia*. Bulletin of Zoological Nomenclature 60(3): 203–205. [See Opinion 2117, Bulletin of Zoological Nomenclature 62(2): 114–115]

- Merz B, Haenni J-P (2000) Morphology and terminology of adult Diptera. In: Papp L, Darvas B (Eds) Contributions to a Manual of Palaeartic Diptera. Volume 1. Science Herald, Budapest, Hungary, 21–51.
- Saigusa T (2012) A new Asio–Nearctic subgenus of *Rhamphomyia* (Diptera: Empididae: Empidinae). The Canadian Entomologist 144(2): 291–322. doi: 10.4039/tce.2012.28
- Sinclair BJ (2000) Morphology and terminology of Diptera male terminalia. In: Papp L, Darvas B (Eds) Contributions to a Manual of Palaeartic Diptera. Volume 1. Science Herald, Budapest, Hungary, 53–84.
- Sinclair BJ, Cumming JM (2006) The morphology, higher-level phylogeny and classification of the Empidoidea (Diptera). Zootaxa 1180: 1–172.
- Yang D, Zhang K, Yao G, Zhang J (2007) World Catalog of Empididae (Insecta: Diptera). China Agricultural University Press, Beijing, 599 pp.