

# Young Diagrams, Standard Monomials and Invariant Theory

Claudio Procesi

The topics I want to treat have various sources: (1) the fundamental theorems of invariant theory of the classical groups, (2) the representation theory of classical groups, (3) the geometry of the Grassmann and flag varieties.

The initial interest in these questions came for me from the solution given by Formanek and myself, [8], of Mumford's conjecture for the linear group, [25], obtained independently from the general solution due to Haboush [9].

Our approach consisted in relating the geometric reductivity of the group  $\mathrm{Gl}(V)$  with a property of the symmetric group  $S_m$  acting on the tensor space  $V^{\otimes m}$ .

In fact, in characteristic zero, one classical approach to the representation theory of  $\mathrm{Gl}(V)$ , and in particular to the proof that it is linearly reductive, is the following [27], [29]:

The algebras of operators spanned by the symmetric group and by  $\mathrm{Gl}(V)$  on  $V^{\otimes m}$  are each the full centralizer of the other. In characteristic zero then one can apply the theory of semisimple algebras.

In positive characteristic, if we consider the algebra  $\Sigma_m$  spanned by  $S_m$  on  $V^{\otimes m}$ , we have that  $\Sigma_m$  is no more semisimple but nevertheless the limit  $\Sigma = \lim_{m \rightarrow \infty} \Sigma_m$  is semisimple in the sense of Jacobson and this suffices to prove the geometric reductivity of  $\mathrm{Gl}(V)$ .

This analysis led naturally to the question of extending the classical relations between  $\mathrm{Gl}(V)$  and  $S_m$  to all characteristics; in particular to prove that  $\Sigma_m = \mathrm{End}_{\mathrm{Gl}(V)}(V^{\otimes m})$ .

It was well known that this equality is a simple consequence of a statement in invariant theory, the so-called first fundamental theorem, to which it is equivalent

in characteristic zero. Similar questions arise for the other classical groups which require a similar (or sometimes more complicated) analysis [29].

The relation with invariant theory comes from the identification  $\text{End}(V^{\otimes m}) \cong (V^{\otimes m} \otimes V^{*\otimes m})^*$  and hence the identification of  $\text{End}_{\text{Gl}(V)}(V^{\otimes m})$  with the multilinear invariant polynomials in  $m$  vector and  $m$  covector variables.

It turns out, in positive characteristic, that it is possible to study first all the invariant polynomials and then deduce the form of the multilinear ones.

**1. Fundamental theorems of classical invariant theory.** Let us state, in a form slightly different from the usual one, the fundamental theorems of invariant theory of the classical groups, (see [29] for the theory in char 0).

(i)  $G = \text{Gl}(n, k)$ . Consider multiplication of matrices  $\pi: M_{k,n} \times M_{n,h} \rightarrow M_{k,h} \{(A, B) \rightarrow AB\}$ , with image the variety  $V_n$  of  $k \times h$  matrices of rank  $\leq n$ .

**FIRST FUNDAMENTAL THEOREM.** *The coordinate ring of  $V_n$  is the ring of invariants of  $G$  acting on  $M_{k,n} \times M_{n,h}$  by  $X \cdot (A, B) = (AX^{-1}, XB)$ .*

**SECOND FUNDAMENTAL THEOREM.** *The ideal of functions on  $M_{k,h}$  vanishing on  $V_n$  is generated by the  $n+1 \times n+1$  minors.*

We should remark that there exists now in characteristic 0 a beautiful theory of the higher syzygies due to A. Lascoux [15], [22], [23]. Similar theorems hold for the other classical groups.

These theorems, with the exception of the very last, are all classical in characteristic zero and can be found in H. Weyl's book [29].

The second fundamental theorems and various qualitative results on the determinantal varieties have been the object of intensive study by many authors both in positive and zero characteristics [10], [11], [21], [26].

In a joint paper with De Concini [5] we attacked all these problems in the spirit of Igusa's proof of the projective normality for the Grassmann variety [14], again a classical theorem of invariant theory for  $\text{Sl}(n, K)$ . His approach was through a careful analysis of the projective coordinate ring of the Grassmann variety. Such ring, as the one for the flag variety, were always fundamental objects in invariant theory [1] (the classical primary covariants of the Capelli–Deruyts expansion), the explicit bases (standard bases) for such rings were popularized by Hodge [12], [13] (although they seem to have been known to Young [30]).

They give, in algebra, very explicit descriptions of the irreducible representations of the linear groups and the symmetric group (in characteristic zero), in geometry a very thorough understanding of the cellular decomposition of the Grassmann and flag varieties (by Schubert cells).

In char 0 the theory is based on the fact that the irreducible polynomial representations of  $\text{Gl}(V)$  can be indexed, as  $L_\sigma(V)$ , by partitions  $\sigma$ . One has the classical

plethysm formulas:

$$S[V \otimes W] = \bigoplus_{\sigma} L_{\sigma}(V) \otimes L_{\sigma}(W),$$

$$S[S^2V] = \bigoplus L_{\sigma}(V), \sigma \text{ has even columns,}$$

$$S[A^2V] = \bigoplus L_{\sigma}(V), \sigma \text{ has even rows}$$

(nevertheless the space  $L_{\sigma}(V)$  is defined over  $Z!$ ).

**2. Standard bases.** If  $Y=(y_{ij})$  is an  $n \times m$  matrix, we indicate by  $p=(i_k \dots i_2 i_1 | j_1 j_2 \dots j_k)$  the determinant of the minor with rows  $i_i$ 's, columns  $j_u$ 's. If  $p_1, p_2, \dots, p_k$  are minors of size  $\sigma: h_1 \geq h_2 \geq \dots \geq h_k$ , we display their product  $M$  as a double tableau with rows the  $p_i$ 's. We say that  $M$  is a standard monomial (of shape  $\sigma$ ) if the indices are strictly increasing on each row, non decreasing on each column, (separately on the right and on the left) [4], [7]. A similar definition holds for  $Y$  symmetric or antisymmetric.

**THEOREM.** *The standard monomials (in each case) are a  $Z$  basis of  $Z[y_{ij}]$ . (Cf. [4], [7], [12].)*

The theorems stated are very convenient to study the coordinate rings of the various determinantal varieties appearing in the fundamental theorems of invariant theory, in fact such theorems can be proved in a characteristic free way [5], precisely by using such standard bases.

In fact the previous form of the theorems contains explicit algorithms, by use of quadratic equations, to express a non standard product in terms of standard ones.

To a partition  $\sigma: k_1 \geq k_2 \geq \dots \geq k_r$  associate the dual partition  $\check{\sigma}: h_1 \geq h_2 \geq \dots \geq h_l$  with  $h_i = \{\#\{j | k_j \geq i\}$  and a sequence  $\gamma_i(\sigma) = \sum_{j \cong i} h_j$ ; then set  $\sigma \geq \tau$  if  $\gamma_i(\sigma) \geq \gamma_i(\tau)$  for all  $i$ .

A filtration of  $Z[y_{ij}]$  is defined setting  $A_{\sigma}$  = span of all standard monomials of shape  $\geq \sigma$ . It has the following geometric interpretation [3]:  $A_0$  is exactly the ideal of functions vanishing, for each  $i$ , on the variety  $P_i$  of matrices of rank  $i$  to order  $\geq \gamma_{i+1}(\sigma)$ .

The graded space of this  $G$  invariant filtration has again the direct sum decomposition as in the Plethysm formulas.

**3. Admissible pairs.** More or less at the same time that these standard bases were studied for the invariant theory of classical groups, Musili [26] and Seshadri [28] analyzed the relationship between standard bases and Schubert cells having in mind a better understanding of the vanishing theorems (proved in general by Kempf [16]; see also [17]) for the cohomology of line bundles in the positive chamber on a variety  $G/P$ ,  $G$  a reductive group,  $P$  a parabolic subgroup.

If the parabolic  $P$  is associated to a fundamental weight  $\omega$ , with  $L_{\omega}$  the corresponding line bundle, the purpose of the analysis is first of all to understand  $H^0(G/P, L_{\omega}^m)$ . This was done, first of all, when  $\omega$  is minuscule. In this case one has

a basis of  $H^0(G/P, L_\omega)$  by extremal weight vectors which index the Bruhat cells of  $G/P$ .

The face ordering of Bruhat cells induces a partial ordering on these generalized Plücker coordinates which also satisfy quadratic equations, so that:

**THEOREM.** *The standard monomials of degree  $m$  in such coordinates are a basis of  $H^0(G/P, L_\omega^m)$ .*

This theorem includes, by suitable interpretation, the standard basis theory for  $S[V \otimes W]$  and  $S[A^2V]$ .

The case  $S[S^2V]$  can be interpreted as a theorem on  $H^0(G/P, L_\omega^m)$  ( $G$  the symplectic group) but it is related to a non minuscule weight.

From this example Lakshmibai, Musili and Seshadri have been able to formulate [20] and prove [19] a general theorem. The hypothesis is that  $\omega$  is a “classical weight”, this means that intersecting a Schubert variety (closure of a Bruhat cell) with the hyperplane class one obtains the Schubert varieties  $\sigma_i$  faces of  $\sigma$  with multiplicity  $\leq 2$ . We can draw a diagram of the ordered set of Bruhat cells, with a double bond each time that the intersection multiplicity is 2.

One defines an admissible pair of cells as one which can be joined by a sequence of double bonds. The pairs  $(\tau, \sigma)$  are considered admissible and called trivial pairs.

**THEOREM.** (LAKSHMIBAI, MUSILI, SESHADRI [19].)

(a) *There is a basis  $P_{\tau, \sigma}$  of  $H^0(G/P, L_\omega)$  indexed by admissible pairs.  $P_{\tau, \sigma}$  is a weight vector of weight  $-\frac{1}{2}(\tau(\omega) + \sigma(\omega))$ .*

(b) *Define a product  $P_{\tau_1 \sigma_1} P_{\tau_2 \sigma_2} \dots P_{\tau_k \sigma_k}$  to be standard if  $\tau_1 \succ \sigma_1 \succ \tau_2 \succ \sigma_2 \succ \dots \succ \tau_k \succ \sigma_k$ . The standard monomials of degree  $m$  are a basis of  $H^0(G/P, L_\omega^m)$ .*

A similar but more complicated analysis gives, for classical groups, standard bases of  $H^0(G/B, L)$ ,  $B$  a Borel subgroup and  $L$  any positive line bundle.

**4. Combinatorial theory of invariant ideals.** We go back to the decomposition  $A = S[V \otimes W] = \bigoplus L_\sigma(V) \otimes L_\sigma(W)$ . The module  $M_\sigma = L_\sigma(V) \otimes L_\sigma(W)$  is irreducible under  $G = \text{Gl}(V) \times \text{Gl}(W)$ . Any invariant subspace  $I$  of  $A$  is thus of the form  $\bigoplus_{\sigma \in \mathcal{J}} M_\sigma$  for some set  $\mathcal{J}$  of partitions. In particular one may study  $G$  invariant ideals; this has been accomplished in joint work with De Concini and Eisenbud [3]. We summarize the results.

**DEFINITION.** A set  $\mathcal{J}$  of partitions is an “ideal” if:  $\sigma \in \mathcal{J}$  and  $\tau \supseteq \sigma$  implies  $\tau \in \mathcal{J}$ . (If  $\sigma: k_1 \succ k_2 \succ \dots \succ k_r$ ,  $\tau: m_1 \succ \dots \succ m_r$  we say  $\tau \supseteq \sigma$  if  $m_i \succ k_i$  for all  $i$ .)


**THEOREM.** *The decomposition  $I = \bigoplus_{\sigma \in \mathcal{J}} M_\sigma$  establishes a 1-1 correspondence between  $G$ -invariant ideals of  $A$  and “ideals” of partitions.*

**DEFINITION.**  $\sigma \cdot \tau$  is the partition having as columns the sum of the corresponding columns.

- DEFINITION. (a) An “ideal”  $\mathcal{I}$  of partitions is prime if  $\sigma\tau \in \mathcal{I}$ ,  $\sigma \notin \mathcal{I}$  implies  $\tau \in \mathcal{I}$ .  
 (b)  $\sqrt{\mathcal{I}} = \{\sigma \mid \sigma^k \in \mathcal{I} \text{ for some } k\}$ .  
 (c)  $\mathcal{I}$  is primary if  $\sigma\tau \in \mathcal{I}$ ,  $\sigma \notin \sqrt{\mathcal{I}}$  implies  $\tau \in \mathcal{I}$ .

THEOREM. *The 1–1 correspondence previously given preserves the notions of prime, primary and radical.*

As an example let us define  $I_\sigma$  as the ( $G$ -invariant) ideal generated by  $M_\sigma$ .  $I_\sigma$  corresponds to the “principal ideal” of partitions  $(\sigma) = \{\tau \mid \tau \supseteq \sigma\}$ . Then  $I_\sigma$  has a nice primary decomposition: e.g.



$\sigma_1, \sigma_2, \sigma_3, \sigma_4$  the diagrams of maximal rectangles in  $\sigma$ , then:  $I_\sigma = I_{\sigma_1} \cap I_{\sigma_2} \cap I_{\sigma_3} \cap I_{\sigma_4}$ .

- THEOREM. (i)  $A_\sigma = I_{k_1} I_{k_2} \dots I_{k_r}$  ( $\sigma: k_1 \geq k_2 \geq \dots \geq k_r$ ).  
 (ii)  $A_\sigma = I_1^{(y_1(\sigma))} \cap I_2^{(y_2(\sigma))} \cap \dots \cap I_n^{(y_n(\sigma))}$   
 (where  $P^{(m)}$  means the symbolic power) is a primary decomposition.  
 (iii)  $A_\sigma$  is the integral closure of  $I_\sigma$ .

THEOREM. *An integrally closed  $G$ -invariant ideal is of the form  $\Sigma A_{\sigma_i}$  with the restriction:*

*If  $\gamma(\tau) \geq$  convex combination of  $\gamma(\sigma_i)$ , then  $\gamma(\tau) \geq$  some  $\gamma(\sigma_i)$ .*

We want to mention a last result which can be obtained by using all the ingredients of the theory [4]: Let  $I_k^{(n)}$  denote the  $n$ th symbolic power of  $I_k$ . The algebra  $B = \bigoplus_n I_k^{(n)} / I_k^{(n+1)}$  has a very explicit theory of standard monomials and:

- (i)  $B$  is a finitely generated algebra;
- (ii)  $B$  is normal and Cohen–Macaulay;
- (iii)  $B$  is the ring of global functions on the normal bundle (in the affine space of matrices) of the variety of matrices of rank  $k-1$ .

### References

1. A. Capelli, *Lezioni sulla teoria delle forme algebriche*, Libr. Sci. Pellerano, Napoli, 1902.
2. C. De Concini, *Standard symplectic tableaux*, Advances in Math. (to appear).
3. C. De Concini, D. Eisenbud and C. Procesi, *Young diagrams and determinantal varieties* (preprint).
4. *Algebras with a straightening law* (preprint).

5. C. De Concini and C. Procesi, *A characteristic free approach to invariant theory*, Advances in Math. **21** (1976), 330—354.
6. D. Desarmenien, J. P. S. Kung and G. C. Rota, *Invariant theory, Young bitableaux, and combinatorics*, Advances in Math. **27** (1978), 63—92.
7. P. Doubilet, G. C. Rota and J. Stein, *On the foundations of combinatorial theory: IX, combinatorial methods in invariant theory*, Studies in Appl. Math. **103** (1974), 185—216.
8. E. Formanek and C. Procesi, *Mumford's conjecture for the general linear group*, Advances in Math. **19** (1976), 292—305.
9. W. J. Haboush, *Reductive groups are geometrically reductive*, Ann. of Math. **102** (1975), 375—376.
10. M. Hochster, *Grassmannians and their Schubert varieties are arithmetically Cohen-Macaulay*, J. Algebra **25** (1973), 40—57.
11. M. Hochster and J. A. Eagon, *Cohen-Macaulay rings, invariant theory, and the generic perfection of the determinantal loci*, Amer. J. Math. **93** (1971), 1020—1058.
12. W. V. D. Hodge, *Some enumerative results in the theory of forms*, Proc. Cambridge Philos. Soc. **39** (1943), 22—30.
13. W. V. D. Hodge and D. Pedoe, *Methods of algebraic geometry*, Vol. II, Cambridge Univ. Press, London, 1952.
14. J. Igusa, *On the arithmetic normality of the Grassmann variety*, Proc. Nat. Acad. Sci. U. S.A. **40** (1954), 309—313.
15. T. Jozefiak and P. Pragacz, *Szyzigies de Pfaffiens* (preprint).
16. G. R. Kempf, *Linear systems on homogeneous spaces*, Ann. of Math. **103** (1976), 557—591.
17. V. Lakshmibai, C. Musili and C. S. Seshadri, *Cohomology of line bundles on  $G/B$* , Ann. Sci. École Norm. Sup. **7** (1974), 89—138.
18. *Geometry of  $G/P$ . III (Standard monomial theory for quasi-minuscule  $P$ )*, Proc. Indian Acad. Sci. (to appear).
19. *Geometry of  $G/P$ . IV (Standard monomial theory for classical types)* (to appear).
20. V. Lakshmibai and C. S. Seshadri, *Geometry of  $G/P$ . II (The work of De Concini and Procesi and the basic conjectures)*, Proc. Indian Acad. Sci. **87A** (1978), 1—54.
21. D. Laksov, *The arithmetic Cohen-Macaulay character of Schubert schemes*, Acta Math. **129** (1972), 1—9.
22. A. Lascoux, *Szyzigies of determinantal varieties*, Advances in Math. **30** (1978), 202—237.
23. *Szyzigies pour les mineurs de matrices symetriques*, preprint, Paris 1977.
24. D. E. Littlewood, *The theory of group characters and matrix representations of groups*, 2nd ed., Clarendon Press, Oxford, 1950.
25. D. Mumford, *Geometric invariant theory*, Springer-Verlag, New York, 1965.
26. C. Musili, *Postulation formula for Schubert varieties*, J. Indian Math. Soc. **36** (1972), 143—171.
27. I. Schur, *Gesammelte Abhandlungen*, Springer-Verlag, Berlin, 1973.
28. C. S. Seshadri, *Geometry of  $G/P$ . I (Standard monomial theory for a minuscule  $P$ )*, C. P. Ramanujan: A tribute, Tata Institute of Fundamental Research (to appear).
29. H. Weyl, *The classical groups*, Princeton Univ. Press, Princeton, N. J., 1946.
30. A. Young, *On quantitative substitutional analysis (third paper)*, Proc. London Math. Soc. **2** (1927), 255—292.