

# Calcium-Activated Potassium Channels Contribute to Human Coronary Microvascular Dysfunction After Cardioplegic Arrest

Jun Feng, MD, PhD; Yuhong Liu, MD; Richard T. Clements, PhD; Neel R. Sodha, MD; Kamal R. Khabbaz, MD; Venkatachalam Senthilnathan, MD; Katherine K. Nishimura, BA; Seth L. Alper, MD, PhD; Frank W. Sellke, MD

**Background**—Cardioplegic arrest (CP) followed by reperfusion after cardiopulmonary bypass induces coronary microvascular dysfunction. We investigated the role of calcium-activated potassium ( $K_{Ca}$ ) channels in this dysfunction in the human coronary microvasculature.

**Methods and Results**—Human atrial tissue was harvested before CP from a nonischemic segment and after CP from an atrial segment exposed to hyperkalemic cold blood CP (mean CP time, 58 minutes) followed by 10-minute reperfusion. In vitro relaxation responses of precontracted arterioles (80 to 180  $\mu$ m in diameter) in a pressurized no-flow state were examined in the presence of  $K_{Ca}$  channel activators/blockers and several other vasodilators. We also examined expression and localization of  $K_{Ca}$  channel gene products in the coronary microvasculature using reverse transcriptase–polymerase chain reaction, immunoblot, and immunofluorescence photomicroscopy. Post-CP reperfusion relaxation responses to the activator of intermediate and small conductance  $K_{Ca}$  channels ( $IK_{Ca}/SK_{Ca}$ ), NS309 ( $10^{-5}$  M), and to the endothelium-dependent vasodilators, substance P ( $10^{-8}$  M) and adenosine 5' diphosphate ( $10^{-5}$  M), were significantly reduced compared with pre-CP responses ( $P < 0.05$ ,  $n = 8$ /group). In contrast, relaxation responses to the activator of large conductance  $K_{Ca}$  channels ( $BK_{Ca}$ ), NS1619 ( $10^{-5}$  M), and to the endothelium-independent vasodilator, sodium nitroprusside ( $10^{-4}$  M), were unchanged pre- and post-CP reperfusion ( $n = 8$ /group). Endothelial denudation significantly diminished NS309-induced vasodilatation and abolished substance P- or adenosine 5' diphosphate-induced relaxation ( $P < 0.05$ ), but had no effect on relaxation induced by either NS1619 or sodium nitroprusside. The total polypeptide levels of  $BK_{Ca}$ ,  $IK_{Ca}$ , and  $SK_{Ca}$  and the expression of  $IK_{Ca}$  mRNA were not altered post-CP reperfusion.

**Conclusion**—Cardioplegic arrest followed by reperfusion after cardiopulmonary bypass causes microvascular dysfunction associated with and likely in part due to impaired function of  $SK_{Ca}$  and  $IK_{Ca}$  channels in the coronary microcirculation. These results suggest novel mechanisms of endothelial and smooth muscle microvascular dysfunction after cardiac surgery. (*Circulation*. 2008;118[suppl 1]:S46–S51.)

**Key Words:** calcium-activated potassium channels ■ cardioplegia ■ ischemia and reperfusion ■ microcirculation

Microvascular endothelial dysfunction and altered vascular reactivity often occur after ischemic arrest and cardiopulmonary bypass, although cardioplegia has been routinely used for the protection of the myocardium against ischemic injury during cardiac surgery.<sup>1–3</sup> Endothelium has been found to release 3 major endothelium-dependent relaxing factors: nitric oxide, prostacyclin, and endothelium-dependent hyperpolarization factor (EDHF).<sup>4</sup> Impaired endothelial synthesis and release of nitric oxide, prostacyclin, and EDHF associated with cardioplegic ischemia and reperfusion may contribute to microvascular endothelial dysfunction.<sup>1–3</sup> These observations have prompted addition to cardioplegic solutions of nitric oxide donors, exogenous prostacyclin,

and EDHF agonists in attempts to attenuate ischemia/reperfusion-mediated endothelial dysfunction.<sup>5–7</sup> The vasodilatory influence of EDHF is believed predominant in smaller arterioles.

EDHF has been proposed to dilate the coronary arteries through opening of calcium-activated potassium channels ( $K_{Ca}$ ) in endothelial and smooth muscle cells.<sup>4</sup> Recently, we found that bradykinin preconditioning of coronary microvasculature in isolated rabbit hearts undergoing cardioplegic arrest was mediated in part by the activation of  $K_{Ca}$  channels.<sup>8</sup> However, the roles of  $K_{Ca}$  channel activation in cardioplegic arrest (CP) and reperfusion-related microvascular dysfunction in human coronary resistance arterioles and the regula-

From the Division of Cardiothoracic Surgery, Department of Surgery, Divisions of Molecular and Vascular Medicine and Nephrology, Department of Medicine, Beth Israel Deaconess Medical Center, and Harvard Medical School, Boston, Mass.

Presented at the American Heart Association Scientific Sessions, November 4–7, 2007, Orlando, Fla.

Correspondence to Frank W. Sellke, MD, Division of Cardiothoracic Surgery, BIDMC, LMOB 2A, 110 Francis Street, Boston, MA 02215. E-mail fsellke@caregroup.harvard.edu

© 2008 American Heart Association, Inc.

*Circulation* is available at <http://circ.ahajournals.org>

DOI: 10.1161/CIRCULATIONAHA.107.755827

tory properties of these  $K_{Ca}$  channels in this vascular bed have little been investigated.

This study was designed to examine the effect of blood CP and brief reperfusion (Rep) on vascular responses of human atrial microvessels to  $K_{Ca}$  channel openers and other vasoactive substances and to correlate these responses to possible alterations in expression of  $K_{Ca}$  channel mRNAs and polypeptides in human atrial tissue.

## Materials and Methods

### Human Subjects and Tissue Harvesting

Samples of right atrial appendage were harvested from clinically similar patients undergoing coronary artery bypass graft surgery before and after exposure of the heart to blood CP and short-term reperfusion under conditions of cardiopulmonary bypass. Samples were harvested with cold sharp dissection and handled in a nontraumatic fashion. Double 3-0 polypropylene pursestring sutures (Ethicon, Somerville, NJ) were placed in the atrial appendage. During placement of the venous cannula, the first sample of atrial appendage was harvested before cardioplegia and reperfusion (pre-CP Rep). The superior suture was tightened to secure the venous cannula. The inferior suture remained loose to allow this portion of the atrium to be perfused with blood, exposed to blood CP, and reperfused (post-CP Rep) after removal of the aortic crossclamp. An initial 800 to 1000 mL of cold-blood (0°C to 4°C) hyperkalemic (15 mmol/L  $K^+$ ) cardioplegic solution was delivered antegrade into the aortic root. This was followed at 8- to 15-minute intervals with 250 to 300 mL of cold CP solution (15 mmol/L  $K^+$ ). The CP solutions consisted of a mixture of oxygenated blood with crystalloid solution of the following final composition (in mmol/L): 15 KCl, 3.5  $MgSO_4$ , 135NaCl, 1.0  $CaCl_2$ , 11 glucose, 11 mannitol, and 4 tromethamine.

The second sample of atrial appendage was harvested during removal of the venous cannula. Tissue samples for immunoblot analysis assay were immediately frozen in liquid nitrogen. Tissue for immunofluorescent staining was fixed in 10% formalin-buffered solution for 24 hours followed by paraffin mounting and sectioning into 5- $\mu$ m slices. Tissue for microvascular studies was placed in cold (5° to 10°C) Krebs buffer solution. All procedures were approved by the Institutional Review Board of Beth Israel Deaconess Medical Center, Harvard Medical School, and informed consent was obtained from all enrolled patients required by the Institutional Review Board.

### Coronary Microvessel Relaxation Studies

Coronary arterioles (80- to 180- $\mu$ m internal diameters) were dissected from the right atria appendage pre- and post-CP Rep. Microvessel studies were performed by *in vitro* organ bath videomicroscopy as described previously.<sup>3,6</sup> After a 60-minute stabilization period in the organ chamber, the microvessels were precontracted with thromboxane  $A_2$  analog U46619  $\times$  30% to 40% of the baseline diameter. After achievement of this constricted steady state, dose-dependent relaxation was measured in response to (extraluminal) application of the following vasodilators: the activator of intermediate and small conductance  $K_{Ca}$  channels ( $IK_{Ca}$ /SK $Ca$ ), NS309, the activator of large conductance  $K_{Ca}$  channels ( $BK_{Ca}$ ), NS1619 (both from  $10^{-9}$  to  $10^{-5}$  M), sodium nitroprusside (SNP), adenosine 5'-diphosphate (both from  $10^{-9}$  to  $10^{-4}$  M), and substance P ( $10^{-12}$  to  $10^{-7}$  M). After exposure to substance P, the vessel was discarded to avoid tachyphylaxis. One or 3 interventions were performed on each vessel. The order of drug administration was random. Six vessels harvested before CP Rep were pretreated with a mixture of the  $IK_{Ca}$  blocker, TRAM34 ( $10^{-7}$  M), and the SK $Ca$  blocker, apamin ( $10^{-6}$  M), before perfusion with NS309. Six additional vessels were pretreated with the  $BK_{Ca}$  blocker, iberiotoxin ( $10^{-7}$  M) before perfusion with NS1619. In some cases, endothelial denudation was carried out by advancing a human hair into the lumen and gently abrading the luminal surface.

### Immunoblot

Small coronary arteries were dissected and cleaned of connective tissues and solubilized in SDS-PAGE buffer. Total protein (40  $\mu$ g) was fractionated on an 8% to 16% SDS-PAGE and then transferred to a polyvinylidene difluoride membrane (Immobilon-P; Millipore Corporation, Bedford, Mass) as previously described.<sup>3</sup> Membranes were incubated for 1 hour at room temperature with 1:200 dilutions of individual rabbit polyclonal primary antibodies to  $BK_{Ca}$ - $\alpha$ ,  $BK_{Ca}$ - $\beta_1$  (Santa Cruz Biotechnology, Santa Cruz, Calif),  $IK_{Ca}$ , or SK3 $Ca$ - $\alpha$  (Alomone Labs Ltd, Jerusalem, Israel). The membranes were then incubated for 1 hour with horseradish peroxidase-conjugated secondary anti-Ig, washed  $3\times$  in Tris saline buffer, and processed for chemiluminescent detection (Pierce, Rockford, Ill) on x-ray film (Kodak, Rochester, NY). Band intensity was measured by densitometric analysis of autoradiograph films using NIH Image J1.33. Specificities of the anti- $BK_{Ca}$ - $\alpha$ , anti- $BK_{Ca}$ - $\beta_1$ , anti- $IK_{Ca}$ , and anti-SK3 $Ca$  antibodies were demonstrated in the previous studies, respectively.<sup>8-10</sup>

### Reverse Transcriptase-Polymerase Chain Reaction of $IK_{Ca}$ mRNA


Total RNA was prepared with the RNeasy kit (Qiagen, Valencia, Calif) from flash-frozen pre- and post-CP Rep atrial tissue samples. RNA integrity was verified by formaldehyde-agarose gel electrophoresis, and RNA concentration was estimated by absorbance at 260 nm. Reverse transcription was performed with the First Strand cDNA Synthesis Kit (Ambion, Austin, Texas) using equal amounts of total RNA. Five percent of the cDNA mixture was used for polymerase chain reaction with HotStart DNA Polymerase (Qiagen; total reaction volume 20  $\mu$ L in supplier's recommended buffer). Two primer pairs tested yielded nearly identical results: forward primer IK1.F6 5'-GGAAGCTCCGGGAACAAGTG-3' coupled with reverse primer  $IK_{Ca}$ .R3 5'-CTACTTKGACTGCTGGCTKGGTTC-3' (209 bp polymerase chain reaction product) or forward primer IK1.F7 5'-GGTGGACATCTCCAAGATGCAC-3' coupled with the same reverse primer (181 bp polymerase chain reaction product). Complete reaction mixes were incubated 15 minutes at 95°C, then cycled through 45 seconds denaturation at 94°C, 2 minutes annealing at 60°C, and 2 minutes elongation at 72°C. A 7-minute final extension at 72°C was terminated by rapid cooling to room temperature. Polymerase chain reaction products were separated in 1% agarose gels, visualized with ethidium bromide, and imaged (Gel-Doc; Biorad, Hercules, Calif). Polymerase chain reaction cycle number was adjusted for low abundance transcript detection within the range of log linear amplification. Thirty-six cycles was chosen for amplification of  $IK_{Ca}$  transcript and 23 cycles for  $\beta$ actin controls.

### Immunofluorescence Photomicroscopy

Atrial tissue sections from 5 patients were deparaffinized in xylene, rehydrated in graded ethanol and phosphate-buffered saline solution, and antigen-unmasked with sodium citrate (10 mmol/L, pH=6.0) followed by phosphate-buffered saline wash and blocking with 2% bovine serum albumin in phosphate-buffered saline at room temperature for 2 hours. After phosphate-buffered saline wash, overnight incubation with anti- $BK_{Ca}$ - $\alpha$  and  $BK_{Ca}$ - $\beta_1$  (each used 1:200; Santa Cruz Biotechnology) was performed at 4°C. Antimouse,  $\alpha$ -smooth muscle actin (1:1000; Sigma, St Louis, Mo) was used to detect microvascular smooth muscle. Sections were then washed in phosphate-buffered saline and incubated with the appropriate Alexa fluor secondary antibody and mounted using fluorescent mounting medium (Vector Labs, Burlingame, Calif). Tissue was visualized using a Zeiss LSM510 confocal microscope system (Carl Zeiss MicroImaging, Inc, Thornwood, NY). Tissue labeling with secondary antibody alone or with normal rabbit IgG or serum in place of primary antibody served as a negative control.

### Chemicals

NS309, NS1619, U46619, SNP, adenosine 5' diphosphate (ADP), and substance P were obtained from Sigma. SNP, ADP, and substance P were dissolved in ultrapure distilled water and prepared



**Figure 1.** Vasoreactivity of human coronary arterioles to the IK/SK channel activator NS309. A, Dose-dependent relaxation of coronary arterioles ( $n=8$ ) in response to NS309 before CP and Rep and afterward (post-CP Rep);  $*P<0.05$ . B, Dose-dependent NS309-mediated relaxation of pre-CP Rep human coronary arterioles pretreated in the absence or presence of TRAM34 and apamin ( $n=6$ );  $**P<0.001$ . C, Dose-dependent relaxation of pre-CP Rep human coronary arterioles from pre-CP Rep tissues ( $n=6$ ) with intact or denuded endothelium in response to NS309;  $**P<0.001$ . Data are presented as percent relaxation after preconstriction with U46619.

on the day of the study. U46619 was dissolved in ethanol to make a stock solution. NS309 and NS1619 were dissolved in dimethylsulfoxide to make a stock solution. All stock solutions were stored at  $-20^{\circ}\text{C}$ . All dilutions were prepared daily.

### Data Analysis

Data are presented as the mean and SEM. The relaxation responses were expressed as the percentage of relaxation of the U46619-precontracted diameter of the microvessels. Repeated-measures analysis of variance and Student *t* test were used to compare variables between or among vessels. The treatment effects were statistically examined by paired or independent 2-tailed Student *t* test. Statistical significance was taken at a probability value of  $<0.05$ .

### Statement of Responsibility

The authors had full access to the data and take full responsibility for its integrity. All authors have read and agree to the manuscript as written.

## Results

### Patient Characteristics

Tissue samples from 27 patients were studied. Twenty-seven patients (mean age  $67\pm 6$  years) underwent coronary artery bypass graft with duration of cardioplegic arrest of  $58\pm 4$  minutes and cardiopulmonary bypass time of  $71.0\pm 18$  minutes. Twenty-one patients were male and 6 were female. Twenty-two of the 27 patients carried a preoperative diagnosis of hypertension. All patients with preoperative hypertension were on medication ( $\beta$ -blocker, calcium channel blocker, or angiotensin-converting enzyme inhibitor) and received perioperative  $\beta$ -blockade. Diabetes mellitus (type I or type II) was present in 5 of 27 patients.

### Vessel Characteristics

Coronary microvessels ranged from 80 to 180  $\mu\text{m}$  in internal diameter. Diameters in the NS309 groups were  $107\pm 4$   $\mu\text{m}$  in the pre-CP Rep group,  $110\pm 8$   $\mu\text{m}$  in the post-CP Rep group,  $103\pm 7$   $\mu\text{m}$  in the endothelium denudation group, and  $137\pm 9$   $\mu\text{m}$  in the TRAM34/apamin+NS309 group. Diameters in the NS1619 groups were  $125\pm 10$   $\mu\text{m}$  in the pre-CP Rep group,  $114\pm 9$   $\mu\text{m}$  in the post-CP Rep group,  $146\pm 11$   $\mu\text{m}$  in the endothelium denudation group, and  $126\pm 8$   $\mu\text{m}$  in the iberiotoxin+NS1619 group. In the substance P groups, diameters were  $134\pm 6$   $\mu\text{m}$  in the pre-CP


Rep group,  $112\pm 7$   $\mu\text{m}$  in the post-CP Rep group, and  $119\pm 7$   $\mu\text{m}$  in the endothelium denudation group. In the ADP groups, diameters were  $118\pm 10$   $\mu\text{m}$  in the pre-CP Rep group,  $130\pm 12$   $\mu\text{m}$  in the post-CP Rep group, and  $135\pm 7$   $\mu\text{m}$  in the endothelium denudation group. In the SNP groups, diameters were  $116\pm 6$   $\mu\text{m}$  in the pre-CP Rep group,  $108\pm 7$   $\mu\text{m}$  in the pre-CP Rep group, and  $122\pm 11$   $\mu\text{m}$  in the endothelium denudation group. Most intergroup differences of the vessel diameters were statistically insignificant. The degree of precontraction by the thromboxane  $A_2$  analog U46619 was  $29\pm 2\%$  in the pre-CP Rep group and  $31\pm 2\%$  in the post-CP Rep group. Similar U46619 concentrations were required to produce adequate precontraction in both groups.

### Responses to NS309 and NS1619

Both NS309 (Figure 1A) and NS1619 (Figure 2A) induced dose-dependent relaxation of human coronary arterioles. Cardioplegic arrest and reperfusion significantly impaired the relaxation response to NS 309 compared with pre-CP Rep responses ( $P<0.05$ ; Figure 1A). In contrast, the pre-CP Rep responses to NS1619 were unchanged post-CP Rep (Figure 2A). Pretreatment with TRAM34/apamin abolished vessel relaxation induced by NS309 ( $P<0.001$ ; Figure 1B), and NS1619-induced relaxation was inhibited by pretreatment with iberiotoxin ( $P<0.001$ ; Figure 2B). Furthermore, NS309-mediated vasorelaxation of the coronary arterioles was significantly diminished by removal of the endothelium ( $P<0.001$ ; Figure 1C), whereas responses to NS1619 were unaffected (Figure 2C).

### Responses to Adenosine 5' Diphosphate, Substance P, and Sodium Nitroprusside

ADP, substance P, and SNP induced dose-dependent relaxation of coronary arterioles (Figure 3A–C). Cardioplegic arrest and reperfusion remarkably reduced the relaxation responses both to ADP and to substance P compared with pre-CP Rep responses ( $P<0.05$ ; Figure 3A–B). In contrast, pre-CP Rep responses to SNP were unchanged post-CP Rep (Figure 3C). Endothelium denudation abolished responses to both ADP and substance P ( $P<0.001$ ; Figure 3A–B), but left intact SNP-induced relaxation.



**Figure 2.** Vasoreactivity of human coronary arterioles to the BK channel activator NS1619. A, Dose-dependent relaxation of coronary arterioles in response to NS1619 either pre-CP Rep or post-CP Rep ( $n=8$ /group). B, Dose-dependent NS1619-mediated relaxation of human coronary arterioles from pre-CP Rep atrial tissues pretreated in the absence or presence of iberiotoxin ( $n=6$ ;  $**P<0.001$ ). C, Dose-dependent NS1619-induced relaxation of pre-CP Rep human coronary arterioles with intact or denuded endothelium ( $n=6$ /group). Data are presented as percent relaxation after precontraction with U46619.

### Effect of Cardioplegic Arrest Reperfusion on levels of $K_{Ca}$ Polypeptides and $IK_{Ca}$ mRNA

Pre-CP Rep coronary artery levels of the  $K_{Ca}$  channel polypeptides  $BK_{Ca-\alpha}$  and  $BK_{Ca}$   $BK_{Ca-\beta_1}$ ,  $IK_{Ca-\alpha}$ , and  $SK3_{Ca-\alpha}$  were unchanged in the post-CP Rep period as detected by immunoblot (Figure 4A–B). Levels of  $IK_{Ca}$  mRNA were similarly unaltered after cardioplegic ischemia and reperfusion (Figure 4C).

### Immunolocalization of Microvessel $K_{Ca}$ Polypeptides Pre- and Postcardioplegic Arrest Reperfusion


Immunofluorescent staining of coronary microvessels displayed a strong signal for  $BK_{Ca-\alpha}$  (Figure 5A) and  $BK_{Ca-\beta_1}$  (Figure 5B) subunits localized to the microvascular smooth muscle and less abundantly to endothelium. Negative controls documented low-level background fluorescence.

### Discussion


This study presents several novel findings. First, both the  $IK_{Ca}/SK_{Ca}$  activator NS309 and the  $BK_{Ca}$ -specific opener NS1619 induced dose-dependent vasodilatation of human coronary arterioles. Second, the relaxation response to NS309 was significantly impaired after CP and Rep, whereas the response to NS1619 was unaltered. Third, the response to NS309 was significantly diminished after endothelium denu-

ation, whereas the response to NS1619 was unaffected. Fourth, the presence of  $BK_{Ca}$ ,  $IK_{Ca}$ , and  $SK_{Ca}$  polypeptides in human coronary microvasculature was documented by immunoblot and by immunofluorescence microscopy.  $BK_{Ca}$  immunostaining was present predominantly in smooth muscle cells and less apparent in endothelium. However,  $SK_{Ca}$  immunostaining was detected predominantly in endothelium and less in smooth muscle cells.  $IK_{Ca}$  mRNA was also present in human atrial tissue. Finally, cardioplegic ischemia and Rep change neither total polypeptide levels of  $BK_{Ca}$ ,  $IK_{Ca}$ , or  $SK_{Ca}$  polypeptides nor the level of  $IK_{Ca}$  mRNA.

We and several others have previously shown that cardioplegic ischemia and Rep resulted in microvascular endothelial dysfunction and decreased hyperpolarization in animals and humans.<sup>1–3,5</sup> The present study confirmed previous findings indicating that responses to several endothelial-dependent vasodilators such as ADP and substance P were significantly reduced after cardioplegic ischemia and Rep. Recent studies have demonstrated that  $K_{Ca}$  channels contribute to the vasodilatation and hyperpolarization induced by the EDHF stimulators, substance P and bradykinin.<sup>4,10</sup> Three types of  $K_{Ca}$  channels,  $BK_{Ca}$ ,  $IK_{Ca}$ , and  $SK_{Ca}$ , are present in the vasculature.<sup>4,11,12</sup> In the present study, relaxation responses to NS309 were completely blocked in the combined presence of the  $IK_{Ca}$  blocker TRAM 34 and the  $SK_{Ca}$  inhibitor apamin, supporting specificity of NS309 as an activator of both  $IK_{Ca}$



**Figure 3.** In vitro response of precontracted human atrial arterioles to endogenous vasodilators. Arterioles were compared pre-CP Rep and post-CP Rep ( $n=8$ /group). Pre-CP Rep arterioles were compared with intact endothelium and after endothelial denudation ( $n=6$ /group) as indicated. A, Response to the endothelium-dependent vasodilator ADP.  $*P<0.05$  versus pre-CP Rep;  $**P<0.001$  versus intact pre-CP Rep. B, Response to substance P.  $*P<0.05$  versus pre-CP Rep;  $**P<0.001$  versus pre-CP Rep. C, Response to the endothelium-independent vasodilator, SNP.



**Figure 4.** A, top left, Representative immunoblot of human coronary microvessels. Lanes 1 to 4 loaded with 40  $\mu$ g protein were developed for BK $_{Ca}$ - $\alpha$ , BK $_{Ca}$ - $\beta$ 1, IK $_{Ca}$ - $\alpha$ , and SK3 $_{Ca}$ - $\alpha$  polypeptides B, top right, Densitometric evaluation of immunoblot band intensity shows unaltered levels of K $_{Ca}$  polypeptides after CP Rep (n=5/group). C, bottom, Semiquantitative reverse transcriptase–polymerase chain reaction showing steady-state levels of mRNA for IK $_{Ca}$  and  $\beta$ -actin control in human atrial tissue pre- and CP and Rep.

and SK $_{Ca}$  channels. Relaxation responses to NS1619 were abolished in the presence of the BK $_{Ca}$  blocker iberiotoxin, consistent with the specificity of NS1619 as a BK $_{Ca}$  activator. Endothelium denudation significantly diminished NS309-induced relaxation but failed to affect the response to NS1619. These results show that whereas NS309-induced relaxation is endothelium-dependent, relaxation induced by NS1619 is endothelium-independent. Another important finding of the present study is that cardioplegic ischemia and Rep remarkably impaired IK $_{Ca}$ /SK $_{Ca}$  channel activator NS309-induced dose-dependent vasodilatation, but not the BK $_{Ca}$  opener NS1619-induced relaxation, suggesting that the inactivation of IK $_{Ca}$ /SK $_{Ca}$  may be involved in endothelial dysfunction of human coronary microvasculature. All the results presented were obtained in vessels precontracted with a thromboxane agonist. It is thus possible that the proportional vasodilatory contributions of IK $_{Ca}$ /SK $_{Ca}$  and BK $_{Ca}$  channels may vary in coronary microvessels precontracted by other agonists such as endothelin, norepinephrine, or elevated extracellular calcium.

To detect IK $_{Ca}$ /SK $_{Ca}$  and BK $_{Ca}$  gene products in the human coronary microvasculature, we performed immunoblot and immunohistochemistry to localize channel polypeptides. We found that BK $_{Ca}$  was predominantly present in smooth muscle cells. This finding is consistent with the microvessel relax-

ation data, indicating that BK $_{Ca}$  activator-mediated relaxation is endothelium-independent. BK $_{Ca}$ , IK $_{Ca}$ , and SK3 $_{Ca}$  polypeptides were detected by immunoblot of extracts from human coronary microvessels. IK $_{Ca}$  mRNA was detected in human atrial appendage. However, cardioplegic ischemia and Rep did not alter expression of BK $_{Ca}$ , IK $_{Ca}$ , and SK3 $_{Ca}$  polypeptides or IK $_{Ca}$  mRNA levels, suggesting that short periods of intraoperative hypothermic ischemia and reperfusion may modify the functional state of channel protein activation or intracellular distribution rather than the steady-state levels of protein and (in the case of IK $_{Ca}$ ) mRNA.

The molecular mechanisms responsible for the CP Rep-related IK $_{Ca}$ /SK $_{Ca}$  channel dysfunction remain unclear. Ischemia and Rep may induce activation and phosphorylation of several protein and tyrosine kinases, which in turn may phosphorylate IK $_{Ca}$ /SK $_{Ca}$  channels or their regulatory proteins, resulting in channel inactivation<sup>13</sup> either by decreasing open probability or by reduction of channel number at the cell surface. Generation of free radicals and cytokine release during ischemia and Rep may also inhibit K $_{Ca}$  channel activity.<sup>14</sup>


In conclusion, CP followed by Rep after cardiopulmonary bypass causes microvascular dysfunction, likely due in part to impaired function of IK $_{Ca}$ /SK $_{Ca}$  channels in human coronary microcirculation. These results may provide novel mechanisms by which to explain the presently unavoidable endothelial and smooth muscle microvascular dysfunction after cardiac surgery. Pharmacological administration of K $_{Ca}$  openers may represent a novel strategy to improve microvascular protection during cardiac surgery and minimally invasive revascularization.

## Acknowledgments

We thank Dr Boris E. Shmukler for discussion and advice.

## Sources of Funding

This research project was supported in part by the National Heart, Lung, and Blood Institute HL-69024-05 and HL-46716-15 (F.W.S.) and HL-077765 (S.L.A.). Y.L., R.T.C., and N.R.S. were supported by a cardiovascular research training grant (T32) from the National Institutes of Health (HL076130-02 [F.W.S.]).



**Figure 5.** Immunolocalization of K $_{Ca}$  channel polypeptides in human coronary microvessels. Vessels were costained for smooth muscle  $\alpha$ -actin and either (A) BK $_{Ca}$ - $\alpha$  or (B) BK $_{Ca}$ - $\beta$ 1. Matched negative controls are displayed below each row of primary antibody staining as indicated.

### Disclosures

F.W.S. has received grant support from Ikaria and Orthologic and is on the steering committee for Novo Nordisk and on the DSMB for Dyax.

### References

- Sellke FW, Boyle EM Jr, Verrier ED. Endothelial cell injury in cardiovascular surgery the pathophysiology of vasomotor dysfunction. *Ann Thorac Surg.* 1996;62:1222–1228.
- Metias C, Li J, Simons M, Sellke FW. Serotonin-induced coronary contraction increases after blood cardioplegia-reperfusion: role of COX-2 expression. *Circulation.* 1999;100(suppl II):II328–334.
- Feng J, Bianchi C, Sandmeyer JL, Li JY, Sellke FW. Molecular indices of apoptosis after intermittent blood and crystalloid cardioplegic ischemia. *Circulation.* 2005;112(suppl I):I-184–189.
- Ledoux J, Werner ME, Brayden JE, Nelson MT. Calcium-activated potassium channels and the regulation of vascular tone. *Physiology.* 2006;21:69–78.
- Feng J, Li HL, Rosenkranz ER. Bradykinin protects the rabbit heart after cardioplegia via NO-dependent pathways. *Ann Thorac Surg.* 2000;70:2119–2124.
- Feng J, Liu RY, Wu GZ, Tang SB. PGE<sub>1</sub> reduces cardiac-derived TXA<sub>2</sub> in ischemic arrest in isolated working rat heart. *Int J Cardiol.* 1996;55:265–270.
- Feng J, Sellke ME, Ramlawi B, Boodhwani M, Clements R, Li JY, Bianchi C, Sellke FW. Bradykinin induces microvascular preconditioning via opening of calcium-activated potassium channels. *Surgery.* 2006;140:192–197.
- Si H, Heyken WT, Wolfle SE, Tysiac M, Schubert R, Grgic I, Vilianovich L, Giebing G, Maier T, Gross V, Bader M, de Wit C, Hoyer J, Kohler R. Impaired endothelium-derived hyperpolarizing factor-mediated dilations and increased blood pressure in mice deficient of the intermediate-conductance Ca<sup>2+</sup>-activated K<sup>+</sup> channel. *Circ Res.* 2006;99:537–544.
- Ambroisine ML, Favre J, Oliviero P, Rodriguez C., Gao J, Thuillez C, Samuel J, Richard LV, Delcayre C. Aldosterone-induced coronary dysfunction in transgenic mice involves the calcium-activated potassium (BKCa) channels of vascular smooth muscle cells. *Circulation.* 2007;116:2435–2443.
- Weston AH, Absi M, Ward DT, Ohanian J, Dodd RH, Dauban P, Petrel C, Ruat M, Edwards G. Evidence in favor of a calcium-sensing receptor in arterial endothelial cells: studies with calindol and Calhex 231. *Circ Res.* 2005;19:97:391–398.
- Wei AD, Gutman GA, Aldrich R, Chandy KG, Grissmer S, Wulff H. International Union of Pharmacology. Nomenclature and molecular relationships of calcium-activated potassium channels. *Pharmacol Rev.* 2005;5:463–472.
- Begenisich T, Nakamoto T, Ovitt CE, Nehrke K, Brugnara C, Alper SL, Melvin JE. Physiological roles of the intermediate conductance, Ca<sup>2+</sup>-activated potassium channel K<sub>cnj4</sub>. *J Biol Chem.* 2004;279:47681–47687.
- Schubert R, Nelson MT. Protein kinases: tuners of the BK<sub>Ca</sub> channel in smooth muscle. *Trends Pharmacol Sci.* 2001;22:505–512.
- Liu Y, Terata K, Chai Q, Li H, Kleinman LH, Gutterman DD. Peroxynitrite inhibits Ca<sup>2+</sup>-activated K<sup>+</sup> channel activity in smooth muscle of human coronary arterioles. *Circ Res.* 2002;9:1070–1076.

## Calcium-Activated Potassium Channels Contribute to Human Coronary Microvascular Dysfunction After Cardioplegic Arrest

Jun Feng, Yuhong Liu, Richard T. Clements, Neel R. Sodha, Kamal R. Khabbaz, Venkatachalam Senthilnathan, Katherine K. Nishimura, Seth L. Alper and Frank W. Sellke

*Circulation*. 2008;118:S46-S51

doi: 10.1161/CIRCULATIONAHA.107.755827

*Circulation* is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2008 American Heart Association, Inc. All rights reserved.

Print ISSN: 0009-7322. Online ISSN: 1524-4539

The online version of this article, along with updated information and services, is located on the World Wide Web at:

[http://circ.ahajournals.org/content/118/14\\_suppl\\_1/S46](http://circ.ahajournals.org/content/118/14_suppl_1/S46)

**Permissions:** Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

**Reprints:** Information about reprints can be found online at:  
<http://www.lww.com/reprints>

**Subscriptions:** Information about subscribing to *Circulation* is online at:  
<http://circ.ahajournals.org/subscriptions/>