

Enhanced miR-210 expression promotes the pathogenesis of endometriosis through activation of signal transducer and activator of transcription 3

M. Okamoto¹, K. Nasu^{1,2,*}, W. Abe¹, Y. Aoyagi¹, Y. Kawano¹, K. Kai¹, M. Moriyama³, and H. Narahara¹

¹Department of Obstetrics and Gynecology, Faculty of Medicine, Oita University, Hasama-machi, Yufu-shi, Oita 879-5593, Japan

²Division of Obstetrics and Gynecology, Support System for Community Medicine, Faculty of Medicine, Oita University, Yufu-shi, Oita 879-5593, Japan ³Department of Molecular Pathology, Faculty of Medicine, Oita University, Yufu-shi, Oita 879-5593, Japan

*Correspondence address. Tel: +81-97-586-5922; Fax: +81-97-586-6687; E-mail: nasu@oita-u.ac.jp

Submitted on May 10, 2014; resubmitted on November 18, 2014; accepted on November 25, 2014

STUDY QUESTION: What are the roles of the microRNA miR-210—an miRNA that is up-regulated in endometriotic cyst stromal cells (ECSCs)—in the pathogenesis of endometriosis?

SUMMARY ANSWER: Up-regulated miR-210 expression in ECSCs is involved in their proliferation, resistance to apoptosis and angiogenesis through signal transducer and activator of transcription (STAT) 3.

WHAT IS KNOWN ALREADY: In the pathogenesis of endometriosis, a number of roles for microRNAs (miRNAs) are becoming apparent.

STUDY DESIGN, SIZE, DURATION: ECSCs and normal endometrial stromal cells (NESCs) were isolated from ovarian endometriotic tissues (patients aged 24–40 years undergoing salpingo-oophorectomy or evisceration for the treatment of ovarian endometriotic cysts, $n = 10$) and the eutopic endometrial tissues without endometriosis (premenopausal patients aged 35–45 years undergoing hysterectomies for subserosal leiomyoma, $n = 13$), respectively.

PARTICIPANTS/MATERIALS, SETTING, METHODS: We used a global gene expression microarray technique to identify downstream targets of miR-210, and we assessed the functions of miR-210 in the pathogenesis of endometriosis by using the miR-210-transfected NESCs.

MAIN RESULTS AND THE ROLE OF CHANCE: Gene expression microarray analysis revealed that one of the key target molecules of miR-210 is STAT3. In the NESCs, in comparison to the control, miR-210 transfection resulted in the induction of cell proliferation ($P < 0.0005$), the production of vascular endothelial cell growth factor (VEGF) ($P < 0.0005$) and the inhibition of apoptosis ($P < 0.05$) through STAT3 activation [increased levels of mRNA ($P < 0.0005$), and protein ($P < 0.005$)]. In the ECSCs, inhibitors of STAT3 inhibited the cell proliferation and VEGF production ($P < 0.05$), and induced the apoptosis of these cells ($P < 0.05$).

LIMITATIONS, REASONS FOR CAUTION: The roles of aberrant miR-210 expression were investigated only in the stromal component of ectopic and eutopic endometrium. Control endometrial tissues were obtained from premenopausal patients who had subserosal leiomyoma and NESC gene expression patterns may be altered in these women. Furthermore, the effects of STAT3 inhibitors were evaluated only in ECSCs and not in NESCs.

WIDER IMPLICATIONS OF THE FINDINGS: The present findings indicate that miR-210 induces NESCs to differentiate into the endometriotic phenotype and we speculate that up-regulated miR-210 expression in ECSCs is involved in the creation of the endometriosis-specific cellular dysfunctions through epigenetic mechanisms. The data indicate that STAT3 inhibitors may be promising candidates for the treatment of endometriosis.

STUDY FUNDING/COMPETING INTEREST(S): This work was supported in part by Grants-in-Aid for Scientific Research from the Japan Society for the Promotion of Science (no. 13237327 to K.N., no. 25861500 to Y.K. and no. 23592407 to H.N.). There are no conflicts of interest to declare.

Key words: endometriosis / microRNA / signal transducer and activator of transcription 3 / vascular endothelial growth factor

Introduction

Endometriosis is an estrogen-dependent disease exhibiting the benign ectopic growth of proliferative endometrial tissue. As a disease most frequently observed in women of reproductive age, endometriosis is most commonly based in the peritoneum, ovaries and rectovaginal septum (Giudice and Kao, 2004). Dysmenorrhea, chronic pelvic pain, subfertility and/or dyspareunia are the main symptoms, often greatly decreasing the quality of life of the affected women (Giudice and Kao, 2004).

Although endometriotic tissues share many histological characteristics with normal proliferative endometrial tissues (Giudice and Kao, 2004), they show several interesting molecular differences such as those concerning gene expression and protein production, synthesis and responsiveness to steroids and cytokines, immune components, adhesion molecules, and proteolytic enzymes and their inhibitors; endometriotic tissues also have a tissue structure and cell proliferation rates that differ from those of normal endometrial tissues (Nasu et al., 2011a,b). The occurrence and development of endometriosis may be based on these aberrations in molecular processes, which have been rather stable and consistent in endometriosis, and some sort of unknown trigger(s) that initiate these molecular alterations might exist. To identify the responsible mechanism(s) of the pathogenesis of endometriosis, our research has been focused on the dysregulation of microRNA (miRNA) expression (Abe et al., 2013), histone modification (Nasu et al., 2014) and DNA methylation (Nasu et al., 2011a) in endometriotic cells from the viewpoint of epigenetics. We hypothesized that the acquisition of aberrant gene expression by epigenetic mechanisms may induce the endometrial cells to differentiate into an endometriotic phenotype (Nasu et al., 2014).

miRNAs are short RNAs that comprise a class of regulatory genes characterized as endogenous, single-stranded, non-coding RNA (Bartel, 2004). They function by regulating the translation of specific targeted protein-coding genes (Bartel, 2004). It has been estimated that, as components of epigenetic mechanisms, miRNAs regulate the expression of 50–60% of the human genes without changing DNA sequences (Pillai, 2005; Engels and Hutvagner, 2006). Dozens of genes can be targeted by a single miRNA and, by the perfect or partial base-pairing with the 3'-untranslated region (UTR) of the target mRNAs, various cellular functions are induced or promoted (Pillai, 2005; Engels and Hutvagner, 2006). A broad range of physiological and pathological processes have been shown to involve one or more essential roles for miRNAs: the entire cell cycle (including embryogenesis, development, differentiation and proliferation), metabolism, cell–cell communication, cell survival and apoptosis, immune responses and oncogenesis (Bartel, 2004; Engels and Hutvagner, 2006; Burney et al., 2009; Ohlsson Teague et al., 2009, 2010).

A group of miRNAs that are differentially expressed among normal endometrium without endometriosis, eutopic endometrial tissues with endometriosis and endometriotic lesions have been demonstrated in microarray studies (Pan et al., 2007; Toloubeydokhti et al., 2008; Burney et al., 2009; Ohlsson Teague et al., 2009, 2010; Filigheddu

et al., 2010; Kuokkanen et al., 2010; Hawkins et al., 2011; Braza-Boils et al., 2014), indicating the importance of miRNAs in the pathogenesis of endometriosis. In our recent miRNA microarray analysis (Abe et al., 2013), we identified a number of miRNAs that are aberrantly expressed in human endometriotic cyst stromal cells (ECSCs) compared with human normal endometrial stromal cells (NESC) in primary culture. We found that miR-196b, one of the repressed miRNAs in the ECSCs, had anti-proliferative and pro-apoptotic functions in these cells.

We designed the present study to evaluate the role of miR-210, one of the up-regulated miRNAs in ECSCs (Abe et al., 2013), in the pathogenesis of endometriosis. Using the miR-210-transfected NESC, we observed the proliferative, anti-apoptotic and angiogenic functions of miR-210 and the possible downstream targets of this miRNA. We also evaluated the efficacy of signal transducer and activator of transcription (STAT) 3 inhibitors as promising drugs for the treatment of endometriosis.

Materials and Methods

ECSC and NESC isolation procedure and cell culture conditions

Endometriotic tissues were obtained from patients with regular menstrual cycles who had undergone a salpingo-oophorectomy or ovisceration for the treatment of ovarian endometriotic cysts ($n = 10$, aged 24–40 years), as described (Nishida et al., 2004). For ethical reasons, it was difficult to obtain endometrial tissues from healthy women; therefore, eutopic endometrial tissues were obtained from premenopausal patients who had undergone hysterectomies for subserous leiomyoma and had no evidence of endometriosis ($n = 13$, aged 35–45 years), as described (Nishida et al., 2004). None of the patients had received any hormonal treatments for at least 2 years prior to the operation. All of the specimens were confirmed as being in the mid- to late-proliferative phases according to pathological observation and/or menstrual cycles. The patients in the present study were chosen without randomization. This study was approved by the institutional review board (IRB) of the Faculty of Medicine, Oita University, and written informed consent was obtained from all patients.

ECSCs and NESC were isolated from ovarian endometriotic tissues and the eutopic endometrial tissues, respectively, by enzymatic digestion with collagenase as previously described (Nishida et al., 2004). Isolated ECSCs and NESC were cultured in Dulbecco's modified eagle medium supplemented with 100 IU/ml of penicillin, 50 mg/ml of streptomycin and 10% heat-inactivated fetal bovine serum (all obtained from Gibco-BRL, Gaithersburg, MD, USA) at 37°C in 5% CO₂ in air. ECSCs and NESC in the monolayer culture after the third passage were >99% pure as determined by immunocytochemical staining with antibodies to vimentin, CD10, cytokeratin, factor VIII and leukocyte common antigen (Nishida et al., 2004). Each experiment was performed in triplicate and repeated at least three times with the cells from at least three separate patients, except for gene expression microarray study.

Reverse transfection of miRNA precursors

NESC were transfected with precursor hsa-miR-210 (Pre-miR miRNA precursor-hsa-miR-210, Ambion, Austin, TX, USA) or negative control

precursor miRNA (Pre-miR miRNA precursor-negative control #1, Ambion) at a final concentration of 10 nM, using Lipofectamine RNAiMAX (Invitrogen, Carlsbad, CA, USA) using the reverse transfection method, as previously described (Abe et al., 2013).

Isolation of total RNA and the gene expression microarray analysis

Forty-eight hours after transfection, total RNA from cultured NESCs transfected with precursor hsa-miR-210 ($n = 3$) and NESCs ($n = 3$) transfected with negative control precursor miRNA was extracted with an RNeasy Mini kit (Qiagen, Valencia, CA, USA). The quality of the extracted RNA was confirmed by measuring the absorbance at 230, 260, and 280 nm using a spectrophotometer (NanoDrop 2000, Thermo Scientific, Wilmington, DE, USA) and by an Experion System (Bio-Rad Laboratories, Hercules, CA, USA). The samples were then subjected to a gene expression microarray analysis with a commercially available human mRNA microarray (G4845A, Human Gene Expression 4 × 44 K v2, Agilent Technologies, Santa Clara, CA, USA), which consists of 44 000 probes for 27 958 human RNAs, based on RefSeq Build 36.3, Ensemble Release 52, Unigene Build 216 and GenBank (April 2009).

Briefly, the total RNA was amplified, labeled and hybridized to a 44 K Agilent 60-mer oligomicroarray according to the manufacturer's instructions. All hybridized microarray slides were scanned by an Agilent scanner. Relative hybridization intensities and background hybridization values were calculated using Agilent Feature Extraction Software (9.5.1.1). Raw signal intensities and Flags for each probe were calculated from hybridization intensities (gProcessedSignal), and spot information (glsSaturated), according to the procedures recommended by Agilent Technologies. The flag criteria on the GeneSpring Software were, Absent (A), 'Feature is not positive and significant' and 'Feature is not above background'; Marginal (M), 'Feature is not Uniform', 'Feature is Saturated', and 'Feature is a population outlier'; Present (P), others. The raw signal intensities of six samples were then log₂-transformed and normalized by a quantile algorithm with the 'preprocess-Core' library package on Bioconductor software (Gentleman et al., 2004).

We selected probes that call the 'P' flag in both of two samples. To identify up- or down-regulated genes, we calculated Z-scores and ratios (non-log scaled fold-change) from the normalized signal intensities of each probe for comparison between NESCs transfected with precursor hsa-miR-210 and NESCs transfected with negative control precursor miRNA (Quackenbush, 2002). We then established the criteria for regulated genes: Z-score ≥ 2.0 and ratio ≥ 1.5 -fold for up-regulated genes and Z-score less than or equal to -2.0 and ratio ≤ 0.66 for down-regulated genes. All data are available at Gene Expression Omnibus via the National Center for Biotechnology Information under Accession No. GSE56854 (<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE56854>).

Ingenuity pathways analysis

To investigate possible biological interactions of differentially expressed genes, we imported data sets representing genes with an altered expression profile derived from the microarray analyses into the ingenuity pathways analysis (IPA) software (Ingenuity Systems, Redwood City, CA, USA) with the IPA knowledgebase (IPA Winter Release 2012).

Assessment of the cell viability of NESCs after miR-210 transfection and ECSCs after treatment with STAT3 inhibitors

WPI066 (573097; Merck Millipore, Darmstadt, Germany), S31-201 (573102; Merck Millipore) and cryptotanshinone (79852; Sigma-Aldrich Co., St Louis, MO, USA) were chosen as the representative STAT3 inhibitors in the present study. These STAT3 inhibitors were dissolved in dimethyl

sulfoxide (DMSO, Wako Pure Chemical, Osaka, Japan) at the concentration of 100 mM as the stock solutions. The cell viability of NESCs after miRNA transfection and ECSCs after treatment with STAT3 inhibitors [WPI066 (1–8 μ M), S31-201 (20–160 μ M) and cryptotanshinone (8–64 μ M)] was determined by a modified methylthiazolotetrazolium (MTT) assay using the CellTiter 96[®] AQ_{ueous} One Solution Cell Proliferation Assay (Promega, Madison, WI, USA). Briefly, 5×10^3 NESCs transfected with precursor hsa-miR-210 or negative control precursor miRNA by reverse transfection method were placed on 96-well flat-bottomed microplates (Corning, New York, NY, USA) and incubated for 72 h, as previously described (Abe et al., 2013). ECSCs (5×10^3 cells) were placed in 96-well flat-bottomed microplates and incubated with STAT3 inhibitors and 0.1% DMSO for 72 h. Thereafter, 20 μ l of CellTiter 96[®] AQ_{ueous} One Solution Reagent was added to each well and the cells were further incubated for 1 h. Cell viability was determined by measuring absorbance at 490 nm.

Assessment of the apoptosis of NESCs after miR-210 transfection and ECSCs after treatment with STAT3 inhibitors

We determined the apoptosis of NESCs after miRNA transfection and ECSCs after treatment with STAT3 inhibitors [WPI066 (1–8 μ M), S31-201 (20–160 μ M) and cryptotanshinone (8–64 μ M)] by the direct determination of nucleosomal DNA fragmentation using an enzyme-linked immunosorbent assay (ELISA) (Cell Death Detection ELISA, Roche Diagnostics) (Abe et al., 2013). NESCs after miRNA transfection (5×10^3 cells) were placed on 96-well flat-bottomed microplates (Corning). After 72 h of culture, the cells were lysed according to the manufacturer's instructions, followed by centrifugation (200g, 5 min). The mono- and oligonucleosomes contained in the supernatants were determined using an anti-histone-biotin antibody. The concentration of nucleosomes-antibody was evaluated by measuring the absorbance at 405 nm using 2,2'-azino-di(3-ethylbenzthiazolinesulphonate) as a substrate. The effects of STAT3 inhibitors on ECSCs were also examined.

Assessment of the activities of caspase-3 and caspase-7 in NESCs after miR-210 transfection and ECSCs after treatment with STAT3 inhibitors

The apoptosis of NESCs after miRNA transfection and ECSCs after treatment with STAT3 inhibitors [WPI066 (1–8 μ M), S31-201 (20–160 μ M) and cryptotanshinone (8–64 μ M)] was evaluated by the Caspase-Glo 3/7 Assay (Promega) as described (Abe et al., 2013). The assay is a luminescent assay that measures the activities of caspase-3 and caspase-7. NESCs after miRNA transfection (5×10^3 cells) were placed on 96-well flat-bottomed microplates (Promega). After 72 h of culture, Caspase-Glo 3/7 reagent was added to each well. The plates were gently shaken for 60 min at room temperature, and then we measured the luminescence in a plate-reading luminometer. The effects of STAT3 inhibitors on ECSCs were also examined.

Assessment of the effects of miR-210 on the STAT3 and VEGF-A mRNA expression in NESCs

The effects of miR-210 on the expressions of possible downstream target genes in NESCs were evaluated by quantitative RT-PCR. The STAT3 and vascular endothelial growth factor A (VEGF-A) were chosen as candidate molecules for evaluation. Briefly, NESCs transfected with precursor hsa-miR-210 or negative control precursor miRNA were cultured in 10-cm culture dishes (Corning). Forty-eight hours after incubation, total RNA was extracted from the cultured NESCs using an miRNeasy Mini kit (Qiagen),

as described above. cDNA was then synthesized from 1 µg of total RNA using the Reverse Transcription System (Promega).

The quantitative RT-PCR was carried out with a LightCycler 480 (Roche Diagnostics GmbH, Penzberg, Germany) using TaqMan Universal PCR Master Mix II with specific primers for STAT3 (Assay ID: Hs00374280_m1, Applied Biosystems, Carlsbad, CA, USA), VEGF-A (Assay ID: Hs00900055_m1, Applied Biosystems), or glyceraldehyde 3-phosphate dehydrogenase (GAPDH) (Assay ID: Hs02758991_g1, Applied Biosystems), as described (Abe et al., 2013). The expression levels of STAT3 and VEGF-A mRNA relative to GAPDH mRNA were calculated from a standard curve.

Assessment of the effects of miR-210 on the phosphorylated and total STAT3 protein levels in NESC

The effects of miR-210 on the phosphorylated and total STAT3 protein levels in NESC were also evaluated, using the Cell-Based Human/Mouse/Rat STAT3 (Tyr705) Phosphorylation ELISA Kit (RayBiotech, Inc., Norcross, GA, USA). Briefly, 3×10^4 cells transfected with precursor hsa-miR-210 or negative control precursor miRNA were placed on a white-walled 96-well microplate (RayBiotech). After 48 h of culture, the plates were processed to measure the absorbance at 405 nm, according to the manufacturer's instructions.

Assessment of the VEGF-A protein levels in the supernatant of NESC after miR-210 transfection and ECSC after treatment with STAT3 inhibitors

The VEGF-A protein levels in the supernatant of NESC after miRNA transfection and those of ECSC after treatment with STAT3 inhibitors were determined by ELISA.

Briefly, subconfluent NESC after miRNA transfection or ECSC after treatment with STAT3 inhibitors were cultured in 24-well culture plates (Corning). After a 24-h culture, the supernatants were then collected and stored at -70°C until assay. The concentration of VEGF-A was determined using commercially available ELISA kits (Human VEGF Immunoassay, R&D systems, Minneapolis, MN, USA). The sensitivity of the assay for VEGF was 9.0 pg/ml.

Statistical analysis

Data were obtained from triplicate samples and are presented as percentages relative to the corresponding controls, as mean \pm SD, and were appropriately analyzed by the Bonferroni test and the Student t-test with Sigmaplot 11.2 (Systat Software, San Jose, CA, USA). Values of $P < 0.05$ were considered significant.

Results

Identification of candidate genes regulated by miR-210 transfection in NESC

Among the 27 958 mRNAs examined by the gene expression microarray, 94 up-regulated mRNAs and 229 down-regulated mRNAs were identified statistically by using the criteria described above. Using the IPA software to find the known target and candidate downstream signaling networks of miR-210, we identified 29 downstream molecules (Table I).

Atrophin 1 (ATN1), elongation factor, RNA polymerase II, 2 (ELL2), histone deacetylase 2 (HDAC2), STAT3 and STAT6 were detected as the transcription factors regulated by miR-210 (Table I). Of these

Table I Regulation of candidate miR-210-target molecules in NESC detected by microarray and ingenuity pathways analyses.

Gene family	Gene symbol	Regulation ^a	Z-score	Ratio
Transcription regulator	ATN1	↓	-3.29	0.36
	ELL2	↓	-3.05	0.49
	HDAC2	↑	3.95	3.61
	STAT3	↑	2.14	2.01
	STAT6	↓	-2.45	0.51
Growth factor	PGF	↑	6.18	7.19
Cytokine	IL11	↑	3.56	3.15
Kinase	ERBB2	↓	-2.27	0.52
	MAP4K4	↓	-2.47	0.45
	RPS6KA5	↑	3.58	2.53
	PTPRF	↓	-2.59	0.55
Peptidase	PLAU	↑	4.77	1.98
Transmembrane receptor	THBD	↑	3.56	2.84
Enzyme	GNA13	↑	2.21	1.9
	HMGCR	↑	2.56	1.93
	IRS2	↓	-3.39	0.38
	MX1	↓	-3.0	0.45
	PNPLA3	↑	2.87	2.3
	PPAT	↑	2.51	2.05
Other	ANGPTL2	↑	2.52	1.96
	ANGPTL4	↑	3.75	2.77
	COL8A1	↓	-2.66	0.51
	DAB2	↓	-2.49	0.57
	IGFBP3	↓	-2.58	0.41
	NF2	↓	-2.72	0.57
	SKP2	↑	3.18	2.79
	SYNPO	↓	-3.65	0.34
	THBS1	↓	-4.24	0.47
U2AF2	↓	-3.78	0.42	

ANGPTL2, angiopoietin-like 2; ANGPTL4, angiopoietin-like 4; ATN1, atrophin 1; ELL2, elongation factor, RNA polymerase II, 2; COL8A1, collagen, type VIII, α 1; DAB2, disabled homolog 2; ERBB2, v-erb-b2 erythroblastic leukemia viral oncogene homolog 2; GNA13, guanine nucleotide binding protein, α 13; HDAC2, histone deacetylase 2; IGFBP3, insulin-like growth factor binding protein 3; IL11, interleukin 11; IRS2, insulin receptor substrate 2; HMGCR, 3-hydroxy-3-methylglutaryl-CoA reductase; MAP4K4, mitogen-activated protein kinase kinase kinase 4; MX1, myxovirus resistance 1, interferon-inducible protein p78; NF2, neurofibromin 2; PGF, placental growth factor; PLAU, plasminogen activator, urokinase; PNPLA3, patatin-like phospholipase domain containing 3; PPAT, phosphoribosyl pyrophosphate amidotransferase; PTPRF, protein tyrosine phosphatase, receptor type, F; RPS6KA5, ribosomal protein S6 kinase, 90 kDa, polypeptide 5; SKP2, S-phase kinase-associated protein 2; STAT, signal transducer and activator of transcription; SYNPO, synaptopodin; THBD, thrombomodulin; THBS1, thrombospondin 1; U2AF2, U2 small nuclear RNA auxiliary factor 2.

^aGenes with Z-score ≥ 2.0 and ratio ≥ 1.5 -fold were defined as up-regulated, and those with Z-score less than or equal to -2.0 and ratio ≤ 0.66 were defined as down-regulated.

transcription factors, we focused on STAT3 as a key molecule regarding the pathogenesis of endometriosis (Fig. 1) and used it in the further experiments. Downstream targets of STAT3 include collagen type IV, interleukin 11 (IL11), myxovirus resistance 1, interferon-inducible protein p78 (MX1), plasminogen activator, urokinase (PLAU), ribosomal protein S6 kinase, 90 kDa, polypeptide 5 (RPS6KA5), sterol regulatory element binding transcription factor 1 (SREBF1), synaptopodin (SYNPO), thrombospondin 1 (THBS1) and VEGF.




Figure 1 The downstream network of the micro RNA (miR)-210-STAT3 signaling pathway detected by Ingenuity pathways analysis. The genes that are shaded were determined to be significant from the statistical analysis. The genes in red are up-regulated and those in green are down-regulated. The intensity of the shading shows to what degree each gene was up- or down-regulated. A solid line represents a direct interaction between the two gene products, and a dotted line means that there is an indirect interaction. ANGPTL2, angiotensin-like 2; ANGPTL4, angiotensin-like 4; Fgfr, fibroblast growth factor receptor; GNA13, guanine nucleotide binding protein, α 13; HDAC2, histone deacetylase 2; IGFBP3, insulin-like growth factor binding protein 3; IL11, interleukin 11; IRS2, insulin receptor substrate 2; MAP4K4 Δ , mitogen-activated protein kinase kinase kinase 4; Mmp, matrix metalloproteinase; MX1, myxovirus resistance 1, interferon-inducible protein p78; NF2, neurofibromin 2; PGF, placental growth factor; PLAU, plasminogen activator, urokinase; PNPLA3, patatin-like phospholipase domain containing 3; PPAT, phosphoribosyl pyrophosphate amidotransferase; PTPase, phosphotyrosine phosphatase; PTPN1, protein tyrosine phosphatase, non-receptor type 1; PTPRF, protein tyrosine phosphatase, receptor type, F; RPS6KA5, ribosomal protein S6 kinase, 90 kDa, polypeptide 5; SKP2, S-phase kinase-associated protein 2; SRC, v-src sarcoma; SREBF1, sterol regulatory element binding transcription factor 1; SYNPO, synaptopodin; THBD, thrombomodulin; THBS1, thrombospondin 1; U2AF2, U2 small nuclear RNA auxiliary factor 2; VEGF, vascular endothelial cell growth factor.

The mRNA expression of some molecules, such as early growth response 2 (EGR2), homeobox A1 (HOXA1), sterol regulatory element binding transcription factor 1 (SREBF1) and protein tyrosine phosphatase, non-receptor type 1 (PTPN1), seem to be unaffected by miR-210 transfection. It is speculated that miR-210 may affect the function of these downstream target molecules without changing their mRNA expression.

Enhanced cell viability of NESC by miR-210 transfection

The effects of miR-210 on the cell viability of NESC were evaluated by a modified MTT assay. As shown in Fig. 2A, the viable cell number was significantly increased by the transfection of miR-210 precursor.

Inhibition of apoptosis of NESC by miR-210 transfection

The anti-apoptotic effects of miR-210 on NESC were determined by the Cell Death Detection ELISA assay. As shown in Fig. 2B, the transfection of miR-210 precursor significantly inhibited the apoptosis of ECSCs.

Inhibition of caspase-3/7 activity in NESC by miR-210 transfection

The anti-apoptotic effects of miR-210 on ECSCs were also assessed by evaluating the activities of caspase-3 and caspase-7. As shown in Fig. 2C, the transfection of miR-210 precursor significantly inhibited the activities of caspase-3 and caspase-7 in ECSCs.




Figure 2 The effects of miR-210 transfection on human NESC cells. (A) cell viability, (B) apoptotic cells, (C) caspase-3/7 activity, (D) VEGF mRNA levels, (E) VEGF protein levels, (F) STAT3 mRNA levels, (G) total STAT3 protein levels and (H) phosphorylated STAT3 protein levels of NESC cells after miR-210 transfection ($n = 3$). The data are presented as percentages relative to the values of NESC cells transfected with negative control precursor miRNA ($n = 3$). * $P < 0.05$, ** $P < 0.005$, *** $P < 0.0005$ versus negative controls (Student's t -test). Representative results are shown.

Enhanced VEGF mRNA and protein expression in NESC cells by miR-210 transfection

The effects of miR-210 transfection on the VEGF mRNA and protein expression in NESC cells were assessed by RT-PCR and ELISA, respectively. The transfection of miR-210 precursor significantly enhanced the VEGF mRNA and protein expression in NESC cells (Fig. 2D and E).

Induction of STAT3 expression and activation in NESC cells by miR-210 transfection

The effects of miR-210 transfection on the STAT3 mRNA and protein expression and STAT3 activation in NESC cells were assessed. As shown in Fig. 2F–H, the transfection of miR-210 precursor significantly enhanced the STAT3 mRNA and protein expression and STAT3 activity in NESC cells.

Suppression of cell viability of ECSCs by STAT3 inhibitors

The effects of STAT3 inhibitors on the cell viability of ECSCs were evaluated by a modified MTT assay. As shown in Fig. 3A, the viable

cell number was significantly decreased by the addition of STAT3 inhibitors.

Induction of apoptosis of ECSCs by STAT3 inhibitors

The effects of STAT3 inhibitors on the apoptosis of ECSCs were determined by the Cell Death Detection ELISA assay. As shown in Fig. 3B, the STAT3 inhibitors significantly induced the apoptosis of ECSCs.

The pro-apoptotic effects of STAT3 inhibitors on ECSCs were also assessed by evaluating the activities of caspase-3 and caspase-7. The STAT3 inhibitors significantly enhanced the activities of caspase-3 and caspase-7 in ECSCs (Fig. 3C).

Suppression of VEGF production in ECSCs by STAT3 inhibitors

The effects of STAT3 inhibitors on the VEGF protein secretion of ECSCs were assessed by ELISA. As shown in Fig. 3D, the STAT3 inhibitors




Figure 3 The effects of STAT3 inhibitors on human ECSCs. **(A)** cell viability, **(B)** apoptotic cells, **(C)** caspase-3/7 activity and **(D)** VEGF protein levels of ECSCs treated with STAT3 inhibitors [WP1066 (1–8 μM), S3I-201 (20–160 μM) and cryptotanshinone (8–64 μM)] ($n = 3$). The data are presented as percentages relative to the values of untreated ECSCs ($n = 3$). * $P < 0.05$, ** $P < 0.005$, *** $P < 0.0005$ versus negative controls (Bonferroni's test). Representative results are shown.

significantly decreased the VEGF protein levels in the supernatant of ECSC culture.

Discussion

We conducted the present study to identify the role of miR-210, one of the up-regulated miRNAs in ECSCs (Abe et al., 2013), in the pathogenesis of endometriosis. By examining the miR-210 transfection in NESC, the gene expression microarray technique and an IPA, we found a variety of candidate molecules as the downstream targets of miR-210. We then focused on STAT3 and performed further functional experiments. We found that miR-210 promoted the proliferation, resistance to apoptosis and VEGF production through STAT3 activation in NESC, whereas STAT3 inhibitors inhibited the proliferation and VEGF production of ECSCs and induced the apoptosis of these cells. These findings suggest that miR-210 induces NESC to differentiate into the endometriotic phenotype, which is characterized by proliferative, anti-apoptotic and angiogenic features. We also suggest that up-regulated miR-210 expression in ECSCs is involved in the creation of the endometriosis-specific cellular dysfunctions as part of epigenetic mechanisms. It is considered that activation of the miR-210-STAT3-VEGF axis is important in the pathogenesis of endometriosis. Based on our findings, we speculate that STAT3 inhibitors could be promising for the treatment of endometriosis.

As summarized in Table II, the roles of several miRNAs in the pathogenesis of endometriosis have been demonstrated (Lin et al., 2012, 2014; Abe et al., 2013; Adamek et al., 2013; Shen et al., 2013; Hsu et al., 2014; Shi et al., 2014). We showed that miR-196b, one of the repressed miRNAs in ECSCs, has anti-proliferative and pro-apoptotic functions in these cells by targeting c-myc and Bcl-2 (Abe et al., 2013). Given the diverse roles that miRNAs play in numerous aspects of cellular functions, it is not surprising that they play key regulatory roles in the pathogenesis of endometriosis. All the miRNAs listed in Table II belong to a different cluster from miR-210. VEGFA was the only target molecule shared with miR-210 and miR-199a-5p. Evaluations of the functions of each aberrantly expressed miRNA are necessary in further research on endometriosis.

Several studies showed that in various cell types, hypoxia was followed by the induction of miR-210 expression through hypoxia-inducible factors (Crosby et al., 2009; Zhang et al., 2009; Huang et al., 2010). A variety of tumors exhibit increased miR-210 expression (Gee et al., 2010; Stephen and Joseph, 2010; Buffa et al., 2011; Yuk et al., 2012). The identified functions of miR-210 include the modulation of cell proliferation, differentiation, cell cycle arrest, cell migration, DNA repair, chromatin remodeling, apoptosis, angiogenesis and metabolism (Fasanaro et al., 2008; Crosby et al., 2009; Zhang et al., 2009; Huang et al., 2010). A large number of target molecules of miR-210 have been reported: apoptosis-inducing factor, mitochondrion-associated, 3 (AIFM3), caspase-8 associated protein-2 (CASP8AP2), death-associated

Table II Reported function of aberrantly expressed miRNAs in endometriosis.

miRNA	Expression	Target gene expression	Function	References
miR-20a	Up-regulated in ESCs	Down-regulation of DUSP2	Augmentation of PGE2-induced FGF-9 expression, induction of prolonged ERK phosphorylation and induction of EGR-1, CYR61 and osteopontin	Lin <i>et al.</i> (2012)
miR-23a/b	Down-regulated in endometriotic tissues and eutopic endometrium of endometriosis patients	Down-regulation of steroidogenic factor-1	Inhibition of estrogen biosynthesis	Shen <i>et al.</i> (2013)
miR-145	Down-regulated in endometriotic tissues	Down-regulation of fascin-1, SOX2, MSI2, OCT4, KLF4, podocalyxin, JAM-A, and PAI-1 Up-regulation of ACTG2 and transgelin	Inhibition of proliferation Induction of invasiveness Reduction of side population and aldehyde dehydrogenase-1 activity	Adammek <i>et al.</i> (2013)
miR-183	Down-regulated in endometriotic tissues	ND	Induction of apoptosis, Inhibition of invasiveness	Shi <i>et al.</i> (2014)
miR-196b	Down-regulated in ESCs	Suppression of c-myc and bcl-2	Inhibition of proliferation Induction of apoptosis	Abe <i>et al.</i> (2013)
miR-199a-5p	Down-regulated in serum of endometriotic patients	VEGFA	Inhibition of proliferation, motility and angiogenesis	Hsu <i>et al.</i> (2014)
miR-302a	Up-regulated in ESCs	Suppression of COUP-TFII	Induction of COX-2	Lin <i>et al.</i> (2014)

ACTG2, smooth-muscle actin isoform γ 2; COUP-TFII, chicken ovalbumin upstream promoter-transcription factor II; COX-2, cyclooxygenase-2; CYR61, cysteine-rich angiogenic inducer 61; DUSP2, dual-specificity phosphatase-2; EGR-1, early growth response protein-1; ESCs, endometriotic stromal cells; FGF-9, fibroblast growth factor-9; JAM-A, junctional adhesion molecule A; KLF4, Kruppel-like-factor 4; MSI2, Musashi-2; ND, not described; OCT4, octamer 4; PAI-1, plasminogen activator inhibitor 1; PGE2, prostaglandin E2; SOX2, SRY-box 2; VEGFA, vascular endothelial growth factor-A.

protein kinase I (DAPK1), E2F transcription factor 3 (E2F3), ephrin-A3 (EFNA3), fibroblast growth factor receptor-like 1 (FGFRL1), HOXA1, HOXA3, iron-sulfur cluster scaffold proteins (ISCU), MAX binding protein (MNT), PTPN1, RAD52 homolog (RAD52), tumor protein p53-inducible protein 11 (TP53I11), VEGF and VEGF receptor 2 (VEGFR2) (Fasanaro *et al.*, 2008; Chan *et al.*, 2009; Crosby *et al.*, 2009; Zhang *et al.*, 2009; Huang *et al.*, 2010; Noman *et al.*, 2012). Since miR-210 is well known as an angiogenic miRNA, we focused on the regulatory mechanism of VEGF expression by miR-210. In this setting, we found miR-210-STAT3-VEGF axis to be an important pathway (Fig. 1). Interestingly, STAT3 has been also shown to regulate cell proliferation and apoptosis (Siveen *et al.*, 2014).

The treatment of endometriosis has been a challenge. Non-steroidal anti-inflammatory agents, agonists of GnRH, progestogens, androgens and contraceptive steroids have all been tried for patients with endometriosis (Committee of the American Society for Reproductive Medicine, 2004), and several surgical and medical strategies have been conducted, with varying degrees of efficacy. The results of the present study suggest that STAT3 inhibitors have potential as a treatment for endometriosis. In fact, several STAT3 inhibitors are now in preclinical use as anticancer drugs (Furqan *et al.*, 2013). BBI608 (Langleben *et al.*, 2013) is now in phase III trial, whereas OPB-31121 (ClinicalTrials.gov Identifier: NCT00955812, Food and Drug Administration, USA) and BBI503 (Laurie *et al.*, 2014) are in phase I trial.

Siveen *et al.* (2014) demonstrated that the activation of STAT3 inhibits apoptosis and induces cell proliferation, angiogenesis, metastasis and invasion. The STAT family of transcription factors, in their inactive form, is initially located in the cytoplasm of the cell. With the stimulation provided by extracellular signals, such as cytokines, Janus kinases (JAKs), growth factors

and hormones are activated thereby inducing the phosphorylation of STAT3 on a tyrosine residue in its COOH terminus (Siveen *et al.*, 2014). Phosphorylated STAT3 proteins activate the transcription of their target genes after they dimerize, translocate into the nucleus and bind with DNA. As shown in Fig. 1, it is suggested that miR-210 induces STAT3 mRNA and protein expression/phosphorylation through protein tyrosine phosphatase, non-receptor type I (PTPN1). Interestingly, as shown in Table I, PTPN1 mRNA expression was not affected by miR-210 transfection in NESCs. Further examinations are necessary to elucidate the precise mechanisms. In the present study, we chose three representative STAT3 inhibitors with diverse mechanisms of action. WP1066 inhibits STAT3 activity by inhibiting its upstream transcription factor JAK (Ferrajoli *et al.*, 2007), whereas S31-201 inhibits STAT3 activity by inhibiting STAT3 dimerization (Fletcher *et al.*, 2009). Cryptotanshinone inhibits STAT3 activity by inhibiting its phosphorylation (Lu *et al.*, 2013). These STAT3 inhibitors show stronger effects in comparison with miR-210 transfection, suggesting that STAT3 can regulate cell viability by a mechanism independent of miR-210. Additional studies are necessary to fully understand the action of STAT3 inhibitors.

Another promising potential strategy for treating endometriosis is miRNA-targeting therapeutics in part because the use of small miRNA-inhibitor oligomers (e.g. DNA, DNA analogs and RNA) can readily accomplish miRNA inhibition (Hemida *et al.*, 2010; Kota and Balasubramanian, 2010). In addition, the DNA analogs known as peptide nucleic acids (PNAs)—in which the sugarphosphate backbone is replaced by *N*-(2-aminoethyl)glycine units (Gambari *et al.*, 2011)—efficiently hybridize with their target miRNAs and inhibit their functioning. The *in vitro* and *in vivo* effects of PNAs targeting miR-210 have been described (Gambari *et al.*, 2011).

Angiogenesis is a critical step in the establishment of endometriosis, and endometriotic lesions are highly vascularized (Donnez et al., 1998). Endothelial cell proliferation and migration for neovascularization are stimulated by VEGF, the main sources of which are endometriotic tissue and peritoneal macrophages, and mesothelial cells (Donnez et al., 1998). As yet another endometriosis treatment strategy, anti-angiogenic therapy against VEGF holds great promise. miR-210-targeting molecules and STAT3 inhibitors can also act as anti-angiogenic agents by suppressing ECSCs' production of VEGF.

For ethical reasons, it is difficult to obtain endometrial tissues from healthy women. Instead, control endometrial tissues were obtained from premenopausal patients who had undergone hysterectomies for subserosal leiomyoma and had no evidence of endometriosis. Gene expression patterns of NESCs may be altered by the presence of leiomyoma (Karmon et al., 2014). Further limitations of the present study are that we have not performed a loss-of-function study using ECSCs transfected with miR-210-antagonist and the effects of STAT3 inhibitors were evaluated only in ECSCs and not in NESCs.

In conclusion, by using miR-210 transfection in NESCs, a gene expression microarray technique and an IPA, we found that STAT3 is one of the key target molecules of miR-210. The miR-210 transfection directed the induction of cell proliferation and VEGF production and the inhibition of apoptosis in NESCs through STAT3 activation, whereas STAT3 inhibitors blocked the proliferation and VEGF production of ECSCs and induced apoptosis of these cells. These findings suggest that up-regulated miR-210 expression in human ECSCs is involved in the creation of cellular dysfunctions that are disease-specific features of endometriosis, and we propose these may involve epigenetic mechanisms. In addition, STAT3 inhibitors are promising candidates for the treatment of endometriosis. Further studies on the repertoire of aberrantly expressed miRNAs, interacting miRNA–target mRNA associations and the regulation and mechanisms of action of miRNA may provide useful information about the pathogenesis of endometriosis.

Acknowledgements

We are indebted to Dr Chisato Nakada and Dr Yoshiyuki Tsukamoto, Department of Molecular Pathology, Faculty of Medicine, Oita University, Japan, for the review of the work.

Authors' roles

K.N., M.M. and H.N. participated in the study design, analysis and manuscript drafting. M.O., W.A., Y.A., Y.K. and K.K. executed the study.

Funding

This work was supported in part by Grants-in-Aid for Scientific Research from the Japan Society for the Promotion of Science (no. 13237327 to K.N., no. 25861500 to Y.K. and no. 23592407 to H.N.).

Conflict of interest

None declared.

References

- Abe W, Nasu K, Nakada C, Kawano Y, Moriyama M, Narahara H. miR-196b targets c-myc and Bcl-2 expression, inhibits proliferation and induces apoptosis in endometriotic stromal cells. *Hum Reprod* 2013;**28**:750–761.
- Adammek M, Greve B, Kässens N, Schneider C, Brüggemann K, Schüring AN, Starzinski-Powitz A, Kiesel L, Götte M. MicroRNA miR-145 inhibits proliferation, invasiveness, and stem cell phenotype of an in vitro endometriosis model by targeting multiple cytoskeletal elements and pluripotency factors. *Fertil Steril* 2013;**99**:1346–1355.e5.
- Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 2004;**116**:281–297.
- Braza-Boils A, Mari-Alexandre J, Gilabert J, Sánchez-Izquierdo D, España F, Estellés A, Gilabert-Estellés J. MicroRNA expression profile in endometriosis: its relation to angiogenesis and fibrinolytic factors. *Hum Reprod* 2014;**29**:978–988.
- Buffa FM, Camps C, Winchester L, Snell CE, Gee HE, Sheldon H, Taylor M, Harris AL, Ragoussis J. microRNA-associated progression pathways and potential therapeutic targets identified by integrated mRNA and microRNA expression profiling in breast cancer. *Cancer Res* 2011;**71**:5635–5645.
- Burney RO, Hamilton AE, Aghajanova L, Vo KC, Nezhat CN, Lessey BA, Giudice LC. MicroRNA expression profiling of eutopic secretory endometrium in women with versus without endometriosis. *Mol Hum Reprod* 2009;**15**:625–631.
- Chan SY, Zhang YY, Hemann C, Mahoney CE, Zweier JL, Loscalzo J. MicroRNA-210 controls mitochondrial metabolism during hypoxia by repressing the iron–sulfur cluster assembly proteins ISCU1/2. *Cell Metab* 2009;**10**:273–284.
- Committee of the American Society for Reproductive Medicine. Endometriosis and infertility. *Fertil Steril* 2004;**81**:1441–1446.
- Crosby ME, Kulshreshtha R, Ivan M, Glazer PM. MicroRNA regulation of DNA repair gene expression in hypoxic stress. *Cancer Res* 2009;**69**:1221–1229.
- Donnez J, Smoes P, Gillerot S, Casanas-Roux F, Nisolle M. Vascular endothelial growth factor (VEGF) in endometriosis. *Hum Reprod* 1998;**13**:1686–1690.
- Engels BM, Hutvagner G. Principles and effects of microRNA-mediated post-transcriptional gene regulation. *Oncogene* 2006;**25**:6163–6169.
- Fasanaro P, D'Alessandra Y, Di Stefano V, Melchionna R, Romani S, Pompilio G, Capogrossi MC, Martelli F. MicroRNA-210 modulates endothelial cell response to hypoxia and inhibits the receptor tyrosine kinase ligand Ephrin-A3. *J Biol Chem* 2008;**283**:15878–15883.
- Ferrajoli A, Faderl S, Van Q, Koch P, Harris D, Liu Z, Hazan-Halevy I, Wang Y, Kantarjian HM, Priebe W et al. WPI066 disrupts Janus kinase-2 and induces caspase-dependent apoptosis in acute myelogenous leukemia cells. *Cancer Res* 2007;**67**:11291–11299.
- Filigheddu N, Gregnanin I, Porporato PE, Surico D, Perego B, Galli L, Patrignani C, Graziani A, Surico N. Differential expression of microRNAs between eutopic and ectopic endometrium in ovarian endometriosis. *J Biomed Biotech* 2010;**2010**:369549.
- Fletcher S, Singh J, Zhang X, Yue P, Page BD, Sharmeen S, Shahani VM, Zhao W, Schimmer AD, Turkson J et al. Disruption of transcriptionally active Stat3 dimers with non-phosphorylated, salicylic acid-based small molecules: potent in vitro and tumor cell activities. *Chembiochem* 2009;**10**:1959–1964.
- Furqan M, Akinleye A, Mukhi N, Mittal V, Chen Y, Liu D. STAT inhibitors for cancer therapy. *J Hematol Oncol* 2013;**6**:90.
- Gambari R, Fabbri E, Borgatti M, Lampronti I, Finotti A, Brognara E, Bianchi N, Manicardi A, Marchelli R, Corradini R. Targeting microRNAs involved in human diseases: a novel approach for modification of gene expression and drug development. *Biochem Pharmacol* 2011;**82**:1416–1429.

- Gee HE, Camps C, Buffa FM, Patiar S, Winter SC. hsa-mir-210 is a marker of tumor hypoxia and a prognostic factor in head and neck cancer. *Cancer* 2010; **116**:2148–2158.
- Gentleman RC, Carey VJ, Bates DM, Bolstad B, Dettling M, Dudoit S, Ellis B, Gautier L, Ge Y, Gentry J et al. Bioconductor: open software development for computational biology and bioinformatics. *Genome Biol* 2004; **5**:R80.
- Giudice LC, Kao LC. Endometriosis. *Lancet* 2004; **364**:1789–1799.
- Hawkins SM, Creighton CJ, Han DY, Zariff A, Anderson ML, Gunaratne PH, Matzuk MM. Functional MicroRNA involved in endometriosis. *Mol Endocrinol* 2011; **25**:821–832.
- Hemida MG, Ye X, Thair S, Yang D. Exploiting the therapeutic potential of microRNAs in viral diseases: expectations and limitations. *Mol Diagn Ther* 2010; **14**:271–282.
- Hsu CY, Hsieh TH, Tsai CF, Tsai HP, Chen HS, Chang Y, Chuang HY, Lee JN, Hsu YL, Tsai EM. miRNA-199a-5p regulates VEGFA in endometrial mesenchymal stem cells and contributes to the pathogenesis of endometriosis. *J Pathol* 2014; **232**:330–343.
- Huang X, Le Q-T, Giaccia AJ. MiR-210—micromanager of the hypoxia pathway. *Trends Mol Med* 2010; **16**:230–237.
- Karmon AE, Cardozo ER, Rueda BR, Styer AK. MicroRNAs in the development and pathobiology of uterine leiomyomata: does evidence support future strategies for clinical intervention? *Hum Reprod Update* 2014; **20**:670–687.
- Kota SK, Balasubramanian S. Cancer therapy via modulation of micro RNA levels: a promising future. *Drug Discov Today* 2010; **15**:733–740.
- Kuokkanen S, Chen B, Ojalvo L, Benard L, Santoro N, Pollard JW. Genomic profiling of microRNAs and messenger RNAs reveals hormonal regulation in microRNA expression in human endometrium. *Biol Reprod* 2010; **82**:791–801.
- Langleben A, Supko JG, Hotte SJ, Batist G, Hirte HW, Rogoff H, Li Y, Li W, Kerstein D, Leggett D et al. A dose-escalation phase I study of a first-in-class cancer stemness inhibitor in patients with advanced malignancies. *J Clin Oncol* 2013; **31**(Suppl):abstr 2542.
- Laurie SA, Jonker DJ, Edenfield WJ, Stephenson J, Keller D, Hitron M, Li W, Li Y, Gada K, Gao Y et al. A phase I dose-escalation study of BBI503, a first-in-class cancer stemness kinase inhibitor in adult patients with advanced solid tumors. *J Clin Oncol* 2014; **32**(suppl):abstr 2527.
- Lin S-C, Li Y-H, Wu M-H, Chang Y-F, Lee D-K, Tsai SY, Tsai M-J, Tsai S-J. Suppression of COUP-TFII by Proinflammatory Cytokines Contributes to the Pathogenesis of Endometriosis. *J Clin Endocrinol Metab* 2014; **99**:E427–E437.
- Lin SC, Wang CC, Wu MH, Yang SH, Li YH, Tsai SJ. Hypoxia-induced microRNA-20a expression increases ERK phosphorylation and angiogenic gene expression in endometriotic stromal cells. *J Clin Endocrinol Metab* 2012; **97**:E1515–E1523.
- Lu L, Li C, Li D, Wang Y, Zhou C, Shao W, Peng J, You Y, Zhang X, Shen X. Cryptotanshinone inhibits human glioma cell proliferation by suppressing STAT3 signaling. *Mol Cell Biochem* 2013; **381**:273–282.
- Nasu K, Kawano Y, Tsukamoto Y, Takano M, Takai N, Li H, Furukawa Y, Abe W, Moriyama M, Narahara H. Aberrant DNA methylation status of endometriosis: epigenetics as the pathogenesis, biomarker and therapeutic target. *J Obstet Gynaecol Res* 2011a; **37**:683–695.
- Nasu K, Nishida M, Kawano Y, Tsuno A, Abe W, Yuge A, Takai N, Narahara H. Aberrant expression of apoptosis-related molecules in endometriosis: a possible mechanism underlying the pathogenesis of endometriosis. *Reprod Sci* 2011b; **18**:206–218.
- Nasu K, Kawano Y, Kai K, Aoyagi Y, Abe W, Okamoto M, Narahara H. Aberrant histone modification in endometriosis. *Front Biosci* 2014; **19**:1202–1214.
- Nishida M, Nasu K, Fukuda J, Kawano Y, Narahara H, Miyakawa I. Down regulation of interleukin-1 receptor expression causes the dysregulated expression of CXC chemokines in endometriotic stromal cells: a possible mechanism for the altered immunological functions in endometriosis. *J Clin Endocrinol Metab* 2004; **89**:5094–5100.
- Noman MZ, Buart S, Romero P, Ketari S, Janji B, Mari B, Mami-Chouaib F, Chouaib S. Hypoxia-inducible miR-210 regulates the susceptibility of tumor cells to lysis by cytotoxic T cells. *Cancer Res* 2012; **72**:4629–4641.
- Ohlsson Teague EMC, Van der Hoek KH, Van der Hoek MB, Perry N, Wagaarachchi P, Robertson SA, Print CG, Hull ML. MicroRNA-regulated pathways associated with endometriosis. *Mol Endocrinol* 2009; **23**:265–275.
- Ohlsson Teague EMC, Print CG, Hull ML. The role of microRNAs in endometriosis and associated reproductive conditions. *Hum Reprod Update* 2010; **16**:142–165.
- Pan Q, Luo X, Toloubeydokhti T, Chegini N. The expression profile of micro-RNA in endometrium and endometriosis and the influence of ovarian steroids on their expression. *Mol Hum Reprod* 2007; **13**:797–806.
- Pillai RS. MicroRNA function: multiple mechanisms for a tiny RNA? *RNA* 2005; **11**:1753–1761.
- Quackenbush J. Microarray data normalization and transformation. *Nat Genet* 2002; **32**(Suppl):496–501.
- Shen L, Yang S, Huang W, Xu W, Wang Q, Song Y, Liu Y. MicroRNA23a and microRNA23b deregulation derepresses SF-1 and upregulates estrogen signaling in ovarian endometriosis. *J Clin Endocrinol Metab* 2013; **98**:1575–1582.
- Shi XY, Gu L, Chen J, Guo XR, Shi YL. Downregulation of miR-183 inhibits apoptosis and enhances the invasive potential of endometrial stromal cells in endometriosis. *Int J Mol Med* 2014; **33**:59–67.
- Siveen KS, Sikka S, Surana R, Dai X, Zhang J, Kumar AP, Tan BK, Sethi G, Bishayee A. Targeting the STAT3 signaling pathway in cancer: role of synthetic and natural inhibitors. *Biochim Biophys Acta* 2014; **1845**:136–154.
- Stephen YC, Joseph L. MicroRNA-210: a unique and pleiotropic hypoxamir. *Cell Cycle* 2010; **9**:1072–1083.
- Toloubeydokhti T, Pan Q, Luo X, Bukulmez O, Chegini N. The expression and ovarian steroid regulation of endometrial micro-RNAs. *Reprod Sci* 2008; **15**:993–1001.
- Yuk CC, Jaideep B, Sang YC, Chandan KS. miR-210: the master hypoxamir. *Microcirculation* 2012; **19**:215–223.
- Zhang Z, Sun H, Dai H, Walsh RM, Imakura M, Schelter J, Burchard J, Dai X, Chang AN, Diaz RL et al. MicroRNA miR-210 modulates cellular response to hypoxia through the MYC antagonist MNT. *Cell Cycle* 2009; **8**:2756–2768.