
Beyond Totally Reflexive Modules and Back

A Survey on Gorenstein Dimensions

Lars Winther Christensen¹, Hans-Bjørn Foxby², and Henrik Holm³

¹ Department of Mathematics and Statistics, Texas Tech University, Mail Stop 1042, Lubbock, TX 79409, U.S.A. lars.w.christensen@ttu.edu

² Department of Mathematical Sciences, University of Copenhagen, Universitetsparken 5, DK-2100 København Ø, Denmark foxby@math.ku.dk

³ Department of Basic Sciences and Environment, University of Copenhagen, Thorvaldsensvej 40, DK-1871 Frederiksberg C, Denmark hholm@life.ku.dk

Summary. Starting from the notion of totally reflexive modules, we survey the theory of Gorenstein homological dimensions for modules over commutative rings. The account includes the theory's connections with relative homological algebra and with studies of local ring homomorphisms. It ends close to the starting point: with a characterization of Gorenstein rings in terms of total acyclicity of complexes.

Key words: Auslander categories, Bass class, Chouinard formula, Frobenius endomorphism, G-dimension, Gorenstein dimension, Gorenstein ring, quasi-Cohen-Macaulay homomorphism, quasi-Gorenstein homomorphism, totally acyclic complex, totally reflexive module.

2000 Mathematics Subject Classification: 13-02, 13D05, 13H10.

1	G-dimension of Finitely Generated Modules	6
2	Gorenstein Projective Dimension	12
3	Gorenstein Injective Dimension	15
4	Gorenstein Flat Dimension	20
5	Relative Homological Algebra	24
6	Modules over Local Homomorphisms	29
7	Local Homomorphisms of Finite G-dimension	30
8	Reflexivity and Finite G-dimension	32
9	Detecting Finiteness of Gorenstein Dimensions	34

Introduction

An important motivation for the study of homological dimensions dates back to 1956, when Auslander and Buchsbaum [7] and Serre [95] proved:

Theorem A. Let R be a commutative Noetherian local ring with residue field k . Then the following conditions are equivalent.

- (i) R is regular.
- (ii) k has finite projective dimension.
- (iii) Every R -module has finite projective dimension.

This result opened to the solution of two long-standing conjectures by Krull. Moreover, it introduced the theme that finiteness of a homological dimension for all modules characterizes rings with special properties. Later work has shown that modules of finite projective dimension over a general ring share many properties with modules over a regular ring. This is an incitement to study homological dimensions of individual modules.

In line with these ideas, Auslander and Bridger [6] introduced in 1969 the G-dimension. It is a homological dimension for finitely generated modules over a Noetherian ring, and it gives a characterization of Gorenstein local rings (Theorem 6), which is similar to Theorem A. Indeed, R is Gorenstein if k has finite G-dimension, and only if every finitely generated R -module has finite G-dimension.

In the early 1990s, the G-dimension was extended beyond the realm of finitely generated modules over a Noetherian ring. This was done by Enochs and Jenda who introduced the notion of Gorenstein projective modules [40]. With the Gorenstein projective dimension at hand, a perfect parallel to Theorem A becomes available (Theorem 12). Subsequent work has shown that modules of finite Gorenstein projective dimension over a general ring share many properties with modules over a Gorenstein ring.

Classical Homological Algebra as Precedent

The notions of injective dimension and flat dimension for modules also have Gorenstein counterparts. It was Enochs and Jenda who introduced Gorenstein injective modules [40] and, in collaboration with Torrecillas, Gorenstein flat modules [46]. The study of Gorenstein dimensions is often called *Gorenstein homological algebra*; it has taken directions from the following principle.

Leitmotif. Every result in classical homological algebra should have a counterpart in Gorenstein homological algebra.

To make this concrete, we review some classical results on homological dimensions and point to their Gorenstein counterparts within the main text. In the balance of this introduction, R is assumed to be a commutative Noetherian local ring with maximal ideal \mathfrak{m} and residue field $k = R/\mathfrak{m}$.

Depth and Finitely Generated Modules

The projective dimension of a finitely generated R -module is closely related to its depth. This is captured by the *Auslander–Buchsbaum Formula* [8]:

Theorem B. For every finitely generated R -module M of finite projective dimension there is an equality $\text{pd}_R M = \text{depth } R - \text{depth}_R M$.

The Gorenstein counterpart (Theorem 5) actually strengthens the classical result; this is a recurring theme in Gorenstein homological algebra.

The injective dimension of a non-zero finitely generated R -module is either infinite or it takes a fixed value:

Theorem C. For every non-zero finitely generated R -module M of finite injective dimension there is an equality $\text{id}_R M = \text{depth } R$.

This result of Bass [19] has its Gorenstein counterpart in Theorem 21.

Characterizations of Cohen–Macaulay Rings

Existence of special modules of finite homological dimension characterizes Cohen–Macaulay rings. The equivalence of (i) and (iii) in the theorem below is still referred to as *Bass’ conjecture*, even though it was settled more than 20 years ago. Indeed, Peskine and Szpiro proved in [83] that it follows from the New Intersection Theorem, which they proved *ibid.* for equicharacteristic rings. In 1987 Roberts [84] settled the New Intersection Theorem completely.

Theorem D. The following conditions on R are equivalent.

- (i) R is Cohen–Macaulay.
- (ii) There is a non-zero R -module of finite length and finite projective dim.
- (iii) There is a non-zero finitely generated R -module of finite injective dim.

A Gorenstein counterpart to this characterization is yet to be found; see Questions 3 and 6.

Gorenstein rings are also characterized by existence of special modules of finite homological dimension. The equivalence of (i) and (ii) below is due to Peskine and Szpiro [83]. The equivalence of (i) and (iii) was conjectured by Vasconcelos [104] and proved by Foxby [55]. The Gorenstein counterparts are given in Theorems 20 and 31; see also Question 9.

Theorem E. The following conditions on R are equivalent.

- (i) R is Gorenstein.
- (ii) There is a non-zero cyclic R -module of finite injective dimension.
- (iii) There is a non-zero finitely generated R -module of finite projective dimension and finite injective dimension.

Local Ring Homomorphisms

The Frobenius endomorphism detects regularity of a local ring of positive prime characteristic. The next theorem collects results of Avramov, Iyengar, and Miller [17], Kunz [80], and Rodicio [86]. The counterparts in Gorenstein homological algebra to these results are given in Theorems 39 and 40.

Theorem F. Let R be of positive prime characteristic, and let ϕ denote its Frobenius endomorphism. Then the following conditions are equivalent.

- (i) R is regular.
- (ii) R has finite flat dimension as an R -module via ϕ^n for some $n \geq 1$.
- (iii) R is flat as an R -module via ϕ^n for every $n \geq 1$.
- (iv) R has finite injective dimension as an R -module via ϕ^n for some $n \geq 1$.
- (v) R has injective dimension $\dim R$ as an R -module via ϕ^n for every $n \geq 1$.

Let (S, \mathfrak{n}) be yet a local ring. A ring homomorphism $\varphi: R \rightarrow S$ is called *local* if $\varphi(\mathfrak{m}) \subseteq \mathfrak{n}$. A classical chapter of local algebra, initiated by Grothendieck, studies transfer of ring theoretic properties along such homomorphisms. If φ is flat, then it is called *Cohen–Macaulay* or *Gorenstein* if its closed fiber $S/\mathfrak{m}S$ is, respectively, a Cohen–Macaulay ring or a Gorenstein ring. These definitions have been extended to homomorphisms of finite flat dimension. The theorem below collects results of Avramov and Foxby from [12] and [14]; the Gorenstein counterparts are given in Theorems 45 and 47.

Theorem G. Let $\varphi: R \rightarrow S$ be a local homomorphism and assume that S has finite flat dimension as an R -module via φ . Then the following hold:

- (a) S is Cohen–Macaulay if and only if R and φ are Cohen–Macaulay.
- (b) S is Gorenstein if and only if R and φ are Gorenstein.

Vanishing of Cohomology

The projective dimension of a module M is at most n if and only if the absolute cohomology functor $\text{Ext}^{n+1}(M, -)$ vanishes. Similarly (Theorem 37), M has Gorenstein projective dimension at most n if and only if the relative cohomology functor $\text{Ext}_{\text{GP}}^{n+1}(M, -)$ vanishes. Unfortunately, the similarity between the two situations does not run too deep. We give a couple of examples:

The absolute Ext is balanced, that is, it can be computed from a projective resolution of M or from an injective resolution of the second argument. In general, however, the only known way to compute the relative Ext is from a (so-called) proper Gorenstein projective resolution of M .

Secondly, if M is finitely generated, then the absolute Ext commutes with localization, but the relative Ext is not known to do so, unless M has finite Gorenstein projective dimension.

Such considerations motivate the search for an alternative characterization of modules of finite Gorenstein projective dimension, and this has been a driving force behind much research on Gorenstein dimensions within the past 15 years. What follows is a brief review.

Equivalence of Module Categories

For a finitely generated R -module, Foxby [56] gave a resolution-free criterion for finiteness of the Gorenstein projective dimension; that is, one that can be verified without construction of a Gorenstein projective resolution. This result from 1994 is Theorem 49. In 1996, Enochs, Jenda, and Xu [48] extended Foxby’s criterion to non-finitely generated R -modules, provided that R is Cohen–Macaulay with a dualizing module D . What they proved is that the modules of finite Gorenstein projective dimension are exactly those in the so-called *Auslander class* with respect to D (Definition 24). This category had been introduced by Foxby [53] already in 1972 to extend a theorem of Sharp [97]. An important special case of this extension reads:

Theorem H. Let R be Cohen–Macaulay with a dualizing module D . Then the horizontal arrows below are equivalences of categories of R -modules.

$$\begin{array}{ccc}
 \mathcal{A}(R) & \begin{array}{c} \xrightarrow{D \otimes_R -} \\ \xleftarrow{\text{Hom}_R(D, -)} \end{array} & \mathcal{B}(R) \\
 \updownarrow & & \updownarrow \\
 \{A \mid \text{pd}_R A \text{ finite}\} & \begin{array}{c} \xrightarrow{D \otimes_R -} \\ \xleftarrow{\text{Hom}_R(D, -)} \end{array} & \{B \mid \text{id}_R B \text{ finite}\}
 \end{array}$$

Here $\mathcal{A}(R)$ is the Auslander class and $\mathcal{B}(R)$ is the *Bass class* (Definition 25).

The upper level equivalence in Theorem H is the Gorenstein counterpart of the lower level equivalence. Indeed, the modules in $\mathcal{B}(R)$ are exactly those of finite Gorenstein injective dimension; this is also proved in [48].

A commutative Noetherian ring has a dualizing complex \mathbf{D} if and only if it is a homomorphic image of a Gorenstein ring of finite Krull dimension; see Kawasaki [77]. For such rings, a result similar to Theorem H was proved by Avramov and Foxby [13] in 1997, and connections (Theorems 52 and 53) between Gorenstein dimensions and the Auslander and Bass classes with respect to \mathbf{D} were established by Christensen, Frankild, and Holm [30] in 2006. Testimony to the utility of these results is the frequent occurrence—e.g. in Theorems 17, 24, 29, 32, 34, and 40—of the assumption that the ground ring “is a homomorphic image of a Gorenstein ring of finite Krull dimension”. Recently, Esmkhani and Tousi [51] and Christensen and Sather-Wagstaff [34] have used the results to establish resolution-free criteria for finiteness of Gorenstein dimensions over general local rings (Theorems 13, 30, and 56).

Scope and Organization

A survey of this modest length is a portrait painted with broad pen strokes. Inevitably, many details are omitted, and some generality has been traded in for simplicity. We have chosen to focus on modules over commutative, and often Noetherian, rings. Much of Gorenstein homological algebra, though, works flawlessly over non-commutative rings, and there are statements in this sur-

vey about Noetherian rings that remain valid for coherent rings. Furthermore, most statements about modules remain valid for complexes of modules. The reader will have to consult the references to qualify these claims.

In most sections, the opening paragraph introduces the main references on the topic in question. We strive to cite the strongest results available and, outside of this introduction, we do not attempt to trace the history of individual results. In notes, placed at the end of sections, we give pointers to the literature on directions of research—often new ones—that are not included in this survey. Even within the scope of this paper, there are open ends, and more than a dozen questions and problems are found throughout the text.

From this point, R denotes a commutative ring. Any extra assumptions on R are explicitly stated. We say that R is *local* if it is Noetherian and has a unique maximal ideal. We use the shorthand (R, \mathfrak{m}, k) for a local ring R with maximal ideal \mathfrak{m} and residue field $k = R/\mathfrak{m}$.

Acknowledgments

It is a pleasure to thank Nanqing Ding, his colleagues, and the students who attended the 2008 Summer School at Nanjing University. The choice of material and the organization of this survey is strongly influenced by the summer school participants' reactions to lectures by L.W.C. We also thank Edgar Enochs, Srikanth Iyengar, and Peter Jørgensen for helpful discussions on their past and current work related to Gorenstein homological algebra. In the preparation of this version, we have been helped and challenged by detailed and thoughtful comments from our colleagues. We thank Sean Sather-Wagstaff and everyone else who gave their comments on earlier versions of the paper: Driss Bennis, Jesse Burke, Nanqing Ding, Najib Mahdou, Ryo Takahashi, and Oana Veliche.

This paper was started and finalized during visits by L.W.C. to Copenhagen; the hospitality and support of the University of Copenhagen is gratefully acknowledged.

1 G-dimension of Finitely Generated Modules

The topic of this section is Auslander and Bridger's notion of G-dimension for finitely generated modules over a Noetherian ring. The notes [5] from a seminar by Auslander outline the theory of G-dimension over commutative Noetherian rings. In [6] Auslander and Bridger treat the G-dimension within a more abstract framework. Later expositions of the commutative theory are given by Christensen [27] and by Mašek [82].

A *complex* \mathbf{M} of modules is (in homological notation) an infinite sequence

$$\mathbf{M} = \cdots \xrightarrow{\partial_{i+1}} M_i \xrightarrow{\partial_i} M_{i-1} \xrightarrow{\partial_{i-1}} \cdots$$

such that $\partial_i \partial_{i+1} = 0$ for every i in \mathbb{Z} . The i th *homology module* of \mathbf{M} is $H_i(\mathbf{M}) = \text{Ker } \partial_i / \text{Im } \partial_{i+1}$. We call \mathbf{M} *acyclic* if $H_i(\mathbf{M}) = 0$ for all $i \in \mathbb{Z}$.

Lemma 1. *Let \mathbf{L} be an acyclic complex of finitely generated projective R -modules. The following conditions on \mathbf{L} are equivalent.*

- (i) *The complex $\mathrm{Hom}_R(\mathbf{L}, R)$ is acyclic.*
- (ii) *The complex $\mathrm{Hom}_R(\mathbf{L}, Q)$ is acyclic for every projective R -module Q .*
- (iii) *The complex $E \otimes_R \mathbf{L}$ is acyclic for every injective R -module E .*

Proof. The Lemma is proved in [27], but here is a cleaner argument: Let Q be a projective module and E be an injective module. As \mathbf{L} consists of finitely generated projective modules, there is an isomorphism of complexes

$$\mathrm{Hom}_R(\mathrm{Hom}_R(\mathbf{L}, Q), E) \cong \mathrm{Hom}_R(Q, E) \otimes_R \mathbf{L}.$$

It follows from this isomorphism, applied to $Q = R$, that (i) implies (iii). Applied to a faithfully injective module E , it shows that (iii) implies (ii), as $\mathrm{Hom}_R(Q, E)$ is an injective module. It is evident that (ii) implies (i).

The following nomenclature is due to Avramov and Martsinkovsky [18]; Lemma 2 clarifies the rationale behind it.

Definition 1. A complex \mathbf{L} that satisfies the conditions in Lemma 1 is called *totally acyclic*. An R -module M is called *totally reflexive* if there exists a totally acyclic complex \mathbf{L} such that M is isomorphic to $\mathrm{Coker}(L_1 \rightarrow L_0)$.

Note that every finitely generated projective module L is totally reflexive, as the complex $0 \rightarrow L \xrightarrow{=} L \rightarrow 0$, with L in homological degrees 0 and -1 , is totally acyclic.

Example 1. If there exist elements x and y in R such that $\mathrm{Ann}_R(x) = (y)$ and $\mathrm{Ann}_R(y) = (x)$, then the complex

$$\dots \xrightarrow{x} R \xrightarrow{y} R \xrightarrow{x} R \xrightarrow{y} \dots$$

is totally acyclic. Thus, (x) and (y) are totally reflexive R -modules. For instance, if X and Y are non-zero non-units in a domain D , then their residue classes x and y in $R = D/(XY)$ generate totally reflexive R -modules.

An elementary construction of rings of this kind shows that non-projective totally reflexive modules may exist over a variety of rings; see also Problem 2.

Example 2. Let S be a commutative ring, and let $m > 1$ be an integer. Set $R = S[X]/(X^m)$, and denote by x the residue class of X in R . Then for every integer n between 1 and $m - 1$, the module (x^n) is totally reflexive.

From Lemma 1 it is straightforward to deduce:

Proposition 1. *Let S be an R -algebra of finite flat dimension. For every totally reflexive R -module G , the module $S \otimes_R G$ is totally reflexive over S .*

For example, Proposition 1 applies to $S = R/(\mathbf{x})$, where \mathbf{x} is an R -regular sequence. If (R, \mathfrak{m}) is local, then it also applies to the \mathfrak{m} -adic completion $S = \widehat{R}$.

Noetherian Rings

Recall that a finitely generated R -module M is called *reflexive* if the canonical map from M to $\text{Hom}_R(\text{Hom}_R(M, R), R)$ is an isomorphism. The following characterization of totally reflexive modules goes back to [6, 4.11].

Lemma 2. *Let R be Noetherian. A finitely generated R -module G is totally reflexive if and only if it is reflexive and for every $i \geq 1$ one has*

$$\text{Ext}_R^i(G, R) = 0 = \text{Ext}_R^i(\text{Hom}_R(G, R), R).$$

Definition 2. An (augmented) G -resolution of a finitely generated module M is an exact sequence $\cdots \rightarrow G_i \rightarrow G_{i-1} \rightarrow \cdots \rightarrow G_0 \rightarrow M \rightarrow 0$, where each module G_i is totally reflexive.

Note that if R is Noetherian, then every finitely generated R -module has a G -resolution, indeed a resolution by finitely generated free modules.

Definition 3. Let R be Noetherian. For a finitely generated R -module $M \neq 0$ the G -dimension, denoted by $\text{G-dim}_R M$, is the least integer $n \geq 0$ such that there exists a G -resolution of M with $G_i = 0$ for all $i > n$. If no such n exists, then $\text{G-dim}_R M$ is infinite. By convention, $\text{G-dim}_R 0 = -\infty$.

In [6, ch. 3] one finds the next theorem and its corollary; see also [27, 1.2.7].

Theorem 1. *Let R be Noetherian and M be a finitely generated R -module of finite G -dimension. For every integer $n \geq 0$ the next conditions are equivalent.*

- (i) $\text{G-dim}_R M \leq n$.
- (ii) $\text{Ext}_R^i(M, R) = 0$ for all $i > n$.
- (iii) $\text{Ext}_R^i(M, N) = 0$ for all $i > n$ and all R -modules N with $\text{pd}_R N$ finite.
- (iv) In every augmented G -resolution

$$\cdots \rightarrow G_i \rightarrow G_{i-1} \rightarrow \cdots \rightarrow G_0 \rightarrow M \rightarrow 0$$

the module $\text{Coker}(G_{n+1} \rightarrow G_n)$ is totally reflexive.

Corollary 1. *Let R be Noetherian. For every finitely generated R -module M of finite G -dimension there is an equality*

$$\text{G-dim}_R M = \sup\{i \in \mathbb{Z} \mid \text{Ext}_R^i(M, R) \neq 0\}.$$

Remark 1. Examples due to Jorgensen and Şega [75] show that in Corollary 1 one cannot avoid the *a priori* condition that $\text{G-dim}_R M$ is finite.

Remark 2. For M as in Corollary 1, the small finitistic projective dimension of R is an upper bound for $\text{G-dim}_R M$; cf. Christensen and Iyengar [32, 3.1.(a)].

A standard argument, see [6, 3.16] or [18, 3.4], yields:

Proposition 2. *Let R be Noetherian. If any two of the modules in an exact sequence $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ of finitely generated R -modules have finite G-dimension, then so has the third.*

The following quantitative comparison establishes the G-dimension as a refinement of the projective dimension for finitely generated modules. It is easily deduced from Corollary 1; see [27, 1.2.10].

Proposition 3. *Let R be Noetherian. For every finitely generated R -module M one has $\text{G-dim}_R M \leq \text{pd}_R M$, and equality holds if $\text{pd}_R M$ is finite.*

By [6, 4.15] the G-dimension of a module can be measured locally:

Proposition 4. *Let R be Noetherian. For every finitely generated R -module M there is an equality $\text{G-dim}_R M = \sup\{\text{G-dim}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} \mid \mathfrak{p} \in \text{Spec } R\}$.*

For the projective dimension even more is known: Bass and Murthy [20, 4.5] prove that if a finitely generated module over a Noetherian ring has finite projective dimension locally, then it has finite projective dimension globally—even if the ring has infinite Krull dimension. This leads to the following:

Question 1. Let R be Noetherian and M be a finitely generated R -module. Suppose that for every prime ideal \mathfrak{p} in R , the module $M_{\mathfrak{p}}$ has finite G-dimension over $R_{\mathfrak{p}}$. Does then M have finite G-dimension over R ?

Recall that a local ring is called *Gorenstein* if it has finite self-injective dimension. A Noetherian ring is Gorenstein if its localization at each prime ideal is a Gorenstein local ring, that is, $\text{id}_{R_{\mathfrak{p}}} R_{\mathfrak{p}}$ is finite for every prime ideal \mathfrak{p} in R . Consequently, the self-injective dimension of a Gorenstein ring equals its Krull dimension; that is $\text{id}_R R = \dim R$. The next result follows from [6, 4.20] in combination with Proposition 4.

Theorem 2. *Let R be Noetherian and n be a non-negative integer. Then R is Gorenstein with $\dim R \leq n$ if and only if one has $\text{G-dim}_R M \leq n$ for every finitely generated R -module M .*

By a result of Goto [62] also Gorenstein rings of infinite Krull dimension are characterized by finiteness of G-dimension.

Theorem 3. *Let R be Noetherian. Then R is Gorenstein if and only if every finitely generated R -module has finite G-dimension.*

Recall that the *grade* of a finitely generated module M over a Noetherian ring R can be defined as follows:

$$\text{grade}_R M = \inf\{i \in \mathbb{Z} \mid \text{Ext}_R^i(M, R) \neq 0\}.$$

Foxby [54] makes the following:

Definition 4. Let R be Noetherian. A finitely generated R -module M is called *quasi-perfect* if it has finite G-dimension equal to $\text{grade}_R M$.

The next theorem applies to $S = R/\widehat{\langle \mathbf{x} \rangle}$, where \mathbf{x} is an R -regular sequence. If R is local, then it also applies to $S = \widehat{R}$. Special (local) cases of the theorem are due to Golod [61] and to Avramov and Foxby [13, 7.11]. Christensen's proof [28, 6.5] establishes the general case.

Theorem 4. *Let R be Noetherian and S be a commutative Noetherian module-finite R -algebra. If S is a quasi-perfect R -module of grade g with $\text{Ext}_R^g(S, R) \cong S$, then the next equality holds for every finitely generated S -module N ,*

$$\text{G-dim}_R N = \text{G-dim}_S N + \text{G-dim}_R S.$$

Note that an S -module has finite G-dimension over R if and only if it has finite G-dimension over S ; see also Theorem 44. The next question is raised in [13].

Question 2. Let R be Noetherian, let S be a commutative Noetherian module-finite R -algebra, and let N be a finitely generated S -module. If $\text{G-dim}_S N$ and $\text{G-dim}_R S$ are finite, is then $\text{G-dim}_R N$ finite?

This is known as the *Transitivity Question*. By [13, 4.7] and [28, 3.15 and 6.5] it has an affirmative answer if $\text{pd}_S N$ is finite; see Theorem 42.

Local Rings

Before we proceed with results on G-dimension of modules over local rings, we make a qualitative comparison to the projective dimension. Theorem 4 reveals a remarkable property of the G-dimension, one that has almost no counterpart for the projective dimension. Here is an instance:

Example 3. Let (R, \mathfrak{m}, k) be local of positive depth. Pick a regular element x in \mathfrak{m} and set $S = R/\langle x \rangle$. Then one has $\text{grade}_R S = 1 = \text{pd}_R S$ and $\text{Ext}_R^1(S, R) \cong S$, but $\text{pd}_S N$ is infinite for every S -module N such that x is in $\mathfrak{m} \text{Ann}_R N$; see Shamash [96, §3]. In particular, if R is regular and x is in \mathfrak{m}^2 , then S is not regular, so $\text{pd}_S k$ is infinite while $\text{pd}_R k$ is finite; see Theorem A.

If G is a totally reflexive R -module, then every R -regular element is G -regular. A strong converse holds for modules of finite projective dimension; it is (still) referred to as *Auslander's zero-divisor conjecture*: let R be local and $M \neq 0$ be a finitely generated R -module with $\text{pd}_R M$ finite. Then every M -regular element is R -regular; see Roberts [85, 6.2.3]. An instance of Example 1 shows that one can *not* relax the condition on M to finite G-dimension:

Example 4. Let k be a field and consider the local ring $R = k[[X, Y]]/(XY)$. Then the residue class x of X generates a totally reflexive module. The element x is (x) -regular but nevertheless a zero-divisor in R .

While a tensor product of projective modules is projective, the next example shows that totally reflexive modules do not have an analogous property.

Example 5. Let R be as in Example 4. The R -modules (x) and (y) are totally reflexive, but $T = (x) \otimes_R (y)$ is not. Indeed, T is not a submodule of a free R -module as it has torsion: T is annihilated by the R -regular element $x + y$.

The next result [6, 4.13] is parallel to Theorem B in the Introduction; it is known as the Auslander–Bridger Formula.

Theorem 5. *Let R be local. For every finitely generated R -module M of finite G-dimension there is an equality*

$$\mathrm{G-dim}_R M = \mathrm{depth} R - \mathrm{depth}_R M.$$

In [82] Mašek corrects the proof of [6, 4.13]. Proofs can also be found in [5] and [27].

By Lemma 2 the G-dimension is preserved under completion:

Proposition 5. *Let R be local. For every finitely generated R -module M there is an equality $\mathrm{G-dim}_R M = \mathrm{G-dim}_{\widehat{R}}(\widehat{R} \otimes_R M)$.*

The following main result from [5, §3.2] is akin to Theorem A, but it differs in that it only deals with finitely generated modules.

Theorem 6. *For a local ring (R, \mathfrak{m}, k) the next conditions are equivalent.*

- (i) R is Gorenstein.
- (ii) $\mathrm{G-dim}_R k$ is finite.
- (iii) $\mathrm{G-dim}_R M$ is finite for every finitely generated R -module M .

It follows that non-projective totally reflexive modules exist over any non-regular Gorenstein local ring. On the other hand, Example 2 shows that existence of such modules does not force a local ring to be, say, Cohen–Macaulay.

A theorem of Christensen, Piepmeyer, Striuli, and Takahashi [33, 4.3] shows that fewness of totally reflexive modules comes in two distinct flavors:

Theorem 7. *Let R be local. If there are only finitely many indecomposable totally reflexive R -modules, up to isomorphism, then R is Gorenstein or every totally reflexive R -module is free.*

This dichotomy brings two problems to light:

Problem 1. Let R be a local ring that is not Gorenstein and assume that there exists a non-free totally reflexive R -module. Find constructions that produce infinite families of non-isomorphic indecomposable totally reflexive modules.

Problem 2. Describe the local rings over which every totally reflexive module is free.

While the first problem is posed in [33], the second one was already raised by Avramov and Martsinkovsky [18], and it is proved *ibid.* that over a Golod local ring that is not Gorenstein, every totally reflexive module is free. Another partial answer to Problem 2 is obtained by Yoshino [112], and by Christensen and Veliche [35]. The problem is also studied by Takahashi in [101].

Finally, Theorem D in the Introduction motivates:

Question 3. Let R be a local ring. If there exists a non-zero R -module of finite length and finite G-dimension, is then R Cohen–Macaulay?

A partial answer to this question is obtained by Takahashi [98, 2.3].

Notes

A topic that was only treated briefly above is constructions of totally reflexive modules. Such constructions are found in [16] by Avramov, Gasharov and Peeva, in work of Takahashi and Watanabe [102], and in Yoshino’s [112].

Hummel and Marley [71] extend the notion of G-dimension to finitely presented modules over coherent rings and use it to define and study coherent Gorenstein rings.

Gerko [60, §2] studies a dimension—the PCI-dimension or CI_* -dimension—based on a subclass of the totally reflexive modules. Golod [61] studies a generalized notion of G-dimension: the G_C -dimension, based on total reflexivity with respect to a semidualizing module C . These studies are continued by, among others, Gerko [60, §1], and Salarian, Sather-Wagstaff, and Yassemi [88]; see also the notes in Section 8.

An approach to homological dimensions that is not treated in this survey is based on so-called quasi-deformations. Several authors—among them Avramov, Gasharov, and Peeva [16] and Veliche [105]—take this approach to define homological dimensions that are intermediate between the projective dimension and the G-dimension for finitely generated modules. Gerko [60, §3] defines a Cohen–Macaulay dimension, which is a refinement of the G-dimension. Avramov [10, §8] surveys these dimensions.

2 Gorenstein Projective Dimension

To extend the G-dimension beyond the realm of finitely generated modules over Noetherian rings, Enochs and Jenda [40] introduced the notion of Gorenstein projective modules. The same authors, and their collaborators, studied these modules in several subsequent papers. The associated dimension, which is the focus of this section, was studied by Christensen [27] and Holm [65]. In organization, this section is parallel to Section 1.

Definition 5. An R -module A is called *Gorenstein projective* if there exists an acyclic complex \mathbf{P} of projective R -modules such that $\text{Coker}(P_1 \rightarrow P_0) \cong A$ and such that $\text{Hom}_R(\mathbf{P}, Q)$ is acyclic for every projective R -module Q .

It is evident that every projective module is Gorenstein projective.

Example 6. Every totally reflexive module is Gorenstein projective; this follows from Definition 1 and Lemma 1.

Basic categorical properties are recorded in [65, §2]:

Proposition 6. *The class of Gorenstein projective R -modules is closed under direct summands and coproducts.*

Every projective module is a direct summand of a free one. A parallel result for Gorenstein projective modules, Theorem 8 below, is due to Bennis and Mahdou [23, §2]; as substitute for free modules they define:

Definition 6. An R -module A is called *strongly Gorenstein projective* if there exists an acyclic complex \mathbf{P} of projective R -modules, in which all the differentials are identical, such that $\text{Coker}(P_1 \rightarrow P_0) \cong A$, and such that $\text{Hom}_R(\mathbf{P}, Q)$ is acyclic for every projective R -module Q .

Theorem 8. *An R -module is Gorenstein projective if and only if it is a direct summand of a strongly Gorenstein projective R -module.*

Definition 7. An (augmented) *Gorenstein projective resolution* of a module M is an exact sequence $\cdots \rightarrow A_i \rightarrow A_{i-1} \rightarrow \cdots \rightarrow A_0 \rightarrow M \rightarrow 0$, where each module A_i is Gorenstein projective.

Note that every module has a Gorenstein projective resolution, as a free resolution is trivially a Gorenstein projective one.

Definition 8. The *Gorenstein projective dimension* of an R -module $M \neq 0$, denoted by $\text{Gpd}_R M$, is the least integer $n \geq 0$ such that there exists a Gorenstein projective resolution of M with $A_i = 0$ for all $i > n$. If no such n exists, then $\text{Gpd}_R M$ is infinite. By convention, $\text{Gpd}_R 0 = -\infty$.

In [65, §2] one finds the next standard theorem and corollary.

Theorem 9. *Let M be an R -module of finite Gorenstein projective dimension. For every integer $n \geq 0$ the following conditions are equivalent.*

- (i) $\text{Gpd}_R M \leq n$.
- (ii) $\text{Ext}_R^i(M, Q) = 0$ for all $i > n$ and all projective R -modules Q .
- (iii) $\text{Ext}_R^i(M, N) = 0$ for all $i > n$ and all R -modules N with $\text{pd}_R N$ finite.
- (iv) In every augmented Gorenstein projective resolution

$$\cdots \rightarrow A_i \rightarrow A_{i-1} \rightarrow \cdots \rightarrow A_0 \rightarrow M \rightarrow 0$$

the module $\text{Coker}(A_{n+1} \rightarrow A_n)$ is Gorenstein projective.

Corollary 2. *For every R -module M of finite Gorenstein projective dimension there is an equality*

$$\text{Gpd}_R M = \sup\{i \in \mathbb{Z} \mid \text{Ext}_R^i(M, Q) \neq 0 \text{ for some projective } R\text{-module } Q\}.$$

Remark 3. For every R -module M as in the corollary, the finitistic projective dimension of R is an upper bound for $\text{Gpd}_R M$; see [65, 2.28].

The next result [65, 2.24] extends Proposition 2.

Proposition 7. *If in an exact sequence $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ any two of the modules have finite Gorenstein projective dimension, then so has the third.*

The Gorenstein projective dimension is a refinement of the projective dimension; this follows from Corollary 2:

Proposition 8. *For every R -module M one has $\text{Gpd}_R M \leq \text{pd}_R M$, and equality holds if M has finite projective dimension.*

Supplementary information comes from Holm [66, 2.2]:

Proposition 9. *If M is an R -module of finite injective dimension, then there is an equality $\text{Gpd}_R M = \text{pd}_R M$.*

The next result of Foxby is published in [31, Ascent table II(b)].

Proposition 10. *Let S be an R -algebra of finite projective dimension. For every Gorenstein projective R -module A , the module $S \otimes_R A$ is Gorenstein projective over S .*

Noetherian Rings

Finiteness of the Gorenstein projective dimension characterizes Gorenstein rings. The next result of Enochs and Jenda [42, 12.3.1] extends Theorem 2.

Theorem 10. *Let R be Noetherian and n be a non-negative integer. Then R is Gorenstein with $\dim R \leq n$ if and only if $\text{Gpd}_R M \leq n$ for every R -module M .*

The next result [27, 4.2.6] compares the Gorenstein projective dimension to the G-dimension.

Proposition 11. *Let R be Noetherian. For every finitely generated R -module M there is an equality $\text{Gpd}_R M = \text{G-dim}_R M$.*

The Gorenstein projective dimension of a module can not be measured locally; that is, Proposition 4 does not extend to non-finitely generated modules. As a consequence of Proposition 10, though, one has the following result.

Proposition 12. *Let R be Noetherian of finite Krull dimension. For every R -module M and every prime ideal \mathfrak{p} in R one has $\text{Gpd}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} \leq \text{Gpd}_R M$.*

Theorem E and Proposition 9 yield:

Theorem 11. *Let R be Noetherian and M be a finitely generated R -module. If $\text{Gpd}_R M$ and $\text{id}_R M$ are finite, then $R_{\mathfrak{p}}$ is Gorenstein for every \mathfrak{p} in $\text{Supp}_R M$.*

Local Rings

The next characterization of Gorenstein local rings—akin to Theorem A in the Introduction—follows from Theorems 6 and 10 via Proposition 11.

Theorem 12. *For a local ring (R, \mathfrak{m}, k) the next conditions are equivalent.*

- (i) R is Gorenstein.
- (ii) $\text{Gpd}_R k$ is finite.
- (iii) $\text{Gpd}_R M$ is finite for every R -module M .

The inequality in the next theorem is a consequence of Proposition 10. The second assertion is due to Esmkhani and Tousi [51, 3.5], cf. Christensen, Frankild, and Holm [30, 4.1]. The result should be compared to Proposition 5.

Theorem 13. *Let R be local and M be an R -module. Then one has*

$$\text{Gpd}_{\widehat{R}}(\widehat{R} \otimes_R M) \leq \text{Gpd}_R M,$$

and if $\text{Gpd}_{\widehat{R}}(\widehat{R} \otimes_R M)$ is finite, then so is $\text{Gpd}_R M$.

Notes

Holm and Jørgensen [68] extend Golod’s [61] notion of G_C -dimension to non-finitely generated modules in the form of a C -Gorenstein projective dimension. Further studies of this dimension are made by White [108].

3 Gorenstein Injective Dimension

The notion of Gorenstein injective modules is (categorically) dual to that of Gorenstein projective modules. The two were introduced in the same paper by Enochs and Jenda [40], and investigated in subsequent works by the same authors, by Christensen and Sather-Wagstaff [34], and by Holm [65]. This section is structured parallelly to the previous ones.

Definition 9. An R -module B is called *Gorenstein injective* if there exists an acyclic complex \mathbf{I} of injective R -modules such that $\text{Ker}(I^0 \rightarrow I^1) \cong B$, and such that $\text{Hom}_R(E, \mathbf{I})$ is acyclic for every injective R -module E .

It is clear that every injective module is Gorenstein injective.

Example 7. Let \mathbf{L} be a totally acyclic complex of finitely generated projective modules, see Definition 1, and let I be an injective module. Then the acyclic complex $\mathbf{I} = \text{Hom}_R(\mathbf{L}, I)$ consists of injective modules, and it follows from Lemma 1 that the complex $\text{Hom}_R(E, \mathbf{I}) \cong \text{Hom}_R(E \otimes_R \mathbf{L}, I)$ is acyclic for every injective module E . Thus, for every totally reflexive module G and every injective module I , the module $\text{Hom}_R(G, I)$ is Gorenstein injective.

Basic categorical properties are established in [65, 2.6]:

Proposition 13. *The class of Gorenstein injective R -modules is closed under direct summands and products.*

Under extra assumptions on the ring, Theorem 17 gives more information.

Definition 10. An (augmented) *Gorenstein injective resolution* of a module M is an exact sequence $0 \rightarrow M \rightarrow B^0 \rightarrow \dots \rightarrow B^{i-1} \rightarrow B^i \rightarrow \dots$, where each module B^i is Gorenstein injective.

Note that every module has a Gorenstein injective resolution, as an injective resolution is trivially a Gorenstein injective one.

Definition 11. The *Gorenstein injective dimension* of an R -module $M \neq 0$, denoted by $\text{Gid}_R M$, is the least integer $n \geq 0$ such that there exists a Gorenstein injective resolution of M with $B^i = 0$ for all $i > n$. If no such n exists, then $\text{Gid}_R M$ is infinite. By convention, $\text{Gid}_R 0 = -\infty$.

The next standard theorem is [65, 2.22].

Theorem 14. *Let M be an R -module of finite Gorenstein injective dimension. For every integer $n \geq 0$ the following conditions are equivalent.*

- (i) $\text{Gid}_R M \leq n$.
- (ii) $\text{Ext}_R^i(E, M) = 0$ for all $i > n$ and all injective R -modules E .
- (iii) $\text{Ext}_R^i(N, M) = 0$ for all $i > n$ and all R -modules N with $\text{id}_R N$ finite.
- (iv) In every augmented Gorenstein injective resolution

$$0 \rightarrow M \rightarrow B^0 \rightarrow \dots \rightarrow B^{i-1} \rightarrow B^i \rightarrow \dots$$

the module $\text{Ker}(B^n \rightarrow B^{n+1})$ is Gorenstein injective.

Corollary 3. *For every R -module M of finite Gorenstein injective dimension there is an equality*

$$\text{Gid}_R M = \sup\{i \in \mathbb{Z} \mid \text{Ext}_R^i(E, M) \neq 0 \text{ for some injective } R\text{-module } E\}.$$

Remark 4. For every R -module M as in the corollary, the finitistic injective dimension of R is an upper bound for $\text{Gid}_R M$; see [65, 2.29].

The next result [65, 2.25] is dual to Proposition 7.

Proposition 14. *If in an exact sequence $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ any two of the modules have finite Gorenstein injective dimension, then so has the third.*

The Gorenstein injective dimension is a refinement of the injective dimension; this follows from Corollary 3:

Proposition 15. *For every R -module M one has $\text{Gid}_R M \leq \text{id}_R M$, and equality holds if M has finite injective dimension.*

Supplementary information comes from Holm [66, 2.1]:

Proposition 16. *If M is an R -module of finite projective dimension, then there is an equality $\text{Gid}_R M = \text{id}_R M$. In particular, one has $\text{Gid}_R R = \text{id}_R R$.*

In [31] Christensen and Holm study (co)base change of modules of finite Gorenstein homological dimension. The following is elementary to verify:

Proposition 17. *Let S be an R -algebra of finite projective dimension. For every Gorenstein injective R -module B , the module $\text{Hom}_R(S, B)$ is Gorenstein injective over S .*

For a conditional converse see Theorems 22 and 56.

The next result of Bennis, Mahdou, and Ouarghi [24, 2.2] should be compared to Theorems 10, 11, 16, and 19. It is a perfect Gorenstein counterpart to a classical result due to Faith and Walker among others; see e.g. [107, 4.2.4].

Theorem 15. *The following conditions on R are equivalent.*

- (i) R is quasi-Frobenius.
- (ii) Every R -module is Gorenstein projective.
- (iii) Every R -module is Gorenstein injective.
- (iv) Every Gorenstein projective R -module is Gorenstein injective.
- (v) Every Gorenstein injective R -module is Gorenstein projective.

Noetherian Rings

Finiteness of the Gorenstein injective dimension characterizes Gorenstein rings; this is due to Enochs and Jenda [41, 3.1]:

Theorem 16. *Let R be Noetherian and n be a non-negative integer. Then R is Gorenstein with $\dim R \leq n$ if and only if $\text{Gid}_R M \leq n$ for every R -module M .*

A ring is Noetherian if and only if every coproduct of injective modules is injective. The “if” part has a perfect Gorenstein counterpart:

Proposition 18. *If every coproduct of Gorenstein injective R -modules is Gorenstein injective, then R is Noetherian.*

Proof. It is sufficient to see that every coproduct of injective R -modules is injective. Let $\{E_j\}$ be a family of injective modules. By assumption, the module $\coprod E_j$ is Gorenstein injective; in particular, there is an epimorphism $\varphi: I \rightarrow \coprod E_j$ such that I is injective and $\text{Hom}_R(E, \varphi)$ is surjective for every injective R -module E . Applying this to $E = E_j$ it is elementary to verify that φ is a split epimorphism.

Christensen, Frankild, and Holm [30, 6.9] provide a partial converse:

Theorem 17. *Assume that R is a homomorphic image of a Gorenstein ring of finite Krull dimension. Then the class of Gorenstein injective modules is closed under direct limits; in particular, it is closed under coproducts.*

As explained in the Introduction, the hypothesis on R in this theorem ensures the existence of a dualizing R -complex and an associated Bass class, cf. Section 9. These tools are essential to the known proof of Theorem 17.

Question 4. Let R be Noetherian. Is then every direct limit of Gorenstein injective R -modules Gorenstein injective?

Next follows a Gorenstein version of Chouinard's formula [26, 3.1]; it is proved in [34, 3.2]. Recall that the *width* of a module M over a local ring (R, \mathfrak{m}, k) is defined as

$$\text{width}_R M = \inf\{i \in \mathbb{Z} \mid \text{Tor}_i^R(k, M) \neq 0\}.$$

Theorem 18. *Let R be Noetherian. For every R -module M of finite Gorenstein injective dimension there is an equality*

$$\text{Gid}_R M = \{\text{depth } R_{\mathfrak{p}} - \text{width}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} \mid \mathfrak{p} \in \text{Spec } R\}.$$

Let M be an R -module, and let \mathfrak{p} be a prime ideal in R . Provided that $\text{Gid}_{R_{\mathfrak{p}}} M_{\mathfrak{p}}$ is finite, the inequality $\text{Gid}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} \leq \text{Gid}_R M$ follows immediately from the theorem. However, the next question remains open.

Question 5. Let R be Noetherian and B be a Gorenstein injective R -module. Is then $B_{\mathfrak{p}}$ Gorenstein injective over $R_{\mathfrak{p}}$ for every prime ideal \mathfrak{p} in R ?

A partial answer is known from [30, 5.5]:

Proposition 19. *Assume that R is a homomorphic image of a Gorenstein ring of finite Krull dimension. For every R -module M and every prime ideal \mathfrak{p} there is an inequality $\text{Gid}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} \leq \text{Gid}_R M$.*

Theorem E and Proposition 16 yield:

Theorem 19. *Let R be Noetherian and M be a finitely generated R -module. If $\text{Gid}_R M$ and $\text{pd}_R M$ are finite, then $R_{\mathfrak{p}}$ is Gorenstein for every \mathfrak{p} in $\text{Supp}_R M$.*

Local Rings

The following theorem of Foxby and Frankild [57, 4.5] generalizes work of Peskine and Szpiro [83], cf. Theorem E.

Theorem 20. *A local ring R is Gorenstein if and only if there exists a non-zero cyclic R -module of finite Gorenstein injective dimension.*

Theorems 16 and 20 yield a parallel to Theorem 6, akin to Theorem A.

Corollary 4. *For a local ring (R, \mathfrak{m}, k) the next conditions are equivalent.*

- (i) R is Gorenstein.
- (ii) $\text{Gid}_R k$ is finite.
- (iii) $\text{Gid}_R M$ is finite for every R -module M .

The first part of the next theorem is due to Christensen, Frankild, and Iyengar, and published in [57, 3.6]. The equality in Theorem 21—the Gorenstein analogue of Theorem C in the Introduction—is proved by Khatami, Tousi, and Yassemi [78, 2.5]; see also [34, 3.3].

Theorem 21. *Let R be local and $M \neq 0$ be a finitely generated R -module. Then $\text{Gid}_R M$ and $\text{Gid}_{\widehat{R}}(\widehat{R} \otimes_R M)$ are simultaneously finite, and when they are finite, there is an equality*

$$\text{Gid}_R M = \text{depth } R.$$

Remark 5. Let R be local and $M \neq 0$ be an R -module. If M has finite length and finite G-dimension, then its Matlis dual has finite Gorenstein injective dimension, cf. Example 7. See also Takahashi [100].

This elementary fact and Theorem D from the Introduction motivate:

Question 6. Let R be local. If there exists a non-zero finitely generated R -module of finite Gorenstein injective dimension, is then R Cohen–Macaulay?

A partial answer to this question is given by Yassemi [111, 1.3].

Esmkhani and Tousi [52, 2.5] prove the following conditional converse to Proposition 17. Recall that an R -module M is said to be *cotorsion* if $\text{Ext}_R^1(F, M) = 0$ for every flat R -module F .

Theorem 22. *Let R be local. An R -module M is Gorenstein injective if and only if it is cotorsion and $\text{Hom}_R(\widehat{R}, M)$ is Gorenstein injective over \widehat{R} .*

The example below demonstrates the necessity of the cotorsion hypothesis. Working in the derived category one obtains a stronger result; see Theorem 56.

Example 8. Let (R, \mathfrak{m}) be a local domain which is not \mathfrak{m} -adically complete. Aldrich, Enochs, and López-Ramos [1, 3.3] show that the module $\text{Hom}_R(\widehat{R}, R)$ is zero and hence Gorenstein injective over \widehat{R} . However, $\text{Gid}_R R$ is infinite if R is not Gorenstein, cf. Proposition 16.

Notes

Dual to the notion of strongly Gorenstein projective modules, see Definition 6, Ben-nis and Mahdou [23] also study strongly Gorenstein injective modules.

Several authors—Asadollahi, Sahandi, Salarian, Sazeeleh, Sharif, and Yassemi—have studied the relationship between Gorenstein injectivity and local cohomology; see [3], [87], [93], [94], and [111].

4 Gorenstein Flat Dimension

Another extension of the G-dimension is based on Gorenstein flat modules—a notion due to Enochs, Jenda, and Torrecillas [46]. Christensen [27] and Holm [65] are other main references for this section. The organization follows the pattern from Sections 1–3.

Definition 12. An R -module A is called *Gorenstein flat* if there exists an acyclic complex \mathbf{F} of flat R -modules such that $\text{Coker}(F_1 \rightarrow F_0) \cong A$, and such that $E \otimes_R \mathbf{F}$ is acyclic for every injective R -module E .

It is evident that every flat module is Gorenstein flat.

Example 9. Every totally reflexive module is Gorenstein flat; this follows from Definition 1 and Lemma 1.

Here is a direct consequence of Definition 12:

Proposition 20. *The class of Gorenstein flat R -modules is closed under coproducts.*

See Theorem 24 for further categorical properties of Gorenstein flat modules.

Definition 13. An (augmented) *Gorenstein flat resolution* of a module M is an exact sequence $\cdots \rightarrow A_i \rightarrow A_{i-1} \rightarrow \cdots \rightarrow A_0 \rightarrow M \rightarrow 0$, where each module A_i is Gorenstein flat.

Note that every module has a Gorenstein flat resolution, as a free resolution is trivially a Gorenstein flat one.

Definition 14. The *Gorenstein flat dimension* of an R -module $M \neq 0$, denoted by $\text{Gfd}_R M$, is the least integer $n \geq 0$ such that there exists a Gorenstein flat resolution of M with $A_i = 0$ for all $i > n$. If no such n exists, then $\text{Gfd}_R M$ is infinite. By convention, $\text{Gfd}_R 0 = -\infty$.

The next duality result is an immediate consequence of the definitions.

Proposition 21. *Let M be an R -module. For every injective R -module E there is an inequality $\text{Gid}_R \text{Hom}_R(M, E) \leq \text{Gfd}_R M$.*

Recall that an R -module E is called *faithfully injective* if it is injective and $\text{Hom}_R(M, E) = 0$ only if $M = 0$. The next question is inspired by the classical situation. It has an affirmative answer for Noetherian rings; see Theorem 25.

Question 7. Let M and E be R -modules. If E is faithfully injective and the module $\text{Hom}_R(M, E)$ is Gorenstein injective, is then M Gorenstein flat?

A straightforward application of Proposition 21 shows that the Gorenstein flat dimension is a refinement of the flat dimension; cf. Bennis [22, 2.2]:

Proposition 22. *For every R -module M one has $\text{Gfd}_R M \leq \text{fd}_R M$, and equality holds if M has finite flat dimension.*

The following result is an immediate consequence of Definition 12. Over a local ring a stronger result is available; see Theorem 30.

Proposition 23. *Let S be a flat R -algebra. For every R -module M there is an inequality $\text{Gfd}_S(S \otimes_R M) \leq \text{Gfd}_R M$.*

Corollary 5. *Let M be an R -module. For every prime ideal \mathfrak{p} in R there is an inequality $\text{Gfd}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} \leq \text{Gfd}_R M$.*

Noetherian Rings

Finiteness of the Gorenstein flat dimension characterizes Gorenstein rings; this is a result of Enochs and Jenda [41, 3.1]:

Theorem 23. *Let R be Noetherian and n be a non-negative integer. Then R is Gorenstein with $\dim R \leq n$ if and only if $\text{Gfd}_R M \leq n$ for every R -module M .*

A ring is coherent if and only if every product of flat modules is flat. We suggest the following problem on products of Gorenstein flat modules.

Problem 3. Describe the rings over which every product of Gorenstein flat modules is Gorenstein flat.

Some insight into the problem comes from the second assertion in Theorem 24; it is due to Christensen, Frankild, and Holm [30, 5.7]. The first assertion follows from work of Enochs, Jenda, and López-Ramos [45, 2.1] and [39, 3.3].

Theorem 24. *Let R be Noetherian. Then the class of Gorenstein flat R -modules is closed under direct summands and direct limits.*

If R is a homomorphic image of a Gorenstein ring of finite Krull dimension, then the class of Gorenstein flat R -modules is closed under products.

A result of Govorov [63] and Lazard [81, 1.2] asserts that a module is flat if and only if it is a direct limit of finitely generated projective modules. For Gorenstein flat modules, the picture is muddled:

Remark 6. Let R be Noetherian. It follows from Example 9 and Theorem 24 that a direct limit of totally reflexive modules is Gorenstein flat. If R is Gorenstein of finite Krull dimension, then every Gorenstein flat R -module can be written as a direct limit of totally reflexive modules; see Enochs and Jenda [42, 10.3.8]. If R is not Gorenstein, this conclusion may fail; see Beligiannis and Krause [21, 4.2 and 4.3] and Theorem 32.

The next result [27, 6.4.2] gives a partial answer to Question 7.

Theorem 25. *Let R be Noetherian, and let M and E be R -modules. If E is faithfully injective, then there is an equality*

$$\mathrm{Gid}_R \mathrm{Hom}_R(M, E) = \mathrm{Gfd}_R M.$$

Theorem 26 and its corollary are found in [65, 3.14]. They can be obtained by application of Theorem 25 to the corresponding results for the Gorenstein injective dimension; that is, Theorem 14 and Corollary 3.

Theorem 26. *Let R be Noetherian, and let M be an R -module of finite Gorenstein flat dimension. For every integer $n \geq 0$ the following are equivalent.*

- (i) $\mathrm{Gfd}_R M \leq n$.
- (ii) $\mathrm{Tor}_i^R(E, M) = 0$ for all $i > n$ and all injective R -modules E .
- (iii) $\mathrm{Tor}_i^R(N, M) = 0$ for all $i > n$ and all R -modules N with $\mathrm{id}_R N$ finite.
- (iv) In every augmented Gorenstein flat resolution

$$\cdots \rightarrow A_i \rightarrow A_{i-1} \rightarrow \cdots \rightarrow A_0 \rightarrow M \rightarrow 0$$

the module $\mathrm{Coker}(A_{n+1} \rightarrow A_n)$ is Gorenstein flat.

Corollary 6. *Let R be Noetherian. For every R -module M of finite Gorenstein flat dimension there is an equality*

$$\mathrm{Gfd}_R M = \sup\{i \in \mathbb{Z} \mid \mathrm{Tor}_i^R(E, M) \neq 0 \text{ for some injective } R\text{-module } E\}.$$

Remark 7. For every R -module M as in the corollary, the finitistic flat dimension of R is an upper bound for $\mathrm{Gfd}_R M$; see [65, 3.24].

The next result [65, 3.15] follows by Theorem 25 and Proposition 14.

Proposition 24. *Let R be Noetherian. If any two of the modules in an exact sequence $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ have finite Gorenstein flat dimension, then so has the third.*

A result of Holm [66, 2.6] supplements Proposition 22:

Proposition 25. *Let R be Noetherian of finite Krull dimension. For every R -module M of finite injective dimension there is an equality $\mathrm{Gfd}_R M = \mathrm{fd}_R M$.*

Recall that the *depth* of a module M over a local ring (R, \mathfrak{m}, k) is given as

$$\mathrm{depth}_R M = \inf\{i \in \mathbb{Z} \mid \mathrm{Ext}_R^i(k, M) \neq 0\}.$$

Theorem 27 is a Gorenstein version of Chouinard's formula [26, 1.2]. It follows from [65, 3.19] and [29, 2.4(b)]; see also Iyengar and Sather-Wagstaff [74, 8.8].

Theorem 27. *Let R be Noetherian. For every R -module M of finite Gorenstein flat dimension there is an equality*

$$\mathrm{Gfd}_R M = \{\mathrm{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} \mid \mathfrak{p} \in \mathrm{Spec} R\}.$$

The next two results compare the Gorenstein flat dimension to the Gorenstein projective dimension. The inequality in Theorem 28 is [65, 3.4], and the second assertion in this theorem is due to Esmkhani and Tousi [51, 3.4].

Theorem 28. *Let R be Noetherian of finite Krull dimension, and let M be an R -module. Then there is an inequality*

$$\text{Gfd}_R M \leq \text{Gpd}_R M,$$

and if $\text{Gfd}_R M$ is finite, then so is $\text{Gpd}_R M$.

It is not known whether the inequality in Theorem 28 holds over every commutative ring. For finitely generated modules one has [27, 4.2.6 and 5.1.11]:

Proposition 26. *Let R be Noetherian. For every finitely generated R -module M there is an equality $\text{Gfd}_R M = \text{Gpd}_R M = \text{G-dim}_R M$.*

The next result [30, 5.1] is related to Theorem 25; the question that follows is prompted by the classical situation.

Theorem 29. *Assume that R is a homomorphic image of a Gorenstein ring of finite Krull dimension. For every R -module M and every injective R -module E there is an inequality*

$$\text{Gfd}_R \text{Hom}_R(M, E) \leq \text{Gid}_R M,$$

and equality holds if E is faithfully injective.

Question 8. Let R be Noetherian, and let M and E be R -modules. If M is Gorenstein injective and E is injective, is then $\text{Hom}_R(M, E)$ Gorenstein flat?

Local Rings

Over a local ring there is a stronger version [51, 3.5] of Proposition 23:

Theorem 30. *Let R be local. For every R -module M there is an equality*

$$\text{Gfd}_{\widehat{R}}(\widehat{R} \otimes_R M) = \text{Gfd}_R M.$$

Combination of [66, 2.1 and 2.2] with Theorem E yields the next result. Recall that a non-zero finitely generated module has finite depth.

Theorem 31. *For a local ring R the following conditions are equivalent.*

- (i) R is Gorenstein.
- (ii) There is an R -module M with $\text{depth}_R M$, $\text{fd}_R M$, and $\text{Gid}_R M$ finite.
- (iii) There is an R -module M with $\text{depth}_R M$, $\text{id}_R M$, and $\text{Gfd}_R M$ finite.

We have for a while been interested in:

Question 9. Let R be local. If there exists an R -module M with $\text{depth}_R M$, $\text{Gfd}_R M$, and $\text{Gid}_R M$ finite, is then R Gorenstein?

A theorem of Jørgensen and Holm [70] brings perspective to Remark 6.

Theorem 32. *Assume that R is a homomorphic image of a Gorenstein local ring and that R is Henselian. If every Gorenstein flat R -module is a direct limit of totally reflexive modules, then R is Gorenstein or every totally reflexive module is free.*

Notes

Parallel to the notion of strongly Gorenstein projective modules, see Definition 6, Bennis and Mahdou [23] also study strongly Gorenstein flat modules. A different notion of strongly Gorenstein flat modules is studied by Ding, Li, and Mao in [37].

5 Relative Homological Algebra

Over a Gorenstein local ring, the totally reflexive modules are exactly the maximal Cohen–Macaulay modules, and their representation theory is a classical topic. Over rings that are not Gorenstein, the representation theory of totally reflexive modules was taken up by Takahashi [99] and Yoshino [112]. Conclusive results have recently been obtained by Christensen, Piepmeyer, Striuli, and Takahashi [33] and by Holm and Jørgensen [70]. These results are cast in the language of precovers and preenvelopes; see Theorem 33.

Relative homological algebra studies dimensions and (co)homology functors based on resolutions that are constructed via precovers or preenvelopes. Enochs and Jenda and their collaborators have made extensive studies of the precovering and preenveloping properties of the classes of Gorenstein flat and Gorenstein injective modules. Many of their results are collected in [42]. An important subsequent result establishes the existence of Gorenstein flat covers for every module over a Noetherian ring; see Theorem 36.

Terminology

Let \mathcal{H} be a class of R -modules. Recall that an \mathcal{H} -precover of an R -module M is a homomorphism $\varphi: H \rightarrow M$ with H in \mathcal{H} such that

$$\text{Hom}_R(H', \varphi): \text{Hom}_R(H', H) \longrightarrow \text{Hom}_R(H', M)$$

is surjective for every H' in \mathcal{H} . That is, every homomorphism from a module in \mathcal{H} to M factors through φ . Dually one defines \mathcal{H} -preenvelopes.

Remark 8. If \mathcal{H} contains all projective modules, then every \mathcal{H} -precover is an epimorphism. Thus, Gorenstein projective/flat precovers are epimorphisms.

If \mathcal{H} contains all injective modules, then every \mathcal{H} -preenvelope is a monomorphism. Thus, every Gorenstein injective preenvelope is a monomorphism.

Fix an \mathcal{H} -precover φ . It is called *special* if it is an epimorphism and the cohomology module $\text{Ext}_R^1(H', \text{Ker } \varphi)$ vanishes for every module H' in \mathcal{H} . It is called a *cover* if in every factorization $\varphi = \varphi\psi$, the map $\psi: H \rightarrow H$ is an automorphism. If \mathcal{H} is closed under extensions, then every epi \mathcal{H} -cover is a special precover. This is known as Wakamatsu’s lemma; see Xu [109, 2.1.1]. Dually one defines special \mathcal{H} -preenvelopes and \mathcal{H} -envelopes.

Remark 9. Let \mathbf{I} be a complex of injective modules as in Definition 9. Then every differential in \mathbf{I} is a special injective precover if its image. Similarly, in a complex \mathbf{P} of projective modules as in Definition 5, every differential ∂_i is a special projective preenvelope of the cokernel of the previous differential ∂_{i+1} .

Totally Reflexive Covers and Envelopes

The next result of Avramov and Martsinkovsky [18, 3.1] corresponds over a Gorenstein local ring to the existence of maximal Cohen–Macaulay approximations in the sense of Auslander and Buchweitz [9].

Proposition 27. *Let R be Noetherian. For every finitely generated R -module M of finite G -dimension there is an exact sequence $0 \rightarrow K \rightarrow G \rightarrow M \rightarrow 0$, where G is totally reflexive and $\text{pd}_R K = \max\{0, G\text{-dim}_R M - 1\}$. In particular, every finitely generated R -module of finite G -dimension has a special totally reflexive precover.*

An unpublished result of Auslander states that every finitely generated module over a Gorenstein local ring has a totally reflexive cover; see [49] for a generalization. A strong converse is contained in the next theorem, which combines Auslander’s result with recent work of several authors; see [33] and [70].

Theorem 33. *For a local ring (R, \mathfrak{m}, k) the next conditions are equivalent.*

- (i) *Every finitely generated R -module has a totally reflexive cover.*
- (ii) *The residue field k has a special totally reflexive precover.*
- (iii) *Every finitely generated R -module has a totally reflexive envelope.*
- (iv) *Every finitely generated R -module has a special totally reflexive preenvelope.*
- (v) *R is Gorenstein or every totally reflexive R -module is free.*

If R is local and Henselian (e.g. complete), then existence of a totally reflexive precover implies existence of a totally reflexive cover; see [99, 2.5]. In that case one can drop “special” in part (ii) above. In general, though, the next question from [33] remains open.

Question 10. Let (R, \mathfrak{m}, k) be local. If k has a totally reflexive precover, is then R Gorenstein or every totally reflexive R -module free?

Gorenstein Projective Precovers

The following result is proved by Holm in [65, 2.10].

Proposition 28. *For every R -module M of finite Gorenstein projective dimension there is an exact sequence $0 \rightarrow K \rightarrow A \rightarrow M \rightarrow 0$, where A is Gorenstein projective and $\text{pd}_R K = \max\{0, \text{Gpd}_R M - 1\}$. In particular, every R -module of finite Gorenstein projective dimension has a special Gorenstein projective precover.*

Under an extra assumption on the ring, Jørgensen [76] proves existence of Gorenstein projective precovers for all modules:

Theorem 34. *If R is a homomorphic image of a Gorenstein ring of finite Krull dimension, then every R -module has a Gorenstein projective precover.*

Remark 10. Actually, the argument in Krause's proof of [79, 7.12.(1)] applies to the setup in [76] and yields existence of a *special* Gorenstein projective precover for every module over a ring as in Theorem 34.

Over any ring, every module has a special projective precover; hence:

Problem 4. Describe the rings over which every module has a (special) Gorenstein projective precover.

Gorenstein Injective Preenvelopes

In [65, 2.15] one finds:

Proposition 29. *For every R -module M of finite Gorenstein injective dimension there is an exact sequence $0 \rightarrow M \rightarrow B \rightarrow C \rightarrow 0$, where B is Gorenstein injective and $\text{id}_R C = \max\{0, \text{Gid}_R M - 1\}$. In particular, every R -module of finite Gorenstein injective dimension has a special Gorenstein injective preenvelope.*

Over Noetherian rings, existence of Gorenstein injective preenvelopes for all modules is proved by Enochs and López-Ramos in [50]. Krause [79, 7.12] proves a stronger result:

Theorem 35. *If R is Noetherian, then every R -module has a special Gorenstein injective preenvelope.*

Over Gorenstein rings, Enochs, Jenda, and Xu [47, 6.1] prove more:

Proposition 30. *If R is Gorenstein of finite Krull dimension, then every R -module has a Gorenstein injective envelope.*

Over any ring, every module has an injective envelope; this suggests:

Problem 5. Describe the rings over which every module has a Gorenstein injective (pre)envelope.

Gorenstein Flat Covers

The following existence result is due to Enochs and López-Ramos [50, 2.11].

Theorem 36. *If R is Noetherian, then every R -module has a Gorenstein flat cover.*

Remark 11. Let R be Noetherian, and let M be a finitely generated R -module. If M has finite G-dimension, then by Proposition 27 it has a finitely generated Gorenstein projective/flat precover, cf. Proposition 26. If M has infinite G-dimension, then it still has a Gorenstein projective/flat precover by Theorems 34 and 36, but by Theorem 33 this need *not* be finitely generated.

Over any ring, every module has a flat cover, as proved by Bican, El Bashir, and Enochs [25]. This motivates:

Problem 6. Describe the rings over which every module has a Gorenstein flat (pre)cover.

Over a Noetherian ring, every module has a flat preenvelope; cf. Enochs [38, 2.1]. A Gorenstein version of this result follows from Theorem 24 and [50, 2.5]:

Proposition 31. *If R is a homomorphic image of a Gorenstein ring of finite Krull dimension, then every R -module has a Gorenstein flat preenvelope.*

Relative Cohomology via Gorenstein Projective Modules

The notion of a proper resolution is central in relative homological algebra. Here is a special case:

Definition 15. An augmented Gorenstein projective resolution,

$$\mathbf{A}^+ = \dots \xrightarrow{\varphi_{i+1}} A_i \xrightarrow{\varphi_i} A_{i-1} \xrightarrow{\varphi_{i-1}} \dots \xrightarrow{\varphi_1} A_0 \xrightarrow{\varphi_0} M \longrightarrow 0$$

of an R -module M is said to be *proper* if for every Gorenstein projective module A' , the complex $\text{Hom}_R(A', \mathbf{A}^+)$ is acyclic.

Remark 12. Assume that every R -module has a Gorenstein projective precover. Then every R -module has a proper Gorenstein projective resolution constructed by taking as φ_0 a Gorenstein projective precover of M and as φ_i a Gorenstein projective precover of $\text{Ker } \varphi_{i-1}$ for $i > 0$.

Definition 16. The *relative Gorenstein projective dimension* of an R -module $M \neq 0$, denoted by $\text{rel-Gpd}_R M$, is the least integer $n \geq 0$ such that there exists a proper Gorenstein projective resolution of M with $A_i = 0$ for all $i > n$. If no such n (or no such resolution) exists, then $\text{rel-Gpd}_R M$ is infinite. By convention, $\text{rel-Gpd}_R 0 = -\infty$.

The following result is a consequence of Proposition 28.

Proposition 32. *For every R -module M one has $\text{rel-Gpd}_R M = \text{Gpd}_R M$.*

It is shown in [42, §8.2] that the next definition makes sense.

Definition 17. Let M and N be R -modules and assume that M has a proper Gorenstein projective resolution \mathbf{A} . The i th relative cohomology module $\text{Ext}_{\text{GP}}^i(M, N)$ is $H^i(\text{Hom}_R(\mathbf{A}, N))$.

Remark 13. Let R be Noetherian, and let M and N be finitely generated R -modules. Unless M has finite G-dimension, it is not clear whether the cohomology modules $\text{Ext}_{\text{GP}}^i(M, N)$ are finitely generated, cf. Remark 11.

Vanishing of relative cohomology Ext_{GP}^i characterizes modules of finite Gorenstein projective dimension. The proof is standard; see [67, 3.9].

Theorem 37. *Let M be an R -module that has a proper Gorenstein projective resolution. For every integer $n \geq 0$ the following conditions are equivalent.*

- (i) $\text{Gpd}_R M \leq n$.
- (ii) $\text{Ext}_{\text{GP}}^i(M, -) = 0$ for all $i > n$.
- (iii) $\text{Ext}_{\text{GP}}^{n+1}(M, -) = 0$.

Remark 14. Relative cohomology based on totally reflexive modules is studied in [18]. The results that correspond to Proposition 32 and Theorem 37 in that setting are contained in [18, 4.8 and 4.2].

Notes

Based on a notion of coproper Gorenstein injective resolutions, one can define a relative Gorenstein injective dimension and cohomology functors Ext_{GI}^i with properties analogous to those of Ext_{GP}^i described above. There is, similarly, a relative Gorenstein flat dimension and a relative homology theory based on proper Gorenstein projective/flat resolutions. The relative Gorenstein injective dimension and the relative Gorenstein flat dimension were first studied by Enochs and Jenda [41] for modules over Gorenstein rings. The question of balancedness for relative (co)homology is treated by Enochs and Jenda [42], Holm [64], and Iacob [72].

In [18] Avramov and Martsinkovsky also study the connection between relative and Tate cohomology for finitely generated modules. This study is continued by Veliche [106] for arbitrary modules, and a dual theory is developed by Asadollahi and Salarian [4]. Jørgensen [76] and Krause [79] study connections between Gorenstein relative cohomology and generalized notions of Tate cohomology.

Sather-Wagstaff and White [92] use relative cohomology to define an Euler characteristic for modules of finite G-dimension. In collaboration with Sharif, they study cohomology theories related to generalized Gorenstein dimensions [90].

6 Modules over Local Homomorphisms

In this section $\varphi: (R, \mathfrak{m}) \rightarrow (S, \mathfrak{n})$ is a local homomorphism, that is, $\varphi(\mathfrak{m}) \subseteq \mathfrak{n}$. The topic is Gorenstein dimensions over R of finitely generated S -modules. The utility of this point of view is illustrated by a generalization, due to Christensen and Iyengar [32, 4.1], of the Auslander–Bridger Formula (Theorem 5):

Theorem 38. *Let N be a finitely generated S -module. If N has finite Gorenstein flat dimension as an R -module via φ , then there is an equality*

$$\text{Gfd}_R N = \text{depth } R - \text{depth}_R N.$$

For a finitely generated S -module N of finite flat dimension over R , this equality follows from work of André [2, II.57]. For a finitely generated S -module of finite injective dimension over R , an affirmative answer to the next question is already in [103, 5.2] by Takahashi and Yoshino.

Question 11. Let N be a non-zero finitely generated S -module. If N has finite Gorenstein injective dimension as an R -module via φ , does then the equality $\text{Gid}_R N = \text{depth } R$ hold?

The Frobenius Endomorphism

If R has positive prime characteristic, we denote by ϕ the Frobenius endomorphism on R and by ϕ^n its n -fold composition. The next two theorems are special cases of [74, 8.14 and 8.15] by Iyengar and Sather-Wagstaff and of [57, 5.5] by Foxby and Frankild.

Theorem 39. *Let R be local of positive prime characteristic. The following conditions are equivalent.*

- (i) R is Gorenstein.
- (ii) R has finite Gorenstein flat dim. as an R -module via ϕ^n for some $n \geq 1$.
- (iii) R is Gorenstein flat as an R -module via ϕ^n for every $n \geq 1$.

Theorem 40. *Let R be local of positive prime characteristic, and assume that it is a homomorphic image of a Gorenstein ring. The following are equivalent:*

- (i) R is Gorenstein.
- (ii) R has finite Gorenstein injective dim. as R -module via ϕ^n for some $n \geq 1$.
- (iii) R has Gorenstein injective dimension equal to $\dim R$ as an R -module via ϕ^n for every $n \geq 1$.

Part (iii) above is actually not included in [57, 5.5]. Part (iii) follows, though, from part (i) by Corollary 4 and Theorem 18.

G-dimension over a Local Homomorphism

The homomorphism φ fits in a commutative diagram of local homomorphisms:

$$\begin{array}{ccccc} & & R' & & \\ & \nearrow \varphi & & \searrow \varphi' & \\ R & \xrightarrow{\varphi} & S & \xrightarrow{\iota} & \widehat{S} \end{array}$$

where φ is flat with regular closed fiber $R'/\mathfrak{m}R'$, the ring R' is complete, and φ' is surjective. Set $\hat{\varphi} = \iota\varphi$; a diagram as above is called a *Cohen factorization* of $\hat{\varphi}$. This is a construction due to Avramov, Foxby, and Herzog [15, 1.1].

The next definition is due to Iyengar and Sather-Wagstaff [74, §3]; it is proved *ibid.* that it is independent of the choice of Cohen factorization.

Definition 18. Choose a Cohen factorization of $\hat{\varphi}$ as above. For a finitely generated S -module N , the *G-dimension of N over φ* is given as

$$\mathrm{G-dim}_{\varphi} N = \mathrm{G-dim}_{R'}(\widehat{S} \otimes_S N) - \mathrm{edim}(R'/\mathfrak{m}R').$$

Example 10. Let k be a field and let φ be the extension from k to the power series ring $k[[x]]$. Then one has $\mathrm{G-dim}_{\varphi} k[[x]] = -1$.

Christensen and Sather-Wagstaff [34, 2.4] prove:

Theorem 41. *A finitely generated S -module N has finite G-dimension over φ if and only if it has finite Gorenstein flat dimension as an R -module via φ .*

It is clear from Example 10 that $\mathrm{Gfd}_R N$ and $\mathrm{G-dim}_{\varphi} N$ need not be equal.

7 Local Homomorphisms of Finite G-dimension

This section treats transfer of ring theoretic properties along a local homomorphism of finite G-dimension. Our focus is on the Gorenstein property, which was studied by Avramov and Foxby in [13], and the Cohen–Macaulay property, studied by Frankild in [58].

As in Section 6, $\varphi: (R, \mathfrak{m}) \rightarrow (S, \mathfrak{n})$ is a local homomorphism. In view of Definition 18, a notion from [13, 4.3] can be defined as follows:

Definition 19. Set $\mathrm{G-dim} \varphi = \mathrm{G-dim}_{\varphi} S$; the homomorphism φ is said to be of *finite G-dimension* if this number is finite.

Notice that by Theorem 41 the homomorphism φ has finite G-dimension if and only if S has finite Gorenstein flat dimension as an R -module via φ .

The next descent result is [34, 2.5].

Theorem 42. *Let φ be of finite G-dimension. For every S -module N one has:*

- (a) *If $\text{fd}_S N$ is finite then $\text{Gfd}_R N$ is finite.*
- (b) *If $\text{id}_S N$ is finite then $\text{Gid}_R N$ is finite.*

It is not known if the composition of two local homomorphisms of finite G-dimension has finite G-dimension; it would follow from an affirmative answer to Question 2, cf. [13, 4.8]. Some insight is provided by Theorem 46 and the next result, which is due to Iyengar and Sather-Wagstaff [74, 5.2].

Theorem 43. *Let $\psi: S \rightarrow T$ be a local homomorphism such that $\text{fd}_S T$ is finite. Then $\text{G-dim } \psi\varphi$ is finite if and only if $\text{G-dim } \varphi$ is finite.*

Quasi-Gorenstein Homomorphisms

Let M be a finitely generated module over a local ring (R, \mathfrak{m}, k) . For every integer $i \geq 0$ the i th Bass number $\mu_R^i(M)$ is the dimension of the k -vector space $\text{Ext}_R^i(k, M)$.

Definition 20. The homomorphism φ is called *quasi-Gorenstein* if it has finite G-dimension and for every $i \geq 0$ there is an equality of Bass numbers

$$\mu_R^{i+\text{depth } R}(R) = \mu_S^{i+\text{depth } S}(S).$$

Example 11. If R is Gorenstein, then the natural surjection $R \twoheadrightarrow k$ is quasi-Gorenstein. More generally, if φ is surjective and S is quasi-perfect as an R -module via φ , then φ is quasi-Gorenstein if and only if there is an isomorphism $\text{Ext}_R^g(S, R) \cong S$ where $g = \text{G-dim}_R S$ with; see [13, 6.5, 7.1, 7.4].

Several characterizations of the quasi-Gorenstein property are given in [13, 7.4 and 7.5]. For example, it is sufficient that $\text{G-dim } \varphi$ is finite and the equality of Bass numbers holds for some $i > 0$.

The next ascent-descent result is [34, 2.6].

Theorem 44. *Let φ be quasi-Gorenstein. For every S -module N the following assertions hold.*

- (a) *$\text{Gfd}_S N$ is finite if and only if $\text{Gfd}_R N$ is finite.*
- (b) *$\text{Gid}_S N$ is finite if and only if $\text{Gid}_R N$ is finite.*

Ascent and descent of the Gorenstein property is described by [13, 7.7.2]. It should be compared to part (b) in Theorem G.

Theorem 45. *The following conditions on φ are equivalent.*

- (i) *R and S are Gorenstein.*
- (ii) *R is Gorenstein and φ is quasi-Gorenstein.*
- (iii) *S is Gorenstein and φ is of finite G-dimension.*

The following (de)composition result is [13, 7.10, 8.9, and 8.10]. It should be compared to Theorem 4.

Theorem 46. *Assume that φ is quasi-Gorenstein, and let $\psi: S \rightarrow T$ be a local homomorphism. The following assertions hold.*

- (a) $\text{G-dim } \psi\varphi$ is finite if and only if $\text{G-dim } \psi$ is finite.
- (b) $\psi\varphi$ is quasi-Gorenstein if and only if ψ is quasi-Gorenstein.

Quasi-Cohen–Macaulay Homomorphisms

The next definition from [58, 5.8 and 6.2] uses terminology from Definition 4.

Definition 21. The homomorphism φ is *quasi-Cohen–Macaulay*, for short *quasi-CM*, if φ has a Cohen factorization where \widehat{S} is a quasi-perfect R' -module.

If φ is quasi-CM, then \widehat{S} is a quasi-perfect R' -module in every Cohen factorization of φ ; see [58, 5.8]. The following theorem is part of [58, 6.7]; it should be compared to part (a) in Theorem G.

Theorem 47. *The following assertions hold.*

- (a) *If R is Cohen–Macaulay and φ is quasi-CM, then S is Cohen–Macaulay.*
- (b) *If S is Cohen–Macaulay and $\text{G-dim } \varphi$ is finite, then φ is quasi-CM.*

In view of Theorem 46, Frankild’s work [58, 6.4 and 6.5] yields:

Theorem 48. *Assume that φ is quasi-Gorenstein, and let $\psi: S \rightarrow T$ be a local homomorphism. Then $\psi\varphi$ is quasi-CM if and only if ψ is quasi-CM.*

8 Reflexivity and Finite G-dimension

In this section R is Noetherian. Let M be a finitely generated R -module. If M is totally reflexive, then the cohomology modules $\text{Ext}_R^i(M, R)$ vanish for all $i > 0$. The converse is true if M is known *a priori* to have finite G-dimension, cf. Corollary 1. In general, though, one can not infer from such vanishing that M is totally reflexive—explicit examples to this effect are constructed by Jorgensen and Şega in [75]—and this motivated a search for alternative criteria for finiteness of G-dimension.

Reflexive Complexes

One such criterion was given by Foxby and published in [110]. Its habitat is the derived category $\mathcal{D}(R)$ of the category of R -modules. The objects in $\mathcal{D}(R)$ are R -complexes, and there is a canonical functor F from the category of R -complexes to $\mathcal{D}(R)$. This functor is the identity on objects and it maps

homology isomorphisms to isomorphisms in $\mathcal{D}(R)$. The restriction of F to modules gives an embedding of the module category into $\mathcal{D}(R)$.

The homology $H(\mathbf{M})$ of an R -complex \mathbf{M} is a (graded) R -module, and \mathbf{M} is said to have *finitely generated homology* if this module is finitely generated. For R -modules M and N , the (co)homology of the derived Hom and tensor product complexes gives the classical Ext and Tor modules:

$$\text{Ext}_R^i(M, N) = H^i(\mathbf{RHom}_R(M, N)) \quad \text{and} \quad \text{Tor}_i^R(M, N) = H_i(M \otimes_R^{\mathbf{L}} N).$$

Definition 22. An R -complex \mathbf{M} is called *reflexive* if \mathbf{M} and $\mathbf{RHom}_R(\mathbf{M}, R)$ have finitely generated homology and the canonical morphism

$$\mathbf{M} \longrightarrow \mathbf{RHom}_R(\mathbf{RHom}_R(\mathbf{M}, R), R)$$

is an isomorphism in the derived category $\mathcal{D}(R)$. The full subcategory of $\mathcal{D}(R)$ whose objects are the reflexive R -complexes is denoted by $\mathcal{R}(R)$.

Theorem 49 below is Foxby’s criterion for finiteness of G-dimension of a finitely generated module [110, 2.7]. It differs significantly from Definition 3 as it does not involve construction of a G-resolution of the module.

Theorem 49. *Let R be Noetherian. A finitely generated R -module has finite G-dimension if and only if it belongs to $\mathcal{R}(R)$.*

If R is local, then the next result is [27, 2.3.14]. In the generality stated below it follows from Theorems 3 and 49: the implication (ii) \Rightarrow (iii) is the least obvious, it uses [27, 2.1.12].

Corollary 7. *Let R be Noetherian. The following conditions are equivalent.*

- (i) R is Gorenstein.
- (ii) Every R -module is in $\mathcal{R}(R)$.
- (iii) Every R -complex with finitely generated homology is in $\mathcal{R}(R)$.

G-dimension of Complexes

Having made the passage to the derived category, it is natural to consider G-dimension for complexes. For every R -complex \mathbf{M} with finitely generated homology there exists a complex \mathbf{G} of finitely generated free R -modules, which is isomorphic to \mathbf{M} in $\mathcal{D}(R)$; see [11, 1.7.(1)]. In Christensen’s [27, ch. 2] one finds the next definition and the two theorems that follow.

Definition 23. Let \mathbf{M} be an R -complex with finitely generated homology. If $H(\mathbf{M}) \neq 0$, then the *G-dimension* of \mathbf{M} is the least integer n such that there exists a complex \mathbf{G} of totally reflexive R -modules which is isomorphic to \mathbf{M} in $\mathcal{D}(R)$ and has $G_i = 0$ for all $i > n$. If no such integer n exists, then $\text{G-dim}_R \mathbf{M}$ is infinite. If $H(\mathbf{M}) = 0$, then $\text{G-dim}_R \mathbf{M} = -\infty$ by convention.

Note that this extends Definition 3. As a common generalization of Theorem 49 and Corollary 1 one has [27, 2.3.8]:

Theorem 50. *Let R be Noetherian. An R -complex M with finitely generated homology has finite G-dimension if and only if it is reflexive. Furthermore, for every reflexive R -complex M there is an equality*

$$\mathrm{G}\text{-dim}_R M = \sup\{i \in \mathbb{Z} \mid H^i(\mathbf{R}\mathrm{Hom}_R(M, R)) \neq 0\}.$$

In broad terms, the theory of G-dimension for finitely generated modules extends to complexes with finitely generated homology. One example is the next extension [27, 2.3.13] of the Auslander–Bridger Formula (Theorem 5).

Theorem 51. *Let R be local. For every complex M in $\mathcal{R}(R)$ one has*

$$\mathrm{G}\text{-dim}_R M = \mathrm{depth} R - \mathrm{depth}_R M.$$

Here the *depth* of a complex M over a local ring (R, \mathfrak{m}, k) is defined by extension of the definition for modules, that is,

$$\mathrm{depth}_R M = \inf\{i \in \mathbb{Z} \mid H^i(\mathbf{R}\mathrm{Hom}_R(k, M)) \neq 0\}.$$

Notes

In [27, ch. 2] the theory of G-dimension for complexes is developed in detail.

Generalized notions of G-dimension—based on reflexivity with respect to semi-dualizing modules and complexes—are studied by Christensen [28], Frankild and Sather-Wagstaff [59], Gerko [60], Golod [61], Holm and Jørgensen [69], by Salarian, Sather-Wagstaff, and Yassemi [88], and White [108]. See also the notes in Section 1.

9 Detecting Finiteness of Gorenstein Dimensions

In the previous section, we discussed a resolution-free characterization of modules of finite G-dimension (Theorem 49). The topic of this section is similar characterizations of modules of finite Gorenstein projective/injective/flat dimension. By work of Christensen, Frankild, and Holm [30], appropriate criteria are available for modules over a Noetherian ring that has a dualizing complex (Theorems 52 and 53). As mentioned in the Introduction, a Noetherian ring has a dualizing complex if and only if it is a homomorphic image of a Gorenstein ring of finite Krull dimension. For example, every complete local ring has a dualizing complex by Cohen’s structure theorem.

Auslander Categories

The next definition is due to Foxby; see [13, 3.1] and [53, §2].

Definition 24. Let R be Noetherian and assume that it has a dualizing complex \mathbf{D} . The *Auslander class* $\mathcal{A}(R)$ is the full subcategory of the derived category $\mathcal{D}(R)$ whose objects \mathbf{M} satisfy the following conditions.

- (1) $H_i(\mathbf{M}) = 0$ for $|i| \gg 0$.
- (2) $H_i(\mathbf{D} \otimes_R^{\mathbf{L}} \mathbf{M}) = 0$ for $i \gg 0$.
- (3) The natural map $\mathbf{M} \rightarrow \mathbf{RHom}_R(\mathbf{D}, \mathbf{D} \otimes_R^{\mathbf{L}} \mathbf{M})$ is an isomorphism in $\mathcal{D}(R)$.

The relation to Gorenstein dimensions is given by [30, 4.1]:

Theorem 52. *Let R be Noetherian and assume that it has a dualizing complex. For every R -module M , the following conditions are equivalent.*

- (i) M has finite Gorenstein projective dimension.
- (ii) M has finite Gorenstein flat dimension.
- (iii) M belongs to $\mathcal{A}(R)$.

Remark 15. Every complete local ring has a dualizing complex, so in view of Theorems 13 and 30 the result above gives a resolution-free characterization of modules of finite Gorenstein projective/flat dimension over any local ring.

The next definition is in [13, 3.1]; the theorem that follows is [30, 4.4].

Definition 25. Let R be Noetherian and assume that it has a dualizing complex \mathbf{D} . The *Bass class* $\mathcal{B}(R)$ is the full subcategory of the derived category $\mathcal{D}(R)$ whose objects \mathbf{M} satisfy the following conditions.

- (1) $H^i(\mathbf{M}) = 0$ for $|i| \gg 0$.
- (2) $H^i(\mathbf{RHom}_R(\mathbf{D}, \mathbf{M})) = 0$ for $i \gg 0$.
- (3) The natural map $\mathbf{D} \otimes_R^{\mathbf{L}} \mathbf{RHom}_R(\mathbf{D}, \mathbf{M}) \rightarrow \mathbf{M}$ is an isomorphism in $\mathcal{D}(R)$.

Theorem 53. *Let R be Noetherian and assume that it has a dualizing complex. For every R -module M , the following conditions are equivalent.*

- (i) M has finite Gorenstein injective dimension.
- (ii) M belongs to $\mathcal{B}(R)$.

From the two theorems above and from Theorems 16 and 23 one gets:

Corollary 8. *Let R be Noetherian and assume that it has a dualizing complex. The following conditions are equivalent.*

- (i) R is Gorenstein.
- (ii) Every R -complex \mathbf{M} with $H_i(\mathbf{M}) = 0$ for $|i| \gg 0$ belongs to $\mathcal{A}(R)$.
- (iii) Every R -complex \mathbf{M} with $H_i(\mathbf{M}) = 0$ for $|i| \gg 0$ belongs to $\mathcal{B}(R)$.

Gorenstein Dimensions of Complexes

It turns out to be convenient to extend the Gorenstein dimensions to complexes; this is illustrated by Theorem 56 below.

In the following we use the notion of a semi-projective resolution. Every complex has such a resolution, by [11, 1.6], and a projective resolution of a module is semi-projective. In view of this and Theorem 9, the next definition, which is due to Veliche [106, 3.1 and 3.4], extends Definition 8.

Definition 26. Let \mathbf{M} be an R -complex. If $H(\mathbf{M}) \neq 0$, then the *Gorenstein projective dimension* of \mathbf{M} is the least integer n such that $H_i(\mathbf{M}) = 0$ for all $i > n$ and there exists a semi-projective resolution \mathbf{P} of \mathbf{M} for which the module $\text{Coker}(P_{n+1} \rightarrow P_n)$ is Gorenstein projective. If no such n exists, then $\text{Gpd}_R \mathbf{M}$ is infinite. If $H(\mathbf{M}) = 0$, then $\text{Gpd}_R \mathbf{M} = -\infty$ by convention.

In the next theorem, which is due to Iyengar and Krause [73, 8.1], the *unbounded Auslander class* $\widehat{\mathcal{A}}(R)$ is the full subcategory of $\mathcal{D}(R)$ whose objects satisfy conditions (2) and (3) in Definition 24.

Theorem 54. *Let R be Noetherian and assume that it has a dualizing complex. For every R -complex \mathbf{M} , the following conditions are equivalent.*

- (i) \mathbf{M} has finite Gorenstein projective dimension.
- (ii) \mathbf{M} belongs to $\widehat{\mathcal{A}}(R)$.

One finds the next definition in [4, 2.2 and 2.3] by Asadollahi and Salarian. It uses the notion of a semi-injective resolution. Every complex has such a resolution, by [11, 1.6], and an injective resolution of a module is semi-injective. In view of Theorem 14, the following extends Definition 11.

Definition 27. Let \mathbf{M} be an R -complex. If $H(\mathbf{M}) \neq 0$, then the *Gorenstein injective dimension* of \mathbf{M} is the least integer n such that $H^i(\mathbf{M}) = 0$ for all $i > n$ and there exists a semi-injective resolution \mathbf{I} of \mathbf{M} for which the module $\text{Ker}(I^n \rightarrow I^{n+1})$ is Gorenstein injective. If no such integer n exists, then $\text{Gid}_R \mathbf{M}$ is infinite. If $H(\mathbf{M}) = 0$, then $\text{Gid}_R \mathbf{M} = -\infty$ by convention.

In the next result [73, 8.2], the *unbounded Bass class* $\widehat{\mathcal{B}}(R)$ is the full subcategory of $\mathcal{D}(R)$ whose objects satisfy conditions (2) and (3) in Definition 25.

Theorem 55. *Let R be Noetherian and assume that it has a dualizing complex. For every R -complex \mathbf{M} , the following conditions are equivalent.*

- (i) \mathbf{M} has finite Gorenstein injective dimension.
- (ii) \mathbf{M} belongs to $\widehat{\mathcal{B}}(R)$.

The next result of Christensen and Sather-Wagstaff [34, 1.6] should be compared to Theorem 22.

Theorem 56. *Let R be local. For every R -module M there is an equality*

$$\text{Gid}_R M = \text{Gid}_{\widehat{R}} \mathbf{R}\text{Hom}_R(\widehat{R}, M).$$

Remark 16. Via this result, Theorem 55 gives a resolution-free characterization of modules of finite Gorenstein injective dimension over any local ring.

Acyclicity Versus Total Acyclicity

The next results characterize Gorenstein rings in terms of the complexes that define Gorenstein projective/injective/flat modules. The first one is [73, 5.5].

Theorem 57. *Let R be Noetherian and assume that it has a dualizing complex. Then the following conditions are equivalent.*

- (i) R is Gorenstein.
- (ii) For every acyclic complex \mathbf{P} of projective R -modules and every projective R -module Q , the complex $\text{Hom}_R(\mathbf{P}, Q)$ is acyclic.
- (iii) For every acyclic complex \mathbf{I} of injective R -modules and every injective R -module E , the complex $\text{Hom}_R(E, \mathbf{I})$ is acyclic.

In the terminology of [73], part (ii)/(iii) above says that every acyclic complex of projective/injective modules is totally acyclic.

The final result is due to Christensen and Veliche [36]:

Theorem 58. *Let R be Noetherian and assume that it has a dualizing complex. Then there exist acyclic complexes \mathbf{F} and \mathbf{I} of flat R -modules and injective R -modules, respectively, such that the following conditions are equivalent.*

- (i) R is Gorenstein.
- (ii) For every injective R -module E , the complex $E \otimes_R \mathbf{F}$ is acyclic.
- (iii) For every injective R -module E , the complex $\text{Hom}_R(E, \mathbf{I})$ is acyclic.

It is not known, in general, if there exists an acyclic complex \mathbf{P} of projective R -modules such that R is Gorenstein if $\text{Hom}_R(\mathbf{P}, Q)$ is acyclic for every projective R -module Q .

Notes

In broad terms, the theory of Gorenstein dimensions for modules extends to complexes. It is developed in detail by Asadollahi and Salarian [4], Christensen, Frankild, and Holm [30] and [31], Christensen and Sather-Wagstaff [34], and by Veliche [106].

Objects in Auslander and Bass classes with respect to semi-dualizing complexes have interpretations in terms of generalized Gorenstein dimensions; see [43] and [44] by Enochs and Jenda, [68] by Holm and Jørgensen, and [89] by Sather-Wagstaff.

Sharif, Sather-Wagstaff, and White [91] study totally acyclic complexes of Gorenstein projective modules. They show that the cokernels of the differentials in such complexes are Gorenstein projective. That is, a “Gorenstein Gorenstein projective” module is Gorenstein projective.

References

1. Stephen T. Aldrich, Edgar E. Enochs, and Juan A. López-Ramos, *Derived functors of Hom relative to flat covers*, Math. Nachr. **242** (2002), 17–26. MR1916846
2. Michel André, *Homologie des algèbres commutatives*, Springer-Verlag, Berlin, 1974, Die Grundlehren der mathematischen Wissenschaften, Band 206. MR0352220
3. Javad Asadollahi and Shokrollah Salarian, *Cohomology theories based on Gorenstein injective modules*, Trans. Amer. Math. Soc. **358** (2006), no. 5, 2183–2203. MR2197453
4. ———, *Gorenstein injective dimension for complexes and Iwanaga-Gorenstein rings*, Comm. Algebra **34** (2006), no. 8, 3009–3022. MR2250584
5. Maurice Auslander, *Anneaux de Gorenstein, et torsion en algèbre commutative*, Secrétariat mathématique, Paris, 1967, Séminaire d'Algèbre Commutative dirigé par Pierre Samuel, 1966/67. Texte rédigé, d'après des exposés de Maurice Auslander, par Marquerite Mangeney, Christian Peskine et Lucien Szpiro. École Normale Supérieure de Jeunes Filles. Available from <http://www.numdam.org>. MR0225844
6. Maurice Auslander and Mark Bridger, *Stable module theory*, Memoirs of the American Mathematical Society, No. 94, American Mathematical Society, Providence, R.I., 1969. MR0269685
7. Maurice Auslander and David A. Buchsbaum, *Homological dimension in Noetherian rings*, Proc. Nat. Acad. Sci. U.S.A. **42** (1956), 36–38. MR0075190
8. ———, *Homological dimension in local rings*, Trans. Amer. Math. Soc. **85** (1957), 390–405. MR0086822
9. Maurice Auslander and Ragnar-Olaf Buchweitz, *The homological theory of maximal Cohen-Macaulay approximations*, Mém. Soc. Math. France (N.S.) (1989), no. 38, 5–37, Colloque en l'honneur de Pierre Samuel (Orsay, 1987). MR1044344
10. Luchezar L. Avramov, *Homological dimensions and related invariants of modules over local rings*, Representations of algebra. Vol. I, II, Beijing Norm. Univ. Press, Beijing, 2002, pp. 1–39. MR2067368
11. Luchezar L. Avramov and Hans-Bjørn Foxby, *Homological dimensions of unbounded complexes*, J. Pure Appl. Algebra **71** (1991), no. 2-3, 129–155. MR1117631
12. ———, *Locally Gorenstein homomorphisms*, Amer. J. Math. **114** (1992), no. 5, 1007–1047. MR1183530
13. ———, *Ring homomorphisms and finite Gorenstein dimension*, Proc. London Math. Soc. (3) **75** (1997), no. 2, 241–270. MR1455856
14. ———, *Cohen-Macaulay properties of ring homomorphisms*, Adv. Math. **133** (1998), no. 1, 54–95. MR1492786
15. Luchezar L. Avramov, Hans-Bjørn Foxby, and Bernd Herzog, *Structure of local homomorphisms*, J. Algebra **164** (1994), no. 1, 124–145. MR1268330
16. Luchezar L. Avramov, Vesselin N. Gasharov, and Irena V. Peeva, *Complete intersection dimension*, Inst. Hautes Études Sci. Publ. Math. (1997), no. 86, 67–114 (1998). MR1608565
17. Luchezar L. Avramov, Srikanth Iyengar, and Claudia Miller, *Homology over local homomorphisms*, Amer. J. Math. **128** (2006), no. 1, 23–90. MR2197067

18. Luchezar L. Avramov and Alex Martsinkovsky, *Absolute, relative, and Tate cohomology of modules of finite Gorenstein dimension*, Proc. London Math. Soc. (3) **85** (2002), no. 2, 393–440. MR1912056
19. Hyman Bass, *On the ubiquity of Gorenstein rings*, Math. Z. **82** (1963), 8–28. MR0153708
20. Hyman Bass and M. Pavaman Murthy, *Grothendieck groups and Picard groups of abelian group rings*, Ann. of Math. (2) **86** (1967), 16–73. MR0219592
21. Apostolos Beligiannis and Henning Krause, *Thick subcategories and virtually Gorenstein algebras*, preprint, 2006, arXiv:math/0608710 [math.RA], to appear in Illinois J. Math.
22. Driss Bennis, *A note on gorenstein flat dimension*, preprint, 2008, arXiv:0811.2650v1 [math.AC], to appear in Algebra Colloq.
23. Driss Bennis and Najib Mahdou, *Strongly Gorenstein projective, injective, and flat modules*, J. Pure Appl. Algebra **210** (2007), no. 2, 437–445. MR2320007
24. Driss Bennis, Najib Mahdou, and Khalid Ouarghi, *Rings over which all modules are gorenstein (resp., strongly gorenstein) projective*, preprint, 2007, arXiv:0712.0127v2 [math.AC].
25. Ladislav Bican, Robert El Bashir, and Edgar E. Enochs, *All modules have flat covers*, Bull. London Math. Soc. **33** (2001), no. 4, 385–390. MR1832549
26. Leo G. Chouinard, II, *On finite weak and injective dimension*, Proc. Amer. Math. Soc. **60** (1976), 57–60 (1977). MR0417158
27. Lars Winther Christensen, *Gorenstein dimensions*, Lecture Notes in Mathematics, vol. 1747, Springer-Verlag, Berlin, 2000. MR1799866
28. ———, *Semi-dualizing complexes and their Auslander categories*, Trans. Amer. Math. Soc. **353** (2001), no. 5, 1839–1883. MR1813596
29. Lars Winther Christensen, Hans-Bjørn Foxby, and Anders Frankild, *Restricted homological dimensions and Cohen-Macaulayness*, J. Algebra **251** (2002), no. 1, 479–502. MR1900297
30. Lars Winther Christensen, Anders Frankild, and Henrik Holm, *On Gorenstein projective, injective and flat dimensions—A functorial description with applications*, J. Algebra **302** (2006), no. 1, 231–279. MR2236602
31. Lars Winther Christensen and Henrik Holm, *Ascent properties of Auslander categories*, Canad. J. Math., online 17 August 2008, 33 pp. Preprint, arXiv:math/0509570v2 [math.AC].
32. Lars Winther Christensen and Srikanth Iyengar, *Gorenstein dimension of modules over homomorphisms*, J. Pure Appl. Algebra **208** (2007), no. 1, 177–188. MR2269838
33. Lars Winther Christensen, Greg Piepmeyer, Janet Striuli, and Ryo Takahashi, *Finite Gorenstein representation type implies simple singularity*, Adv. Math. **218** (2008), no. 4, 1012–1026. MR2419377
34. Lars Winther Christensen and Sean Sather-Wagstaff, *Transfer of Gorenstein dimensions along ring homomorphisms*, preprint, 2008, arXiv:0807.070v2 [math.AC], 13 pp.
35. Lars Winther Christensen and Oana Veliche, *Acyclicity over local rings with radical cube zero*, Illinois J. Math. **51** (2007), no. 4, 1439–1454. MR2417436
36. ———, *A test complex for Gorensteinness*, Proc. Amer. Math. Soc. **136** (2008), no. 2, 479–487. MR2358487
37. Nanqing Ding, Yuanlin Li, and Lixin Mao, *Strongly Gorenstein flat modules*, preprint, 2007, to appear in J. Aust. Math. Soc.

38. Edgar E. Enochs, *Injective and flat covers, envelopes and resolvents*, Israel J. Math. **39** (1981), no. 3, 189–209. MR636889
39. Edgar E. Enochs and Overtoun M. G. Jenda, *Balanced functors applied to modules*, J. Algebra **92** (1985), no. 2, 303–310. MR778450
40. ———, *Gorenstein injective and projective modules*, Math. Z. **220** (1995), no. 4, 611–633. MR1363858
41. ———, *Gorenstein injective and flat dimensions*, Math. Japon. **44** (1996), no. 2, 261–268. MR1416263
42. ———, *Relative homological algebra*, de Gruyter Expositions in Mathematics, vol. 30, Walter de Gruyter & Co., Berlin, 2000. MR1753146
43. ———, *Ω -Gorenstein projective and flat covers and Ω -Gorenstein injective envelopes*, Comm. Algebra **32** (2004), no. 4, 1453–1470. MR2100367
44. ———, *Gorenstein and Ω -Gorenstein injective covers and flat preenvelopes*, Comm. Algebra **33** (2005), no. 2, 507–518. MR2124340
45. Edgar E. Enochs, Overtoun M. G. Jenda, and Juan A. López-Ramos, *The existence of Gorenstein flat covers*, Math. Scand. **94** (2004), no. 1, 46–62. MR2032335
46. Edgar E. Enochs, Overtoun M. G. Jenda, and Blas Torrecillas, *Gorenstein flat modules*, Nanjing Daxue Xuebao Shuxue Bannian Kan **10** (1993), no. 1, 1–9. MR1248299
47. Edgar E. Enochs, Overtoun M. G. Jenda, and Jinzhong Xu, *Covers and envelopes over Gorenstein rings*, Tsukuba J. Math. **20** (1996), no. 2, 487–503. MR1422636
48. ———, *Foxby duality and Gorenstein injective and projective modules*, Trans. Amer. Math. Soc. **348** (1996), no. 8, 3223–3234. MR1355071
49. ———, *A generalization of Auslander’s last theorem*, Algebr. Represent. Theory **2** (1999), no. 3, 259–268. MR1715748
50. Edgar E. Enochs and Juan A. López-Ramos, *Kaplansky classes*, Rend. Sem. Mat. Univ. Padova **107** (2002), 67–79. MR1926201
51. Mohammad Ali Esmkhani and Massoud Tousei, *Gorenstein homological dimensions and Auslander categories*, J. Algebra **308** (2007), no. 1, 321–329. MR2290924
52. ———, *Gorenstein injective modules and Auslander categories*, Arch. Math. (Basel) **89** (2007), no. 2, 114–123. MR2341722
53. Hans-Bjørn Foxby, *Gorenstein modules and related modules*, Math. Scand. **31** (1972), 267–284 (1973). MR0327752
54. ———, *Quasi-perfect modules over Cohen-Macaulay rings*, Math. Nachr. **66** (1975), 103–110. MR0376663
55. ———, *Bounded complexes of flat modules*, J. Pure Appl. Algebra **15** (1979), no. 2, 149–172. MR0535182
56. ———, *Gorenstein dimension over Cohen-Macaulay rings*, Proceedings of international conference on commutative algebra (W. Bruns, ed.), Universität Osnabrück, 1994.
57. Hans-Bjørn Foxby and Anders J. Frankild, *Cyclic modules of finite Gorenstein injective dimension and Gorenstein rings*, Illinois J. Math. **51** (2007), no. 1, 67–82. MR2346187
58. Anders Frankild, *Quasi Cohen–Macaulay properties of local homomorphisms*, J. Algebra **235** (2001), no. 1, 214–242. MR1807663

59. Anders Frankild and Sean Sather-Wagstaff, *Reflexivity and ring homomorphisms of finite flat dimension*, Comm. Algebra **35** (2007), no. 2, 461–500. MR2294611
60. Aleksandr A. Gerko, *On homological dimensions*, Mat. Sb. **192** (2001), no. 8, 79–94. MR1862245
61. Evgeniy S. Golod, *G-dimension and generalized perfect ideals*, Trudy Mat. Inst. Steklov. **165** (1984), 62–66, Algebraic geometry and its applications. MR752933
62. Shiro Goto, *Vanishing of $\text{Ext}_A^i(M, A)$* , J. Math. Kyoto Univ. **22** (1982/83), no. 3, 481–484. MR674605
63. V. E. Govorov, *On flat modules*, Sibirsk. Mat. Ž. **6** (1965), 300–304. MR0174598
64. Henrik Holm, *Gorenstein derived functors*, Proc. Amer. Math. Soc. **132** (2004), no. 7, 1913–1923. MR2053961
65. ———, *Gorenstein homological dimensions*, J. Pure Appl. Algebra **189** (2004), no. 1-3, 167–193. MR2038564
66. ———, *Rings with finite Gorenstein injective dimension*, Proc. Amer. Math. Soc. **132** (2004), no. 5, 1279–1283. MR2053331
67. ———, *Relative Ext groups, resolutions, and Schanuel classes*, Osaka J. Math. **45** (2008), no. 3, 719–735.
68. Henrik Holm and Peter Jørgensen, *Semi-dualizing modules and related Gorenstein homological dimensions*, J. Pure Appl. Algebra **205** (2006), no. 2, 423–445. MR2203625
69. ———, *Cohen-Macaulay homological dimensions*, Rend. Semin. Mat. Univ. Padova **117** (2007), 87–112. MR2351787
70. ———, *Rings without a Gorenstein analogue of the Govorov-Lazard theorem*, preprint, 2008, arXiv:0810.3168v2 [math.AC].
71. Livia Hummel and Thomas Marley, *The Auslander-Bridger formula and the Gorenstein property for coherent rings*, preprint, 2008, arXiv:0809.1595v1 [math.AC].
72. Alina Iacob, *Absolute, Gorenstein, and Tate torsion modules*, Comm. Algebra **35** (2007), no. 5, 1589–1606. MR2317632
73. Srikanth Iyengar and Henning Krause, *Acyclicity versus total acyclicity for complexes over Noetherian rings*, Doc. Math. **11** (2006), 207–240. MR2262932
74. Srikanth Iyengar and Sean Sather-Wagstaff, *G-dimension over local homomorphisms. Applications to the Frobenius endomorphism*, Illinois J. Math. **48** (2004), no. 1, 241–272. MR2048224
75. David A. Jorgensen and Liana M. Şega, *Independence of the total reflexivity conditions for modules*, Algebr. Represent. Theory **9** (2006), no. 2, 217–226. MR2238367
76. Peter Jørgensen, *Existence of Gorenstein projective resolutions and Tate cohomology*, J. Eur. Math. Soc. (JEMS) **9** (2007), no. 1, 59–76. MR2283103
77. Takesi Kawasaki, *On arithmetic Macaulayfication of Noetherian rings*, Trans. Amer. Math. Soc. **354** (2002), no. 1, 123–149. MR1859029
78. Leila Khatami, Massoud Tousei, and Siamak Yassemi, *Finiteness of Gorenstein injective dimension of modules*, preprint, 2008, to appear in Proc. Amer. Math. Soc.
79. Henning Krause, *The stable derived category of a Noetherian scheme*, Compos. Math. **141** (2005), no. 5, 1128–1162. MR2157133
80. Ernst Kunz, *Characterizations of regular local rings for characteristic p* , Amer. J. Math. **91** (1969), 772–784. MR0252389

81. Daniel Lazard, *Autour de la platitude*, Bull. Soc. Math. France **97** (1969), 81–128. MR0254100
82. Vladimir Mašek, *Gorenstein dimension and torsion of modules over commutative Noetherian rings*, Comm. Algebra **28** (2000), no. 12, 5783–5811, Special issue in honor of Robin Hartshorne. MR1808604
83. Christian Peskine and Lucien Szpiro, *Dimension projective finie et cohomologie locale. Applications à la démonstration de conjectures de M. Auslander, H. Bass et A. Grothendieck*, Inst. Hautes Études Sci. Publ. Math. (1973), no. 42, 47–119. MR0374130
84. Paul Roberts, *Le théorème d'intersection*, C. R. Acad. Sci. Paris Sér. I Math. **304** (1987), no. 7, 177–180. MR880574
85. Paul C. Roberts, *Multiplicities and Chern classes in local algebra*, Cambridge Tracts in Mathematics, vol. 133, Cambridge University Press, Cambridge, 1998. MR1686450
86. Antonio G. Rodicio, *On a result of Avramov*, Manuscripta Math. **62** (1988), no. 2, 181–185. MR963004
87. Parviz Sahandi and Tirdad Sharif, *Dual of the Auslander-Bridger formula and GF-perfectness*, Math. Scand. **101** (2007), no. 1, 5–18. MR2353238
88. Shokrollah Salarian, Sean Sather-Wagstaff, and Siamak Yassemi, *Characterizing local rings via homological dimensions and regular sequences*, J. Pure Appl. Algebra **207** (2006), no. 1, 99–108. MR2244383
89. Sean Sather-Wagstaff, *Complete intersection dimensions and Foxby classes*, preprint, 2008, arXiv:0709.2442v3 [math.AC].
90. Sean Sather-Wagstaff, Tirdad Sharif, and Diana White, *Comparison of relative cohomology theories with respect to semidualizing modules*, preprint, 2007, arXiv:0706.3635v1 [math.AC] 28 pp., to appear in Math. Z.
91. ———, *Stability of Gorenstein categories*, J. Lond. Math. Soc. (2) **77** (2008), no. 2, 481–502. MR2400403
92. Sean Sather-Wagstaff and Diana White, *An Euler characteristic for modules of finite G-dimension*, Math. Scand. **102** (2008), no. 2, 206–230. MR2437187
93. Reza Sazeedeh, *Gorenstein injective modules and local cohomology*, Proc. Amer. Math. Soc. **132** (2004), no. 10, 2885–2891. MR2063107
94. ———, *Gorenstein injective, Gorenstein flat modules and the section functor*, J. Pure Appl. Algebra **211** (2007), no. 3, 773–783. MR2344228
95. Jean-Pierre Serre, *Sur la dimension homologique des anneaux et des modules noethériens*, Proceedings of the international symposium on algebraic number theory, Tokyo & Nikko, 1955 (Tokyo), Science Council of Japan, 1956, pp. 175–189. MR0086071
96. Jack Shamash, *The Poincaré series of a local ring*, J. Algebra **12** (1969), 453–470. MR0241411
97. Rodney Y. Sharp, *Finitely generated modules of finite injective dimension over certain Cohen-Macaulay rings*, Proc. London Math. Soc. (3) **25** (1972), 303–328. MR0306188
98. Ryo Takahashi, *Some characterizations of Gorenstein local rings in terms of G-dimension*, Acta Math. Hungar. **104** (2004), no. 4, 315–322. MR2082781
99. ———, *On the category of modules of Gorenstein dimension zero*, Math. Z. **251** (2005), no. 2, 249–256. MR2191025
100. ———, *The existence of finitely generated modules of finite Gorenstein injective dimension*, Proc. Amer. Math. Soc. **134** (2006), no. 11, 3115–3121. MR2231892

101. ———, *On G -regular local rings*, preprint, 2007, to appear in *Comm. Algebra*.
102. Ryo Takahashi and Kei-ichi Watanabe, *Totally reflexive modules constructed from smooth projective curves of genus $g \geq 2$* , *Arch. Math. (Basel)* **89** (2007), no. 1, 60–67. MR2322781
103. Ryo Takahashi and Yuji Yoshino, *Characterizing Cohen-Macaulay local rings by Frobenius maps*, *Proc. Amer. Math. Soc.* **132** (2004), no. 11, 3177–3187. MR2073291
104. Wolmer V. Vasconcelos, *Divisor theory in module categories*, North-Holland Publishing Co., Amsterdam, 1974, North-Holland Mathematics Studies, No. 14, Notas de Matemática No. 53. [Notes on Mathematics, No. 53]. MR0498530
105. Oana Veliche, *Construction of modules with finite homological dimensions*, *J. Algebra* **250** (2002), no. 2, 427–449. MR1899298
106. ———, *Gorenstein projective dimension for complexes*, *Trans. Amer. Math. Soc.* **358** (2006), no. 3, 1257–1283. MR2187653
107. Charles A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994. MR1269324
108. Diana White, *Gorenstein projective dimension with respect to a semidualizing module*, preprint, 2006, [arXiv:math/0611711](https://arxiv.org/abs/math/0611711) [math.AC].
109. Jinzhong Xu, *Flat covers of modules*, Lecture Notes in Mathematics, vol. 1634, Springer-Verlag, Berlin, 1996. MR1438789
110. Siamak Yassemi, *G -dimension*, *Math. Scand.* **77** (1995), no. 2, 161–174. MR1379262
111. ———, *A generalization of a theorem of Bass*, *Comm. Algebra* **35** (2007), no. 1, 249–251. MR2287566
112. Yuji Yoshino, *Modules of G -dimension zero over local rings with the cube of maximal ideal being zero*, *Commutative algebra, singularities and computer algebra (Sinaia, 2002)*, NATO Sci. Ser. II Math. Phys. Chem., vol. 115, Kluwer Acad. Publ., Dordrecht, 2003, pp. 255–273. MR2030276