

ORDER-THEORETICAL CONNECTIVITY

T.A. RICHMOND

Department of Mathematics
Western Kentucky University
Bowling Green KY 42101, U. S. A.

and

R. VAINIO

Department of Mathematics
Åbo Akademi
SF-20500 Åbo, Finland

(Received December 29, 1989 and in revised form February 10, 1990)

ABSTRACT. Order-theoretically connected posets are introduced and applied to create the notion of T -connectivity in ordered topological spaces. As special cases T -connectivity contains classical connectivity, order-connectivity, and link-connectivity.

KEY WORDS AND PHRASES: Topologies on posets, links, maximal chains, order-theoretical connectivity, order-connectivity, link-connectivity, and T -connectivity.

AMS SUBJECT CLASSIFICATION CODES: 54D05, 54F05.

INTRODUCTION

In the literature on order and topology (ordered topological spaces, topological lattices, order-determined topologies, etc.) connectivity has generally been introduced as a *topological property*. Given some natural compatibility between topology and order, much is known about the impact different connectivity properties have on the order in question. Classical theory (in case of total order), as well as [2], [4], and [5], have revealed the close relationship between topological connectivity on one hand and the order-theoretical properties "conditionally complete" and "order dense" on the other. The aim of this note is to define a concept of *order-theoretical connectivity* (using conditional completeness and order density), and then to apply this concept to the theory of ordered topological spaces. First, some relevant remarks on (partial) order and connectivity will be given.

A triple $S \equiv (S, \tau, o)$ consisting of a set, an order o and a topology τ on that set is called an *ordered topological space*. The space S is called *order-connected* (cf. [1]) if all order-preserving continuous maps from S to the space $\mathbf{2}$ are constants ($\mathbf{2} \equiv$ the set $\{0, 1\}$ endowed with discrete topology and natural order); *connected* (*path-connected*) if it is connected (*path-connected*) as a topological space; and *link-connected* if any two points of S can be joined by a finite set of connected links of S (link between x and $y \equiv$ maximal chain of $[x, y]$). A class \mathcal{C} of subsets of a given set S is called a *connectivity system* (cf. [2]) if any set $A \subseteq S$ is a member of \mathcal{C} whenever for all $x, y \in A$ there are sets $C_1, C_2, \dots, C_n \in \mathcal{C}$ in A with $x \in C_1, y \in C_n$, and $C_i \cap C_{i+1} \neq \emptyset$ for $i = 1, 2, \dots, n-1$. The classes of order-connected, or connected, or path-connected, or link-connected sets constitute connectivity systems.

An ordered topological space is a T_i -ordered space (an i -space) if every maximal chain inherits a topology which is finer than or equal to (which coincides with) its own interval topology (see [2], [5]). Note that for any lattice all topologies between the interval topology and Birkhoff's order topology are i -topologies (cf. [5]).

"Conditionally complete" and "order dense" will be abbreviated "cc" and "od", respectively. A subchain of an ordered set is called cc (od) if it is cc (od) regarded as an ordered space in its own right. It is important to recall that

- (A) a chain is connected in its interval topology iff it is od and cc;
- (B) an ordered set is od (cc) iff all maximal chains of it are od (cc);
- (C) an od and cc chain between two points of a given ordered set is always a complete link.

1. ORDER-THEORETICAL CONNECTIVITY

Let P be an arbitrary poset. Regard the equivalence relation \sim^{ord} defined in P by $a \sim^{\text{ord}} b \iff a$ and b are joined by a finite set of od and cc links in P .

The \sim^{ord} classes are called the *order-theoretical components* in P . In case of only one equivalence class, P is *order-theoretically connected*. The order-theoretically connected subsets (defined in the obvious way) of any poset are a connectivity system in the sense of [2]. Moreover, a poset is order-theoretically connected iff it is link-connected in its interval topology (or equivalently, in any of its i -structures).

PROPOSITION 1. *For any $a \in P$, the order-theoretical component of a equals the maximal order-theoretically connected subset of P containing a .*

LEMMA 2. *In any product poset $P \equiv \prod P_i$, all projections of an od and cc link J are od and cc links.*

PROOF. For a topological proof, endow P and the P_i 's with Birkhoff's order topology o , note that J is a connected set in $o(P)$ and that all projections are continuous maps $o(P) \rightarrow o(P_i)$. Thus $pr_i(J)$ is a connected chain in $o(P_i)$, and since $pr_i(J)$ carries a structure which is finer than its own interval topology it is an od and cc chain. •

PROPOSITION 3. *The product poset $P \equiv P_1 \times P_2 \times \dots \times P_n$ is order-theoretically connected iff every factor is. The order-theoretical component of $a \equiv (a_1, a_2, \dots, a_n) \in P$ equals the product of the order-theoretical components of the a_k 's ($k = 1, 2, \dots, n$).*

PROOF. To see the first part for $n = 2$, take $a_2 \in P_2$, express the product set $P_1 \times P_2$ as $\bigcup_{a_1 \in P_1} [(P_1 \times \{a_2\}) \cup (\{a_1\} \times P_2)]$, and use Lemma 2. Then proceed via induction over the natural numbers. The remaining part is straightforward. •

EXAMPLE 4 (denumerable products). Consider the Euclidean plane, let l_k denote the line segment ($y = \frac{1}{k}, 0 \leq x \leq 1$) and m_k the line segment connecting the right endpoint of l_k and the left endpoint of l_{k+1} . Define the lattice L to be the union of all the l_k 's and m_k 's ($k \in \mathbf{Z}_+$) endowed with the natural order of the plane. Obviously, L is order-theoretically connected. To see that the product lattice $P \equiv L^{\mathbf{Z}^+}$ is not, take pairs of points a_n, b_n in L which can be joined by a set of no less than n od and cc links of L and note $(a_n) \sim (b_n)$ is not satisfied in P (use Lemma 2).

EXAMPLE 5 (basic remarks). The od and cc lattices form a class of order-theoretically connected structures which is closed under formation of arbitrary products. A lattice is od and cc iff all maximal chains (or all links) are order-theoretically connected. As was demonstrated in Lemma 2, projections preserve order-theoretical connectivity. However,

a lattice with prime ideals (for instance any distributive lattice) allows a lattice-morphism onto the lattice $\mathbf{2}$. This shows that order-theoretical connectivity cannot be characterized via morphisms (and, of course, that morphisms do not preserve order-theoretical connectivity). Cantor's teepee (cf. [3]) with base interval $[0, 1]$ and top $(1, 1)$ is a connected subset of \mathbf{R}^2 ; nevertheless, all its order-theoretical components are singletons (which is a quite natural order-theoretical property of this structure). Finally, note that the topological closure of the lattice L of Example 4 (the Euclidean topology) consists of two order-theoretical components, although L is order-theoretically connected.

In ordered topological spaces S , order-theoretical connectedness relates to path-connectedness, as demonstrated below.

PROPOSITION 6. *Suppose each $x \in S$ has an order-theoretically connected neighborhood. Then S is order-theoretically connected if S is path-connected.*

PROOF: Suppose $x, y \in S$ and $p : \mathbf{I} \rightarrow S$ is a path between x and y . For each $z \in p(\mathbf{I})$, let N_z be an order-theoretically connected neighborhood of z , and let $M_z \subseteq N_z$ be an open neighborhood of z . Consider $\{N_i\}_{i=1}^n$ where $\{M_i\}_{i=1}^n$ is a subcover of $p(\mathbf{I})$ from $\{M_z : z \in p(\mathbf{I})\}$. Now $(\cup_{i=1}^k N_i) \cap (\cup_{i=k+1}^n N_i) \neq \emptyset$ for any $1 \leq k \leq n$, for otherwise $p^{-1}(\cup_{i=1}^k M_i) \cup p^{-1}(\cup_{i=k+1}^n M_i)$ would separate \mathbf{I} for some k . Since the order-theoretically connected subsets of S form a connectivity system, it follows that x and y are in the same order-theoretical component and thus S is order-theoretically connected. •

An obvious sufficient condition for order-theoretical connectedness to imply path-connectedness is that every cc and od link from x to y be a path from x to y . Applying this and the proposition we finally get

COROLLARY 7. *An open subset of \mathbf{R}^n is path-connected iff it is order-theoretically connected, given the usual topology and order on \mathbf{R}^n .*

2. T-CONNECTIVITY

Order-theoretical connectedness is now employed to create a natural connectivity concept in the category of (partially) ordered topological spaces and order-preserving continuous maps (here called *morphisms*).

Let T denote an ordered topological space.

The ordered topological space S is called *T-connected* if for every morphism $\varphi : S \rightarrow T$ the image $\varphi(S)$ is an order-theoretically connected set in T . A subset of an arbitrary ordered topological space is a *T-connected set* if it is a *T-connected space* (regarded as a subspace). Every ordered topological space can be partitioned into maximal *T-connected sets*, the *T-components* of the space.

Obviously, the *T-connected sets* in any ordered topological space form a connectivity system in the sense of [2]. *T-connectivity* provides the realm for several well-known connectivity concepts. For instance, an ordered topological space is order-connected iff it is $\mathbf{2}$ -connected. As a special case of this, note that a topological space is connected in the classical sense iff it is a $\mathbf{2}$ -connected ordered topological space when equipped with the discrete order. A connected space endowed with any order is always an \mathbf{R} -connected ordered topological space.

PROPOSITION 8. *Let S be a T_i -ordered topological space. Consider the following statements*

- (1) S is link-connected;
- (2) S is T -connected for all T_i -ordered topological spaces T ;
- (3) S is S -connected;
- (4) S is order-theoretically connected.

Implications (1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (4) are always true. If S is an i -space, then (4) \Rightarrow (1).

PROPOSITION 9. *Everything stated in Proposition 8 holds for T -connectivity; moreover, T -connectivity is preserved by morphisms (the choice of T is arbitrary, throughout).*

EXAMPLE 10 (*denumerable products*). The cc lattice L of Example 4 endowed with Euclidean topology is T -connected for all T_i -ordered topological spaces T (cf. Proposition 8). The product lattice $P \equiv L^{\mathbb{Z}^+}$ endowed with product topology is an i -space. Since the space P cannot be link-connected (pick elements $(a_n), (b_n) \in P$ as in Example 4), Proposition 8 guarantees the existence of at least one T_i -ordered space T , for which P is not T -connected.

EXAMPLE 11 (*basic remarks*). The set L of the previous example is an \mathbf{R}^2 -connected set in the Euclidean plane. However, the closure of L (denoted by \bar{L}) consists of two \mathbf{R}^2 -components, one of which fails to be a closed set in the space \bar{L} . Also, note that any $T_{3,5}$ topological space endowed with discrete order is totally \mathbf{R}^2 -disconnected (i.e. all \mathbf{R}^2 -components consist of only one point).

Again let T be any ordered topological space.

PROPOSITION 12. *A connected ordered topological space S , in which every point has a T -connected neighborhood, is T -connected.*

PROOF. For $x, y \in S$, let $x \sim^T y$ denote that for all morphisms $\varphi : S \rightarrow T$ the points $\varphi(x), \varphi(y)$ are joined by a finite set of od and cc links in $\varphi(S)$. Take $a \in S$, define the subset A of S by $x \in A \iff x \sim^T a$, and prove $A = S$ by showing A is open and closed as follows. First, take $b \in A$ and let V be a T -connected neighborhood of b . For any $x \in V$, $x \sim^T b \sim^T a$, and thus $x \in A$. Then, for b in the closure of A and V as above, $A \cap V \neq \emptyset$ and every $x \in A \cap V$ satisfies $b \sim^T x \sim^T a$, which implies $b \in A$. •

REFERENCES

1. D.C.J. BURGESS and S.D. MCCARTAN, *Order-continuous functions and order-connected spaces*, **Proc. Camb. Phil. Soc.** **68** (1970), pp 27-31.
2. M. ERNÉ and R. VAINIO, *Connectivity in lattice-ordered spaces*, to appear in **Math. Nachr.**
3. L.A. STEEN and J.A. SEEBACH, *Counterexamples in topology*, Holt - Rinehart - Winston, New York 1968.
4. R. VAINIO, *Connectedness properties of lattices*, **Canad. Math. Bull.** **29** (1986), pp 314-320.
5. R. VAINIO, *A maximal chain approach to topology and order*, **Internat. J. Math. and Math. Sci.** **11** no. 3 (1988), pp 465-472.

Special Issue on Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mssklai@cityu.edu.hk