

Regulatory Genes Controlling Fatty Acid Catabolism and Peroxisomal Functions in the Filamentous Fungus *Aspergillus nidulans*[†]

Michael J. Hynes,* Sandra L. Murray, Anna Duncan,‡ Gillian S. Khew, and Meryl A. Davis

Department of Genetics, University of Melbourne, Parkville 3010, Australia

Received 16 December 2005/Accepted 20 February 2006

The catabolism of fatty acids is important in the lifestyle of many fungi, including plant and animal pathogens. This has been investigated in *Aspergillus nidulans*, which can grow on acetate and fatty acids as sources of carbon, resulting in the production of acetyl coenzyme A (CoA). Acetyl-CoA is metabolized via the glyoxalate bypass, located in peroxisomes, enabling gluconeogenesis. Acetate induction of enzymes specific for acetate utilization as well as glyoxalate bypass enzymes is via the Zn₂-Cys₆ binuclear cluster activator FacB. However, enzymes of the glyoxalate bypass as well as fatty acid beta-oxidation and peroxisomal proteins are also inducible by fatty acids. We have isolated mutants that cannot grow on fatty acids. Two of the corresponding genes, *farA* and *farB*, encode two highly conserved families of related Zn₂-Cys₆ binuclear proteins present in filamentous ascomycetes, including plant pathogens. A single ortholog is found in the yeasts *Candida albicans*, *Debaryomyces hansenii*, and *Yarrowia lipolytica*, but not in the *Ashbya*, *Kluyveromyces*, *Saccharomyces* lineage. Northern blot analysis has shown that deletion of the *farA* gene eliminates induction of a number of genes by both short- and long-chain fatty acids, while deletion of the *farB* gene eliminates short-chain induction. An identical core 6-bp *in vitro* binding site for each protein has been identified in genes encoding glyoxalate bypass, beta-oxidation, and peroxisomal functions. This sequence is overrepresented in the 5' region of genes predicted to be fatty acid induced in other filamentous ascomycetes, *C. albicans*, *D. hansenii*, and *Y. lipolytica*, but not in the corresponding genes in *Saccharomyces cerevisiae*.

It has become increasingly clear that the breakdown of fatty acids is important in the metabolism, development, and pathogenicity of many fungi. Catabolism occurs via the beta-oxidation pathway, in which fatty acids are activated to the corresponding acyl coenzyme A (CoA) and then oxidation by a series of enzyme steps releases acetyl-CoA and an acyl-CoA shortened by two carbons, which can undergo additional cycles of beta-oxidation. In mammals, beta-oxidation of long-chain fatty acids occurs in peroxisomes, while medium- and short-chain fatty acids undergo beta-oxidation in the mitochondria (reviewed in references 16 and 84). In contrast, in *Saccharomyces cerevisiae* fatty acids are metabolized entirely in peroxisomes (reviewed in reference 29). In fungi, where fatty acids can serve as sole sources of carbon and energy, the acetyl-CoA must be converted to C₄ compounds via the glyoxalate bypass, comprising the enzymes isocitrate lyase and malate synthase, allowing gluconeogenesis (40, 64). Isocitrate lyase and malate synthase are usually, but not always, located in peroxisomes. It has been found that mutations affecting isocitrate lyase, malate synthase, and peroxisomal functions can affect the pathogenicity of both plant and animal pathogens (33, 37, 45, 46, 66). Furthermore it has been found that genes encoding enzymes for fatty acid catabolism are up regulated during infection. For example, microarray analysis of genes during infection of macrophages by *Candida albicans* showed increased expression of

a number of genes involved in fatty acid utilization (47). In the plant pathogen *Magnaporthe grisea*, the generation of turgor pressure required for penetration of the appressorium has been proposed to depend on the conversion of lipids to glycerol via beta-oxidation and the glyoxalate bypass (69, 85). In addition, fatty acid metabolism has been implicated in secondary metabolism and development in fungi (7, 59, 72). Therefore, the understanding of the regulation of these metabolic pathways in fungi is of considerable significance.

In *S. cerevisiae*, oleate utilization depends on peroxisomal beta-oxidation. A wide range of peroxisomal proteins and the enzymes necessary for complete conversion of oleate to acetyl-CoA are induced by oleate (29). Acetyl-CoA produced in the peroxisome is converted to acetyl-carnitine for export to the cytosol and the mitochondrion (18, 39, 63, 79, 80). The related transcriptional activators Oaf1 and Pip2 containing DNA binding domains of the Zn₂-Cys₆ binuclear class together with Adr1, a C₂H₂ zinc finger protein, are responsible for this induction via the *cis*-acting elements (ORE) and UAS1, respectively (4, 25, 34, 35, 48, 57, 58). The genes for metabolism of the resulting acetyl-CoA by the glyoxalate bypass and gluconeogenesis is controlled by the Zn₂-Cys₆ binuclear proteins Cat8 and Sip4 acting at carbon source-responsive elements (CSREs) in the 5' region of the relevant genes (26, 27, 54, 55, 56, 81). Some of these genes are also regulated by Adr1 (86). Growth on ethanol or acetate as sole carbon sources is also dependent on the Cat8, Sip4, and Adr1 activators as well as the Snf1 kinase (82, 86). In the presence of glucose the Mig1 repressor represses the expression of these genes (23). Growth on acetate depends on acetyl-CoA synthetase-dependent formation of acetyl-CoA in the cytosol. This is converted to acetyl-carnitine by a cytosolic acetyl-carnitine transferase and transported

* Corresponding author. Mailing address: Department of Genetics, University of Melbourne, Parkville, Victoria 3010, Australia. Phone: 61 3 83446239. Fax: 61 3 83445139. E-mail: mjhynes@unimelb.edu.au.

[†] Supplemental material for this article may be found at <http://ec.asm.org/>.

[‡] Present address: Department of Physiology, University of Melbourne, Parkville 3010, Australia.

into the mitochondrion, where it is converted back to acetyl-CoA by a mitochondrial acetyl-carnitine transferase for metabolism by the tricarboxylic acid cycle (18, 62, 80).

In the filamentous fungus *Aspergillus nidulans*, mutations in the *facA* gene, encoding acetyl-CoA synthetase, and in *facC*, encoding cytosolic acetyl-carnitine transferase, result in an inability to use acetate as a sole carbon source. These genes are induced by acetate via the FacB activator, which is related to the Cat8 and Sip4 proteins of *S. cerevisiae* (10, 31, 61, 67, 74, 76, 77). Mutations in *acuD* and *acuE*, which encode the glyoxalate bypass enzymes isocitrate lyase and malate synthase, respectively, are also unable to use acetate, but in addition are unable to use fatty acids as sole carbon sources (2, 3, 21, 36, 61). These genes are also regulated by acetate induction via FacB, but *facB* mutations do not prevent growth on fatty acids, and these genes are, in addition, regulated by fatty acid induction (2, 5). Similarly, *acuJ* encoding an acetyl-carnitine transferase located in both mitochondria and peroxisomes is required for growth on both acetate and fatty acids because it is necessary for the shuttling of acetyl-CoA via the transporter AcuH (14, 50, 67). The *acuJ* gene is also subject to FacB-mediated acetate induction as well as fatty acid induction (67; M. J. Hynes, A. Andrianopoulos, S. Delimitrou, S. L. Murray, H. Sealy-Lewis, and M. A. Davis, unpublished data). This is also the case for *idpA*, the gene for NADP-isocitrate dehydrogenase located in peroxisomes and in mitochondria (68).

A. nidulans is able to utilize short-chain and long-chain fatty acids, and recently it has been shown that beta-oxidation pathways exist in both mitochondria and peroxisomes, as is the case for mammals and plants. There is evidence for fatty acid induction of some of the relevant enzymes in *A. nidulans* (49, 78) and in *Neurospora crassa* (28, 38). Nothing is known about the mechanism or the regulatory genes involved. It is therefore predicted that there are three classes of genes: (i) acetate specific, regulated by FacB; (ii) acetate and fatty acid induced and controlled by FacB and unknown fatty acid regulators, and (iii) fatty acid specific induced only by the fatty acid regulators. This latter class is predicted to include enzymes required for fatty acid breakdown via beta-oxidation as well as proteins involved in peroxisome biogenesis and function.

We have sought to investigate this regulatory circuit by isolating mutants unable to use fatty acids as carbon sources but still capable of growth on acetate and by cloning the corresponding genes. Among the genes identified are ones predicted to encode Zn₂-Cys₆ binuclear proteins. These have been shown to control the expression of genes involved in fatty acid catabolism but not to affect expression of genes involved specifically in acetate utilization. One of these genes affects induction by short-chain (C₂ to C₆) fatty acids and is not highly conserved in other fungi. Two other genes, one affecting short-chain induction and one affecting induction by all chain lengths, encode related proteins with highly conserved orthologs found in other euascomycetes and a single ortholog in the hemiascomycetes *Candida albicans*, *Debaromyces hansenii*, and *Yarrowia lipolytica* but not in other hemiascomycete lineages. These proteins have been found to bind in vitro to sequences found to be overrepresented in the 5' region of a large number of genes predicted to be involved in fatty acid breakdown in the species which contain predicted orthologs but not in the 5' region of the corresponding genes in *S.*

TABLE 1. Primers used in this study

Primer name	Sequence
acuJ upper	GACACTAGTATATGATCGGAC
acuJ lower	AACTGCAGCGGAGGGACGA
LT1:RT5'	AACCTTCTCTGGATGGATGC
LT1:RT3'	AGAGCACTGAAACAATCCGG
BUZ7:RT5'	ATGACACGCAGTCGATTTGG
BUZ7:RT3'	TTTCTGACCTGCCCAAGC
BUX75'-RT	GCATTGAAAGTATTGTGCGG
BUX7RTPCR3'	GAAGATCTACTTTCACGCAGCGTCTCTT

cerevisiae. We have therefore identified a highly conserved regulatory circuit controlling the use of fatty acids in ascomycete fungi.

MATERIALS AND METHODS

A. nidulans strains, media, and transformation. Media and conditions for growth of *A. nidulans* were as described by Cove (11). Carbon and nitrogen sources were added as appropriate to minimal salts. The pH of these was adjusted to 6.5 where necessary, and long-chain fatty acids added were usually dispersed in the medium by the addition of 0.5 to 1% Tergitol (NP-40; Sigma) before melting or autoclaving. All strains were derived from the original Glasgow strain and contained the *velA1* mutation to promote uniform flat conidiation, and standard genetic manipulations were as previously described (8, 9). Preparation of protoplasts and transformation were as described elsewhere (1).

Molecular techniques. Standard methods for DNA manipulations, RNA isolation, nucleic acid blotting, and hybridization have been described previously (12, 60, 75). The sequences of oligonucleotides used are given in Table 1.

Isolation and assays of *acuJ-lacZ* reporter strains. A PCR fragment from the 5' region of the *acuJ* gene (corresponding to the predicted gene AN6279.2 in the genome database [http://www.broad.mit.edu/annotation/fungi/aspergillus/]), which has been shown to encode a peroxisomal-mitochondrial carnitine acetyltransferase (Hynes et al., unpublished), was generated using the primers *acuJ* upper (containing a SpeI site) and *acuJ* lower (containing a PstI site). This fragment contained sequences from -82 to -349 of the 5' region (relative to the ATG) and was cloned into SpeI-PstI-digested pANXKSK+, a vector containing a minimal promoter (gpd-mini) driving expression of the *Escherichia coli lacZ* gene as well as a mutated *A. nidulans argB* gene that allows site-specific integration at the *argB* locus when transformed into *argB2*-containing strains (53). Transformation of the vector alone has resulted in a strain with a single plasmid copy integrated, and assays of β -galactosidase showed low levels of expression under the conditions used in this paper (M. J. Hynes, unpublished data). Transformation of the *acuJ-lacZ* plasmid into an *argB2* strain yielded a transformant, MH9975, shown by Southern blotting to have a single plasmid integrated, in which β -galactosidase activities were regulated by fatty acids. Crosses generated *acuJ-lacZ* strains in various genetic backgrounds. Extracts of mycelium were prepared and assayed for β -galactosidase as described elsewhere (13).

Isolation of mutants. A conidial suspension of strain MH9975 was treated with diepoxyoctane as described previously (32), and dilutions plated on glucose-minimal ammonium tartrate medium with 0.08% sodium deoxycholate were added to restrict the colony size. Colonies were velvet replicated to minimal medium containing 20 mM sodium butyrate as the sole carbon source with 10 mM ammonium chloride as the nitrogen source. Potential mutants affected in butyrate utilization were isolated and tested for their ability to use L-glutamate (50 mM) and acetate (50 mM) as carbon sources. Mutants specifically affected in butyrate utilization were isolated for further study. This yielded strains containing the *scfA7* and *farB7* mutations. One of the butyrate-nonutilizing mutants (like a number of others) was completely inhibited in growth on medium containing fatty acids. This mutant subsequently was shown to have a mutation in a gene (*pexF*) encoding an ortholog of the peroxin Pex6 (42, 72, 83). This strain was inoculated onto minimal medium containing 0.5% lactose and 0.2% Tween 80, as a source of the fatty acid oleate, and ammonium chloride. Spontaneous resistant sectors appearing after incubation for 3 to 5 days were isolated. Outcrossing of these yielded isolates separated from the original *pexF* mutation that were unable to use fatty acids as sole carbon sources but did not show the extreme *pexF* phenotype. This yielded a strain with the *farA1* mutation.

Molecular cloning and sequencing of genes. Strains containing the proposed regulatory mutations were crossed to a strain containing the *pyrG89* mutation to




FIG. 1. Growth of regulatory gene mutants on fatty acids as sole carbon sources. The following carbon sources were added to minimal medium with 10 mM ammonium chloride as the nitrogen source: glucose (1%); acetate (50 mM); butyrate and valerate (10 mM); hexanoate (5 mM), propionate, Tween 20, and Tween 80 (0.1%); lauric, myristic, palmitic, heptadecanoic, stearic, oleate, elaidic, and erucic acids (2.5 mM). Growth was for 2 to 3 days at 37°C.

generate PyrG⁻ double mutants. These strains were used for cloning the genes by functional complementation by transforming with a genomic library in the autonomously replicating vector pRG3AMA1 (51). Transformants selected for uracil-uridine prototrophy were velvet replicated to medium containing butyrate as the sole carbon source, complementing transformants were selected and purified, and genomic DNA was prepared. Plasmid DNA was recovered by transforming this DNA into *E. coli*, selecting for ampicillin resistance. Alternatively, direct PCR on genomic DNA was used to recover insert DNA from complementing plasmids. Subcloning, sequencing, and comparison with the *A. nidulans* database (<http://www.broad.mit.edu/annotation/fungi/aspergillus/>) showed that *scfA* corresponded to the annotated gene AN1303.2, *farA* corresponded to AN7050.2, and *farB* corresponded to AN1425.2.

cDNA corresponding to the genes was synthesized from total RNA using gene-specific primer pairs LT1:RT5' and LT1:RT3', BUZ7RT5' and BUZ7RT3', and BUX75'RT and BUX7RTPCR3' for *farA*, *farB*, and *scfA*, respectively, using the Superscript II (Invitrogen) one-step reverse transcriptase PCR kit. Products were cloned into the plasmid pGEMTEasy (Promega Corp.) and sequenced. This led to a reannotation of intron positions in the genes (see Fig. 2 and 7, below).

Generation of gene deletions. Restriction fragments containing the *A. nidulans* *pyrG* gene were inserted into the plasmids pSM5953 (*farA*), pAD5875 (*farB*), and pSM5831 (*scfA*), replacing sequences with coordinates relative to the genes +225 to +2760, -277 to +2368, and -295 to +1470, respectively. Linear fragments generated by restriction digests were transformed into the *pyrG89* strain MH10076, selecting for uracil-uridine prototrophy. Transformants showing the phenotypes of the corresponding original mutants were analyzed by Southern blot analysis of genomic DNA to confirm precise gene replacement events and further confirmed by complementation of phenotypes by transforming deletion mutants with plasmids containing wild-type genes.

Expression of MBP fusion proteins. Fragments of *farA*, *farB*, and *scfA* genes were expressed as maltose binding protein (MBP) fusion proteins using the pMAL system (New England BioLabs). The cDNA clones of *farA*, *farB*, and *scfA*, pSM6128, pSM6123, and pSM5777, respectively, were used as templates for PCRs using primers to introduce a restriction site enabling the cloning of complete cDNAs of *farA*, *farB*, and *scfA* into the pMALC2 polylinker in the same reading frame as *malE*. The resulting plasmids, pSM6350 (encoding MBP fused to amino acids 1 to 399 of FarA), pSM6351 (encoding MBP fused to amino acids 1 to 514 of FarB), and pSM6229 (encoding MBP fused to amino acids 1 to 410 of ScfA), together with the previously described pRT2013 (encoding MBP fused to amino acids 4 to 417 of FacB [74]) were transformed into *E. coli* strain BL21-DE3 or Rosetta (Novagen) for expression. An overnight culture was diluted 1:100 in M9ZB medium, with the addition of 1 mM MgSO₄, 0.4% glucose, and grown at 37°C with aeration to an optical density at 600 nm of 0.6. Expression of fusion plasmids was induced by the addition of isopropyl-β-D-thiogalactopyranoside to a 1 mM final concentration for 2 h. Cells were harvested, and crude extracts were produced as described in the pilot experiment of the pMAL protein fusion and purification system manual (New England BioLabs). Protein concentrations were determined using the Bio-Rad protein assay reagent (Bio-Rad). Expression of fusion proteins of the predicted sizes was visualized by sodium dodecyl sulfate-polyacrylamide gel electrophoresis.

Electrophoretic mobility shift assays (EMSAs). Oligonucleotide probes were prepared, and binding reactions were carried out as previously described except that no poly(dI-dC) or Mg²⁺ was added (74). Electrophoresis of samples occurred on a nondenaturing 5% polyacrylamide gel in 0.5× Tris-borate-EDTA at

100 to 150 V. Gels were blotted onto 3MM paper, dried in a vacuum dryer, and exposed to X-ray film.

Nucleotide sequence accession numbers. The sequences determined in this study have been deposited in GenBank and assigned the accession numbers DQ386636 (*scfA*), DQ386637 (*farA*), and DQ386638 (*farB*).

RESULTS

Isolation of *farA* and *farB* mutants and cloning of the corresponding genes. Mutants unable to grow on medium containing butyrate as a sole carbon source were isolated by replica plating from glucose plates containing colonies derived from diluted mutagenized conidiospores. Those specifically unable to grow on butyrate but capable of growth on glutamate and acetate as sole carbon sources were isolated for further analysis. Studies on mutants found to be affected in a variety of beta-oxidation and peroxisomal functions will be reported elsewhere. One of these mutants was completely inhibited on medium containing fatty acids but not acetate. This mutant was found to contain a mutation in the ortholog of the *PEX6* gene of *S. cerevisiae*, which encodes a peroxin required for the import of proteins into the peroxisomal matrix (42, 72, 83). It was thought that this toxicity resulted from mislocalization of peroxisomal beta-oxidation enzymes. Therefore, it was proposed that mutations affecting fatty acid induction would result in resistance. This mutant was used to isolate revertants capable of growth on medium containing 0.5% lactose (to provide a nonrepressing carbon source) and 0.1% Tween 80, which contains principally oleate. The revertants showed weak growth on this medium compared to the parent strain, which was completely inhibited, consistent with growth on lactose as the carbon source and resistance to the effects of oleate catabolism. Outcrossing of one of these mutants allowed the separation of the new mutation from the original peroxisome defect mutant. Strains containing the new mutation were unable to use all fatty acids tested as sole carbon sources, and this was shown to be due to a mutation in a single gene, designated *farA* (Fig. 1). A further two alleles of this gene were isolated in this revertant screen.

A second mutant was found to contain a mutation in a single gene, designated *farB*, resulting in loss of the ability to use the short-chain fatty acids butyrate, valerate, and hexanoate as sole carbon sources but capable of growth on longer-chain-length fatty acids (lauric acid, C₁₂ and above) (Fig. 1A). As reported




FIG. 2. Structures of *farA* and *farB* genes and comparisons with orthologs in other fungi. (A) The *farA* gene, showing intron positions and the sequences coding for the indicated domains. *farA*Δ was generated by replacing sequences from +225 to +2670 (relative to the start codon) with the *A. nidulans pyrG* gene followed by gene replacement. (B) Structure of the *farB* gene, showing intron positions and the sequences coding for the indicated domains. *farB*Δ was generated by replacing sequences from -227 to +2368 (relative to the start codon) with the *A. nidulans pyrG* gene followed by gene replacement. (C) Comparison of the Zn₂-Cys₆ binuclear cluster domains of the FarA and FarB proteins in filamentous fungi and the orthologs in hemiascomycetes compared to *A. nidulans* FarA (Yeast group). Identical residues present in at least 60% of the sequences are indicated by black boxes, whereas gray shading represents similar residues. Overall comparisons of the proteins, database accession numbers, and intron

elsewhere, C₇ to C₁₀ fatty acids were found to be toxic to all strains (49).

The *farA* and *farB* genes were cloned by complementation by the isolation of double mutants containing the *pyrG89* mutation, transforming with a genomic library in the autonomously replicating vector pRG3AMA1 (51), selecting for PyrG⁺ transformants, and replica plating to medium with butyrate as the sole carbon source. Colonies capable of growth were purified and DNA isolated. The corresponding autonomously replicating plasmids were recovered by transformation into *E. coli*. Sequencing of insert DNA in the recovered plasmids showed that *farA* corresponded to the annotated gene AN7050.2, while *farB* corresponded to AN1425.2 in the genome sequence (<http://www.broad.mit.edu/annotation/fungi/aspergillus/>). cDNA of each gene, made by reverse transcription-PCR, was cloned and sequenced, leading to a reannotation of intron positions in each gene (Fig. 2A and B). Each gene was deleted by replacement of sequences with the *pyrG* gene as shown in Fig. 2. The generated mutations segregated as single genes, and each deletion mutant was complemented by subclones of the relevant cloned genes, indicating that no additional events occurred during the isolation of the deletions.

The corresponding phenotypes with respect to fatty acid utilization of each gene deletion were identical to those of the original isolated mutants (Fig. 1B). More extensive testing of the deletion mutants showed that *farB*Δ resulted in loss of utilization of short-chain fatty acids but not inhibition of growth and did not affect growth on long-chain fatty acids. The *farA* deletion resulted in loss of the utilization of short-chain fatty acids but also resulted in loss of utilization and inhibition on all fatty acids tested longer than C₁₀ (lauric acid) up to C₂₂ (erucic acid), including the odd-chain-length heptadecanoate (C₁₇) and the unsaturated oleic and elaidic acids (C₁₈). The loss-of-function *facB101* mutation (2, 76) did not affect growth on fatty acids apart from a slight effect on butyrate. Growth on propionate (C₃) was inhibited in the mutants as well as the wild type and the *facB101* mutant was resistant, consistent with previous results showing that propionyl-CoA generated from propionate is inhibitory to growth (6).

The *farA* and *farB* genes encode conserved proteins containing fungal-specific transcription factor domains. Each of the predicted proteins encoded by the two genes were found to contain both a Zn₂-Cys₆ binuclear cluster domain and a conserved central domain characteristic of fungal transcription factors containing this DNA binding domain (Fig. 2A, B, and C) (62, 73). Database searches revealed predicted conserved orthologs corresponding to each of these genes in filamentous euascomycetes, including a number of plant pathogens (see

positions in the genes are presented in the Fig. S1 to S3 in the supplemental material. (D) Unrooted neighbor joining distance tree showing the relationships of the FarA, FarB, and yeast proteins to each other. Sequences were aligned using ClustalW (70) and distance matrices were calculated (19), which were used to create the tree in NJPlot (52). Species abbreviations: A.n, *Aspergillus nidulans*; A.f, *Aspergillus fumigatus*; S.n, *Stagonospora nodorum*; F.g, *Fusarium graminearum*; N.h, *Nectria haematococca*; M.g, *Magnaporthe grisea*; N.c, *Neurospora crassa*; C.g, *Chaetomium globosum*; B.c, *Botrytis cineria*; C.a, *Candida albicans*; D.h, *Debaryomyces hansenii*; Y.l, *Yarrowia lipolytica*.

Fig. S1 and S2 in the supplemental material). In the case of *farA* the first four introns, including two introns within the sequence encoding the Zn₂-Cys₆ domain, were found to be conserved in all genes and to have led to extensive misannotation in the databases (see Fig. S4 in the supplemental material). The FarA and FarB proteins showed similarity to each other, and each of the species also had a single copy of each gene, with the exception of *Aspergillus fumigatus*, which was found to have two predicted copies of the *farB* gene, apparently arising due to a segmental duplication of approximately 22.6 kbp encoding seven hypothetical annotated genes from chromosome 8 to chromosome 1 (see Fig. S2 in the supplemental material). A single copy of a related gene was found in the hemiascomycete species *Candida albicans*, *Debaryomyces hansenii*, and *Yarrowia lipolytica*, but not in *Ashbya (Eremothecium) gossypii* nor in *S. cerevisiae* (see Fig. S3 in the supplemental material). *Y. lipolytica* had one of the introns conserved in other species, while the *C. albicans* and *D. hansenii* genes lacked introns (see Fig. S4 in the supplemental material). The neighbor joining tree constructed showed that the hemiascomycete proteins were most closely related to the FarA group, which were tightly clustered in the euascomycetes. The FarB group was more divergent (Fig. 2D).

In the filamentous ascomycetes, the proposed Zn₂-Cys₆ binuclear DNA binding domains of these proteins were found to be extraordinarily conserved, particularly within the FarA group, and to be related between groups and with the corresponding domains in the hemiascomycete proteins (Fig. 2C). Of particular note was the finding that the predicted proteins were highly similar to the cutinase transcription factors CTF α and CTF β of *Nectria hematococca*. These proteins have been found to bind to sequences in the 5' region of cutinase genes in this plant pathogen (43, 44). Cutin is a polymer containing hydroxy-oleate, strongly suggesting a connection with fatty acid breakdown.

Effects of the *farA* and *farB* genes on induction of an *acuJ-lacZ* reporter construct. The *acuJ* gene has been found to encode a carnitine-acetyltransferase located to peroxisomes and mitochondria and is regulated by FacB-dependent acetate induction and induced by fatty acids (50, 67; Hynes et al., unpublished). A sequence from -82 to -349 from the *acuJ* gene was inserted upstream of the minimal *gpd* promoter driving *lacZ* expression (53), and this was inserted in single copy at the *argB* locus (Fig. 3A). For studies of induction, proline was used as a noninducing limiting carbon source control. The original *farA* and *farB* mutations as well as the deletions were found to affect induction with *farB*, reducing induction by short-chain fatty acids but not by oleate or Tween 20 (a source of lauric, palmitic, and myristic acids), while the *farA* mutations led to elevated levels of noninduced expression but reduced induction by all fatty acids tested (Fig. 3B, C, and H). This was also observed for the *farA* Δ *farB* Δ double mutant (Fig. 3G). In short time course induction experiments, butyrate and oleate induction was detectable at about 60 min (Fig. 3D and E). Acetate and propionate were weak inducers, and this induction was not affected by the *facB101* mutation. In fact, stronger induction in this background as well as in a *facA303* background (lacking acetyl-CoA synthetase) was observed (Fig. 3G). This suggested that loss of activation of acetate and propionate to the respective acyl-CoA might enhance the weak

induction observed. Induction was observed in the presence of 1% glucose, and a loss of function mutation in the *creA* gene, encoding the major glucose repressor protein (15), only slightly increased the response to an inducer (Fig. 3F).

Northern blot analysis of fatty acid induction. RNA isolated from induced and noninduced (proline) strains was analyzed using probes derived from relevant genes (Fig. 4). Expression of the acetate-specific genes *facA* and *facC* was induced by acetate and butyrate, and this was dependent on FacB. Butyrate induction was lost in the *farA* Δ and *farB* Δ strains, which we interpret as indicating that butyrate must be metabolized to acetyl-CoA to result in induction via FacB. Long-chain (Tween 20 and oleate) induction was not observed. Similarly, genes required for both acetate and fatty acid utilization, *acuD*, *acuE*, *acuH*, and *acuJ*, were induced by acetate and butyrate in a FacB-dependent fashion. However, these genes were also long-chain fatty acid inducible, dependent on FarA. The other genes investigated were predicted to be involved specifically in fatty acid utilization, and these were not induced by acetate but were induced by butyrate (FarB and FarA dependent) and Tween 20 and oleate (FarA dependent).

Detection of DNA binding by expressed FarA and FarB fusion proteins. FarA and FarB proteins containing the proposed DNA binding domains were expressed in *E. coli* as fusion proteins with *E. coli* MBP (see Materials and Methods) for use in EMSAs. Because of the high degree of similarity between the proposed DNA binding domains of FarA and FarB with the *N. hematococca* cutinase transcription factors (Fig. 2C), the choice of DNA probes for analysis was guided by studies with these proteins (43, 44). As summarized in Fig. 5A, binding of these proteins to a sequence with a core CCGAGG but not to mutant versions of this sequence were previously observed, and this was consistent with methylation protection experiments (44). Two sequences containing this core were observed in the *acuJ* 5' fragment that conferred fatty acid induction on the *lacZ* reporter (Fig. 3A). Two probes, J1-2 and J7-8, based on the *acuJ* 5' sequence (Fig. 5B), were bound by the MBP-FarA and MBP-FarB extracts but not by the MBP extract (Fig. 5C). In addition, J1-2 and J7-8 were weakly bound by MBP-FacB (74). Both of these probes contain the sequence CCAN5CCG (where N = A, G, C, or T), a sequence related to the *S. cerevisiae* CSRE element bound by Sip4/Cat8 (81). The similarity of the FacB DNA binding domain to that of Cat8/Sip4 (77) presumably accounts for this weak binding. However, this is unlikely to be physiologically significant, as indicated by the lack of effect of the *facB101* mutation on expression of the *acuJ-lacZ* reporter (Fig. 3G).

Additional probes for the EMSA were based on the presence of the core CCGAGG (complement, CCTCGG) in the 5' region of other genes predicted to be regulated by fatty acids. The gene designated as *pexK* (corresponding to the annotated gene AN1921) is predicted to encode the peroxin Pex11 involved in peroxisome biogenesis (42, 72, 83). This was found to be induced by fatty acids (Fig. 4). The probe K13-14 containing the core sequence was found to be bound by MBP-FarA and MBP-FarB (Fig. 5D). Similarly, the probe EC9-10, based on a 5' sequence upstream of the *echA* gene encoding a mitochondrial enoyl-CoA hydratase (49), also bound MBP-FarA and MBP-FarB (Fig. 5D). However, probe AE11-12, based on a sequence from the *acuE* 5' region, was found to only weakly




FIG. 3. Effects of regulatory gene mutations on expression of an *acuJ-lacZ* reporter. (A) A fragment of the *acuJ* 5' region was inserted upstream of a minimal promoter (*gpd mini*) driving *lacZ* expression. The black bars represent CCGAGG sequences present in this sequence. (B to H) Mycelia of strains of the indicated genotypes were grown for 16 to 18 h in 1% glucose–10 mM ammonium tartrate medium and then transferred to minimal medium with the indicated carbon sources together with 10 mM ammonium chloride for 6 h or for the indicated times (D and E) before harvesting. Mycelia were extracted and assayed for β -galactosidase as previously described (13). The specific activity is expressed in Miller units per minute per milligram of protein. The bars represent standard errors. The concentrations of carbon sources were 10 mM for proline, acetate, butyrate, valerate, and oleate, 5 mM for hexanoate, 0.1% for propionate and Tween 20, and 2.5 mM for myristic, palmitic, heptadecanoic, elaidic, stearic, and erucic acids.

bind to MBP-FarB and not MBP-FarA (Fig. 5D). The biological significance of this result remains to be determined.

Unlabeled DNA (J3-4) containing the mutant sequence CCcgGG replacing the core sequence showed a loss of competition for binding of MBP-FarA and MBP-FarB with the J7-8 probe, although some competition for MBP-FarB was observed (Fig. 5E). Overall the results showed, in agreement with previous observations for the *N. hematococca* CTF proteins, that the sequence CCGAGG (complement, CCTCGG) is the core binding sequence for these conserved proteins. It was therefore predicted that this sequence would be found in the 5' regions of genes induced by fatty acids.

Analysis of 5' sequences of fungal genes for predicted binding sites. Sequences from the 5' region (1 kb upstream of the start codon) of a large number of genes from the *A. nidulans*

genome were scanned for the CCTCGG sequence. One or more sequences were found in nearly all genes predicted to be involved in fatty acid metabolism (see Table S1 in the supplemental material). These included genes determining proteins involved in peroxisome biogenesis or function, enzymes of beta-oxidation, including those predicted to be in mitochondria or peroxisomes, metabolite transporters, and glutathione peroxidases (but not catalases) involved in dealing with oxidative stress. The motif was found to occur in either orientation, and at least one copy was most commonly found within 300 bp of the start codon.

As part of the comparative analysis of the genomes of *A. nidulans*, *Aspergillus oryzae*, and *A. fumigatus* (22), a three-way alignment of 1 kb upstream of orthologous genes using Mlgan was performed. The CCTCGG motif was found to commonly




FIG. 4. Analysis of the effects of regulatory gene mutations on expression of various genes. RNAs were extracted from strains of the indicated genotypes grown for 16 h in 1% glucose–10 mM ammonium tartrate medium and then transferred to medium with the indicated carbon sources for 6 h. Abbreviations: pro, 10 mM L-proline; ace, 10 mM acetate; but, 10 mM n-butyrate; ole, 10 mM oleate; Tw20, 0.1% Tween 20. The results represent three different Northern blot membranes probed multiple times with sequences corresponding to the indicated genes with their proposed functions. Probes were made by labeling either gel-purified restriction fragments or PCR products generated using primers based on genome sequences. H3 represents an EcoRI fragment of the histone H3 clone (17), and rRNA refers to the large rRNA species observed by ethidium bromide staining.

occur in high-scoring conserved sequences upstream of relevant genes (these are indicated in Table S1 in the supplemental material). Furthermore, analysis of the patterns of occurrence of these conserved sequences (“conpat” analysis) identified the CCTCGG sequence to be enriched in genes for lipid metabolism and peroxisome localization (see Table 3 in reference 22).

Analysis of the 5' regions of genes required for acetate utilization showed the presence of FacB sites in all genes, while the CCTCGG motif was present only in genes also regulated by fatty acids (Fig. 6). A more limited analysis of 1 kb upstream of genes from other species containing *farA* and *farB* orthologs showed that the CCTCGG motif was overrepresented in those genes that might be expected to be regulated by fatty acids but

not in acetate-specific genes (Table 2; see also Table S1 in the supplemental material). In the case of the plant pathogen *M. grisea*, this included members of the cutinase gene family, and the motif was also found upstream of cutinase genes as well as a predicted lipase in *A. nidulans*. *S. cerevisiae* lacks *farA* and *farB* orthologs, and fatty acid induction is dependent on the Oaf1/Pip2 heterodimer (29, 34, 35, 58). Consistent with this, the CCTCGG motif was not overrepresented in the 5' region of relevant genes from *S. cerevisiae* (Table 2).

A third Zn₂-Cys₆ domain-containing protein involved in induction by short-chain fatty acids. Another mutant, isolated by virtue of lack of growth on butyrate, was found to contain a mutation in a single gene, designated *scfA*, and resulted in loss





FIG. 6. Positions of FacB binding sites and CCTCGG motifs in the 5' regions of genes involved in acetate utilization in *A. nidulans*. FacB binding sites are based on those previously proposed (67, 74) or by sequence inspection in the case of *acuJ*. Vertical lines represent 100-bp intervals upstream of the start codon for each gene.

of the ability to use the short-chain fatty acids butyrate, valerate, and hexanoate as sole carbon sources but was able to grow on longer-chain-length fatty acids (lauric acid [C_{12}] and above) (Fig. 1A). The *scfA* gene was cloned by complementation by the same method used for the other genes. Sequencing showed that this gene corresponded to the annotated gene AN1303.2 (<http://www.broad.mit.edu/annotation/fungi/aspergillus/>). However, isolation and sequencing of cDNA showed that reassignment of intron positions was required. This resulted in the prediction that *scfA* encoded a protein containing a Zn_2 -Cys₆ binuclear DNA binding motif as well as a fungal-specific transcription factor central domain (Fig. 7A). In marked contrast to FarA and FarB, ScfA was not highly conserved in other fungi, with two possible orthologs with similar DNA binding domains being found only in *A. fumigatus* and *N. crassa* (Fig. 7B), and the overall similarity was not high (see Fig. S5 in the supplemental material). The *scfA* gene was deleted by replacement with *pyrG* (Fig. 7A), and the phenotypes of *scfA* Δ strains were found to be the same as for *scfA7* mutant strains and very similar to

FIG. 5. EMSA analysis of DNA binding by expressed MBP fusion proteins. (A) Summary of the key DNA binding studies of cutinase transcription factors in *N. haematococca* (43, 44). Binding was detected to the probe Pal2 but not Pal2M1 or Pal2M2. Methylation protection studies showed the protection of the G residues, indicated by asterisks. (B) Probes used in EMSAs in this study. Complementary pairs of oligonucleotides were designed based on sequences from the 5' regions of the genes *acuJ*, *echA*, *acuE*, and *pexK*, with *SpeI/XbaI*- or *BamHI/BglII*-compatible ends added (in lowercase letters). The database designation of these genes is given in Table S1 in the supplemental material. The conserved core CCGAGG sequence is indicated in bold, and only one strand is shown. Probe J3-4 is identical to J1-2 except for the modification of the core CCGAGG to CCgGG. (C and D) EMSA of extracts (20 μ g of protein added to binding reaction mixtures) to the indicated probes together with no-extract controls. Free probe is designated as F. Binding is represented by the decreased mobility of the probe in the presence of protein, compared to the free probe. (E) Specific competition of binding of FarB and FarA to probe J7-8. Binding was performed with MBP-FarA and MBP-FarB extracts (20 μ g of protein) in the presence of increasing amounts of unlabeled competitor DNA. Competitors were the unlabeled probes J1-2 and J3-4, increasing from 10 to 100 to 500 times the concentration of labeled probe.

TABLE 2. Presence of CCTCGG motif in the 5' region of relevant genes from fungal species^a

Protein function	Position relative to start codon in:						
	<i>A. nidulans</i>	<i>N. crassa</i>	<i>M. grisea</i>	<i>Y. lipolytica</i>	<i>C. albicans</i>	<i>D. hansenii</i>	<i>S. cerevisiae</i>
Acetyl-CoA synthetase		-796					-170
Carnitine acetyl transferase (cytosol)		-577	-420	-697			
			-503				
Carnitine acetyl transferase (peroxisome/mitochondrion)	-175	-955	-578	-138	-397	-132	
	-338			-661	-983	-167	
				-994		-806	
Isocitrate lyase	-432	-291	-471	-291	-338	-193	
		-718	-639	-908	-623	-236	
		-770	-793			-589	
Malate synthase	-603	-164	^b	-663	-470	-402	-429
	-738	-643					
Peroxin (PEX11)	-280	-471	-224	-115	-308	-183	
			-364				
			-624				

^a Positions of CCTCCG or the complement CGGAGG were obtained by scanning 1 kb upstream of the proposed start codon. Genes for *M. grisea*, *N. crassa*, and *A. nidulans* were from annotated sequences obtained from <http://www.broad.mit.edu/annotation/fgi/>. Reannotation was carried out as necessary. Genes from *Y. lipolytica* and *D. hansenii* were from <http://www.ncbi.nlm.nih.gov/>; genes from *C. albicans* were from <http://www.candidagenome.org/>, and genes from *S. cerevisiae* were from <http://www.yeastgenome.org/>.

^b Only approximately 400 bp was available because of the position at end of the contig.

those for *farBΔ* strains (Fig. 1B). Furthermore, *scfAΔ* affected short-chain but not long-chain fatty acid induction of the *acuJ-lacZ* reporter (Fig. 3) as well as showing a similar pattern to the effects of *farBΔ* on transcription of the genes studied by Northern blotting (Fig. 4). Expression of ScfA as an MBP fusion protein followed by EMSA with the probes used to detect FarA and FarB binding failed to give binding (Fig. 5). Additional probes based on sequences present in the *acuJ* -82 to -349 region have so far also failed to give significant binding (results not shown).

DISCUSSION

The starting point for this study was the finding that genes specific for acetate utilization were only controlled by FacB-dependent acetate induction, while those required for both acetate and fatty acid utilization were regulated both by FacB and independently by fatty acids. This is consistent with the finding that *facB* mutations lead to loss of the ability to use acetate as a carbon source but do not affect growth on fatty acids requiring beta-oxidation. The approach of isolating mutants able to grow on acetate but not on butyrate has been successful in leading to the discovery of regulatory genes involved in fatty acid induction of genes regulated also by FacB as well as genes only regulated by fatty acids. Like *facB*, the *farA* and *farB* genes encode proteins containing Zn₂-Cys₆ binuclear DNA binding domains.

The FarA and FarB proteins are highly conserved in filamentous ascomycetes and have related DNA binding domains. The previously described cutinase transcription factors from *N. hematococca* (43, 44), which were defined by DNA binding studies using 5' sequences from cutinase genes, are clearly orthologs and, in agreement with these studies, we have shown that expressed fusion proteins of FarA and FarB bind in vitro to sequences containing the core CCTCGG sequence. Analysis of fungal genomes has shown that this sequence is present in one or more copies within 1 kb of the predicted start codon of a large number of genes predicted to encode proteins involved in fatty acid metabolism, providing strong support for these regulatory genes being involved in induction by fatty acids. Furthermore, comparative analyses of the genomes of *Aspergillus* species indicate a conservation of this sequence in the upstream region of orthologous genes enriched for lipid metabolism and peroxisomal functions (22). Our findings on the in vivo functions of FarA and FarB provide a striking validation of the value of the phylogenetic footprinting methods used to determine potential functional motifs. The ease of detection of this simple sequence makes it possible to predict in many




FIG. 7. Structure of the *scfA* gene. (A) The *scfA* gene, showing intron positions and the sequences coding for the indicated domains. *scfAΔ* was generated by replacing sequences from -295 to +1470 (relative to the start codon) with the *A. nidulans pyrG* gene followed by gene replacement. (B) Comparison of the Zn₂-Cys₆ binuclear cluster domains of *scfA* with proposed orthologs in *N. crassa* and *A. fumigatus*. Identical residues present in at least 60% of the sequences are indicated by black boxes, whereas gray shading represents similar residues. Overall comparisons of the proteins and database accession numbers are presented in Fig. S5 in the supplemental material.

ascomycete species those genes likely to be regulated by fatty acid induction via these proteins.

In addition to the glyoxalate cycle enzymes, we have found direct evidence for FarA and FarB regulating genes involved in the shuttling of acetyl-CoA between peroxisomes and mitochondria (*acuJ* and *acuH*) as well as genes encoding both mitochondrial and peroxisomal beta-oxidation enzymes. We have also found that an ortholog of *S. cerevisiae* *PEX11*, predicted to encode a peroxin required for peroxisome proliferation (42, 72, 83), is regulated by FarA and FarB, and we have shown that deletion of this gene in *A. nidulans* results in reduced growth on fatty acids (G. S. Khew and M. J. Hynes, unpublished data). The CCTCGG motif was found upstream of a number of predicted orthologs for key peroxisomal proteins, including Pex3, which is absolutely required for de novo peroxisome formation (30, 41), as well as Pex1, Pex6, and Pex5, which are required for protein import into the peroxisomal matrix. We have isolated strains containing mutations in these genes and found that the resulting loss of peroxisomal functions leads to mild developmental defects (e.g., reduced sporulation) and complete inhibition of growth by both long- and short-chain fatty acids (Khew and Hynes, unpublished). The *farA*Δ and *farB*Δ strains do not show these phenotypes. This indicates that the Pex phenotypes result from mislocalization of induced fatty acid metabolic activities, and these effects are not observed when induction is lost. In addition, it is possible that induction of these peroxisomal functions is not absolutely required for expression but rather serves to increase the capacity for peroxisome formation and function.

Filamentous ascomycetes are very versatile in their ability to use diverse carbon sources. The finding that the conserved CCTCGG motif is present in the upstream region of the cutinase gene families of the plant pathogens *M. grisea* and *N. hematoococca* as well as the saprophyte *A. nidulans* indicates that regulation of these genes is via fatty acid induction. We have not extensively scanned potential lipase genes, but in one case in *A. nidulans* we found the 5' motif. In an expressed sequence tag analysis of the insect pathogen *Metarhizium anisopliae* grown under conditions of maximal cuticle degradation, lipases and phospholipases were detected and, significantly, an ortholog of the *N. hematoococca* cutinase transcription factor was found (20). This result is now understandable, since it is highly likely that this fungal species uses orthologs of *farA* and *farB* to control enzymes involved in lipid degradation. A broad role for these transcription factors in controlling the expression of enzymes for degradation of sources of fatty acids in filamentous fungi is strongly suggested.

At this stage many questions relating to the mechanism of action of these transcription factors remain to be answered. Deletion of *farA* results in loss of the ability to use all fatty acids tested and loss of induction by fatty acids. However, the noninduced activity of the *acuJ-lacZ* reporter was found to be elevated, implying a repressing function for FarA in the absence of inducer. Deletion of *farB* abolishes growth on and induction by short-chain fatty acids but does not affect long-chain induction. Somewhat increased induction by long-chain fatty acids of the *acuJ-lacZ* reporter was also observed in the *farB*Δ background (Fig. 3). Given that FarA and FarB recognize the same DNA sequence, this may indicate competition for binding between inducer-activated and nonactivated pro-

teins. Deletion of *scfA* gives a very similar phenotype to that of the *farB*Δ mutant in causing loss of growth on and induction by short-chain fatty acids but does not result in increased long-chain fatty acid induction. We have so far been unable to determine an ScfA binding site, and it is puzzling that, in stark contrast to FarA and FarB, highly conserved orthologs of this protein have not been found in other species. The possibilities that ScfA acts upstream of FarB by controlling the production of the short-chain inducing signal or that it is required for FarB expression are currently being examined. The nature of the inducing signal and the discrimination between short-chain and long-chain inducers is of considerable interest. Response of the *acuJ-lacZ* reporter to inducer is relatively rapid (Fig. 3), perhaps indicating that the fatty acid inducer does not require extensive metabolism to generate the signal. In *S. cerevisiae* it has been proposed that the activators Oaf1 and Pip2 respond directly to oleate as inducer, but direct evidence is lacking (29). Zn₂-Cys₆ binuclear cluster proteins have been proposed to commonly respond directly to inducing metabolites (65). It should also be noted that as beta-oxidation of long-chain fatty acids proceeds, the shortening of chain length might result in an increasing level of short-chain induction. Analysis of the *A. nidulans* genome reveals multiple genes potentially carrying out enzymatic steps in beta-oxidation, and the substrate and induction specificity of these may be complex.

The finding that the hemiascomycete yeasts *C. albicans*, *D. hansenii*, and *Y. lipolytica* contain orthologs of FarA is intriguing. Other hemiascomycetes have orthologs of the well-studied Oaf1 and Pip2 genes of *S. cerevisiae*. The overrepresentation of the CCTCGG motif in the 5' region of relevant genes in these species, but not in *S. cerevisiae*, provides strong circumstantial evidence for the functional role of these FarA orthologs. The phylogenetic analysis presented (Fig. 2D) suggests one scenario in which a FarA orthog involved in long-chain induction was present in a progenitor of the ascomycetes. In filamentous euascomycetes, this was duplicated and diverged, resulting in the FarB genes being involved in a greater metabolic capacity to respond to short-chain substrates. FarA may have been lost and replaced by Oaf1/Pip2 orthologs in the other hemiascomycetes. Peroxisome biogenesis and alkane utilization in *Y. lipolytica* have been the subject of much study (e.g., references 24, 71, and 72), and investigation of the regulatory role of the FarA ortholog in this species is required.

Recent studies have suggested that peroxisomal functions and lipid metabolism are important for fungal pathogenesis (33, 37, 45–47, 66, 69, 85). Our discovery in plant and animal pathogens of transcription factors that are highly likely to be involved in controlling these processes is therefore significant. Pathogens may use these regulatory proteins to respond to host fatty acid signals to upregulate functions necessary for the establishment of infection. Deletion of the genes in these species and testing pathogenicity will therefore be of considerable interest.

ACKNOWLEDGMENTS

This work was supported by a grant from the Australian Research Council.

Assistance by Khanh Nguyen, Rosemary Genovese, Sophie Delimitrou, and Oliver Evans is acknowledged.

REFERENCES

- Andrianopoulos, A., and M. J. Hynes. 1988. Cloning and analysis of the positively acting regulatory gene *amdR* from *Aspergillus nidulans*. *Mol. Cell. Biol.* **8**:3532–3541.
- Armitt, S., W. McCullough, and C. F. Roberts. 1976. Analysis of acetate non-utilizing (*acu*) mutants in *Aspergillus nidulans*. *J. Gen. Microbiol.* **92**: 263–282.
- Ballance, D. J., and G. Turner. 1986. Gene cloning in *Aspergillus nidulans*: isolation of the isocitrate lyase gene (*acuD*). *Mol. Gen. Genet.* **202**:271–275.
- Baumgartner, U., B. Hamilton, M. Piskacek, H. Ruis, and H. Rottensteiner. 1999. Functional analysis of the Zn_2Cys_6 transcription factors Oaf1p and Pip2p. Different roles in fatty acid induction of beta-oxidation in *Saccharomyces cerevisiae*. *J. Biol. Chem.* **274**:22208–22216.
- Bowyer, P., J. R. De Lucas, and G. Turner. 1994. Regulation of the expression of the isocitrate lyase gene (*acuD*) of *Aspergillus nidulans*. *Mol. Gen. Genet.* **242**:484–489.
- Brock, M., and W. Buckel. 2004. On the mechanism of action of the anti-fungal agent propionate. *Eur. J. Biochem.* **271**:3227–3241.
- Calvo, A. M., H. W. Gardner, and N. P. Keller. 2001. Genetic connection between fatty acid metabolism and sporulation in *Aspergillus nidulans*. *J. Biol. Chem.* **276**:25766–25774.
- Clutterbuck, A. J. 1974. *Aspergillus nidulans* genetics, p. 447–510. In R. C. King (ed.), *Handbook of genetics*, vol. 1. Plenum Publishing Corp., New York, N.Y.
- Clutterbuck, A. J. 1994. Linkage map and locus list, p. 791–795. In S. D. Martinelli and J. R. Kinghorn (ed.), *Aspergillus: 50 years on*. Elsevier, Amsterdam, The Netherlands.
- Connerton, I. F., J. R. S. Fincham, R. A. Sandeman, and M. J. Hynes. 1990. Comparison and cross-species expression of the acetyl-CoA synthetase genes of the ascomycete fungi, *Aspergillus nidulans* and *Neurospora crassa*. *Mol. Microbiol.* **4**:451–460.
- Cove, D. J. 1966. The induction and repression of nitrate reductase in the fungus *Aspergillus nidulans*. *Biochim. Biophys. Acta* **113**:51–56.
- Davis, M. A., M. C. Askin, and M. J. Hynes. 2005. Amino acid catabolism by an *areA*-regulated gene encoding an L-amino acid oxidase with broad substrate specificity in *Aspergillus nidulans*. *Appl. Environ. Microbiol.* **71**:3551–3555.
- Davis, M. A., C. S. Cobbett, and M. J. Hynes. 1988. An *amdS-lacZ* fusion for studying gene regulation in *Aspergillus*. *Gene* **63**:199–212.
- De Lucas, J. R., A. I. Dominguez, S. Valenciano, G. Turner, and F. Laborda. 1999. The *acuH* gene of *Aspergillus nidulans*, required for growth on acetate and long-chain fatty acids, encodes a putative homologue of the mammalian carnitine/acylcarnitine carrier. *Arch. Microbiol.* **171**:386–396.
- Dowzer, C. E. A., and J. M. Kelly. 1991. Analysis of the *creA* gene, a regulator of carbon catabolite repression in *Aspergillus nidulans*. *Mol. Cell. Biol.* **11**: 5701–5709.
- Eaton, S., K. Bartlett, and M. Pourfarzam. 1996. Mammalian mitochondrial beta-oxidation. *Biochem. J.* **320**:345–357.
- Ehinger, A., S. H. Denison, and G. S. May. 1990. Sequence, organization and expression of the core histone genes of *Aspergillus nidulans*. *Mol. Gen. Genet.* **222**:416–424.
- Elgersma, Y., C. W. van Roermund, R. J. Wanders, and H. F. Tabak. 1995. Peroxisomal and mitochondrial carnitine acetyltransferases of *Saccharomyces cerevisiae* are encoded by a single gene. *EMBO J.* **14**:3472–3479.
- Felsenstein, J. 1989. PHYLIP: phylogeny inference package (version 3.2). *Cladistics* **5**:164–166.
- Freimoser, F. M., S. Screen, S. Bagga, G. Hu, and R. J. St. Leger. 2003. Expressed sequence tag (EST) analysis of two subspecies of *Metarhizium anisopliae* reveals a plethora of secreted proteins with potential activity in insect hosts. *Microbiology* **149**:239–247.
- Gainey, L. D., I. F. Connerton, E. H. Lewis, G. Turner, and D. J. Balance. 1992. Characterization of the glyoxysomal isocitrate lyase genes of *Aspergillus nidulans* (*acuD*) and *Neurospora crassa* (*acu-3*). *Curr. Genet.* **21**:43–47.
- Galagan, J. E., S. E. Calvo, C. Cuomo, L.-J. Ma, J. Wortman, S. Batzoglou, S.-I. Lee, M. Batürkmen, C. C. Spevak, J. Clutterbuck, V. Kapitonov, J. Jurka, C. Scacciochio, M. Farman, J. Butler, S. Purcell, S. Harris, G. H. Braus, O. Draht, S. Busch, C. D'Enfert, C. Bouchier, G. H. Goldman, D. Bell-Pedersen, S. Griffiths-Jones, J. H. Doonan, J. Yu, K. Veenken, A. Pain, M. Freitag, E. U. Selker, D. B. Archer, M. Á. Peñalva, B. R. Oakley, M. Momany, T. Tanaka, T. Kumagai, K. Asai, M. Machida, W. C. Nierman, D. W. Denning, M. Caddick, M. Hynes, M. Paoletti, R. Fischer, B. Miller, P. Dyer, M. S. Sachs, S. A. Osmani, and B. Birren. 2005. Sequencing of *Aspergillus nidulans* and comparative analysis with *A. fumigatus* and *A. oryzae*. *Nature* **438**:1105–1115.
- Gancedo, J. M. 1998. Yeast carbon catabolite repression. *Microbiol. Mol. Biol. Rev.* **62**:334–361.
- Guo, T., Y. Y. Kit, J. M. Nicaud, M. T. Le Dall, S. K. Sears, H. Vali, H. Chan, R. A. Rachubinski, and V. I. Titorenko. 2003. Peroxisome division in the yeast *Yarrowia lipolytica* is regulated by a signal from inside the peroxisome. *J. Cell Biol.* **162**:1255–1266.
- Gurvitz, A., J. K. Hiltunen, R. Erdmann, B. Hamilton, A. Hartig, H. Ruis, and H. Rottensteiner. 2001. *Saccharomyces cerevisiae* Adr1p governs fatty acid beta-oxidation and peroxisome proliferation by regulating POX1 and PEX11. *J. Biol. Chem.* **276**:31825–31830.
- Haurie, V., M. Perrot, T. Mini, P. Jenö, F. Saggiocco, and H. Boucherie. 2001. The transcriptional activator Cat8p provides a major contribution to the reprogramming of carbon metabolism during the diauxic shift in *Saccharomyces cerevisiae*. *J. Biol. Chem.* **276**:76–85.
- Hedges, D., M. Proft, and K. D. Entian. 1995. CAT8, a new zinc cluster-encoding gene necessary for derepression of gluconeogenic enzymes in the yeast *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **15**:1915–1922.
- Hii, V., and J. B. Courtright. 1982. Induction of acyl coenzyme A synthetase and hydroxyacyl coenzyme A dehydrogenase during fatty acid degradation in *Neurospora crassa*. *J. Bacteriol.* **150**:981–983.
- Hiltunen, J. K., A. M. Mursula, H. Rottensteiner, R. K. Wierenga, A. J. Kastaniotis, and A. Gurvitz. 2003. The biochemistry of peroxisomal beta-oxidation in the yeast *Saccharomyces cerevisiae*. *FEMS Microbiol. Rev.* **27**: 35–64.
- Hoepfner, D., D. Schildknecht, I. Braakman, P. Philippsen, and H. F. Tabak. 2005. Contribution of the endoplasmic reticulum to peroxisome formation. *Cell* **122**:85–95.
- Hynes, M. J. 1977. Induction of the acetamidase of *Aspergillus nidulans* by acetate metabolism. *J. Bacteriol.* **131**:770–775.
- Hynes, M. J. 1979. Fine-structure mapping of the acetamidase structural gene and its controlling region in *Aspergillus nidulans*. *Genetics* **91**:381–392.
- Idnurm, A., and B. J. Howlett. 2002. Isocitrate lyase is essential for pathogenicity of the fungus *Leptosphaeria maculans* to canola (*Brassica napus*). *Eukaryot. Cell* **1**:719–724.
- Karpichev, I. V., Y. Luo, R. C. Marians, and G. M. Small. 1997. A complex containing two transcription factors regulates peroxisome proliferation and the coordinate induction of beta-oxidation enzymes in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **17**:69–80.
- Karpichev, I. V., and G. M. Small. 1998. Global regulatory functions of Oaf1p and Pip2p (Oaf2p), transcription factors that regulate genes encoding peroxisomal proteins in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **18**:6560–6570.
- Katz, M. E., and M. J. Hynes. 1989. Isolation and analysis of the acetate regulatory gene, *facB*, from *Aspergillus nidulans*. *Mol. Cell. Biol.* **9**:5696–5701.
- Kimura, A., Y. Takano, I. Furusawa, and T. Okuno. 2001. Peroxisomal metabolic function is required for appressorium-mediated plant infection by *Colletotrichum lagenarium*. *Plant Cell* **13**:1945–1958.
- Kionka, C., and W. H. Kunau. 1985. Inducible beta-oxidation pathway in *Neurospora crassa*. *J. Bacteriol.* **161**:153–157.
- Kispal, G., B. Sumegi, K. Kietmeier, I. Bock, G. Gajdos, T. Tomcsanyi, and A. Sandor. 1993. Cloning and sequencing of a cDNA encoding *Saccharomyces cerevisiae* carnitine acetyltransferase. *J. Biol. Chem.* **268**:1824–1829.
- Kornberg, H. L. 1966. The role and control of the glyoxylate cycle in *Escherichia coli*. *Biochem. J.* **99**:1–11.
- Kragt, A., T. Voorn-Brouwer, M. van den Berg, and B. Distel. 2005. Endoplasmic reticulum-directed Pex3p routes to peroxisomes and restores peroxisome formation in a *Saccharomyces cerevisiae* *pex3Δ* strain. *J. Biol. Chem.* **280**:34350–34357.
- Lazarow, P. B. 2003. Peroxisome biogenesis: advances and conundrums. *Curr. Opin. Cell Biol.* **15**:489–497.
- Li, D., and P. E. Kolattukudy. 1997. Cloning of cutinase transcription factor 1, a transactivating protein containing Cys_8Zn_2 binuclear cluster DNA-binding motif. *J. Biol. Chem.* **272**:12462–12467.
- Li, D., T. Sirakova, L. Rogers, W. F. Ettinger, and P. E. Kolattukudy. 2002. Regulation of constitutively expressed and induced cutinase genes by different zinc finger transcription factors in *Fusarium solani* f. sp. *pisi* (*Nectria haematococca*). *J. Biol. Chem.* **277**:7905–7912.
- Lorenz, M. C., and G. R. Fink. 2001. The glyoxylate cycle is required for fungal virulence. *Nature* **412**:83–86.
- Lorenz, M. C., and G. R. Fink. 2002. Life and death in a macrophage: role of the glyoxylate cycle in virulence. *Eukaryot. Cell* **1**:657–662.
- Lorenz, M. C., J. A. Bender, and G. R. Fink. 2004. Transcriptional response of *Candida albicans* upon internalization by macrophages. *Eukaryot. Cell* **3**:1076–1087.
- Luo, Y., L. V. Karpichev, R. A. Kohanski, and G. M. Small. 1996. Purification, identification and properties of a *Saccharomyces cerevisiae* oleate-activated upstream activating sequence-binding protein that is involved in the activation of *POX1*. *J. Biol. Chem.* **271**:12068–12075.
- Maggio-Hall, L. A., and N. P. Keller. 2004. Mitochondrial beta-oxidation in *Aspergillus nidulans*. *Mol. Microbiol.* **54**:1173–1185.
- Midgley, M. 1993. Carnitine acetyltransferase is absent from *acuJ* mutants of *Aspergillus nidulans*. *FEMS Microbiol. Lett.* **108**:7–10.
- Oshover, N., and G. May. 2000. Conidial germination in *Aspergillus nidulans* requires RAS signaling and protein synthesis. *Genetics* **155**:647–656.
- Perrière, G., and M. Gouy. 1996. WWW-Query: an on-line retrieval system for biological sequence banks. *Biochimie* **78**:364–369.
- Punt, P. J., A. Kuyvenhoven, and C. A. van den Hondel. 1995. A mini-

- promoter *lacZ* gene fusion for the analysis of fungal transcription control sequences. *Gene* **158**:119–123.
54. **Rahner, A., A. Scholer, E. Martens, B. Gollwitzer, and H. J. Schuller.** 1996. Dual influence of the yeast Cat1p (Snf1p) protein kinase on carbon source-dependent transcriptional activation of gluconeogenic genes by the regulatory gene CAT8. *Nucleic Acids Res.* **24**:2331–2337.
 55. **Roth, S., and H.-J. Schuller.** 2001. Cat8 and Sip4 mediate regulated transcriptional activation of the yeast malate dehydrogenase gene MDH2 by three carbon source-responsive promoter elements. *Yeast* **18**:151–162.
 56. **Roth, S., J. Kumme, and H.-J. Schüller.** 2004. Transcriptional activators Cat8 and Sip4 discriminate between sequence variants of the carbon source-responsive promoter element in the yeast *Saccharomyces cerevisiae*. *Curr. Genet.* **45**:121–128.
 57. **Rottensteiner, H., A. J. Kal, B. Hamilton, H. Ruis, and H. F. Tabak.** 1997. A heterodimer of the Zn₂Cys₆ transcription factors Pip2p and Oaf1p controls induction of genes encoding peroxisomal proteins in *Saccharomyces cerevisiae*. *Eur. J. Biochem.* **247**:776–783.
 58. **Rottensteiner, H., L. Wabnegger, R. Erdmann, B. Hamilton, H. Ruis, A. Hartig, and A. Gurvitz.** 2003. *Saccharomyces cerevisiae* PIP2 mediating oleic acid induction and peroxisome proliferation is regulated by Adr1p and Pip2p-Oaf1p. *J. Biol. Chem.* **278**:27605–27611.
 59. **Ruprich-Robert, G., V. Berteaux-Lecellier, D. Zickler, A. Panvier-Adoutte, and M. Picard.** 2002. Identification of six loci in which mutations partially restore peroxisome biogenesis and/or alleviate the metabolic defect of *pep2* mutants in *Podospira*. *Genetics* **161**:1089–1099.
 60. **Sambrook, J., E. F. Fritsch, and T. Maniatis.** 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
 61. **Sandeman, R. A., and M. J. Hynes.** 1989. Isolation of the *facA* (acetyl-coenzyme A synthetase) and *acuE* (malate synthase) genes of *Aspergillus nidulans*. *Mol. Gen. Genet.* **218**:87–92.
 62. **Schjerling, P., and S. Holmberg.** 1996. Comparative amino acid sequence analysis of the C6 zinc cluster family of transcriptional regulators. *Nucleic Acids Res.* **24**:4599–4607.
 63. **Schmalix, W., and W. Bandlow.** 1993. The ethanol-inducible *YAT1* gene from yeast encodes a presumptive mitochondrial outer carnitine acetyltransferase. *J. Biol. Chem.* **268**:27428–27439.
 64. **Schuller, H. J.** 2003. Transcriptional control of nonfermentative metabolism in the yeast *Saccharomyces cerevisiae*. *Curr. Genet.* **43**:139–160.
 65. **Sellick, C. A., and R. J. Reece.** 2005. Eukaryotic transcription factors as direct nutrient sensors. *Trends Biochem. Sci.* **30**:405–412.
 66. **Solomon, P. S., R. C. Lee, T. J. Wilson, and R. P. Oliver.** 2004. Pathogenicity of *Stagonospora nodorum* requires malate synthase. *Mol. Microbiol.* **53**:1065–1073.
 67. **Stemple, C. J., M. A. Davis, and M. J. Hynes.** 1998. The *facC* gene of *Aspergillus nidulans* encodes an acetate-inducible carnitine acetyltransferase. *J. Bacteriol.* **180**:6242–6251.
 68. **Szewczyk, E., A. Andrianopoulos, M. A. Davis, and M. J. Hynes.** 2001. A single gene produces mitochondrial, cytoplasmic, and peroxisomal NADP-dependent isocitrate dehydrogenase in *Aspergillus nidulans*. *J. Biol. Chem.* **276**:37722–37729.
 69. **Thines, E., R. W. Weber, and N. J. Talbot.** 2000. MAP kinase and protein kinase A-dependent mobilization of triacylglycerol and glycogen during appressorium turgor generation by *Magnaporthe grisea*. *Plant Cell* **12**:1703–1718.
 70. **Thompson, J. D., D. G. Higgins, and T. J. Gibson.** 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**:4673–4680.
 71. **Titorenko, V. I., J. M. Nicaud, H. Wang, H. Chan, and R. A. Rachubinski.** 2002. Acyl-CoA oxidase is imported as a heteropentameric, cofactor-containing complex into peroxisomes of *Yarrowia lipolytica*. *J. Cell Biol.* **156**:481–494.
 72. **Titorenko, V. I., and R. A. Rachubinski.** 2001. The life cycle of the peroxisome. *Nat. Rev. Mol. Cell Biol.* **2**:357–368.
 73. **Todd, R. B., and A. Andrianopoulos.** 1997. Evolution of a fungal regulatory gene family: the Zn(II)₂Cys₆ binuclear cluster DNA binding motif. *Fungal Genet. Biol.* **21**:388–405.
 74. **Todd, R. B., A. Andrianopoulos, M. A. Davis, and M. J. Hynes.** 1998. FacB, the *Aspergillus nidulans* activator of acetate utilization genes, binds dissimilar DNA sequences. *EMBO J.* **17**:2042–2054.
 75. **Todd, R. B., J. A. Fraser, K.-H. Wong, M. A. Davis, and M. J. Hynes.** 2005. Nuclear accumulation of the GATA factor AreA in response to complete nitrogen starvation by regulation of nuclear export. *Eukaryot. Cell* **4**:1646–1653.
 76. **Todd, R. B., J. M. Kelly, M. A. Davis, and M. J. Hynes.** 1997. Molecular characterization of mutants of the acetate regulatory gene *facB* of *Aspergillus nidulans*. *Fungal Genet. Biol.* **22**:92–102.
 77. **Todd, R. B., R. L. Murphy, H. M. Martin, J. A. Sharp, M. A. Davis, M. E. Katz, and M. J. Hynes.** 1997. The acetate regulatory gene *facB* of *Aspergillus nidulans* encodes a Zn(II)₂Cys₆ transcriptional activator. *Mol. Gen. Genet.* **254**:495–504.
 78. **Valenciano, S., J. R. Lucas, A. Pedregosa, I. F. Monistrol, and F. Laborda.** 1996. Induction of beta-oxidation enzymes and microbody proliferation in *Aspergillus nidulans*. *Arch. Microbiol.* **166**:336–341.
 79. **van Roermund, C. W. T., Y. Elgersma, N. Singh, R. J. A. Wanders, and H. Tabak.** 1995. The membrane of peroxisomes in *Saccharomyces cerevisiae* is impermeable to NAD(H) and acetyl-CoA under in vivo conditions. *EMBO J.* **14**:3480–3486.
 80. **van Roermund, C. W., E. H. Hetteema, M. van den Berg, H. F. Tabak, and R. J. Wanders.** 1999. Molecular characterization of carnitine-dependent transport of acetyl-CoA from peroxisomes to mitochondria in *Saccharomyces cerevisiae* and identification of a plasma membrane carnitine transporter, Agp2p. *EMBO J.* **18**:5843–5852.
 81. **Vincent, O., and M. Carlson.** 1998. Sip4, a Snf1 kinase-dependent transcriptional activator, binds to the carbon source-responsive element of gluconeogenic genes. *EMBO J.* **17**:7002–7008.
 82. **Vincent, O., and M. Carlson.** 1999. Gal83 mediates the interaction of the Snf1 kinase complex with the transcription activator Sip4. *EMBO J.* **18**:6672–6681.
 83. **Voorn-Brouwer, T., I. van der Leij, W. Hemrika, B. Distel, and H. F. Tabak.** 1993. Sequence of the *PAS8* gene, the product of which is essential for biogenesis of peroxisomes in *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta* **1216**:325–328.
 84. **Wanders, R. J., P. Vreken, S. Ferdinandusse, G. A. Jansen, H. R. Waterham, C. W. van Roermund, and E. G. Van Grunsven.** 2001. Peroxisomal fatty acid alpha- and beta-oxidation in humans: enzymology, peroxisomal metabolite transporters and peroxisomal diseases. *Biochem. Soc. Trans.* **29**:250–267.
 85. **Wang, Z. Y., C. R. Thornton, M. J. Kershaw, L. Debaio, and N. J. Talbot.** 2003. The glyoxylate cycle is required for temporal regulation of virulence by the plant pathogenic fungus *Magnaporthe grisea*. *Mol. Microbiol.* **47**:1601–1612.
 86. **Young, E. T., K. M. Dombek, C. Tachibana, and T. Ideker.** 2003. Multiple pathways are co-regulated by the protein kinase Snf1 and the transcription factors Adr1 and Cat8. *J. Biol. Chem.* **278**:26146–26158.