

Regulation of JNK signaling by GSTp

Victor Adler, Zhimin Yin, Serge Y.Fuchs, Miriam Benezra, Lilliam Rosario¹, Kenneth D.Tew¹, Matthew R.Pincus², Mohinder Sardana³, Colin J.Henderson⁴, C.Roland Wolf⁴, Roger J.Davis⁵ and Ze'ev Ronai⁶

The Rutenberg Cancer Center, Mount Sinai School of Medicine, One Gustave L.Levy Place, Box 1130, New York, NY 10029-6574, ¹Department of Pharmacology, Fox Chase Cancer Center, Philadelphia, PA, ²Department of Pathology and Laboratory Medicine, SUNY Health Science Center, Brooklyn, NY, ³Protein Sequencing Facility, Merck Sharp & Dohme, West Point, PA, USA, ⁴Imperial Cancer Research Fund, Molecular Pharmacology Unit, Dundee, DD1 9SY, UK and ⁵Howard Hughes Medical Institute, University of Massachusetts, Worcester, MA, USA

⁶Corresponding author
e-mail: ronaiz01@doc.mssm.edu

Studies of low basal Jun N-terminal kinase (JNK) activity in non-stressed cells led us to identify a JNK inhibitor that was purified and identified as glutathione S-transferase Pi (GSTp) and was characterized as a JNK-associated protein. UV irradiation or H₂O₂ treatment caused GSTp oligomerization and dissociation of the GSTp–JNK complex, indicating that it is the monomeric form of GSTp that elicits JNK inhibition. Addition of purified GSTp to the Jun–JNK complex caused a dose-dependent inhibition of JNK activity. Conversely, immunodepleting GSTp from protein extracts attenuated JNK inhibition. Furthermore, JNK activity was increased in the presence of specific GSTp inhibitors and a GSTp-derived peptide. Forced expression of GSTp decreased MKK4 and JNK phosphorylation which coincided with decreased JNK activity, increased c-Jun ubiquitination and decreased c-Jun-mediated transcription. Co-transfection of MEKK1 and GSTp restored MKK4 phosphorylation but did not affect GSTp inhibition of JNK activity, suggesting that the effect of GSTp on JNK is independent of the MEKK1–MKK4 module. Mouse embryo fibroblasts from GSTp-null mice exhibited a high basal level of JNK activity that could be reduced by forced expression of GSTp cDNA. In demonstrating the relationships between GSTp expression and its association with JNK, our findings provide new insight into the regulation of stress kinases.

Keywords: GSTp/JNK/signaling/stress kinase

Introduction

Jun N-terminal kinases (JNKs) belong to the multi-member family of stress kinases that are activated transiently in response to UV- or X-irradiation, heat shock, osmotic

shock or inflammatory cytokines (Galcheva-Gargova *et al.*, 1994; Kyriakis *et al.*, 1994; Westwick *et al.*, 1994). JNK activation in response to UV irradiation is mediated by upstream signaling components, including cdc42, p21^{PAK}, ASK1, MLK, MEKK1, SEK1/MKK4, MKK7 (Coso *et al.*, 1995; Fanger *et al.*, 1997; Tournier *et al.*, 1997; reviewed by Ip and Davis, 1998) and p21^{ras} (Minden *et al.*, 1994; Adler *et al.*, 1995a, 1996), in concert with nuclear DNA lesions (Adler *et al.*, 1995b). Different forms of stress have been shown to mediate JNK activation via various cellular pathways (Adler *et al.*, 1995c). Activated JNK phosphorylates the transcription factors c-Jun, ATF2, p53 and ELK-1 (Kyriakis *et al.*, 1994; Gupta *et al.*, 1995; Whitmarsh *et al.*, 1995; Adler *et al.*, 1997; Fuchs *et al.*, 1998a). Phosphorylation by JNK has been implicated in stabilization (Fuchs *et al.*, 1997, 1998b) and transcriptional activity (Kyriakis *et al.*, 1994) of its substrates, which directly contribute to the mammalian stress response through changes in the cell cycle, DNA repair or apoptosis (Xia *et al.*, 1995; Fuchs *et al.*, 1998a; Kasibhatla *et al.*, 1998).

Despite the significant advances in identifying the components of stress-activated protein kinases, the mechanisms underlying the regulation of JNK before and immediately after stress are not well understood. In cells maintained under normal growth conditions, the basal activity of JNK is low, although JNK phosphorylation by upstream kinases occurs in response to growth factors (Minden *et al.*, 1994) and hence should also be observed in cells proliferating under normal growth conditions. While some unstressed cell types have been found to contain high levels of JNK activity (Dhar *et al.*, 1996), the reason for this high basal level is unknown. A 2- to 4-fold increase in constitutive JNK activity has been reported and, while this is lower than the activation upon exposure to DNA-damaging agents, it is equivalent to tumor necrosis factor- α (TNF- α) treatment and the degree of activation elicited by multiple exposures to low dose UV-B (Adler *et al.*, 1995d). As a key component in regulating the stability and activity of its substrates (Fuchs *et al.*, 1998c; Kyriakis *et al.*, 1994), changes in JNK's basal activity could influence key cellular functions, including growth, apoptosis and transformation.

Neither transcript nor protein levels of JNK are affected by stress. JNK activation is accomplished upon its phosphorylation by upstream kinases in response to stress (Coso *et al.*, 1995; Tournier *et al.*, 1997). Dephosphorylation of JNK at a later stage (Cavigelli *et al.*, 1996; Hanada *et al.*, 1998) is thought to constitute the primary mechanism underlying the regulation of JNK activity as a kinase. Nevertheless, several studies have demonstrated the existence of JNK inhibitors in normal growing cells. Thus, stress-mediated JNK activation may be independent of its upstream kinases. Among the cellular components

involved in regulating JNK substrates is a 'δ inhibitor' which blocks transactivation of c-Jun by interacting with the δ domain (Baichwal *et al.*, 1991). This is an integral 27 amino acid sequence (30–57) from the N-terminal region of c-Jun required for JNK binding to c-Jun, enabling Jun phosphorylation (Adler *et al.*, 1992) or ubiquitination (Treier *et al.*, 1994; Fuchs *et al.*, 1996). Other JNK regulatory proteins include p21^{cip/waf1}, which has been shown to exert greater inhibition of JNK activity in response to stress (Shim *et al.*, 1996).

Our interest in exploring the nature of JNK's low basal activity originated in the observation that extracts of non-stressed cells efficiently inhibited c-Jun phosphorylation when added to a solid-phase kinase reaction. Our further studies led to the purification, identification and characterization of glutathione *S*-transferase Pi (GSTp) as a JNK inhibitor, described in this report.

GSTs comprise a multigene family, of which GSTp is the most prevalent and ubiquitous non-hepatic isozyme (Jakoby, 1978). Among cellular functions attributed to GSTs are ligand binding and xenobiotic detoxification (Tew, 1994). Reduced glutathione (GSH) binds to the 'G' site of GSTp (and other GST isozymes) and plays an important role in detoxification of reactive oxygen species (ROS) and the maintenance of the cellular redox state (Sato *et al.*, 1989).

Among factors implicated in regulating JNK activity are ROS and altered redox potential (Adler *et al.*, 1995c; Gomez-del-Arco *et al.*, 1996; Cui *et al.*, 1997; Wilhelm *et al.*, 1997). ROS have also been associated with regulation of other signaling cascades, e.g. certain isozymes of protein kinase C (PKC; Konishi *et al.*, 1997) and mitogen-activated protein kinase (MAPK; Guyton *et al.*, 1996). The addition of exogenous oxidants or anti-oxidants has been found to influence the activation of MAPK/JNK (Lo *et al.*, 1996; Wilhelm *et al.*, 1997).

Changes in ROS can also directly influence transcriptional activity, as demonstrated for NF-κB (Wang *et al.*, 1995), Ref-1 (Xanthoudakis *et al.*, 1992) and c-Jun (Gomez-del-Arco *et al.*, 1996). Thus, evidence is accumulating that redox status can play an integral role in kinase-mediated stress response pathways. Our characterization of GSTp as an endogenous regulator of JNK activity provides a novel function for a protein with previously characterized catalytic and ligand-binding properties.

Results

A JNK inhibitor is present in cellular extracts of normally growing cells

JNK activities were measured in solid-phase kinase reactions using Jun as a substrate. Proteins prepared from UV-treated 3T3-4A mouse fibroblast cells exhibited marked phosphorylation of c-Jun. The kinetics of JNK activation in these cells increased within 1–5 min, reaching peak levels within 20 min and declining to basal levels within 3–4 h (Adler *et al.*, 1995b). This is a transient activation that is monitored as a multifold increase of kinase activity relative to the control (untreated) cells. Since the assays are performed under conditions that enable selective binding of JNK isozymes (Adler *et al.*,

1995b), we refer to the Jun-bound kinase as JNK and to the overall complex as Jun–JNK.

When cellular extracts prepared from unstressed cells were added to the pre-formed Jun–JNK complex before adding [γ -³²P]ATP, c-Jun phosphorylation was inhibited by 80% (Figure 1A, compare lanes 1 and 2). To elucidate further the nature of this inhibition, proteins derived from the non-stressed cells were absorbed by pre-incubation with increasing amounts of Jun–JNK. Subsequently, beads-bound c-Jun–JNK complexes were spun and the supernatant, which lacked Jun–JNK-bound proteins, was tested to determine its ability to alter the degree of JNK-mediated Jun phosphorylation. Increasing the amounts of Jun–JNK complexes used to absorb the putative inhibitor resulted in a dose-dependent decrease of inhibitory activity (Figure 1A, lanes 3–6). To determine whether this inhibitor activity was heat stable, the protein fraction that contained the inhibitor activity was heated to 95°C for 5 min. This treatment abolished the fraction's ability to block Jun phosphorylation by JNKs (not shown).

UV irradiation abolishes JNK inhibitor activities

The presence of JNK inhibitory activity in non-stressed cells prompted us to determine possible changes to the inhibitory activity after UV irradiation. As shown in Figure 1B, extracts prepared after UV treatment lacked inhibitory activity. The decrease in JNK inhibitor activity depended on the UV dose. Whereas a 10 J/m² dose (a 6 s exposure) caused a 40% reduction (as quantified using a phospho-imager), a 40 J/m² dose (a 25 s exposure) completely abolished the inhibition. In all instances, analysis was performed using the same amount (20 μg) of protein prepared 30 min after administering UV irradiation (Figure 1B). A correlation between JNK activation and the inactivation of its inhibitory activity was also noted in human melanoma cells, which require a 60 J/m² dose for JNK activation. In melanoma cells, 40 J/m² doses caused a 50% inhibition, whereas a 60 J/m² dose abolished JNK inhibitory activities (not shown).

Purification of the JNK inhibitor

To purify the putative inhibitor, extracts of normally growing cells were subjected to the six steps described below. After each step, the presence of the putative inhibitor (tested by measuring inhibition of c-Jun phosphorylation) was monitored in a solid-phase kinase assay using the _{his}Jun–JNK (1 μg) complex. Proteins from non-stressed cells were first precipitated in ammonium sulfate (50%). After dialysis against kinase buffer, the non-precipitated proteins were concentrated on membranes with various molecular weight cut-offs. Inhibitor activity was found in the flow-through of both 100 and 30 kDa cut-off filters; it was retained on the 10 kDa membrane, indicating that the inhibitor is between 10 and 30 kDa in molecular weight (Figure 1C). The 10–30 kDa fractions were loaded on a gel filtration column; several fractions within the 20–25 kDa range (based on calibration of the gel filtration column with known molecular weight standards) retained inhibitory activity (Figure 1D). Active fractions were pooled and fractionated on a MonoQ anion exchange column; a 70 mM NaCl eluate was found to contain most of the inhibitor activity (Figure 1E). Active MonoQ fractions were then loaded onto a phenyl-




Fig. 1. (A) UV-mediated JNK activation is inhibited by proteins obtained from cells maintained under normal growth conditions. Phosphorylation of Jun by JNK (lane 1) is inhibited when the pre-formed complex (Jun–JNK; total 1 μg at 1:1 ratio) is incubated with proteins (10 μg) from untreated cells before adding the [γ -³²P]ATP (lane 2). Pre-incubation of proteins from non-stressed cells with increasing concentrations of Jun–JNK complex resulted in a dose-dependent decrease of inhibitory activity (lanes 3–6 represent increasing amounts of the pre-formed complex 0.25, 0.5, 1 and 2 μg, respectively). (B) UV irradiation abolishes inhibitor activities. Pre-formed Jun–JNK complex was incubated with whole-cell extract prepared 30 min after sham or UV-C irradiation at the doses indicated prior to addition of [γ -³²P]ATP. (C) Purification of JNK inhibitor on molecular weight cut-off columns. Whole-cell extracts from non-stressed cells were subjected to an ammonium sulfate precipitation and the supernatants were fractionated on the indicated membranes with a cut-off mol. wt of 100, 30 or 10 kDa before being added to pre-formed Jun–JNK complex, followed by addition of [γ -³²P]ATP (the basal activity in UV-treated cells is shown in lane JNK). Shown is an autoradiograph of the phosphorylated c-Jun. (D) Gel filtration of 3–30 kDa fractions. Fractions <30 kDa were concentrated and loaded (50 μl) onto a Superdex 75 column. From each fraction, 20 μl were added to the pre-formed Jun–JNK complex to measure the ability to inhibit JNK activity (the first lane points to JNK activity with UV-treated extract before this inhibition; numbers represent the respective fractions). (E) Fractionation of JNK inhibitor via an anion exchange column. Fractions 18–24 obtained by Superdex 75 separation were pooled and applied to a MonoQ column. The fractions were assayed for inhibitor activities as shown in the inset. (F) Identifying the inhibitor component on the basis of its binding to Jun–JNK complex. his-Jun or his-Jun–JNK complex was incubated with the phenyl-Sepharose eluate (A) or with a purified form of GSTp (B). Beads-bound material was washed and separated on SDS–PAGE. Silver staining of bound proteins is shown in the left panel, whereas the immunoblot of a duplicate gel with antibodies to GSTp is shown on the right. The arrow points to the position of GSTp.

Table I. Purification of JNK inhibitor

| Purification step | Protein (mg) | Total activity (U) ^a | Specific activity ^b (_{his} Jun-JNK) |
|------------------------|--------------|---------------------------------|--|
| Total lysate | 80 | 10 160 | 0.127 |
| Ammonium sulfate (50%) | 32 | 8128 | 0.254 |
| Membrane cut-off | 6.7 | 6096 | 0.91 |
| MonoQ | 2.1 | 5425 | 2.69 |
| Phenyl-Sepharose | 1.4 | 4820 | 3.46 |
| _{his} Jun-JNK | 0.063 | 4511 | 70.70 |

All purification steps were performed using the _{his}Jun-JNK complex (on nickel beads) as a substrate to measure the degree of JNK activity/inhibition (see Materials and methods for details).

^aTotal activity is defined in units: 1 U is the ability to decrease JNK activity by 50%.

^bSpecific activity is calculated as inhibitor activity of 1 µg of protein.

Sepharose column in the presence of 0.6 M ammonium sulfate. Using decreased concentrations of ammonium sulfate, the inhibitor activity was found in 0.2 M eluate fractions (not shown). The active fractions from the phenyl-Sepharose column were incubated in a _{his}Jun-JNK column. Analysis of the _{his}Jun-JNK-bound material revealed a single band on silver-stained gels (Figure 1F). Microsequencing analysis of the corresponding band obtained after a large-scale purification (Table I) revealed a 12 amino acid peptide (Pro-Pro-Tyr-Thr-Val-Val-Tyr-Phe-Pro-Val-Arg-Gly) that exhibited 100% homology to the human form of GSTp. To confirm this protein's identity, a purified form of GSTp was separated parallel to the _{his}Jun-JNK-bound material and subjected to Western blot analysis using polyclonal antibodies to GST. Both migration and immunoreactivity of purified GSTp were identical to those of the _{his}Jun-JNK-bound material (Figure 1F).

GST inhibition of JNK activity does not alter phosphorylation of Jun or JNK

To determine whether GSTp affected the degree of JNK or Jun phosphorylation, increasing concentrations of GSTp (purified form) were added to pre-formed Jun-JNK complex which contained the phosphorylated form of JNK obtained from UV-treated cells. GSTp decreased JNK phosphorylation of c-Jun in a dose-dependent manner (within a range of 0.05–1 µg) (Figure 2A), but it did not decrease the number of phosphate groups on JNK, as revealed by immunoblots with phospho-JNK antibodies (Figure 2B). Dual activity protein phosphatase (Ishibashi *et al.*, 1992) was used as a positive control in these reactions (Figure 2C).

To determine whether GSTp affects the number of phospho groups on c-Jun, cellular extracts from normally growing cells were incubated with pre-formed Jun-JNK complex for the indicated periods of time before or after the phosphorylation step with [γ -³²P]ATP. The extent of c-Jun phosphorylation was not altered when cell extracts were added after the phosphorylation reaction (Figure 2D). This observation suggested that GSTp did not reduce the number of phospho groups on c-Jun.

Effect of different GST isoforms on JNK activity

Incubation of whole-cell extract prepared from non-stressed mouse fibroblasts with the _{his}Jun-JNK complex




Fig. 2. (A) GSTp as a JNK inhibitor. GSTp was added at the indicated concentrations (micrograms) to the pre-formed Jun-JNK complex and the level of Jun phosphorylation was measured by means of autoradiography. (B) GSTp does not affect JNK phosphorylation. To measure the effect of GSTp on JNK phosphorylation, JNK was immunoprecipitated from UV-treated cells and incubated with GSTp (at the microgram concentrations indicated) followed by Western blot with antibodies to phospho-JNK (upper panel). The lower panel depicts the level of JNK on the same blot after re-probing with antibodies to JNK. (C) As a positive control, JNK from UV-treated cells was incubated with no protein (lane C), dual specificity phosphatase (lane PP; Ishibashi *et al.*, 1992) or GSTp (lane GSTp) before carrying out immunoblot analysis with antibodies to phospho-JNK. Quantification via densitometer scanning revealed 35% inhibition of JNK phosphorylation by PP, whereas GSTp did not elicit such inhibition. (D) GSTp does not alter c-Jun phospho groups. The autoradiograph depicts the c-Jun phosphorylation level after incubation with the inhibitor for the indicated time periods (minutes) before or after phosphorylation by JNK.

identified GSTp as the associated protein. A marked decrease in this association was found in proteins prepared after UV irradiation (Figure 3A). In addition to GSTp, isoforms of the GST α and GST μ families were also capable of associating with the Jun-JNK complex *in vitro* (Figure 3A). GSTp exhibited greater JNK inhibitory activity than did GST μ , which was more potent than GST α (Figure 3B). Bacterially expressed GST (GST-2T) also mediated JNK inhibition (Figure 3B). This excludes the

possibility that the inhibitor activity was dependent on any putative GST-associated cellular component.

Dose-dependent inhibition of JNK by GSTp

GSTp inhibition of JNK activities was determined using six concentrations of purified GSTp added to either full-length or N-terminal forms of c-Jun. The addition of purified GSTp to JNK complexed with GST–Jun^{5–89} or hisJun^{full-length} led to a concentration-dependent inhibition of c-Jun phosphorylation (Figure 3C). The greater inhibitory capacity of GSTp observed for GST–Jun^{5–89} compared with hisJun^{full-length}-based JNK complexes can be attributed to the different conformations of the recombinant substrates (N-terminal GST–Jun^{5–89} versus hisJun^{full-length}). A higher degree of inhibition of the GST–Jun^{5–89} fusion protein is not likely to occur as a result of GST–GST interactions, which would require other experimental conditions and result in different kinetics. Nevertheless, using either GST–Jun^{5–89} or hisJun^{full-length} as substrates, the degree of GSTp inhibitory activity appeared to be linear within the range of 25–180 ng of GSTp (per microgram of pre-formed Jun–JNK complex; Figure 3C). GSTp also inhibited JNK activity when added to a soluble form of hisJun–JNK complex (not shown).

Specificity of GST inhibition

To determine the specificity of the JNK inhibitor, purified GSTp was incubated with substrates for other protein kinases. Incubation with PKA, PKC, casein kinase II (CKII) or MAPK revealed 6, 3, 36 and 9% inhibition, respectively. Under the same conditions, GST elicited 86% inhibition of JNK (Figure 3D). In all cases, the activities of the various kinases tested were normalized (c.p.m./μg protein and an equal ratio between the respective substrate and GSTp) to ensure an equal degree of substrate phosphorylation.

GSTp immunodepletion attenuates JNK inhibition

We further elucidated the ability of GSTp to inhibit JNK activity by modulating its levels or activity, or both, *in vitro* and *in vivo*. The *in vitro* kinase assay in which proteins from non-stressed cells were added to the pre-formed Jun–JNK complex revealed a dose-dependent increase in the degree of JNK inhibition (of up to 80%). Immunodepletion of GST from whole-cell extract decreased the degree of JNK inhibition from 80 to 45% (Figure 3E, lane GST-1). Repeated immunodepletion by GST antibodies further decreased the degree of JNK inhibition to 20% (Figure 3E, lane GST-2). Control immunodepletion reactions with normal rabbit serum (NRS) and protein A/G beads did not affect the degree of JNK inhibition (Figure 3E, lanes NRS-1 and NRS-2, respectively).

Specific GSTp inhibitors and a GSTp-derived peptide efficiently alter JNK inhibition

As an independent approach to inhibit GSTp, we used specific inhibitors which were shown to inactivate GSTp enzymatic activity *in vitro* and *in vivo* (Flatgaard *et al.*, 1993). Adding TER-117, a specific *in vitro* GSTp inhibitor, to a Jun–JNK complex at the same time as GSTp prevented GSTp inhibition of JNK activity. Similarly, TER-293, another GSH peptidomimetic related to TER-117, also

blocked GSTp inhibition of JNK activity (Figure 3F). TER-317, structurally similar to TER-117, yet void of GSTp inhibitory properties, failed to affect GSTp inhibition of JNK. Similarly, TER-199, an inactive prodrug of TER-117 (Flatgaard *et al.*, 1993), was also without effect *in vitro*.

Molecular dynamic calculations on the energy-minimized X-ray crystal structure of GST were performed to identify the most flexible regions of the protein, i.e. those that were computed to have the highest degree of fluctuations. These regions are most likely to undergo significant conformational changes during catalysis or ligand binding. Two such flexible domains, corresponding respectively to amino acids 36–50 and 194–201, which can participate in the GST–Jun–JNK association and inhibition, were tested.

Peptides corresponding to each of these domains were synthesized, and their ability to alter GSTp inhibitor activity was determined *in vitro*. As shown, the GSTp-derived peptide from amino acids 194–201 (at the C-terminal domain designated P1) abrogated GSTp inhibition of JNK activity (Figure 3F). Conversely, neither P2 (aa 36–50; Figure 3F) nor three unrelated, non-GST-based peptides of varying lengths (8–24 amino acids; not shown) were capable of affecting GSTp inhibition of JNK. Treatment of cells with TER-199 (the prodrug form of TER-117 which is converted to active form *in vivo* and has been shown to elicit effective inhibition of GST activity *in vivo*; Flatgaard *et al.*, 1993) led to a 2-fold increase in basal JNK activity in non-stressed cells. UV irradiation also increased (~2.5-fold) JNK activation in TER-199-treated cells (Figure 3G). That TER-199 also affected JNK activities after UV irradiation suggests that the dose of UV irradiation was insufficient to mediate complete inactivation of GST as a JNK inhibitor; thus, the presence of a specific GSTp inhibitor led to an additive effect on overall JNK activities in these cells. The effect of these GSH peptidomimetic agents is likely to be mediated either by altered conformation of the GSTp molecule or by a competition for the binding site, as non-active (bacterially produced) GST was also capable of eliciting JNK inhibition (Figure 3B).


Changes in ROS affect GST oligomerization and its association with, and inhibition of, JNK

We next assessed whether modulation of the cellular redox potential would affect JNK inhibition by GSTp. We monitored the GST–JNK complex *in vivo* by means of immunoprecipitations followed by immunoblot analysis. Exposure of mouse fibroblast cells to either UV or H₂O₂ reduced the amount of the JNK–GSTp complex and increased JNK activity, whereas pre-treating cells with the free radical scavengers *N*-acetylcysteine (NAC) or the ethyl-ester of glutathione (eeGSH) prevented JNK dissociation from GSTp and maintained GSTp inhibitory activity (Figure 4A I–III). These observations indicate that GST–JNK–Jun association is inversely correlated with JNK activity. To elucidate further changes to GST that take place upon altered redox conditions, we measured the possible formation of GST–GST dimers/multimers through disulfide bonds on neighboring cysteines. Monitoring the migration of GSTp on non-reducing SDS–PAGE revealed decreased monomer/dimer forms of GSTp in UV- or


H₂O₂-treated cells (Figure 4A IV). Instead, a high molecular weight band corresponding to a multimer form of GSTp was detected (Figure 4A IV, upper arrow in panel a). Pre-treatment with NAC or eeGSH maintained the lower molecular weight forms of GSTp (Figure 4A

IV, the single arrow in panel a points to the dimer and panel b shows the monomer). H₂O₂ treatment is known to cause the formation of intra- or intersubunit disulfide bonds between cysteines at amino acids 47 and 101 of GST (Shen *et al.*, 1993), resulting in a multimerization


A




B




C




D




E



F



G



of GST subunits which no longer accommodates JNK molecules.

Monomer form of GSTp mediates JNK inhibition

The observation of changes in GSTp from monomer/dimer to multimer forms upon exposure to increased ROS prompted us to determine which of the two prevalent GSTp forms, monomer or dimer, exerts inhibitory activity on JNK. A gel filtration column was used to separate the monomer from the dimer form of GSTp (Figure 4B). A purified form of either monomer or dimer was added to a solid-phase kinase reaction in which pre-formed $_{\text{his}}\text{Jun}^{\text{full-length}}$ -JNK complex served as a substrate. As shown, only the monomeric form of GSTp was capable of mediating efficient inhibition of JNK phosphorylation of c-Jun (Figure 4B).

GSTp expression is inversely correlated with JNK activities

In a complementary approach to study GSTp effects on JNK activity, increasing amounts of GSTp cDNA were transiently transfected into mouse fibroblasts. As shown in Figure 5A, GSTp transfection resulted in a dose-dependent decrease in JNK activation by UV. While transfection of 0.5 μg of GSTp caused a 20% decrease, 2 μg led to a 40% and 10 μg to a 50% decrease in JNK activation by UV irradiation. These observations suggest that it is possible to alter the degree of JNK activation by GSTp transfection, albeit within the relatively narrow range of 0.5–2 μg .

Effect of GSTp on phosphorylation of JNK kinase MKK4/JNKK/SEK1

To elucidate further the mechanism underlying GSTp inhibition of JNK activity, we monitored possible changes at the level of MKK4 phosphorylation. *In vitro* phosphorylation of JNK by MKK4 was not inhibited by GSTp (not shown). In non-stressed 3T3 fibroblasts, there is a basal level of MKK4 phosphorylation, detected by MKK4 phospho-antibodies (Figure 5BI). UV irradiation

efficiently increased MKK4 phosphorylation. Conversely, forced expression of GSTp reduced the level of MKK4 phosphorylation in a dose-dependent manner (Figure 5BI). Under the same conditions, GSTp did not alter phosphorylation of ERK1 (Figure 5BII). Treatment of GSTp-transfected cells with TER-199, an inhibitor of GSTp, restored the level of MKK4 phosphorylation (Figure 5BI). The observation that under physiological levels of GSTp expression MKK4 was phosphorylated, although JNK activity was inhibited by GSTp, suggests that GSTp does not affect JNK kinase under non-stressed conditions. GSTp inhibition of JNK could be attributed to the nature of the complex between the two proteins. The finding that overexpression of GSTp reduces MKK4 phosphorylation suggests that an excess of GSTp can also affect the MKK4–JNK module.

Further evaluation of the possible interplay between GSTp and upstream JNK kinases was carried out in cells that had been transfected with ΔMEKK1 , an MKK4 upstream kinase. Forced expression of ΔMEKK1 led to constitutive phosphorylation of MKK4, even when GSTp was co-transfected. Interestingly, although GSTp was not able to reduce MKK4 phosphorylation in the presence of ΔMEKK1 , it efficiently blocked JNK-mediated phosphorylation of c-Jun (Figure 5BIII). These findings point to a selective effect of GSTp on JNK. GSTp inhibition of MKK4 depends on the level of GSTp expression and the signal elicited by the upstream kinase (as shown here for MEKK1). GSTp inhibition of JNK is due primarily to their association, which is released upon the conversion of GSTp from a monomer to a dimer form.

GSTp expression results in higher ubiquitination of c-Jun

Since JNK efficiently targets the ubiquitination of its non-phosphorylated associated proteins c-Jun, ATF2 and p53, we determined the possible effects of GSTp on ubiquitination of JNK substrates in this reaction. Under non-stress growth conditions, c-Jun exhibits a short half-life, which is prolonged upon phosphorylation by JNK (Fuchs *et al.*,

Fig. 3. (A) GSTp associates with Jun–JNK *in vitro*. The pre-formed Jun–JNK complex was incubated with whole-cell extract (10 μg) prepared before (WCE cont) or after (WCE UV) UV irradiation or with purified forms of GST isozymes (Ciaccio *et al.*, 1991), as indicated. Following extensive washes, complex-bound and non-bound (absorbed on Jun–JNK; sup) material was analyzed on immunoblots with polyclonal antibodies that recognize multiple forms of GST (Ramgopal and Tew, 1991). Arrows point to the identified forms of GSTp. (B) Effect of different GST isozymes on JNK activity. Pre-formed $_{\text{his}}\text{Jun}$ –JNK was incubated with the indicated forms of GST (a, α ; m, μ ; p, π ; 2T, bacterially produced form of GST) purified as described in Materials and methods before the addition of $[\gamma\text{-}^{32}\text{P}]\text{ATP}$. Autoradiography demonstrates the degree of c-Jun phosphorylation in the presence of the various GSTs. Quantification of phosphorylation is shown in the graph. (C) Dose-dependent effect of GSTp on JNK kinase activity. The purified form of GSTp was added at the indicated concentrations (per μg of the relevant Jun substrate) to GST–Jun^{5–89}–JNK (2 μg /reaction) or $_{\text{his}}\text{c-Jun}$ –JNK (7 μg /reaction) prior to the addition of $[\gamma\text{-}^{32}\text{P}]\text{ATP}$. The degree of inhibitor activity was calculated based on the ratio between Jun phosphorylation activities in the absence and in the presence of GSTp. The insets show representative autoradiographs of the respective reactions (– reflects the degree of phosphorylation without inhibitor added). Quantification of three independent experiments is shown in the graph. (D) Specificity of GST as a JNK inhibitor. The ability of the purified form of GSTp to inhibit phosphorylation by Src, CKII, MAPK or PKA was tested. Bead-bound substrates were incubated with the respective kinases in the presence of GSTp before $[\gamma\text{-}^{32}\text{P}]\text{ATP}$ was added to initiate the kinase reaction. After phosphorylation, the beads were washed and phosphorylation was quantified. The values shown represent average results of three different reactions. (E) Immunodepletion of GSTp increases JNK activity. $_{\text{his}}\text{Jun}$ –JNK complex (containing JNK purified from UV-treated cells) was incubated with whole-cell extract either from non-stressed cells (10 μg) or subjected to one or two sequential immunodepletions of GSTp (GST-1 or GST-2, respectively) using antibodies to GSTp before addition of $[\gamma\text{-}^{32}\text{P}]\text{ATP}$. The control reactions with whole-cell extracts treated under the same conditions with NRS and protein A/G beads are also shown. The inhibitory activity depicted was calculated based on values of c-Jun phosphorylation. The lower panel shows an immunoblot indicating the GSTp level after each of the immunodepletion reactions. (F) GSTp inhibitor increases JNK activities *in vitro*. $_{\text{his}}\text{Jun}$ –JNK or GST–Jun–JNK was incubated with selective GST inhibitors in the presence of the purified form of GSTp, prior to the addition of $[\gamma\text{-}^{32}\text{P}]\text{ATP}$. The degree of Jun phosphorylation in the presence of each of the inhibitors is shown. Peptides tested in parallel represent two flexible domains from GSTp, corresponding to the N- (P2) and the C-terminal regions (P1). (G) GSTp inhibitor increases JNK activities *in vivo*. Mouse fibroblasts were treated with TER-199, a specific inhibitor of GSTp, for 2 h followed by either sham or UV treatment (50 J/m^2). Whole-cell extract proteins were prepared after 45 min and assayed for JNK activity by means of $_{\text{his}}\text{Jun}$ phosphorylation. The inset shows an autoradiograph of $_{\text{his}}\text{Jun}$ phosphorylation (– without; + in the presence of TER-199), which was quantified as shown in the graph.




Fig. 4. (A) Free radical scavengers prevent UV- or H₂O₂-mediated GSTp–JNK complex dissociation and maintain low basal JNK activity. Immunoprecipitations using antibodies to JNK (clone 333; PharMingen) were performed on proteins prepared before (–) and after UV irradiation (60 J/m²) or H₂O₂ treatment (10 μM added in phosphate-buffered saline (PBS) which was replaced after 5 min with medium for another 30 min) of mouse fibroblasts. Cells were pre-treated with NAC (10 mM) or eeGSH (1 mM added to the medium) as indicated. JNK immunoprecipitates (from 3 mg of whole-cell extract) were examined by immunoblot analysis using antibodies to GSTp (I). (II) The same blot probed with JNK antibody (clone 333; PharMingen). (III) The level of JNK activity in immunoprecipitates monitored by means of hisJun phosphorylation. (IV) A GSTp immunoblot reflecting the changes in the migration of GSTp (immunoprecipitated using antibodies to GSTp from cells treated as indicated in the figure) under non-reducing (without β-mercaptoethanol in sample buffer) versus reducing SDS–PAGE. The blot shown in (a) was subjected to a 1 min exposure, whereas the blot in (b) was exposed for 1 h. Migration of purified GSTp is shown on the right lane marked GSTp. The positions of the dimer (lower arrow), trimer (double arrow) and large complex (possible tetramer or higher) seen under non-reducing SDS–PAGE conditions (upper arrow) are indicated. Molecular weight markers are indicated on the left panel. (B) The monomer form of GST mediates JNK inhibition. GSTp was purified by means of gel filtration (Superdex 75) to dissociate monomer from dimer forms of GSTp (a). By adding monomer or dimer GSTp forms to the hisJun–JNK complex prior to the addition of [³²P]ATP, the monomer form of GSTp was identified as the actual inhibitor of JNK activity (b).

1996, 1997; Musti *et al.*, 1997). Transfection of GSTp cDNA into 3T3 mouse fibroblasts increased the level of c-Jun ubiquitination *in vivo* (Figure 5C). Since the level of ubiquitinated Jun is inversely correlated with its degree of phosphorylation (Fuchs *et al.*, 1996; Musti *et al.*, 1997), the increase in ubiquitinated c-Jun is an expected result of the GSTp inhibition of basal JNK activity, which reduces the number of c-Jun molecules that undergo phosphorylation. The noticeable increase in ubiquitinated c-Jun molecules provides an example of the physiological significance of JNK inhibition under normal growth conditions.


JNK activity in cells of GSTp null mice

We established embryo fibroblast cells from *GSTP1/P2*(^{–/–}) mice [MEF^{GSTP1/P2}(^{–/–})] in order to assess JNK activity in a GSTp-free environment. These cells do not express GSTp as monitored by either RT–PCR (data not shown) or Western blot analysis (Figure 6A). Stress in the form of UV irradiation, sorbitol or anisomycin markedly increased JNK activity in both MEF^{wt} and MEF^{GSTP1/P2}(^{–/–}) cells; of these treatments, UV elicited the greater degree of JNK activation (Figure 6B). Interestingly, a higher basal level of JNK activity was found in the MEF^{GSTP1/P2}(^{–/–}) than in the MEF^{wt} (Figure 6B and C). This high activity could be diminished in a dose-dependent manner upon transfection of GSTp cDNA (Figure 6C). Lesser amounts of GSTp cDNA were required to mediate 80–100% inhibition of UV-mediated JNK activation in the MEF^{GSTP1/P2}(^{–/–}) cells, as compared with the MEF^{wt} cells (not shown). In both types of MEFs, transfection of GSTp caused 80–100% inhibition of UV-mediated JNK activation, whereas in mouse fibroblast 3T3 cells, GSTp inhibition reached only 50%. Such differences may be attributed to the overall amount and form (monomer and dimer) of GSTp and/or the levels of other radical scavenger enzymes expressed in each of the cell lines.

To elucidate further the possible mechanism by which GSTp elicits its inhibition of JNK, we monitored levels of JNK phosphorylation before and after stress using JNK phospho-antibodies. In MEF^{GSTP1/P2}(^{–/–}) cells, a higher basal level of JNK phosphorylation was seen (Figure 6D I), when compared with the MEF^{wt} cells. Forced expression of GSTp in these cells revealed an efficient reduction in the number of phospho groups on JNK, prior to, as well as after, UV irradiation (Figure 6D II). Forced expression of a truncated MEKK1 form (ΔMEKK1), which elicits constitutively high levels of JNK activity, partially restored the levels of JNK phosphorylation (Figure 6D II versus I). These observations are in line with the effect of GSTp on phosphorylation of MKK4, and suggest that GSTp elicits inhibition of JNK *in vivo* in spite of JNK phosphorylation by upstream kinases.

Forced expression of GSTp decreases transactivation of c-Jun in GST null cells

MEFs derived from GSTp null mice [MEF^{GSTP1/P2}(^{–/–})] as well as from GSTp wild-type mice were co-transfected with GSTp and Jun promoter (5× *jun2* target sequence)-driven luciferase constructs. MEF^{GSTP1/P2}(^{–/–}) cells exhibited high basal levels of transcriptional activity mediated by the *Jun2* promoter sequence (Figure 6E, control transfection with empty construct, pcDNA3). This activity



reflected the high basal JNK activity found in these cells (Figure 6B and C). Forced expression of GSTp efficiently reduced high basal Jun-driven transcriptional activity in the MEF^{GSTP1/P2(-/-)} cells (Figure 6E), implicating the capacity of GST to reduce JNK phosphorylation and activity (Figures 6C and D). As the amino acid residues which are important for GST-GST dimerization and enzymatic activity have been identified previously (Shen *et al.*, 1993), we mutated GSTp at the respective sites and evaluated their ability to elicit JNK inhibition. When tested in MEF^{GSTP1/P2(-/-)} cells, GSTp whose cysteines were mutated at amino acids 47 and 101 (which are required for GSTp dimerization) and GST whose tyrosine was mutated at amino acid 7 (which abrogates catalytic proton transfer activity) were as potent as wild-type GSTp in inhibiting high basal JNK activity (not shown) and Jun-mediated transactivation (Figure 6E). The lower basal levels of Jun-mediated transactivation found in MEF^{GST^{wt}} were also reduced by each of these constructs. The capacity of GSTp to reduce transcriptional activity mediated by the Jun promoter could be due to reduced phosphorylation of Jun and/or ATF2 by JNK, increased ubiquitination and degradation of c-Jun, or both. These observations further support the hypothesis that GSTp inhibition of JNK does not require GST enzymatic activity and is mediated by its monomeric form. Forced expression of peptide derived from the GSTp C-terminal domain

Fig. 5. (A) Transfection of GSTp into mouse fibroblasts reduces JNK activities. The figure shows the extent of JNK activation by UV in cells transfected with GSTp cDNA as compared with mock-transfected controls (100%). GSTp cDNA was co-transfected with the β -gal (0.5 μ g) construct into 3T3 cells (via lipofection; lipofectamine, Gibco-BRL). Control empty vector was added to reach a constant amount of transfected DNA (15 μ g). At 48 h post-transfection, cells were subjected to UV treatment (50 J/m²) and whole-cell extract proteins were prepared 45 min later. The inset depicts an immunoblot performed on the same protein extracts, revealing the respective increase in expression of GSTp (- reflects mock-transfected, lanes 2-4 represent 0.5, 2 and 10 μ g of GSTp cDNA, respectively). Whole-cell extracts (normalized to transfection efficiency based on β -galactosidase values) were used for a kinase reaction using the his₆-Jun substrate (2 μ g). **(B) (I)** Forced expression of GSTp reduces MKK4 phosphorylation. Left panel: mouse 3T3 fibroblasts analyzed for MKK4 phosphorylation using MKK4 phospho-antibodies. Forced expression of GSTp was achieved via lipofection of GSTp cDNA at the indicated concentrations (micrograms). TER-199 (at the micromolar concentration indicated in the figure) was added 22 h after transfection (2 h before protein preparations). The lower panel depicts the overall expression level of MKK4 (using non-phospho-antibodies). Right panel: level of MKK4 phosphorylation upon GSTp expression (0.5 μ g) or UV irradiation (30 J/m²). PC represents positive control of phosphorylated MKK4. **(II)** Forced expression of GSTp does not affect ERK1 phosphorylation. Proteins prepared as indicated in (I) were subjected to immunoblot analysis using ERK1 phospho-antibodies. **(III)** Forced expression of Δ MEKK1 blocks GSTp effect on MKK4 but not inhibition of JNK phosphorylation of c-Jun. Mouse 3T3 fibroblasts were co-transfected with Δ MEKK1 cDNA (1 μ g) and empty vector (pcDNA3 to a total of 2.5 μ g) or GSTp cDNA at the indicated concentrations. Proteins prepared 24 h after transfection were subjected to JNK immunokinase reaction (upper panel) using his₆-Jun as a substrate, or to immunoblot analysis using antibodies to the phosphorylated form of MKK4 (middle panel). Analysis using non-phospho-MKK4 antibodies is shown in the lower panel. **(C)** Ubiquitination of c-Jun increases upon GSTp expression. Mouse 3T3 cells were co-transfected with his₆-Jun, GSTp and Ub-HA as indicated in the figure. his₆-Jun was purified on Ni beads as described (Terier *et al.*, 1994) and the degree of ubiquitinated Jun was assessed by immunoblots with antibodies to hemagglutinin (HA). The area reflecting the polyubiquitination is marked on the right panel. The expression of c-Jun is shown on the lower panel.

(which blocked GSTp inhibition *in vitro*) increased basal JNK kinase and c-Jun transcriptional activities, respectively (Z.Yin, V.Adler and Z.Ronai, unpublished observations). By monitoring c-Jun transcriptional activities, these experiments point to the biological significance of GSTp as an inhibitor of JNK.

Discussion

In the present study, we identify and characterize GSTp as a JNK inhibitor. Our data point to an additional cellular mechanism that is involved in the regulation of JNK activity before and after stress. In non-stressed cells, the basal levels of JNK phosphorylation are low, in spite of constitutive stimulation by growth factors and endogenously formed ROS. Maintaining a low basal JNK activity is believed to affect the half-life of JNK substrates,

including c-Jun, ATF2 and p53 (Fuchs *et al.*, 1996, 1998c; Musti *et al.*, 1997), and to play a key role in maintenance of controlled cell growth. Indeed, forced expression of GSTp in 3T3 fibroblasts increased the degree of Jun ubiquitination and decreased Jun-mediated transactivation.

GSTp inhibition of JNK is found primarily in normally growing non-stressed cells. Stress, as shown in the case of UV irradiation, decreases this inhibition in a dose-dependent manner. Important to our understanding of GSTp's ability to elicit association with JNK and inhibition of this kinase is the finding that the inhibitory activities are confined to the monomeric form of GSTp. When first identified, the Jun-JNK-associated protein had a mol. wt of 23 kDa, the characteristic size for the monomeric form of GSTp. Immunoprecipitation reactions always identified the monomeric form of GSTp as the Jun-JNK-associated protein. Similarly, the ability of the monomer (but not the dimer) to elicit JNK inhibitory activity *in vitro* supports the role of the monomer GSTp in JNK inhibition *in vivo*. UV irradiation reduces GSTp-JNK association, probably as a result of the formation of GST-GST dimers and multimers; because of disulfide bond-induced steric constraints, dimers/multimers cannot accommodate the Jun-JNK complex. ROS scavengers, such as NAC or eeGSH, inhibit the formation of GSTp multimers, prevent GSTp dissociation from Jun-JNK and maintain the low basal activity of JNK as a kinase. The switch from a monomer to a dimer/multimer form is likely to provide the underlying mechanism for GSTp's ability to sense and transmit changes in redox potential as a regulator of JNK signaling. Thus, based on its cellular conformation, GSTp dictates the association and inhibition of JNK. Importantly, while our studies demonstrate the effect of ROS elicited by UV and H₂O₂ on GSTp dimerization, certain type of stress




Fig. 6. (A) Expression of GST in GSTp null mice. Expression of GST in MEF^{GSTP1/P2^{-/-}} (lanes A) and MEF^{GSTwt} (lanes B) was determined using specific antibodies to GSTP1-1 (Henderson *et al.*, 1998). (B) Basal and induced JNK activity in GSTp null mice. Proteins prepared from either GSTP1/P2^{-/-} (null) or GST wt MEF (MEF wt) before (C) or after exposure to UV irradiation (UV; 40 J/m²), sorbitol (S; 0.6 M) or anisomycin (A; 10 μg/ml) were subjected to a solid-phase kinase reaction using c-Jun as substrate. Levels of phosphorylated Jun are shown. (C) Basal JNK activity in MEF cells of GSTp null mice can be reduced by forced expression of GSTp. JNK activity was monitored in MEF cells derived from wild-type (MEF wt) or GST null (GST P1/P2^{-/-}) mice transfected via lipofection with either empty vector (4 μg; first lane on left) or GSTp cDNA (0.5, 2 or 4 μg). JNK activity was determined 24 h after transfection into each of the MEF lines. (D) JNK phosphorylation is affected by GST expression. JNK phosphorylation was monitored in MEF^{GSTwt} and MEF^{GSTP1/P2^{-/-}} cells before (C) and after UV irradiation (UV) as well as after transfection with the constitutively active form of MEKK1 (ΔMEKK1). The effect of the empty vector (I) or GSTp cDNA (II-IV) on overall JNK phosphorylation (I and II) is shown. The levels of JNK (III) and GSTp expression (IV) are also shown. JNK phosphorylation was determined 30 min after mock or UV irradiation (which was administered 24 h after transfection) using antibodies to phosphorylated residues 183 and 185 on JNK (P-JNK). The control reactions using antibodies that recognize non-phosphorylated forms are shown in (III). (E) Transactivation of the Jun-driven luciferase construct in GSTP1/P2^{-/-} cells. Forced expression of GSTp or the indicated mutant forms in MEFs of GSTpwt or MEFs of GSTP1/P2^{-/-} mice was used to determine the effect on Jun transcriptional activities. Each of the GSTp constructs was co-transfected with the *Jun2*-luciferase vector (consisting of five repeats of the *Jun2* sequence TGACATCA). The amount of luciferase activity was quantitated 24 h after transfection. Values shown were normalized with respect to transfection efficiency.




Fig. 7. Model of GST inhibition of JNK signaling. Based on our findings, the following model is proposed: under non-stressed conditions, GSTp can be free or part of a complex with Jun–JNK. Upon stress, in which ROS are formed, GSTp forms dimers and larger aggregates which cannot accommodate Jun–JNK, thus enabling JNK phosphorylation of c-Jun, which as a result is a stable and active transcription factor.

(i.e. certain cytokines) are not expected to cause the same degree of ROS, and yet are strong inducers of JNK. The latter suggests that dissociation of GSTp from JNK may also occur due to changes other than disulfide bond-based dimerization. The effect of post-translational modifications of GSTp on its association with JNK is being investigated.

The finding that GSTp transfection reduces basal and UV-inducible levels of JNK phosphorylation *in vivo* provides further insight into the mechanism by which GSTp elicits inhibition of JNK signaling. It is possible to rescue GSTp-mediated inhibition of JNK phosphorylation by transfecting a constitutively active form of MEKK1. Independently, under physiological conditions, GSTp inhibition of JNK takes place in spite of activated MKK4, possibly through direct inhibition of JNK-mediated Jun phosphorylation, supporting our original *in vitro* observation where GSTp inhibited c-Jun phosphorylation by an activated form of JNK. The ability of GSTp to block JNK activity in spite of MKK4 and JNK phosphorylation by MEKK1 suggests that the nature of the GST–JNK association disables JNK activity as a kinase. That endogenous GSTp does not affect the MKK4 phosphorylation level in non-stressed cells provides further evidence for selective inhibition of the Jun–JNK module by physiological levels of GSTp. That high expression of GSTp also reduces MKK4 phosphorylation could be attributed to possible inclusion of MKK4 with the GST–JNK module. Our data suggest that the interplay between upstream MKK4 signaling and the amount of monomer GSTp may determine the effect of GSTp on MKK4. Interplay between MEKK1–MKK4 and MKK4–JNK modules (Xia *et al.*, 1998) and their existence in complex with scaffold protein, which determines the specificity of these signaling cascades (Whitmarsh *et al.*, 1998), may explain the nature of changes elicited with different expression levels of GSTp. The specificity of GSTp-mediated JNK inhibition was demonstrated via comparison of different protein kinases in a solid-phase reaction, as well as *in vivo*,

as forced expression of GSTp did not affect ERK1/2 phosphorylation. Our data also imply that MEKK1 is not a target for GSTp inhibition; other MKK4 upstream kinase(s), which include ASK1, TAKs, MLKs and GCKs, are currently being examined. Of interest is the observation that UV doses (10 J/m^2) which decrease GST inhibitory activities are insufficient for JNK activation, which requires doses $>20 \text{ J/m}^2$. This suggests that conversion of the monomer to the dimer may precede JNK ability to elicit kinase activity MKK4/7, and is in agreement with the finding that GSTp efficiently reduces JNK phosphorylation of c-Jun *in vivo*, even when its upstream kinase MKK4 is active.

That specific inhibitors of GSTp efficiently decrease GSTp inhibition of JNK both *in vitro* (TER-117, TER-287) and *in vivo* (TER-199) suggests that their binding to GSTp is within the domain which is also required for association with Jun–JNK or, alternatively, that they alter the conformation of GSTp, affecting its association with JNK. Our results suggest that GSTp does not require transferase activity to mediate inhibition of JNK, in as much as bacterially produced GST, as well as GSTp mutated on Tyr7, which is essential for its catalytic activity, efficiently inhibited JNK enzyme and Jun transcriptional activities, respectively.

Important confirmation of the finding that GSTp is a regulator of JNK kinase activity comes from the use of cells from GSTP1/P2^(-/-)-null mice (Henderson *et al.*, 1998). GSTP1/P2^(-/-)-derived MEFs revealed a noticeably higher basal level of JNK activity, which was also reflected in JNK's phosphorylation and Jun-driven transcriptional activities. Transfection of GSTp into MEF^{GSTP1/P2^(-/-)} cells caused a decrease in high basal JNK phosphorylation, kinase activity and Jun-mediated transactivation, further supporting the role of GSTp as an inhibitor of JNK signaling.

It is well documented that GSTp levels vary between different cell types (Tew, 1994). This, together with

variable cellular compartmentalization and the expression of other detoxification enzymes, will contribute to variability of GSTp inhibition of JNK activity. Indeed, comparison of 3T3 fibroblasts and MEFs revealed that while GSTp transfection causes 100% inhibition in MEFs, it is limited to 50% in the 3T3 fibroblasts.

Overexpression of GSTp has been associated with transformation to malignancy (Sato, 1989) and acquired resistance to electrophilic anticancer drugs (Nakagawa *et al.*, 1990; Tew, 1994). The finding that GSTp is a modulator of JNK inhibitor and the relationship between expression of this protein and JNK inhibition suggests that cancer cells prone to overexpress GSTp may exhibit high intrinsic JNK inhibitory activity. In this model, tumor cells overexpressing GSTp may escape apoptosis, which has been implicated as one of the end-points of JNK activity (Xia *et al.*, 1995; Kasibhatla *et al.*, 1998).

The emerging model suggests that through its association with the Jun–JNK complex under non-stressed conditions, GSTp inhibits JNK phosphorylation and activity (Figure 7). Changes in the level of ROS elicited by UV irradiation or H₂O₂ treatments decrease the amount of monomeric GSTp, resulting in a reduction of the GST–JNK complex, thereby enabling JNK phosphorylation (Figure 7). The fact that transcriptionally active p53 and c-Jun directly affect transcription of radical scavenging enzymes including GST (Polyak *et al.*, 1997; Komarova *et al.*, 1998) points to the possible existence of an autoregulatory loop for GSTp regulation of JNK. Under such autoregulation, stress-mediated GSTp dimerization/multimerization could enable JNK activation, yielding transcriptionally active p53/Jun and, in turn, newly synthesized GSTp that re-forms a complex with JNK to limit the degree and duration of JNK kinase activity. Our studies support this model, which provides a possible underlying mechanism for the regulation of stress kinases by altered redox potential (Adler *et al.*, 1996; Gomez del Arco, 1996; Kuo *et al.*, 1996; Wilhelm *et al.*, 1997; reviewed by Finkel, 1998). Thus, as a guardian of JNK activities in normally growing cells, GSTp may serve as a sensor of intracellular changes in redox potential that are elicited by various forms of stress.

Materials and methods

Cells and protein preparation

The mouse fibroblast cell line 3T3-4A (Adler *et al.*, 1995a), kindly provided by Dr Claudio Basilico, was maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% calf serum and antibiotics (Gibco-BRL). Cells were grown at 37°C with 5% CO₂. MEFs from GSTP1/P2^(-/-) cells and wild-type control animals were prepared using standard protocols. MEFs were maintained in DMEM supplemented with 10% fetal bovine serum (FBS) for no longer than 2 weeks. Proteins were prepared from cells as previously described (Adler *et al.*, 1995a). In all cases, buffers contained a protease 'cocktail' (1 µg/ml of pepstatin, leupeptin and aprotinin) and phosphatase inhibitors (1 mM sodium orthovanadate and 5 mM sodium fluoride).

Chemicals

H₂O₂, eeGSH, GST and NAC were purchased from Sigma.

Antibodies, immunoprecipitations and immunoblots

Antibodies to c-Jun, PKA and ERK1 were purchased from Santa Cruz. Antibodies to JNK were obtained from PharMingen. Antibodies to CKII were purchased from Upstate Biotechnology and antibodies to phospho-JNK were purchased from Promega. Antibodies to GSTs were as described

previously (Ramgamathan and Tew, 1991; Henderson *et al.*, 1998). Antibodies to the phosphorylated form of MKK4, ERK1/2 were obtained from New England Biolabs.

Immunoprecipitations were carried out using 1 mg of protein extracts and 500 ng of the antibodies, for 16 h at 4°C. Protein A/G beads (Gibco-BRL) were added (15 µl) for 30 min at room temperature before washes were carried out in PBS supplemented with Tween-100 (0.5%). Immunoblot analysis was performed using 50 µg of whole-cell extract that had been separated on SDS–PAGE (10%) followed by electrotransfer to PVDF membrane. Ponceau staining was carried out to confirm equal loading, followed by blocking (5% non-fat milk) and reaction with the appropriate antibodies (diluted 1:3000) for 16 h at 4°C. Reactions were visualized using enhanced chemiluminescence (ECL) reagents (Amersham). Analysis of GSTp migration was performed as indicated (Shen *et al.*, 1993), using material that had been immunoprecipitated (0.3 µg of GSTp antibodies and 1 mg of proteins) from cells. Where indicated, separation under non-reducing conditions was performed in SDS–PAGE without including the reducing agent β-mercaptoethanol in sample buffer.

Peptides and inhibitor

Selective inhibitors of GST, including TER-117, TER-199, TER-291 and TER-317, were synthesized, purified to >93% purity, and kindly provided by Telik Inc. (San Francisco, CA). Peptides corresponding to flexible domains on GST [amino acids 194–201, sequence SSPEHVNR (P1), and amino acids 36–50, sequence TIDTWMQGLLKPTCL (P2)] were synthesized (Peptide Technologies Corp., Gaithersburg, MD) and purified by HPLC to >98.5%.

UV irradiation and H₂O₂ treatment

Cells in the logarithmic growth phase were exposed to UV-C (60 J/m², which requires a 15 s exposure in PBS with the lids off), followed by addition of medium and incubation for 45 min. When indicated, NAC (10 mM) was added to the cells 1 h prior to UV irradiation. For H₂O₂ treatment, medium from the culture dish was mixed with freshly diluted H₂O₂ (10 µM) and immediately applied to the fibroblasts (10⁷ cells). When indicated, H₂O₂ was added in the presence of 1 mM eeGSH (Sigma).

Protein kinase assays

Protein kinase assays were carried out using a fusion protein, GST–Jun (amino acids 5–89; Adler *et al.*, 1992) or his-c-Jun (full-length; Treier *et al.*, 1994) as a substrate. JNK2 was purified from UV-treated 3T3-4A cells as previously described (Adler *et al.*, 1995a). The purity of bacterially produced c-Jun and of 3T3-4A-derived JNK was confirmed by silver-stained SDS–PAGE. Solid-phase JNK assays were carried out as previously described (Adler *et al.*, 1995b, 1996). Briefly, the GST–Jun^{5–89} or hisJun^{full-length} fusion proteins (0.5 µg/assay) were bound to glutathione–Sepharose or nickel beads, before incubation with the purified form of JNK (0.5 µg/assay) in the presence of kinase buffer [20 mM HEPES, pH 7.6, 1 mM EGTA, 1 mM dithiothreitol (DDT), 2 mM MgCl₂, 2 mM MnCl₂, 5 mM NaF, 1 mM NaVO₃, 50 mM NaCl] at room temperature for 15 min. The beads were pelleted and washed extensively with PBST [150 mM NaCl, 16 mM sodium phosphate, pH 7.5, 1% Triton X-100, 2 mM EDTA, 0.1% β-mercaptoethanol, 0.2 mM phenylmethylsulfonyl fluoride (PMSF) and 5 mM benzamide], before they were incubated with [³²P]ATP (50 c.p.m./fmol; Amersham) in the presence of kinase buffer. Following extensive washing, the phosphorylated Jun was boiled in SDS sample buffer and the eluted proteins were run on a 15% SDS–polyacrylamide gel. The gel was dried, and phosphorylation of the Jun substrate was determined by autoradiography, followed by quantification with a phosphorimager (Bio-Rad).

To assay for the presence of the JNK inhibitor, we used Jun–JNK as a pre-formed complex. GST–Jun or hisJun (0.5 µg) was first pre-incubated with the purified form of active JNK (0.5 µg), thus forming a Jun–JNK complex which subsequently was incubated with protein extracts (for 30 min) before addition of [³²P]ATP (for 10 min) at room temperature. Important for the formation of these pre-formed complexes is the ratio between Jun and JNK molecules (1:1 molar ratio) and the saturation of the nickel or glutathione beads with EDTA or glutathione to prevent non-specific binding of cellular proteins. In all cases, the buffers used contained a cocktail of protease and phosphatase inhibitors (Adler *et al.*, 1995c, 1997).

Other kinase assays were performed with peptides that are known to serve as specific substrates for Src tyrosine kinases, MAPKs and for PKA, all of which were purchased from Santa Cruz Biotech. Co. (Santa

Cruz, CA). Histone H1 (Sigma) was used as a substrate for CKII. In all cases, the substrates were covalently bound to beads using the immunolink kit (Pierce) according to the manufacturer's recommendations. In these assays, 20 μ l of bead-coupled peptides (1 μ g) were incubated with the source of kinase in the presence or absence of the inhibitor fraction, followed by extensive washing and quantification.

Purification of JNK inhibitor

Protein extracts prepared from 3T3-4A mouse fibroblast cells (4 mg/ml) were precipitated with ammonium sulfate (50%). The precipitated material was dialyzed against kinase buffer and subjected to fractionation on small concentrators with cut-off membranes of 100 and 30 kDa (Millipore, Bedford, MA). Material that passed through the 100 kDa membrane was placed on a 30 kDa column and the filtrate (<30 kDa) was concentrated further to a volume of 50 μ l using a 10 kDa membrane (Amicon). Concentrated material that was 10–30 kDa in size was separated on a Superdex 75 column (SMART system; Pharmacia) pre-calibrated with respect to the position of the expected molecular weight using a combination of protein standards. Using kinase buffer and a flow rate of 40 μ l/min, fractions (200 μ l) were collected and tested for inhibitor activity. In these assays, 20 μ l fractions were added to pre-formed 32 P-hisJun–JNK complex before addition of [γ - 32 P]ATP in the presence of kinase buffer and a cocktail of protease and phosphatase inhibitors. Active fractions were pooled and applied to a MonoQ column with a gradient of 20–500 mM NaCl in kinase buffer at a flow rate of 75 μ l/min. MonoQ fractions were tested for inhibitory activity, and positive fractions were pooled and loaded on a phenyl-Sepharose column in a buffer consisting of 20 mM K₂HPO₄ pH 7.5, 10% glycerol and 0.6 M ammonium sulfate. Using a gradient elution scheme, the inhibitory component was found in the 0.2 M ammonium sulfate fraction. To adjust for kinase buffer and to concentrate the fraction that contained inhibitory activity, the phenyl-Sepharose eluate was concentrated on a 3 kDa column and applied to a pre-formed 32 P-hisJun–JNK complex. This fraction inhibited JNK activity (Table I). Material that was bound to the 32 P-hisJun–JNK complex revealed a single protein on silver-stained SDS–PAGE.

Microsequencing

A large-scale preparation of the JNK inhibitor (Table I) was subjected to multiple separation steps as described above, followed by SDS–PAGE and blotting onto a PVDF membrane. The purity of the single protein identified by Ponceau staining on the PVDF membrane was confirmed in parallel by silver staining of the same material. The band identified on the PVDF membrane was excised and subjected to N-terminal analysis on an ABI protein sequencer Model 494 equipped with a Model 140C phenylthiohydantoin microanalyzer. The sequence Pro-Pro-Tyr-Thr-Val-Val-Tyr-Phe-Pro-Val-Arg-Gly, which was obtained at the 10 pmol level, has 100% homology to the human GST P1-1.

Purification of GSTp

GSTp was purified from human placenta. GST α and GST μ were purified from human liver as described (Ciaccio *et al.*, 1991), followed by Superdex 75 gel filtration (SMART). Purity was confirmed by silver-stained SDS–PAGE.

Transcription and ubiquitination assays

Transcriptional analysis of Jun was carried out using the 5 \times Jun2-driven luciferase construct as previously reported (van Dam *et al.*, 1998). In all cases, values were normalized with respect to transfection efficiency. *In vivo* ubiquitination assays were carried out by transfection of Jun and HA-tagged ubiquitin into the 3T3 cells as indicated elsewhere (Treier *et al.*, 1994).

Acknowledgements

We are grateful to Andras Schaffer, Mark Bluth, Zheng Dong and Theresa Wood for technical assistance, and to Alla Polotskaya and Serge Khilko for advice. We also thank Claudio Basilico for the 3T3-4A cells, Dirk Bohmann for the 32 P-hisJun and Ub-HA constructs, Craig Monell of PharMingen for the antibodies to JNK, Michael Comb of New England Biolabs for the phosphoantibodies to MKK4 and ERK1/2, and Telik Inc. (San Francisco, CA) for the supply of GSTp inhibitors. The work was supported by NCI grant CA77389 to Z.R.

References

Adler,V., Franklin,C.C. and Kraft,A.S. (1992) Phorbol esters stimulate phosphorylation of c-Jun but not c-Jun: regulation by the N-terminal δ domain. *Proc. Natl Acad. Sci. USA*, **89**, 5341–5345.

- Adler,V., Pincus,M.R., Brandt-Rauf,P. and Ronai,Z. (1995a) Complexes of p21^{ras} with jun N-terminal kinase and Jun proteins. *Proc. Natl Acad. Sci. USA*, **92**, 10585–10589.
- Adler,V., Fuchs,S.Y., Kim,J., Kraft,A., King,M.P., Pelling,J. and Ronai,Z. (1995b) Jun-NH₂-terminal kinase activation mediated by UV-induced DNA lesions in melanoma and fibroblast cells. *Cell Growth Differ.*, **6**, 1437–1446.
- Adler,V., Shaffer,A., Kim,J., Dolan,L. and Ronai,Z. (1995c) UV-irradiation and heat shock mediate JNK activation via alternate pathways. *J. Biol. Chem.*, **270**, 26071–26077.
- Adler,V., Polotskaya,A., Kim,J., Dolan,L., Davis,R., Pincus,M. and Ronai,Z. (1995d) Dose rate and mode of exposure are key factors in JNK activation by UV-irradiation. *Carcinogenesis*, **17**, 2073–2076.
- Adler,V., Pincus,M.R., Polotskaya,A., Montano,X., Friedman,F. and Ronai,Z. (1996) Activation of c-Jun NH₂-kinase by UV irradiation is dependent on p21^{ras}. *J. Biol. Chem.*, **271**, 23304–23309.
- Adler,V. *et al.* (1997) Conformation-dependent phosphorylation of p53. *Proc. Natl Acad. Sci. USA*, **94**, 1686–1691.
- Baichwal,V.R., Park,A. and Tjian,R. (1991) Control of c-Jun activity by interaction of a cell specific inhibitor with regulatory domain delta, differences between c-jun and v-jun. *Nature*, **352**, 165.
- Cavigelli,M., Li,W.W., Lin,A., Su,B., Yoshioka,K. and Karin,M. (1996) The tumor promoter arsenite stimulates AP1 activity by inhibiting a JNK phosphatase. *EMBO J.*, **15**, 6269–6279.
- Ciaccio,P.J., Tew,K.D. and La Creta,F.P. (1991) Enzymatic conjugation of chlorambucil with glutathione by human glutathione S-transferases and inhibition by ethacrynic acid. *Biochem. Pharmacol.*, **42**, 1504–1507.
- Coso,O.A., Chiariello,M., Yu,J.C., Teramoto,H., Crespo,P., Xu,N., Miki,T. and Gutkind,J.S. (1995) The small GTP-binding proteins Rac1 and Cdc42 regulate the activity of the JNK/SAPK signaling pathway. *Cell*, **81**, 1137–1146.
- Cui,X.L. and Douglas,J.G. (1997) Arachidonic acid activates c-jun N-terminal kinase through NADPH oxidase in rabbit proximal tubular epithelial cells. *Proc. Natl Acad. Sci. USA*, **94**, 3771–3776.
- Dhar,V., Adler,V., Lehman,A. and Ronai,Z. (1996) Impaired jun-NH₂-terminal kinase activation by ultraviolet irradiation in fibroblasts of patients with Cockayne syndrome complementation group B. *Cell Growth Differ.*, **7**, 841–846.
- Fanger,G.R., Gerwins,P., Widmann,C., Jarpe,M.B., Johnson,G.L. (1997) MEKKs, GCKs, MLKs, PAKs, TAKs and tpls: upstream regulators of the c-Jun amino terminal kinases? *Curr. Opin. Genet. Dev.*, **7**, 67–74.
- Finkel,T. (1998) Oxygen radicals and signaling. *Curr. Opin. Cell Biol.*, **10**, 248–253.
- Flartaard,J.E., Bauer,K.E. and Kauvar,L.M. (1993) Isozyme specificity of novel glutathione S-transferase inhibitors. *Cancer Chemother. Pharmacol.*, **33**, 63–70.
- Fuchs,S.Y., Dolan L.R., Davis,R.J. and Ronai,Z. (1996) JNK targets the ubiquitination of c-Jun in a phosphorylation-dependent manner. *Oncogene*, **13**, 1531–1535.
- Fuchs,S.Y., Xie,B., Adler,V., Fried,V.A., Davis,R.J. and Ronai,Z. (1997) c-Jun NH₂-terminal kinases target the ubiquitination of their associated transcription factors. *J. Biol. Chem.*, **272**, 32163–32168.
- Fuchs,S.Y., Adler,V., Pincus,M.R. and Ronai,Z. (1998a) MEKK1/JNK stabilizes and activates p53. *Proc. Natl Acad. Sci. USA*, **95**, 10541–10546.
- Fuchs,S.Y., Adler,V., Buschmann,T., Yin,Z., Wu,X., Jones,S.T. and Ronai,Z. (1998b) JNK targets p53 ubiquitination and degradation in nonstressed cells. *Genes Dev.*, **12**, 2658–2663.
- Fuchs,S.Y., Fried,V. and Ronai,Z. (1998c) Stress activated kinase regulates protein stability. *Oncogene*, **17**, 1483–1490.
- Galcheva-Gargova,Z., Derijard,B., Wu,I.-H. and Davis,R.J. (1994) An osmosensing signal transducing pathway in mammalian cells. *Science*, **265**, 806–808.
- Gomez del Arco,P., Martinez-Martinez,S., Calvo,V., Armesilla,A.L. and Rendondo,J.M. (1996) JNK is target for antioxidants in T lymphocytes. *J. Biol. Chem.*, **271**, 26335–26340.
- Gupta,S.D., Campbell,B., Derijard,R. and Davis,J. (1995) Transcription factor ATF2 regulation by the JNK signal transduction pathway. *Science*, **267**, 389–393.
- Guyton,K.Z., Liu,Y., Gorospe,M., Xu,Q. and Holbrook,N.J. (1996) Activation of mitogen-activated protein kinase by H₂O₂. Role in cell survival following oxidant injury. *J. Biol. Chem.*, **271**, 4138–4142.
- Hanada,M. *et al.* (1998) Selective suppression of stress-activated protein kinase pathway by protein phosphatase 2C in mammalian cells. *FEBS Lett.*, **437**, 172–176.

- Ip,Y.T. and Davis,R.J. (1998) Signal transduction by the c-Jun N-terminal kinase (JNK) from inflammation to development. *Opin. Cell Biol.*, **10**, 205–219.
- Henderson,C.J., Smith,A.G., Ure,J., Brown,K., Bacon,E.J. and Wolf,C.R. (1998) Increased skin tumorigenesis in mice lacking pi class glutathione S-transferases. *Proc. Natl Acad. Sci. USA*, **95**, 5275–5280.
- Ip,Y.T. and Davis,R.J. (1998) Signal transduction by the c-Jun. N-terminal kinase (JNK) from inflammation to development. *Opin. Cell Biol.*, **10**, 205–219.
- Ishibashi,T., Bottaro,D.P., Chan,A., Miki,T. and Aaronson,S.A. (1992) Expression cloning of a human dual-specificity phosphatase. *Proc. Natl Acad. Sci. USA*, **89**, 12170–12174.
- Jakoby,W.B. (1978) The glutathione S-transferases: a group of multifunctional detoxification enzymes. *Adv. Enzymol.*, **46**, 383–414.
- Kasibhatla,S., Brunner,T., Genestier,L., Echeverri,F., Mahboubi,A., Green,M. and Douglas,R. (1998) DNA damaging agents induce expression of Fas ligand and subsequent apoptosis in T lymphocytes via the activation of NF- κ B and AP-1. *Mol. Cell*, **1**, 543–551.
- Komarova,E.A., Diatchenko,L., Rokhlin,O.W., Hill,J.E., Wang,Z.J., Krivokrysenko,V.I., Feinstein,E. and Gudkov,A.V. (1998) Stress-induced secretion of growth inhibitors: a novel tumor suppressor function of p53. *Oncogene*, **17**, 1089–1096.
- Konishi,H., Tanaka,M., Takemura,Y., Matsuzaki,H., Ono,Y., Kikkawa,U. and Nishizuka,Y. (1997) Activation of protein kinase C by tyrosine phosphorylation in response to H₂O₂. *Proc. Natl Acad. Sci. USA*, **94**, 11233–11237.
- Kuo,M.L., Meng,T.C. and Lin,J.L. (1996) Involvement of glutathione in induction of c-jun proto-oncogene by methylmethanesulfonate in NIH3T3 cells. *Carcinogenesis*, **17**, 815–820.
- Kyriakis,J.M., Banerjee,P., Nikolakaki,E., Dai,T., Rubie,E.A., Ahmad,M.F. and Avruch,J. (1994) The stress-activated protein kinase subfamily of c-Jun kinases. *Nature*, **369**, 156–160.
- Lo,Y.Y.C., Wong,J.M.S. and Cruz,T.F. (1996) Reactive oxygen species mediate cytokine activation of c-Jun NH₂ terminal kinase. *J. Biol. Chem.*, **271**, 15703–15707.
- Minden,A., Lin,A., McMahon,M., Lange-Carter,C., Derijard,B., Davis,R.J., Johnson,G.L. and Karin,M. (1994) Differential activation of ERK and JNK mitogen-activated protein kinases by Raf-1 and MEKK. *Science*, **266**, 1719–1723.
- Musti,A.M., Treier,M. and Bohmann,D. (1997) Reduced ubiquitin-dependent degradation of c-Jun after phosphorylation by MAP kinases. *Science*, **275**, 400–402.
- Nakagawa,K. et al. (1990) Glutathione-S-transferase pi as a determinant of drug resistance in transfectant cell lines. *J. Biol. Chem.*, **265**, 4296–4301.
- Polyak,K., Xia,Y., Zweier,J.L., Kinzler,K.W. and Vogelstein,B. (1997) A model for p53-induced apoptosis. *Nature*, **389**, 300–305.
- Rangamatha,S. and Tew,K.D. (1991) Immunohistochemical localization of glutathione S-transferases alpha, mu and pi in normal tissue and carcinomas from human colon. *Carcinogenesis*, **12**, 2383–2387.
- Sato,K. (1989) Glutathione transferases as markers of preneoplasia and neoplasia. *Adv. Cancer Res.*, **52**, 205–255.
- Shen,H., Tsuchida,S., Tamai,K. and Sato,K. (1993) Identification of cysteine residues involved in disulfide formation in the inactivation of glutathione transferase P form by hydrogen peroxide. *Arch. Biochem. Biophys.*, **300**, 137–141.
- Shim,J., Lee,H., Park,J., Kim,H and Choi,E.J. (1996) A non-enzymatic p21 protein inhibitor of stress-activated protein kinases. *Nature*, **381**, 804–806.
- Tew,K.D. (1994) Glutathione-associated enzymes in anticancer drug resistance. *Cancer Res.*, **54**, 4313–4320.
- Tournier,C., Whitmarsh,A.J., Cavanagh,J., Barret,T. and Davis,R.J. (1997) Mitogen-activated protein kinase 7 is an activator of the c-Jun NH₂-terminal kinase. *Proc. Natl Acad. Sci. USA*, **94**, 7337–7342.
- Treier,L., Staszewski,M. and Bohmann,D. (1994) Ubiquitin-dependent c-Jun degradation *in vivo* is mediated by the delta domain. *Cell*, **78**, 787–798.
- van Dam,H., Huguier,S., Kooistra,K., Baguet,J., Vial,E., van der Eb,A.J., Herrlich,P., Angel,P. and Castellazzi,M. (1998) Autocrine growth and anchorage independence: two complementing Jun-controlled genetic programs of cellular transformation. *Genes Dev.*, **12**, 1227–1239.
- Wang,G.L., Jiang,B.H., Rue,E.A. and Semenza,G.L. (1995) Hypoxia-inducible factor 1 is a basic-helix-loop-helix-PAS heterodimer regulated by cellular O₂ tension. *Proc. Natl Acad. Sci. USA*, **92**, 5510–5514.
- Westwick,J.K., Weitzel,C., Minden,A., Karin,M. and Brenner,D.A. (1994) Tumor necrosis factor alpha stimulates AP-1 activity through prolonged activation of the c-Jun kinase. *J. Biol. Chem.*, **269**, 26396–26401.
- Whitmarsh,A.J., Shore,A.D., Sharrocks,R. and Davis,R.J. (1995) Integration of MAP kinase signal transduction pathways at the serum response element. *Science*, **269**, 403–407.
- Whitmarsh,A.J., Cavanagh,J., Tournier,C., Yasuda,J. and Davis,R.J. (1998) A mammalian scaffold complex that selectively mediates MAP kinase activation. *Science*, **281**, 1671–1674.
- Wilhelm,D., Bender,K., Knebel,A. and Angel,P. (1997) The level of intracellular glutathione is a key regulator for the induction of stress-activated signal transduction pathways including jun N-terminal protein kinases and p38 kinase by alkylating agents. *Mol. Cell Biol.*, **17**, 4792–4800.
- Xanthoudakis,S., Miao,G., Wang,F., Pan,Y.C.E. and Curran,T. (1992) Redox activation of Fos-Jun DNA binding activity is mediated by a DNA repair enzyme. *EMBO J.*, **11**, 3323–3335.
- Xia,M., Dickens,J., Raingeaud,R., Davis,J. and Greenberg,M.E. (1995) Opposing effects of ERK and JNK-p38 MAP kinases on apoptosis. *Science*, **270**, 1326–1331.
- Xia,Y., Wu,Z., Su, B., Murray,B. and Karin,M. (1998) JNKK1 organizes a MAP kinase module through specific and sequential interactions with upstream and downstream components mediated by its amino-terminal extension. *Genes Dev.*, **12**, 3369–3381.

Received November 19, 1998; revised and accepted January 8, 1999