

# The unpredictability of prolonged activation of stress response pathways

Lilian T. Lamech<sup>1</sup> and Cole M. Haynes<sup>1,2</sup>

<sup>1</sup>Cell Biology Program, Memorial Sloan-Kettering Cancer Center, New York, NY 10065

<sup>2</sup>BCMB Allied Program, Weill Cornell Medical College, New York, NY 10065

In response to stress, cellular compartments activate signaling pathways that mediate transcriptional programs to promote survival and reestablish homeostasis. Manipulation of the magnitude and duration of the activation of stress responses has been proposed as a strategy to prevent or repair the damage associated with aging or degenerative diseases. However, as these pathways likely evolved to respond specifically to transient perturbations, the unpredictability of prolonged activation should be considered.

Cellular stresses, such as unfolded or misfolded protein accumulation and organelle deterioration, are associated with numerous diseases as well as the aging process. Thus, enhanced activation of pathways that have evolved to protect against these defects may protect against degenerative diseases such as Parkinson's and Alzheimer's or the ill effects of normal aging (Powers et al., 2009; Bratic and Larsson, 2013). Stress response pathways are typically maintained in the off state or at a baseline level. Upon organelle perturbation, they are activated to the appropriate magnitude and duration to efficiently promote cellular survival and organelle recovery. Once homeostasis is reestablished, the pathway is down-regulated so that cells can properly respond to future stress (Fig. 1 A).

Manipulations of these pathways can mitigate the intracellular damage that occurs during aging or in degenerative diseases. However, these pathways did not likely evolve to deal with prolonged stress or to be activated for extended periods of time (Fig. 1 B). If continued activation were entirely beneficial, these pathways would likely have evolved to be hardwired into developmental or cell-specific programs rather than to be stress inducible. We hypothesize that prolonged stress response activation has not been subject to evolution, as conditions that cause perpetual activation, such as deleterious gene mutations, result in cellular damage and would be selected against. Thus, the potential outcomes of prolonged stress response activation are difficult to predict. Here, we review the evidence suggesting that stress response pathways evolved to be transiently activated to a precise magnitude to match the level of dysfunction and allow the most efficient recovery, and consider the positive and negative effects of enhanced stress response activation. We also consider approaches to therapeutically engage stress response signaling.

Correspondence to Cole M. Haynes: haynesc@mskcc.org


## Stress detection and matched transcriptional activation

Several organelle or stress-specific stress responses have been identified and are described in more detail elsewhere (Åkerfelt et al., 2010; Walter and Ron, 2011; Jensen and Jasper, 2014). Here, we focus on specific responses that are activated by cytosolic, ER, or mitochondrial stress or dysfunction.

**The heat shock response (HSR).** The HSR is mediated by the transcription factor HSF1 and occurs during conditions that cause an increase in unfolded or misfolded proteins primarily in the cytosol and nucleus, such as increased temperature, oxidative stress, and exposure to heavy metals (Ananthan et al., 1986; Åkerfelt et al., 2010). However, it can also be activated independently of misfolded proteins, as stalled ribosome complexes also activate the response (Brandman et al., 2012). The HSR is a transcriptional program that involves the induction of ~500 genes, including cytosolic and nuclear-localized protein homeostasis (proteostasis) machinery such as molecular chaperones and genes involved in protein synthesis, the cell cycle, and the regulation of cell death (Mendillo et al., 2012; Ryno et al., 2014). While the induction of chaperones and proteases garner much of the attention, the HSR also includes the repression of ~1,000 genes including developmental, immune, apoptotic (Mendillo et al., 2012; Ryno et al., 2014), and cytoskeletal maintenance components (Baird et al., 2014; Fig. 2 A).

Normally, HSF1 is repressed by the cytosolic and nuclear-localized molecular chaperone Hsp90, which binds and maintains the transcription factor at a baseline level (Morimoto, 1998; Zou et al., 1998). As unfolded proteins increase, HSF1 is released, allowing it to bind the heat shock promoter element and regulate transcription (Fig. 2 A; Topol et al., 1985; Morgan et al., 1987). In addition to direct regulation by chaperones, HSF1 is also subject to multiple posttranslational modifications (Anckar and Sistonen, 2011). For example, the magnitude and duration of HSF1 activation are further regulated by the acetyltransferase EP300 and the deacetylase SIRT1. HSF1 acetylation by EP300 controls the quantity of HSF1 available for activation by preventing proteasome-dependent degradation (Raychaudhuri et al., 2014). Conversely, deacetylation of HSF1 by SIRT1 promotes activation of HSF1 during stress (Westerheide et al., 2009), but will eventually lead to HSF1 turnover to down-regulate the response (Raychaudhuri et al., 2014). Further regulation of the HSR occurs at the organismal level via thermosensory

© 2015 Lamech and Haynes This article is distributed under the terms of an Attribution-Noncommercial-Share Alike-No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms>). After six months it is available under a Creative Commons License (Attribution-Noncommercial-Share Alike 3.0 Unported license, as described at <http://creativecommons.org/licenses/by-nc-sa/3.0/>).



**Figure 1. Transient versus perpetual or prolonged stress response activation.** (A) The HSR, UPR, and the UPR<sup>mt</sup> remain in an “off” or “low-activity” state until they are activated by compartment-specific stress. As these pathways are not constitutively active, these stress responses likely evolved to promote survival during temporary stressful conditions and ultimately recover once the condition causing the cellular dysfunction is alleviated. Transient stresses include environmental effects such as temperature shifts, exposure to toxins, or altered nutritional status. (B) Genetic mutations that perturb cytosolic, ER, or mitochondrial proteostasis are typically selected against evolutionarily as they cause cellular dysfunction. However, genotoxic perturbations, or damage that has accrued over long periods of time during aging or disease, may cause prolonged or perpetual activation of the HSR, UPR, and UPR<sup>mt</sup>, as the mutation cannot be rectified. Prolonged or perpetual stress response activation is potentially very different than transient activation as there is never a recovery. As evolution did not select for prolonged stress response activation, we hypothesize that it is difficult to predict the outcome of prolonged stress response activation.

neurons and neuroendocrine signaling. Non-cell-autonomous HSF1 activation presumably allows for more precise matching of the HSR to the behavioral and metabolic status of the organism (Prahlad et al., 2008; Prahlad and Morimoto, 2011).

**The unfolded protein response (UPR).** The ER is the site of protein synthesis and folding for the vast majority of proteins that are secreted or localized within the secretory pathway. In response to increased protein flux through the ER or to conditions that perturb ER protein folding, UPR activation limits the load on the stressed organelle by reducing localized protein synthesis and activating protective ER-specific transcriptional programs to reestablish organelle homeostasis (Walter and Ron, 2011; Fig. 2 B). The most conserved branch of the UPR is regulated by the ER-membrane localized kinase IRE1 and the transcription factor XBP1 (Hac1 in yeast). The UPR is activated when accumulating unfolded proteins directly


interact with the luminal domain of IRE1 (Gardner and Walter, 2011), causing it to oligomerize, activating the cytosolic RNase domain (Korennykh et al., 2009). IRE1 cleaves several ER-localized mRNAs, resulting in their degradation, and thus reducing their translation and the burden on the ER protein-folding environment (Han et al., 2009; Hollien et al., 2009). Concomitantly, IRE1 also cleaves an inhibitory intron from the transcript encoding XBP1, which upon ligation allows the translation of the active transcription factor (Cox and Walter, 1996; Yoshida et al., 2001; Calton et al., 2002). Once translated, XBP1 activates a broad transcriptional response that includes ER-localized components that promote protein folding and quality control, compartmental expansion, and increased ER-Golgi trafficking (Travers et al., 2000; Shoulders et al., 2013; Fig. 2 B). However, if ER stress cannot be rectified, an apoptotic program is engaged to eliminate the unsalvageable cell (Tabas and Ron, 2011; Upton et al., 2012; Lu et al., 2014).

**The mitochondrial UPR (UPR<sup>mt</sup>).** The UPR<sup>mt</sup> is a transcriptional response that occurs specifically during mitochondrial dysfunction to promote survival and recovery of mitochondrial activity. The UPR<sup>mt</sup> in *Caenorhabditis elegans* is regulated by the transcription factor ATFS-1, which is normally imported into mitochondria and degraded (Nargund et al., 2012; Haynes et al., 2013). However, during conditions that impair mitochondrial protein import such as respiratory chain dysfunction, mitochondrial unfolded protein accumulation, or high levels of reactive oxygen species (ROS), general mitochondrial import efficiency is reduced, causing mitochondrial proteins to accumulate in the cytosol (Wright et al., 2001; Nargund et al., 2012; Harbauer et al., 2014). As ATFS-1 has a nuclear localization sequence, some of the cytosolic ATFS-1 pool then traffics to the nucleus to mediate UPR<sup>mt</sup> activation (Nargund et al., 2012; Fig. 2 C). Similar to the HSR and UPR, the UPR<sup>mt</sup> receives non-cell-autonomous regulatory inputs (Durieux et al., 2011; Owusu-Ansah et al., 2013; Taylor and Dillin, 2013).

ATFS-1 activation increases mitochondrial protein folding capacity and promotes mitochondrial recovery by increasing mitochondrial chaperones, proteases, respiratory chain complex assembly factors, import, and fission components (Nargund et al., 2012, 2015). Concomitantly, ATFS-1 limits expression of the tricarboxylic acid (TCA) cycle and respiratory chain components, suggesting that ATFS-1 promotes mitochondrial recovery by increasing protein folding and complex assembly capacity while slowing the rate of respiratory complex biogenesis to match the stressed organelle’s capacity (Nargund et al., 2015; Fig. 2 C). To facilitate organelle repair, ATFS-1 increases expression of all glycolysis components in order to maintain energy levels. Thus, the UPR<sup>mt</sup> includes a shift in cellular metabolism to promote survival during mitochondrial dysfunction that is reminiscent of the metabolism in rapidly dividing cells (Vander Heiden et al., 2009), and which presumably must be down-regulated upon return to homeostasis.

#### Regulation of response duration and recovery

In addition to the magnitude of a specific stress response, which is largely governed by activating mechanisms, the duration of the response must be tightly regulated to match cell physiology and promote efficient recovery. Consistent with the idea that prolonged activation of each pathway is potentially detrimental, multiple mechanisms exist to limit and down-regulate stress response activation. Included in the HSR, UPR and UPR<sup>mt</sup> are



**Figure 2. The heat shock response, the UPR, and the mitochondrial UPR.** (A) The HSR is mediated by the transcription factor HSF1. Normally, HSF1 is repressed or maintained in the “off” state by interacting with the molecular chaperone Hsp90. However, when unfolded proteins accumulate in the cytosol or nucleus, HSF1 dissociates from Hsp90 and binds the promoters of HSR genes. The HSR includes the induction (green) of proteostasis machinery including molecular chaperones as well as the repression (red) of many genes required for development. Once proteostasis is recovered, HSF1 is degraded and the HSR is attenuated. (B) The UPR is mediated by at least three ER stress sensor molecules, the most conserved of which is IRE1. Upon detection of ER stress, IRE1 oligomerizes, activating its cytosolic RNase domain which results in (1) the cleavage and subsequent degradation of ER-localized mRNAs, reducing the incoming protein load on the stressed organelle; and (2) the activation of the transcription factor XBP1, which results in induction of a broad response including ER proteostasis machinery, lipid synthesis to expand ER volume, and increased secretory components. Once ER homeostasis is reestablished, IRE1 signaling is attenuated by association with ER chaperones and XBP1 is degraded. (C) The UPR<sup>mt</sup> is a mitochondrial-specific stress response mediated by ATFS-1. ATFS-1 is activated when mitochondrial protein import is impaired, which can be caused by imbalanced mitochondrial proteostasis or respiratory chain defects. Cytosolic ATFS-1 then traffics to the nucleus and activates the UPR<sup>mt</sup>, which includes an increase (green) in mitochondrial proteostasis machinery such as mitochondrial chaperones and ROS-detoxifying components. The UPR<sup>mt</sup> also involves the repression (red) or limited expression of some of the most highly expressed mitochondrial proteins including components of the TCA cycle and the oxidative phosphorylation (OXPHOS) complexes. Once mitochondrial function is recovered, ATFS-1 is degraded and the UPR<sup>mt</sup> is down-regulated.

components that down-regulate HSF1, XBP1, and ATFS-1, respectively, via negative feedback loops. For example, HSF1 induces expression of Hsp70 and Hsp90, which in addition to promoting efficient protein folding also associates with active HSF1 to dampen the response (Shi et al., 1998). Similarly, XBP1 induces expression of ER-localized chaperones, which associate with IRE1 to attenuate signaling upon proteostasis recovery (Todd-Corlett et al., 2007; Eletto et al., 2014). ATFS-1 also induces multiple components that promote mitochondrial protein import efficiency, which serves to reduce cytosolic and ultimately nuclear accumulation of ATFS-1 (Nargund et al., 2012). Furthermore, all three pathways increase components that target the active transcription factor for degradation, including ubiquitin ligases. In addition to negative regulation of the response regulators, the outputs of the responses are also down-regulated once proteostasis is reestablished. For example, as misfolded or unfolded proteins are depleted, the HSF1-induced chaperone Hsp70 is ubiquitinated by the HSF1-induced ubiquitin ligase CHIP and is degraded by proteasomes, returning the chaperone capacity to baseline levels (Qian et al., 2006).

#### Effects of prolonged activation

HSF1, XBP1, and ATFS-1 have all been shown to be protective during organelle-specific stress, promoting survival and cellular proliferation during conditions that perturb cytosolic and nuclear (Morano et al., 1999; Xiao et al., 1999; Hsu et al., 2003; Morley and Morimoto, 2004), ER (Cox et al., 1993; Shen et al., 2001; Lin et al., 2009; Richardson et al., 2010), or mito-

chondrial proteostasis (Baker et al., 2012; Nargund et al., 2012), respectively. Interestingly, cellular damage that accrues in aging animals activates each pathway when it occurs in young animals. However, all three pathways (Yoneda et al., 2004; Ben-Zvi et al., 2009; David et al., 2010; Durieux et al., 2011) are attenuated and less effective in aging animals, which coincides with a proteostatic collapse (Ben-Zvi et al., 2009), further suggesting that enhanced activation may be beneficial.

Several interesting observations reveal their protective effects against age-associated cellular damage. Impaired insulin signaling, which extends worm lifespan considerably, requires multiple transcription factors including HSF1 and XBP1 (Kimura et al., 1997; Lin et al., 1997; Hsu et al., 2003; Henis-Korenblit et al., 2010). Similarly, modest mitochondrial dysfunction that activates the UPR<sup>mt</sup> also extends lifespan in multiple species including mice, flies, and worms (Dillin et al., 2002b; Durieux et al., 2011; Houtkooper et al., 2013; Owusu-Ansah et al., 2013; Schieber and Chandel, 2014). Thus, pathway activation can mitigate age-associated damage; however, it should be noted that this often comes at the expense of fecundity and normal development (Dillin et al., 2002a).

**HSR.** Impressively, HSF1 activation is sufficient to extend the lifespan of normal worms, indicating that the HSR can be protective over longer periods of time (Hsu et al., 2003; Westerheide et al., 2009). HSF1 activity positively affects proteostasis and reduces aggregation of disease-associated proteins in multiple organisms such as those containing polyglutamine stretches (Calamini et al., 2012; Brunquell et al., 2014),  $\alpha$ -syn-

uclein (Hamamichi et al., 2008), prion protein (PrP; Steele et al., 2008), and  $A\beta^{1-42}$  (Cohen et al., 2006; Calamini et al., 2012).

In addition to these protective effects, accumulating evidence indicates that HSF1 activation can also negatively affect proteostasis. Defects in folding and trafficking of the CFTR protein caused by an amino acid deletion result in cystic fibrosis. While expression of mutated CFTR activates HSF1, it was recently shown that HSF1 inhibition increases CFTR trafficking and function, suggesting that prolonged HSF1 activation creates a maladaptive state (Wang et al., 2006; Roth et al., 2014). Similarly, HSF1 overexpression has been shown to exacerbate aggregation of the polyglutamine protein Huntingtin (Bersuker et al., 2013). Lastly, the HSF1 expression level is associated with poor prognoses in breast cancers, which is consistent with many cancer types requiring HSF1 activity to promote proliferation (Dai et al., 2007; Santagata et al., 2011), highlighting the importance of appropriate HSF1 activation.

**UPR.** Similar to HSF1, expression of XBP1 is sufficient to counteract the secretory pathway dysfunction that occurs during worm aging and results in lifespan extension (Taylor and Dillin, 2013). This suggests that approaches to promote UPR activation may be effective against diseases associated with ER stress, which include neurodegenerative and metabolic diseases as well as those associated with mutations causing expression of terminally misfolded secretory proteins (Ryno et al., 2013). Enhanced UPR activation has been demonstrated to reduce the secretion of a misfolded and dysfunctional variant of rhodopsin that results in photoreceptor cell death (Chiang et al., 2012), promote proper folding and function of mutant lysosomal proteins associated with lysosomal storage disease (Mu et al., 2008), reduce the secretion of amyloidogenic aggregation-prone proteins (Cooley et al., 2014), and limit the neurodegeneration in mouse models of Charcot-Marie Tooth disease and amyotrophic lateral sclerosis (Das et al., 2015).

However, numerous studies indicate that prolonged or inappropriate UPR signaling can be toxic, even if apoptotic induction is avoided (Tabas and Ron, 2011). Phospho-transfer by IRE1's cytosolic kinase domain is not required for activation of the RNase domain; rather, it is required to down-regulate signaling as ER stress is alleviated (Chawla et al., 2011; Rubio et al., 2011). Cells expressing IRE1 with impaired phospho-transfer activity efficiently activate the UPR but are unable to attenuate IRE1 activity. The prolonged UPR activation in these cells fails to return the organelle to proteostasis and is at least partially due to the sustained production of ER-targeted proteins (Rubio et al., 2011), which can lead to developmental arrest or apoptosis (Eletto et al., 2014). Additionally, prolonged turnover of ER-localized mRNAs by IRE1 likely has negative consequences for secretory pathway activity.

**UPR<sup>mt</sup>.** While ATFS-1 is necessary for longevity associated with modest mitochondrial dysfunction (Rea et al., 2007; Schieber and Chandel, 2014), ATFS-1 is not sufficient to promote longevity independent of mitochondrial stress. Mutations in ATFS-1's mitochondrial targeting sequence that cause it to redistribute to the nucleus under normal conditions are quite toxic, impeding development and reducing lifespan (Rauthan et al., 2013). These results may be explained by ATFS-1 functioning as a single component within a broader signaling network that must be integrated to elude protective effects during stress. For example, autophagy (Lapierre et al., 2013), altered protein synthesis (Baker et al., 2012), and additional transcription programs (Lee et al., 2010; Walter et al., 2011) are also

required to promote longevity during mitochondrial dysfunction. Highlighting the protective effects of activating the UPR<sup>mt</sup> to appropriately match the level of mitochondrial dysfunction, worms expressing activated ATFS-1 are resistant when chronically exposed to mitochondrial toxins, statins (Rauthan et al., 2013), or the pathogenic bacteria *Pseudomonas aeruginosa* (Pellegrino et al., 2014), which perturbs mitochondrial function. These results indicate that enhanced UPR<sup>mt</sup> activation can be protective, but the magnitude and duration of the response should be considered as well as other factors that potentially coordinate with the UPR<sup>mt</sup>.

## Conclusions

Manipulations to enhance stress response activation hold promise therapeutically to mitigate the cellular damage that accrues during aging and disease (Calamini et al., 2012; Mouchiroud et al., 2013). Response activation in principle can be achieved by (1) perturbing the protein folding environment, (2) activating the transcription factor directly, or (3) impairing the turnover of the active transcription factors. But the evidence reviewed here suggests that it is difficult to predict the outcome of prolonged activation, as these responses likely evolved to resolve transient proteotoxic stress (Fig. 1 A). Therefore, manipulation of these stress response pathways as a therapeutic measure will require careful consideration of the effects of prolonged activation (Fig. 1 B). Considering the number of transcripts XBP1, ATFS-1, and HSF1 affect in addition to those promoting proteostasis, prolonged activation may alter fundamental aspects of a particular cell. For example, to promote mitochondrial recovery, ATFS-1 activation shifts metabolism to that typically observed in rapidly proliferating cells, which may be detrimental to post-mitotic cells if activation is prolonged (Fig. 2 C).

Despite the challenges in manipulating these pathways to promote organelle recovery and cell survival, we are optimistic that as more knowledge is gained regarding pathway regulation and outputs, therapeutic manipulations can be tailored to limit cellular damage so as to avoid unintended effects of prolonged alterations. A particularly exciting example has been the development of phosphatase inhibitors that partially prolong the effects of the UPR branch that attenuates translation during ER stress (Novoa et al., 2001; Boyce et al., 2005). Impaired activation, or extreme prolonged activation, of the translational control branch of the UPR results in cell death and developmental arrest (Harding et al., 2000, 2009). However, guanabenz inhibits only one of the two phosphatases that attenuate the pathway (Tsaytler et al., 2011). Impressively, guanabenz and, more recently, a related compound have been shown to be protective in a variety of cultured cell lines as well as in mouse models of neurodegeneration (Das et al., 2015; Way et al., 2015).

## Acknowledgements

We apologize to those colleagues whose work on the HSR, UPR, and UPR<sup>mt</sup> we could not include due to space limitations.

This work was supported by the National Institutes of Health (grants R01AG040061 and R01AG047182 to C.M. Haynes).

The authors declare no competing financial interests.

Submitted: 23 March 2015

Accepted: 2 June 2015

## References

- Åkerfelt, M., R.I. Morimoto, and L. Sistonen. 2010. Heat shock factors: integrators of cell stress, development and lifespan. *Nat. Rev. Mol. Cell Biol.* 11:545–555. <http://dx.doi.org/10.1038/nrm2938>
- Ananthan, J., A.L. Goldberg, and R. Voellmy. 1986. Abnormal proteins serve as eukaryotic stress signals and trigger the activation of heat shock genes. *Science.* 232:522–524. <http://dx.doi.org/10.1126/science.3083508>
- Anckar, J., and L. Sistonen. 2011. Regulation of HSF1 function in the heat stress response: implications in aging and disease. *Annu. Rev. Biochem.* 80:1089–1115. <http://dx.doi.org/10.1146/annurev-biochem-060809-095203>
- Baird, N.A., P.M. Douglas, M.S. Simic, A.R. Grant, J.J. Moresco, S.C. Wolff, J.R. Yates III, G. Manning, and A. Dillin. 2014. HSF-1-mediated cytoskeletal integrity determines thermotolerance and life span. *Science.* 346:360–363. <http://dx.doi.org/10.1126/science.1253168>
- Baker, B.M., A.M. Nargund, T. Sun, and C.M. Haynes. 2012. Protective coupling of mitochondrial function and protein synthesis via the eIF2 $\alpha$  kinase GCN-2. *PLoS Genet.* 8:e1002760. <http://dx.doi.org/10.1371/journal.pgen.1002760>
- Ben-Zvi, A., E.A. Miller, and R.I. Morimoto. 2009. Collapse of proteostasis represents an early molecular event in *Caenorhabditis elegans* aging. *Proc. Natl. Acad. Sci. USA.* 106:14914–14919. <http://dx.doi.org/10.1073/pnas.0902882106>
- Bersuker, K., M.S. Hipp, B. Calamini, R.I. Morimoto, and R.R. Kopito. 2013. Heat shock response activation exacerbates inclusion body formation in a cellular model of Huntington disease. *J. Biol. Chem.* 288:23633–23638. <http://dx.doi.org/10.1074/jbc.C113.481945>
- Boyce, M., K.F. Bryant, C. Jousse, K. Long, H.P. Harding, D. Scheuner, R.J. Kaufman, D. Ma, D.M. Coen, D. Ron, and J. Yuan. 2005. A selective inhibitor of eIF2 $\alpha$  dephosphorylation protects cells from ER stress. *Science.* 307:935–939. <http://dx.doi.org/10.1126/science.1101902>
- Brandman, O., J. Stewart-Ornstein, D. Wong, A. Larson, C.C. Williams, G.W. Li, S. Zhou, D. King, P.S. Shen, J. Weibezahn, et al. 2012. A ribosome-bound quality control complex triggers degradation of nascent peptides and signals translation stress. *Cell.* 151:1042–1054. <http://dx.doi.org/10.1016/j.cell.2012.10.044>
- Bratice, A., and N.G. Larsson. 2013. The role of mitochondria in aging. *J. Clin. Invest.* 123:951–957. <http://dx.doi.org/10.1172/JCI64125>
- Brunquell, J., P. Bowers, and S.D. Westerheide. 2014. Fluorodeoxyuridine enhances the heat shock response and decreases polyglutamine aggregation in an HSF-1-dependent manner in *Caenorhabditis elegans*. *Mech. Ageing Dev.* 141:142:1–4. <http://dx.doi.org/10.1016/j.mad.2014.08.002>
- Calamini, B., M.C. Silva, F. Madoux, D.M. Hutt, S. Khanna, M.A. Chalfant, S.A. Saldanha, P. Hodder, B.D. Tait, D. Garza, et al. 2012. Small-molecule proteostasis regulators for protein conformational diseases. *Nat. Chem. Biol.* 8:185–196. <http://dx.doi.org/10.1038/nchembio.763>
- Calfon, M., H. Zeng, F. Urano, J.H. Till, S.R. Hubbard, H.P. Harding, S.G. Clark, and D. Ron. 2002. IRE1 couples endoplasmic reticulum load to secretory capacity by processing the XBP-1 mRNA. *Nature.* 415:92–96. <http://dx.doi.org/10.1038/415092a>
- Chawla, A., S. Chakrabarti, G. Ghosh, and M. Niwa. 2011. Attenuation of yeast UPR is essential for survival and is mediated by IRE1 kinase. *J. Cell Biol.* 193:41–50. <http://dx.doi.org/10.1083/jcb.201008071>
- Chiang, W.C., C. Messah, and J.H. Lin. 2012. IRE1 directs proteasomal and lysosomal degradation of misfolded rhodopsin. *Mol. Biol. Cell.* 23:758–770. <http://dx.doi.org/10.1091/mbc.E11-08-0663>
- Cohen, E., J. Bieschke, R.M. Perciavalle, J.W. Kelly, and A. Dillin. 2006. Opposing activities protect against age-onset proteotoxicity. *Science.* 313:1604–1610. <http://dx.doi.org/10.1126/science.1124646>
- Cooley, C.B., L.M. Ryno, L. Plate, G.J. Morgan, J.D. Hulleman, J.W. Kelly, and R.L. Wiseman. 2014. Unfolded protein response activation reduces secretion and extracellular aggregation of amyloidogenic immunoglobulin light chain. *Proc. Natl. Acad. Sci. USA.* 111:13046–13051. <http://dx.doi.org/10.1073/pnas.1406050111>
- Cox, J.S., and P. Walter. 1996. A novel mechanism for regulating activity of a transcription factor that controls the unfolded protein response. *Cell.* 87:391–404. [http://dx.doi.org/10.1016/S0092-8674\(00\)81360-4](http://dx.doi.org/10.1016/S0092-8674(00)81360-4)
- Cox, J.S., C.E. Shamu, and P. Walter. 1993. Transcriptional induction of genes encoding endoplasmic reticulum resident proteins requires a transmembrane protein kinase. *Cell.* 73:1197–1206. [http://dx.doi.org/10.1016/0092-8674\(93\)90648-A](http://dx.doi.org/10.1016/0092-8674(93)90648-A)
- Dai, C., L. Whitesell, A.B. Rogers, and S. Lindquist. 2007. Heat shock factor 1 is a powerful multifaceted modifier of carcinogenesis. *Cell.* 130:1005–1018. <http://dx.doi.org/10.1016/j.cell.2007.07.020>
- Das, I., A. Krzyzosiak, K. Schneider, L. Wrabetz, M. D'Antonio, N. Barry, A. Sigurdardottir, and A. Bertolotti. 2015. Preventing proteostasis diseases by selective inhibition of a phosphatase regulatory subunit. *Science.* 348:239–242. <http://dx.doi.org/10.1126/science.aaa4484>
- David, D.C., N. Ollikainen, J.C. Trinidad, M.P. Cary, A.L. Burlingame, and C. Kenyon. 2010. Widespread protein aggregation as an inherent part of aging in *C. elegans*. *PLoS Biol.* 8:e1000450. <http://dx.doi.org/10.1371/journal.pbio.1000450>
- Dillin, A., D.K. Crawford, and C. Kenyon. 2002a. Timing requirements for insulin/IGF-1 signaling in *C. elegans*. *Science.* 298:830–834. <http://dx.doi.org/10.1126/science.1074240>
- Dillin, A., A.L. Hsu, N. Arantes-Oliveira, J. Lehrer-Graiwer, H. Hsin, A.G. Fraser, R.S. Kamath, J. Ahringer, and C. Kenyon. 2002b. Rates of behavior and aging specified by mitochondrial function during development. *Science.* 298:2398–2401. <http://dx.doi.org/10.1126/science.1077780>
- Durieux, J., S. Wolff, and A. Dillin. 2011. The cell-non-autonomous nature of electron transport chain-mediated longevity. *Cell.* 144:79–91. <http://dx.doi.org/10.1016/j.cell.2010.12.016>
- Eletto, D., D. Eletto, D. Dersh, T. Gidalevitz, and Y. Argon. 2014. Protein disulfide isomerase A6 controls the decay of IRE1 $\alpha$  signaling via disulfide-dependent association. *Mol. Cell.* 53:562–576. <http://dx.doi.org/10.1016/j.molcel.2014.01.004>
- Gardner, B.M., and P. Walter. 2011. Unfolded proteins are Ire1-activating ligands that directly induce the unfolded protein response. *Science.* 333:1891–1894. <http://dx.doi.org/10.1126/science.1209126>
- Hamamichi, S., R.N. Rivas, A.L. Knight, S. Cao, K.A. Caldwell, and G.A. Caldwell. 2008. Hypothesis-based RNAi screening identifies neuroprotective genes in a Parkinson's disease model. *Proc. Natl. Acad. Sci. USA.* 105:728–733. <http://dx.doi.org/10.1073/pnas.0711018105>
- Han, D., A.G. Lerner, L. Vande Walle, J.P. Upton, W. Xu, A. Hagen, B.J. Backes, S.A. Oakes, and F.R. Papa. 2009. IRE1 $\alpha$  kinase activation modes control alternate endoribonuclease outputs to determine divergent cell fates. *Cell.* 138:562–575. <http://dx.doi.org/10.1016/j.cell.2009.07.017>
- Harbauer, A.B., R.P. Zahedi, A. Sickmann, N. Pfanner, and C. Meisinger. 2014. The protein import machinery of mitochondria—a regulatory hub in metabolism, stress, and disease. *Cell Metab.* 19:357–372. <http://dx.doi.org/10.1016/j.cmet.2014.01.010>
- Harding, H.P., Y. Zhang, A. Bertolotti, H. Zeng, and D. Ron. 2000. Perk is essential for translational regulation and cell survival during the unfolded protein response. *Mol. Cell.* 5:897–904. [http://dx.doi.org/10.1016/S1097-2765\(00\)80330-5](http://dx.doi.org/10.1016/S1097-2765(00)80330-5)
- Harding, H.P., Y. Zhang, D. Scheuner, J.J. Chen, R.J. Kaufman, and D. Ron. 2009. Ppp1r15 gene knockout reveals an essential role for translation initiation factor 2 alpha (eIF2 $\alpha$ ) dephosphorylation in mammalian development. *Proc. Natl. Acad. Sci. USA.* 106:1832–1837. <http://dx.doi.org/10.1073/pnas.0809632106>
- Haynes, C.M., C.J. Fiorese, and Y.F. Lin. 2013. Evaluating and responding to mitochondrial dysfunction: the mitochondrial unfolded-protein response and beyond. *Trends Cell Biol.* 23:311–318. <http://dx.doi.org/10.1016/j.tcb.2013.02.002>
- Henis-Korenblit, S., P. Zhang, M. Hansen, M. McCormick, S.J. Lee, M. Cary, and C. Kenyon. 2010. Insulin/IGF-1 signaling mutants reprogram ER stress response regulators to promote longevity. *Proc. Natl. Acad. Sci. USA.* 107:9730–9735. <http://dx.doi.org/10.1073/pnas.1002575107>
- Hollien, J., J.H. Lin, H. Li, N. Stevens, P. Walter, and J.S. Weissman. 2009. Regulated Ire1-dependent decay of messenger RNAs in mammalian cells. *J. Cell Biol.* 186:323–331. <http://dx.doi.org/10.1083/jcb.200903014>
- Houtkooper, R.H., L. Mouchiroud, D. Ryu, N. Moullan, E. Katsyuba, G. Knott, R.W. Williams, and J. Auwerx. 2013. Mitonuclear protein imbalance as a conserved longevity mechanism. *Nature.* 497:451–457. <http://dx.doi.org/10.1038/nature12188>
- Hsu, A.L., C.T. Murphy, and C. Kenyon. 2003. Regulation of aging and age-related disease by DAF-16 and heat-shock factor. *Science.* 300:1142–1145. <http://dx.doi.org/10.1126/science.1083701>
- Jensen, M.B., and H. Jasper. 2014. Mitochondrial proteostasis in the control of aging and longevity. *Cell Metab.* 20:214–225. <http://dx.doi.org/10.1016/j.cmet.2014.05.006>
- Kimura, K.D., H.A. Tissenbaum, Y. Liu, and G. Ruvkun. 1997. daf-2, an insulin receptor-like gene that regulates longevity and diapause in *Caenorhabditis elegans*. *Science.* 277:942–946. <http://dx.doi.org/10.1126/science.277.5328.942>
- Korennykh, A.V., P.F. Egea, A.A. Korostelev, J. Finer-Moore, C. Zhang, K.M. Shokat, R.M. Stroud, and P. Walter. 2009. The unfolded protein response signals through high-order assembly of Ire1. *Nature.* 457:687–693. <http://dx.doi.org/10.1038/nature07661>
- Lapierre, L.R., C.D. De Magalhaes Filho, P.R. McQuary, C.C. Chu, O. Visvikis, J.T. Chang, S. Gelino, B. Ong, A.E. Davis, J.E. Irazoqui, et al. 2013. The TFEB orthologue HLH-30 regulates autophagy and modulates lon-

- gevity in *Caenorhabditis elegans*. *Nat. Commun.* 4:2267. <http://dx.doi.org/10.1038/ncomms3267>
- Lee, S.J., A.B. Hwang, and C. Kenyon. 2010. Inhibition of respiration extends *C. elegans* life span via reactive oxygen species that increase HIF-1 activity. *Curr. Biol.* 20:2131–2136. <http://dx.doi.org/10.1016/j.cub.2010.10.057>
- Lin, K., J.B. Dorman, A. Rodan, and C. Kenyon. 1997. daf-16: An HNF-3/ forkhead family member that can function to double the life-span of *Caenorhabditis elegans*. *Science.* 278:1319–1322. <http://dx.doi.org/10.1126/science.278.5341.1319>
- Lin, J.H., H. Li, Y. Zhang, D. Ron, and P. Walter. 2009. Divergent effects of PERK and IRE1 signaling on cell viability. *PLoS ONE.* 4:e4170. <http://dx.doi.org/10.1371/journal.pone.00004170>
- Lu, M., D.A. Lawrence, S. Marsters, D. Acosta-Alvear, P. Kimmig, A.S. Mendez, A.W. Paton, J.C. Paton, P. Walter, and A. Ashkenazi. 2014. Cell death. Opposing unfolded-protein-response signals converge on death receptor 5 to control apoptosis. *Science.* 345:98–101. <http://dx.doi.org/10.1126/science.1254312>
- Mendillo, M.L., S. Santagata, M. Koeva, G.W. Bell, R. Hu, R.M. Tamimi, E. Fraenkel, T.A. Ince, L. Whitesell, and S. Lindquist. 2012. HSF1 drives a transcriptional program distinct from heat shock to support highly malignant human cancers. *Cell.* 150:549–562. <http://dx.doi.org/10.1016/j.cell.2012.06.031>
- Morano, K.A., N. Santoro, K.A. Koch, and D.J. Thiele. 1999. A trans-activation domain in yeast heat shock transcription factor is essential for cell cycle progression during stress. *Mol. Cell. Biol.* 19:402–411.
- Morgan, W.D., G.T. Williams, R.I. Morimoto, J. Greene, R.E. Kingston, and R. Tjian. 1987. Two transcriptional activators, CCAAT-box-binding transcription factor and heat shock transcription factor, interact with a human hsp70 gene promoter. *Mol. Cell. Biol.* 7:1129–1138.
- Morimoto, R.I. 1998. Regulation of the heat shock transcriptional response: cross talk between a family of heat shock factors, molecular chaperones, and negative regulators. *Genes Dev.* 12:3788–3796. <http://dx.doi.org/10.1101/gad.12.24.3788>
- Morley, J.F., and R.I. Morimoto. 2004. Regulation of longevity in *Caenorhabditis elegans* by heat shock factor and molecular chaperones. *Mol. Biol. Cell.* 15:657–664. <http://dx.doi.org/10.1091/mbc.E03-07-0532>
- Mouchiroud, L., R.H. Houtkooper, N. Moulhan, E. Katsyuba, D. Ryu, C. Cantó, A. Mottis, Y.S. Jo, M. Viswanathan, K. Schoonjans, et al. 2013. The NAD<sup>+</sup>/Sirtuin pathway modulates longevity through activation of mitochondrial UPR and FOXO signaling. *Cell.* 154:430–441. <http://dx.doi.org/10.1016/j.cell.2013.06.016>
- Mu, T.W., D.S. Ong, Y.J. Wang, W.E. Balch, J.R. Yates III, L. Segatori, and J.W. Kelly. 2008. Chemical and biological approaches synergize to ameliorate protein-folding diseases. *Cell.* 134:769–781. <http://dx.doi.org/10.1016/j.cell.2008.06.037>
- Nargund, A.M., M.W. Pellegrino, C.J. Fiorese, B.M. Baker, and C.M. Haynes. 2012. Mitochondrial import efficiency of ATFS-1 regulates mitochondrial UPR activation. *Science.* 337:587–590. <http://dx.doi.org/10.1126/science.1223560>
- Nargund, A.M., C.J. Fiorese, M.W. Pellegrino, P. Deng, and C.M. Haynes. 2015. Mitochondrial and nuclear accumulation of the transcription factor ATFS-1 promotes OXPHOS recovery during the UPR(mt). *Mol. Cell.* 58:123–133. <http://dx.doi.org/10.1016/j.molcel.2015.02.008>
- Novoa, I., H. Zeng, H.P. Harding, and D. Ron. 2001. Feedback inhibition of the unfolded protein response by GADD34-mediated dephosphorylation of eIF2 $\alpha$ . *J. Cell Biol.* 153:1011–1022.
- Owusu-Ansah, E., W. Song, and N. Perrimon. 2013. Muscle mitohormesis promotes longevity via systemic repression of insulin signaling. *Cell.* 155:699–712. <http://dx.doi.org/10.1016/j.cell.2013.09.021>
- Pellegrino, M.W., A.M. Nargund, N.V. Kirienko, R. Gillis, C.J. Fiorese, and C.M. Haynes. 2014. Mitochondrial UPR-regulated innate immunity provides resistance to pathogen infection. *Nature.* 516:414–417. <http://dx.doi.org/10.1038/nature13818>
- Powers, E.T., R.I. Morimoto, A. Dillin, J.W. Kelly, and W.E. Balch. 2009. Biological and chemical approaches to diseases of proteostasis deficiency. *Annu. Rev. Biochem.* 78:959–991. <http://dx.doi.org/10.1146/annurev.biochem.052308.114844>
- Prahlad, V., and R.I. Morimoto. 2011. Neuronal circuitry regulates the response of *Caenorhabditis elegans* to misfolded proteins. *Proc. Natl. Acad. Sci. USA.* 108:14204–14209. <http://dx.doi.org/10.1073/pnas.1106557108>
- Prahlad, V., T. Cornelius, and R.I. Morimoto. 2008. Regulation of the cellular heat shock response in *Caenorhabditis elegans* by thermosensory neurons. *Science.* 320:811–814. <http://dx.doi.org/10.1126/science.1156093>
- Qian, S.B., H. McDonough, F. Boellmann, D.M. Cyr, and C. Patterson. 2006. CHIP-mediated stress recovery by sequential ubiquitination of substrates and Hsp70. *Nature.* 440:551–555. <http://dx.doi.org/10.1038/nature04600>
- Rauthan, M., P. Ranji, N. Aguilera Pradenas, C. Pitot, and M. Pilon. 2013. The mitochondrial unfolded protein response activator ATFS-1 protects cells from inhibition of the mevalonate pathway. *Proc. Natl. Acad. Sci. USA.* 110:5981–5986. <http://dx.doi.org/10.1073/pnas.1218778110>
- Raychaudhuri, S., C. Loew, R. Körner, S. Pinkert, M. Theis, M. Hayer-Hartl, F. Buchholz, and F.U. Hartl. 2014. Interplay of acetyltransferase EP300 and the proteasome system in regulating heat shock transcription factor 1. *Cell.* 156:975–985. <http://dx.doi.org/10.1016/j.cell.2014.01.055>
- Rea, S.L., N. Ventura, and T.E. Johnson. 2007. Relationship between mitochondrial electron transport chain dysfunction, development, and life extension in *Caenorhabditis elegans*. *PLoS Biol.* 5:e259. <http://dx.doi.org/10.1371/journal.pbio.0050259>
- Richardson, C.E., T. Kooistra, and D.H. Kim. 2010. An essential role for XBP-1 in host protection against immune activation in *C. elegans*. *Nature.* 463:1092–1095. <http://dx.doi.org/10.1038/nature08762>
- Roth, D.M., D.M. Hutt, J. Tong, M. Boucheareilh, N. Wang, T. Seeley, J.F. Dekkers, J.M. Beekman, D. Garza, L. Drew, et al. 2014. Modulation of the maladaptive stress response to manage diseases of protein folding. *PLoS Biol.* 12:e1001998. <http://dx.doi.org/10.1371/journal.pbio.1001998>
- Rubio, C., D. Pincus, A. Korenykh, S. Schuck, H. El-Samad, and P. Walter. 2011. Homeostatic adaptation to endoplasmic reticulum stress depends on Ire1 kinase activity. *J. Cell Biol.* 193:171–184.
- Ryno, L.M., R.L. Wiseman, and J.W. Kelly. 2013. Targeting unfolded protein response signaling pathways to ameliorate protein misfolding diseases. *Curr. Opin. Chem. Biol.* 17:346–352. <http://dx.doi.org/10.1016/j.cbpa.2013.04.009>
- Ryno, L.M., J.C. Genereux, T. Naito, R.I. Morimoto, E.T. Powers, M.D. Shoulders, and R.L. Wiseman. 2014. Characterizing the altered cellular proteome induced by the stress-independent activation of heat shock factor 1. *ACS Chem. Biol.* 9:1273–1283. <http://dx.doi.org/10.1021/cb500062n>
- Santagata, S., R. Hu, N.U. Lin, M.L. Mendillo, L.C. Collins, S.E. Hankinson, S.J. Schnitt, L. Whitesell, R.M. Tamimi, S. Lindquist, and T.A. Ince. 2011. High levels of nuclear heat-shock factor 1 (HSF1) are associated with poor prognosis in breast cancer. *Proc. Natl. Acad. Sci. USA.* 108:18378–18383. <http://dx.doi.org/10.1073/pnas.1115031108>
- Schieber, M., and N.S. Chandel. 2014. TOR signaling couples oxygen sensing to lifespan in *C. elegans*. *Cell Reports.* 9:9–15. <http://dx.doi.org/10.1016/j.celrep.2014.08.075>
- Shen, X., R.E. Ellis, K. Lee, C.Y. Liu, K. Yang, A. Solomon, H. Yoshida, R. Morimoto, D.M. Kurnit, K. Mori, and R.J. Kaufman. 2001. Complementary pathways regulate the unfolded protein response and are required for *C. elegans* development. *Cell.* 107:893–903. [http://dx.doi.org/10.1016/S0092-8674\(01\)00612-2](http://dx.doi.org/10.1016/S0092-8674(01)00612-2)
- Shi, Y., D.D. Mosser, and R.I. Morimoto. 1998. Molecular chaperones as HSF1-specific transcriptional repressors. *Genes Dev.* 12:654–666. <http://dx.doi.org/10.1101/gad.12.5.654>
- Shoulders, M.D., L.M. Ryno, J.C. Genereux, J.J. Moresco, P.G. Tu, C. Wu, J.R. Yates III, A.I. Su, J.W. Kelly, and R.L. Wiseman. 2013. Stress-independent activation of XBP1s and/or ATF6 reveals three functionally diverse ER proteostasis environments. *Cell Reports.* 3:1279–1292. <http://dx.doi.org/10.1016/j.celrep.2013.03.024>
- Steele, A.D., G. Hutter, W.S. Jackson, F.L. Heppner, A.W. Borkowski, O.D. King, G.J. Raymond, A. Aguzzi, and S. Lindquist. 2008. Heat shock factor 1 regulates lifespan as distinct from disease onset in prion disease. *Proc. Natl. Acad. Sci. USA.* 105:13626–13631. <http://dx.doi.org/10.1073/pnas.0806319105>
- Tabas, I., and D. Ron. 2011. Integrating the mechanisms of apoptosis induced by endoplasmic reticulum stress. *Nat. Cell Biol.* 13:184–190. <http://dx.doi.org/10.1038/ncb0311-184>
- Taylor, R.C., and A. Dillin. 2013. XBP-1 is a cell-nonautonomous regulator of stress resistance and longevity. *Cell.* 153:1435–1447. <http://dx.doi.org/10.1016/j.cell.2013.05.042>
- Todd-Corlett, A., E. Jones, C. Seghers, and M.J. Gething. 2007. Lobe IB of the ATPase domain of Kar2p/BiP interacts with Ire1p to negatively regulate the unfolded protein response in *Saccharomyces cerevisiae*. *J. Mol. Biol.* 367:770–787. <http://dx.doi.org/10.1016/j.jmb.2007.01.009>
- Topol, J., D.M. Ruden, and C.S. Parker. 1985. Sequences required for in vitro transcriptional activation of a *Drosophila* hsp 70 gene. *Cell.* 42:527–537. [http://dx.doi.org/10.1016/0092-8674\(85\)90110-2](http://dx.doi.org/10.1016/0092-8674(85)90110-2)
- Travers, K.J., C.K. Patil, L. Wodicka, D.J. Lockhart, J.S. Weissman, and P. Walter. 2000. Functional and genomic analyses reveal an essential coordination between the unfolded protein response and ER-associated degradation. *Cell.* 101:249–258. [http://dx.doi.org/10.1016/S0092-8674\(00\)80835-1](http://dx.doi.org/10.1016/S0092-8674(00)80835-1)
- Tsaytler, P., H.P. Harding, D. Ron, and A. Bertolotti. 2011. Selective inhibition of a regulatory subunit of protein phosphatase 1 restores proteostasis. *Science.* 332:91–94. <http://dx.doi.org/10.1126/science.1201396>

- Upton, J.P., L. Wang, D. Han, E.S. Wang, N.E. Huskey, L. Lim, M. Truitt, M.T. McManus, D. Ruggero, A. Goga, et al. 2012. IRE1 $\alpha$  cleaves select microRNAs during ER stress to derepress translation of proapoptotic Caspase-2. *Science*. 338:818–822. <http://dx.doi.org/10.1126/science.1226191>
- Vander Heiden, M.G., L.C. Cantley, and C.B. Thompson. 2009. Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science*. 324:1029–1033. <http://dx.doi.org/10.1126/science.1160809>
- Walter, P., and D. Ron. 2011. The unfolded protein response: from stress pathway to homeostatic regulation. *Science*. 334:1081–1086. <http://dx.doi.org/10.1126/science.1209038>
- Walter, L., A. Baruah, H.W. Chang, H.M. Pace, and S.S. Lee. 2011. The homeobox protein CEH-23 mediates prolonged longevity in response to impaired mitochondrial electron transport chain in *C. elegans*. *PLoS Biol*. 9:e1001084. <http://dx.doi.org/10.1371/journal.pbio.1001084>
- Wang, X., J. Venable, P. LaPointe, D.M. Hutt, A.V. Koulov, J. Coppinger, C. Gurkan, W. Kellner, J. Matteson, H. Plutner, et al. 2006. Hsp90 cochaperone Aha1 downregulation rescues misfolding of CFTR in cystic fibrosis. *Cell*. 127:803–815. <http://dx.doi.org/10.1016/j.cell.2006.09.043>
- Way, S.W., J.R. Podojil, B.L. Clayton, A. Zaremba, T.L. Collins, R.B. Kunjamma, A.P. Robinson, P. Brugarolas, R.H. Miller, S.D. Miller, and B. Popko. 2015. Pharmaceutical integrated stress response enhancement protects oligodendrocytes and provides a potential multiple sclerosis therapeutic. *Nat. Commun*. 6:6532. <http://dx.doi.org/10.1038/ncomms7532>
- Westerheide, S.D., J. Anckar, S.M. Stevens Jr., L. Sistonen, and R.I. Morimoto. 2009. Stress-inducible regulation of heat shock factor 1 by the deacetylase SIRT1. *Science*. 323:1063–1066. <http://dx.doi.org/10.1126/science.1165946>
- Wright, G., K. Terada, M. Yano, I. Sergeev, and M. Mori. 2001. Oxidative stress inhibits the mitochondrial import of preproteins and leads to their degradation. *Exp. Cell Res*. 263:107–117. <http://dx.doi.org/10.1006/excr.2000.5096>
- Xiao, X., X. Zuo, A.A. Davis, D.R. McMillan, B.B. Curry, J.A. Richardson, and I.J. Benjamin. 1999. HSF1 is required for extra-embryonic development, postnatal growth and protection during inflammatory responses in mice. *EMBO J*. 18:5943–5952. <http://dx.doi.org/10.1093/emboj/18.21.5943>
- Yoneda, T., C. Benedetti, F. Urano, S.G. Clark, H.P. Harding, and D. Ron. 2004. Compartment-specific perturbation of protein handling activates genes encoding mitochondrial chaperones. *J. Cell Sci*. 117:4055–4066. <http://dx.doi.org/10.1242/jcs.01275>
- Yoshida, H., T. Matsui, A. Yamamoto, T. Okada, and K. Mori. 2001. XBP1 mRNA is induced by ATF6 and spliced by IRE1 in response to ER stress to produce a highly active transcription factor. *Cell*. 107:881–891. [http://dx.doi.org/10.1016/S0092-8674\(01\)00611-0](http://dx.doi.org/10.1016/S0092-8674(01)00611-0)
- Zou, J., Y. Guo, T. Guettouche, D.F. Smith, and R. Voellmy. 1998. Repression of heat shock transcription factor HSF1 activation by HSP90 (HSP90 complex) that forms a stress-sensitive complex with HSF1. *Cell*. 94:471–480. [http://dx.doi.org/10.1016/S0092-8674\(00\)81588-3](http://dx.doi.org/10.1016/S0092-8674(00)81588-3)