

## ***Listeria monocytogenes* Mutants Lacking Phosphatidylinositol-specific Phospholipase C Are Avirulent**

By Andrew Camilli, Howard Goldfine, and Daniel A. Portnoy

From the Department of Microbiology, University of Pennsylvania School of Medicine, Philadelphia, Pennsylvania 19104

### **Summary**

A number of bacterial species secrete phosphatidylinositol-specific phospholipase C (PI-PLC). In this report, we show that the facultative intracellular bacterial pathogen, *Listeria monocytogenes*, contains a gene, *plcA*, predicting a polypeptide with 31% amino acid identity to a *Bacillus thuringiensis* PI-PLC. Accordingly, *L. monocytogenes* secretes PI-PLC activity, while a mutant with a transposon insertion in *plcA* lacks detectable PI-PLC activity. In addition, expression of *plcA* in *B. subtilis* resulted in secretion of PI-PLC activity. The *L. monocytogenes* PI-PLC-defective mutant was three logs less virulent for mice and failed to grow in host tissues. The mutant was also defective for in vitro growth in mouse peritoneal macrophages. These results strongly suggest that PI-PLC is an essential determinant of *L. monocytogenes* pathogenesis. Whether the PI-PLC acts on a bacterial or host substrate remains to be determined.

*Listeria monocytogenes*, a facultative intracellular bacterial pathogen, has been extensively studied as a model for infections that are resolved by cell-mediated immunity. Recently, the cell biology of listerial intracellular parasitism has been documented and shown to involve entry of bacteria into the host-cell cytoplasm and cell-to-cell spread (1). Until now, the only proven virulence determinant of *L. monocytogenes* was a secreted pore-forming hemolysin, called listeriolysin O, which has been shown to mediate lysis of the host phagosome (1–3). In a recent study, we reported the isolation of a class of transposon mutants that were normal for hemolysin production and growth in tissue culture cell lines yet were defective for cell-to-cell spread (4). These mutants had insertions in a previously reported open reading frame (5), which we now designate *plcA*. In this report, we show that *plcA* encodes a secreted phosphatidylinositol-specific phospholipase C (PI-PLC), the loss of which results in the inability of the mutant bacterium to propagate in host tissues.

### **Materials and Methods**

**Bacterial Strains.** *L. monocytogenes* 10403S (6) was the wild-type parental strain in these studies. Strain DP-L1054 is a Tn917-LTV3 insertion mutant isolated as a small-plaque former in tissue culture cells (4). Strain DP-L1044 contains a Tn917-LTV3 insertion within the listeriolysin O structural gene (4).


**Construction of *Bacillus subtilis* Strain Expressing Phosphatidylinositol-Specific Phospholipase C.** The *L. monocytogenes plcA* gene was cloned into an asporogenic strain of *B. subtilis* under the control of an isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG)-inducible promoter, essentially as described (2). The *plcA* gene, bases 279–1349 (5), was

amplified from *L. monocytogenes* 10403S chromosomal DNA using PCR, and ligated into the IPTG-inducible promoter using SalI- and XbaI-generated DNA ends. The SalI and XbaI restriction sites of the PCR-amplified product were created as noncomplementary ends of the amplification primers. PCR primers used were 5'-GGG-TCGACTAGGAATAATATATGTTAGTTG-3' and 5'-GGTCTAGATTCTAGTCCTGCTGTCC-3' (Genosys Biotechnologies, Inc., The Woodlands, TX).

**Assay for Phosphatidylinositol-Specific Phospholipase C Activity.** Culture supernatants of *L. monocytogenes* strains were prepared by overnight growth to stationary phase ( $OD_{600} = 2.1$ ) in brain heart infusion (BHI) broth at 38°C with aeration and from *B. subtilis* by growth at 37°C with aeration to early stationary phase ( $OD_{600} = 1.5$ ) in BHI supplemented with 1 mM IPTG. Aliquots of supernatant fluid were then assayed for PI hydrolysis activity essentially as described (7) except that [ $^3$ H-inositol] PI, 0.045  $\mu$ Ci, 17.4 Ci/mmol (Amersham Corp., Arlington Heights, IL), and 13.5  $\mu$ g PI (Sigma Chemical Co., St. Louis, MO) were sonicated in 100  $\mu$ l of 40 mM Tris-HCl, pH 7.2, and 0.2% deoxycholate (per assay), and the PI-sonicate was then incubated with culture supernatant in a final volume of 200  $\mu$ l at 37°C.

**Chromatography of Phosphatidylinositol Hydrolysis Products.** The lipid product of PI (Bovine liver; Sigma Chemical Co.) hydrolysis was chromatographed on Silica gel 60 (E. Merck, Darmstadt, FRG) TLC plates in petroleum ether/ethyl ether/acetic acid, 50:50:1 (vol/vol/vol), and also in hexane/ethyl ether/formic acid 9:6:0.4 (vol/vol/vol). The water-soluble product of PI hydrolysis obtained after partitioning the chloroform-methanol phase with either water or 4% TCA was chromatographed on cellulose (E. Merck) TLC plates in *n*-propanol/ $NH_4OH/H_2O$ ; 5:4:1 (vol/vol/vol) or in 95% ethanol/13.5 M  $NH_4OH$ , 3:2 (vol/vol). The acidified product was also chromatographed on an anion exchange column (8).






**Figure 2.** Growth of wild-type *L. monocytogenes* and DP-L1054 (PI-PLC<sup>-</sup>) in murine liver and spleen. Data points and error bars represent the mean and SD of the number of viable bacteria recovered from whole spleen or liver from five mice.

host tissues was reflected in a three-log increase in LD<sub>50</sub> (Table 1).

**Growth of *L. monocytogenes* in Mouse Peritoneal Macrophages.** Growth of both the parental and mutant strains in vivo was mirrored in vitro in primary cultures of mouse resident peritoneal macrophages (Fig. 3). As previously reported (10) for the parental strain, ~70% of the infecting bacteria were killed by the macrophages followed by subsequent intracellular growth of the survivors. In contrast, a hemolysin-minus mutant, which fails to lyse the phagosomal membrane, showed a continual decline in viable count (Fig. 3). The PI-PLC mutant, DP-L1054, had an intermediate level of viability, but showed no evidence for bacterial growth (Fig. 3). However, microscopic examination of monolayers 5 h after infection with DP-L1054 revealed rare foci of infection resulting from growth of the bacteria in the cytoplasm (data not shown). The most likely explanation for these results is that the PI-PLC mutant was somewhat defective in escaping from the host phagosome although not to the extent seen in the hemolysin-minus mutant.

## Discussion

A number of bacterial species secrete PI-PLCs, including *Staphylococcus aureus*, *Clostridium novyi*, *Bacillus cereus*, and *B. thuringiensis* (13). These enzymes have proven to be valuable reagents for the identification and characterization of a family of eukaryotic membrane proteins anchored to the cell surface by PI-glycan (13). However, a role for PI-PLC in bacterial growth or pathogenicity was previously unknown. The results of this study strongly suggest that the PI-PLC secreted by *L. monocytogenes* is an essential determinant of pathogenicity since a mutant with a transposon insertion in the structural gene encoding PI-PLC had reduced virulence and failed to grow in host tissues. Since *plcA* is monocistronic (5) it is unlikely that the mutation directly affects expression of other genes. In addition, an independent mutant with a transposon insertion in *plcA* had identical characteristics in tissue culture and the same increase in LD<sub>50</sub> as DP-L1054.



**Figure 3.** Growth of wild-type *L. monocytogenes*, DP-L1054 (PI-PLC<sup>-</sup>), and hemolysin-minus mutant DP-L1044 (hly<sup>-</sup>) in murine resident peritoneal macrophages. Data points and error bars represent the mean and SD of the number of viable bacteria recovered from three coverslips.

However, absolute proof for a role of PI-PLC in pathogenesis awaits fulfillment of a molecular version of Koch's postulates, by reintroducing *plcA* into DP-L1054 and showing that virulence is restored (14).

The major question that arises from this study is, what is the role of PI-PLC in pathogenicity and, more specifically, what is the biological substrate(s) for this enzyme? The results presented here indicated that PI-PLC is required for growth in host tissues which may reflect a role in lysis of the macrophage vacuole and, as previously shown (4), subsequent cell-to-cell spread. It is possible that PI-PLC acts in concert with hemolysin and perhaps a distinct PLC secreted by *L. monocytogenes* (15), to mediate entry into the host cytoplasm and to solubilize host membrane(s) during cell-to-cell spread. Thus, the biological substrates for *L. monocytogenes* PI-PLC may be host PI or PI-glycan in the phagosomal membrane and PI or its phosphorylated derivatives in the cytoplasmic membrane. Recognition of the latter substrates is an intriguing possibility because of their roles in signal transduction in eukaryotic cells (13). However, other bacterial PI-PLCs, although capable of hydrolyzing PI and PI-glycan, do not hydrolyze phosphorylated derivatives of PI. The precise substrate-specificity of the *L. monocytogenes* PI-PLC awaits its purification.

Lastly, it is formally possible that the listerial PI-PLC is acting on a bacterial substrate such as a PI-anchored bacterial protein. *L. monocytogenes* has been reported to lack PI (16), and there is no precedence for PI-anchored proteins in prokaryotes. Nevertheless, recent evidence has been provided for the presence of PI-anchored polysaccharides, the lipoarabinomannan and lipomannan in *Mycobacterium tuberculosis* (17). In addition, it has been pointed out that a number of surface proteins from Gram-positive cocci have a conserved hexapeptide with homology to the proposed cleavage and PI-glycan addition site found in some eukaryotic PI-glycan anchored proteins (e.g., the cellular adhesion molecules N-CAM and LFA-3) (18). Thus, there may be a prokaryotic version of a PI-anchor. Whether the *L. monocytogenes* PI-PLC acts on a host or bacterial substrate awaits further study.

---

We thank R. Barry and D. Hinrichs for the LD<sub>50</sub> determinations and R. North for collaborating on the in vivo growth curves.

This work was supported by National Institutes of Health grants AI-26919 and AI-27655 (D. A. Portnoy) and by Biomedical Research Support Grants 2-S07-RR-07083 and S07-RR-05415 (H. Goldfine). A. Camilli was supported by National Institutes of Health Training Grant GM-08216.

Address correspondence to Daniel A. Portnoy, 209 Johnson Pavilion, University of Pennsylvania School of Medicine, Philadelphia, PA 19104-6076.

Received for publication 3 December 1990.

## References

1. Tilney, L.G., and D.A. Portnoy. 1989. Actin filaments and the growth, movement, and spread of the intracellular bacterial parasite, *Listeria monocytogenes*. *J. Cell Biol.* 109:1597.
2. Bielecki, J., P. Youngman, P. Connelly, and D.A. Portnoy. 1990. *Bacillus subtilis* expressing a haemolysin gene from *Listeria monocytogenes*, can grow in mammalian cells. *Nature (Lond.)* 345:175.
3. Gaillard, J.L., P. Berche, J. Mounier, S. Richard, and P. Sansonetti. 1987. In vitro model of penetration and intracellular growth of *Listeria monocytogenes* in the human enterocyte-like cell line Caco-2. *Infect. Immun.* 55:2822.
4. Sun, A.N., A. Camilli, and D.A. Portnoy. 1990. Isolation of *Listeria monocytogenes* small-plaque mutants defective for intracellular growth and cell-to-cell spread. *Infect. Immun.* 58:3770.
5. Mengaud, J., M.F. Vicente, and P. Cossart. 1989. Transcriptional mapping and nucleotide sequence of the *Listeria monocytogenes* *hlyA* region reveal structural features that may be involved in regulation. *Infect. Immun.* 57:3695.
6. Portnoy, D.A., P.S. Jacks, and D.J. Hinrichs. 1988. Role of hemolysin for the intracellular growth of *Listeria monocytogenes*. *J. Exp. Med.* 167:1459.
7. Low, M.G. 1981. Phosphatidylinositol-specific phospholipase C from *Staphylococcus aureus*. *Methods Enzymol.* 71:741.
8. Berridge, M.J., C.P. Downes, and M.R. Hanley. 1982. Lithium amplifies agonist-dependent phosphatidyl responses in brain and salivary glands. *Biochem. J.* 206:587.
9. Newborg, M.F., and R.J. North. 1980. On the mechanism of T cell-dependent anti-listeria resistance in nude mice. *J. Immunol.* 124:571.
10. Portnoy, D.A., R.D. Schreiber, P. Connelly, and L.G. Tilney. 1989.  $\gamma$  Interferon limits access of *Listeria monocytogenes* to the macrophage cytoplasm. *J. Exp. Med.* 170:2141.
11. Lipman, D.J., and W.R. Pearson. 1985. Rapid and sensitive protein similarity searches. *Science (Wash. DC)* 227:1435.
12. Henner, D.J., M. Yang, E. Chen, R. Hellmiss, H. Rodriguez, and M.G. Low. 1988. Sequence of the *Bacillus thuringiensis* phosphatidylinositol specific phospholipase C. *Nucleic Acids Res.* 16:10383.
13. Low, M.G. 1989. The glycosyl-phosphatidylinositol anchor of membrane proteins. *Biochim. Biophys. Acta.* 988:427.
14. Falkow, S. 1988. Molecular Koch's postulates applied to microbial pathogenicity. *Rev. Infect. Dis.* 10:S274.
15. Barclay, R., D.R. Threlfall, and I. Leighton. 1989. Haemolysins and extracellular enzymes of *Listeria monocytogenes* and *L. ivanovii*. *J. Med. Microbiol.* 30:111.
16. Kosaric, N., and K.K. Carroll. 1971. Phospholipids of *Listeria monocytogenes*. *Biochim. Biophys. Acta.* 239:428.
17. Hunter, S.W., and P.J. Brennan. 1990. Evidence for the presence of a phosphatidylinositol anchor on the lipoarabinomannan and lipomannan of *Mycobacterium tuberculosis*. *J. Biol. Chem.* 265:9272.
18. Fischetti, V.A., V. Pancholi, and O. Schneewind. 1990. Conservation of a hexapeptide sequence in the anchor region of surface proteins from gram-positive cocci. *Mol. Microbiol.* 4:1603.