

# Cycles in the chamber homology for $\mathrm{GL}(3)$

Anne-Marie Aubert, Samir Hasan and Roger Plymen

## Abstract

Let  $F$  be a nonarchimedean local field and let  $\mathrm{GL}(N) = \mathrm{GL}(N, F)$ . We prove the existence of parahoric types for  $\mathrm{GL}(N)$ . We construct representative cycles in all the homology classes of the chamber homology of  $\mathrm{GL}(3)$ .

## 1 Introduction

Let  $F$  be a nonarchimedean local field and let  $G = \mathrm{GL}(N) = \mathrm{GL}(N, F)$ . The enlarged building  $\beta^1 G$  of  $G$  is a polysimplicial complex on which  $G$  acts properly. We select a chamber  $C \subset \beta^1 G$ . This chamber is a polysimplex, the product of an  $n$ -simplex by a 1-simplex:

$$C = \Delta_n \times \Delta_1.$$

To this datum we will attach a *homological coefficient system*, see [13, p.11]. To each simplex  $\sigma$  in  $\Delta_n$  we attach the representation ring  $R(G(\sigma))$  of the stabilizer  $G(\sigma)$ , and to each inclusion  $\sigma \rightarrow \tau$  we attach the induction map:

$$\mathrm{Ind}_{G(\sigma)}^{G(\tau)} : R(G(\sigma)) \rightarrow R(G(\tau)).$$

This creates the homology of the simplicial set  $\Delta_n$  with the above coefficient system. The resulting homology groups are denoted  $h_j(G)$ ,  $0 \leq j \leq N - 1$ .

For each point  $\mathfrak{s}$  in the Bernstein spectrum  $\mathfrak{B}(G)$  (see appendix B) we will select an  $\mathfrak{s}$ -type  $(J, \tau)$ . Here,  $J$  denotes a certain compact open subgroup of  $G$ , and  $\tau$  denotes a certain irreducible smooth representation of  $J$ , see [8, 9, 10].

The following result is due to Bushnell-Kutzko [8, 9, 10].

**Theorem 1.** Existence of types. *Let  $\mathfrak{s} \in \mathfrak{B}(G)$ . Then there exists an  $\mathfrak{s}$ -type  $(J, \tau)$ .*

An  $\mathfrak{s}$ -type  $(J, \tau)$  is of *multiplicity 1* if, for each  $\pi \in \text{Irr}(G)$  with inertial support  $\mathfrak{s}$ , the restriction of  $\pi$  to  $J$  contains  $\tau$  with multiplicity 1. The next result is due to Paskunas [16].

**Theorem 2.** Multiplicity one in the supercuspidal case. *Let  $\sigma$  be an irreducible supercuspidal representation of  $G$ , and let  $\mathfrak{s} = [G, \sigma]_G$ . Then there exists an  $\mathfrak{s}$ -type  $(\text{GL}(N, \mathfrak{o}_F), \tau)$ . The representation  $\tau$  is unique up to equivalence, and is of multiplicity 1.*

Let  $\mathfrak{s} = [M, \sigma]_G \in \mathfrak{B}(G)$ . An  $\mathfrak{s}$ -type  $(J, \lambda)$  will be called *parahoric* if  $J$  is a parahoric subgroup of  $G$ .

We present a refinement of Theorem 1.

**Theorem 3.** Existence of parahoric types. *Let  $\mathfrak{s} \in \mathfrak{B}(G)$ . Then there exists a parahoric  $\mathfrak{s}$ -type  $(J^{\mathfrak{s}}, \tau)$ .*

The parahoric subgroup  $J^{\mathfrak{s}}$  only depends on certain invariants attached to  $\mathfrak{s}$ . For details of these invariants, see appendix D.

In the proof of Theorem 3, we have to call upon several of the technical resources developed by Bushnell-Kutzko.

We now specialize to  $\text{GL}(3)$ . In this article, we will explicitly construct representative cycles in *all* the homology classes in  $h_0(G) \oplus h_1(G) \oplus h_2(G)$  when  $G = \text{GL}(3)$ . This allows us to compute the chamber homology groups of  $\text{GL}(3)$  according to the following formulas:

$$H_{\text{ev}}(G; \beta^1 G) = h_0(G) \oplus h_1(G) \oplus h_2(G) = H_{\text{odd}}(G; \beta^1 G).$$

We will demonstrate that each parahoric  $\mathfrak{s}$ -type  $(J^{\mathfrak{s}}, \tau)$  creates finitely many cycles in  $h_0(G) \oplus h_1(G) \oplus h_2(G)$ . To prove that all homology classes in  $h_0(G) \oplus h_1(G) \oplus h_2(G)$  are thereby accounted for, we invoke the  $K$ -theory of the reduced  $C^*$ -algebra  $\mathcal{A} := C_r^*(G)$ . The  $K$ -theory is torsion-free [17].

The abelian groups  $H_{\text{ev/odd}}(G; \beta^1 G)$  and  $K_j(\mathcal{A})$  admit compatible Bernstein decompositions, see appendix B. This leads, for each  $\mathfrak{s} \in \mathfrak{B}(G)$ , to the equalities

$$\text{rank } H_{\text{ev/odd}}(G; \beta^1 G)^{\mathfrak{s}} = \text{rank } K_0(\mathcal{A}^{\mathfrak{s}}) = \text{rank } K_1(\mathcal{A}^{\mathfrak{s}}). \quad (1)$$

The ranks of the finitely generated abelian groups on the right-hand-side are easily computed (see appendix C).

**Theorem 4.** *Let  $G = \text{GL}(3)$ , and let  $\mathfrak{s} = [M, \sigma]_G$ . Each parahoric  $\mathfrak{s}$ -type  $(J^{\mathfrak{s}}, \tau)$  creates finitely many cycles in  $h_0(G) \oplus h_1(G) \oplus h_2(G)$ , and all homology classes in  $h_0(G) \oplus h_1(G) \oplus h_2(G)$  are thereby accounted for. Quite specifically, we have*

- if  $M = \mathrm{GL}(3)$  then

$$\mathrm{H}_{\mathrm{ev}}(G; \beta^1 G)^{\mathfrak{s}} = \mathbb{Z} = \mathrm{H}_{\mathrm{odd}}(G; \beta^1 G)^{\mathfrak{s}}$$

- if  $M = \mathrm{GL}(2) \times \mathrm{GL}(1)$  then

$$\mathrm{H}_{\mathrm{ev}}(G; \beta^1 G)^{\mathfrak{s}} = \mathbb{Z}^2 = \mathrm{H}_{\mathrm{odd}}(G; \beta^1 G)^{\mathfrak{s}}$$

- if  $M = \mathrm{GL}(1) \times \mathrm{GL}(1) \times \mathrm{GL}(1)$  then

$$\mathrm{H}_{\mathrm{ev}}(G; \beta^1 G)^{\mathfrak{s}} = \mathbb{Z}^4 = \mathrm{H}_{\mathrm{odd}}(G; \beta^1 G)^{\mathfrak{s}}$$

From this point of view, the types for  $\mathrm{GL}(3)$  exceed their original expectations. Let  $\widehat{\mathcal{A}}^{\mathfrak{s}}$  denote the dual of the  $C^*$ -algebra  $\mathcal{A}^{\mathfrak{s}}$ . This is a compact Hausdorff space. Since  $K$ -theory for unital  $C^*$ -algebras is compatible with topological  $K$ -theory of compact Hausdorff spaces, we have

$$K_j(\mathcal{A}^{\mathfrak{s}}) \cong K^j(\widehat{\mathcal{A}}^{\mathfrak{s}}).$$

Therefore, the  $\mathfrak{s}$ -type also computes the topological  $K$ -theory of the compact space  $\widehat{\mathcal{A}}^{\mathfrak{s}}$ . The space  $\widehat{\mathcal{A}}^{\mathfrak{s}}$  is precisely the space of all those tempered representations of  $\mathrm{GL}(3)$  which have inertial support  $\mathfrak{s}$ .

Preliminary work in this direction was done with Paul Baum and Nigel Higson, and recorded in [4]. The diagrams in [4] are relevant to the present article. In [4] all computations were in the *tame* case. We confront here the general case: this is much more technical. We require much detailed information in the theory of types; in particular we need detailed information concerning *compact* intertwining sets.

We thank the referees for their detailed and constructive comments.

## 2 General results on types

We will collect here some general results on types which will be used in the paper. In this section  $G$  denotes the group of  $F$ -points of an arbitrary reductive connected algebraic group  $\mathbf{G}$  defined over  $F$ .

Let  $\mathfrak{R}(G)$  denote the category of smooth complex representations of  $G$ . Recall that, for each irreducible smooth representation  $\pi$  of  $G$ , there exists a Levi subgroup  $L$  of a parabolic subgroup  $P$  of  $G$  and an irreducible supercuspidal representation  $\sigma$  of  $L$  such that  $\pi$  is equivalent to a subquotient of the parabolically induced representation  $I_P^G(\sigma)$ . The pair  $(L, \sigma)$  is unique up

to conjugacy and the inertial class  $\mathfrak{s} = [M, \sigma]_G$  (see appendix B) is called the *inertial support* of  $\pi$ .

We have the standard decomposition (see [5, (2.10)])

$$\mathfrak{R}(G) = \prod_{\mathfrak{s} \in \mathfrak{B}(G)} \mathfrak{R}^{\mathfrak{s}}(G) \quad (2)$$

into full sub-categories, where the objects of  $\mathfrak{R}^{\mathfrak{s}}(G)$  are those smooth representations of  $G$  all whose irreducible subquotient have inertial support  $\mathfrak{s}$ .

Let  $\mathfrak{s}$  be a point in the Bernstein spectrum of  $G$ , and let  $(J, \tau)$  be an  $\mathfrak{s}$ -type, *i.e.*,  $\tau$  is an irreducible smooth representation of an open compact subgroup  $J$  of  $G$  such that for any irreducible smooth representation  $\pi$  of  $G$ , the restriction of  $\pi$  to  $J$  contains  $\tau$  if and only if  $\pi$  is an object of  $\mathfrak{R}^{\mathfrak{s}}(G)$ , [9, (4.2)]. When  $G = \mathrm{GL}(N, F)$ , it has been proved [8, 10] that there exists an  $\mathfrak{s}$ -type for each point  $\mathfrak{s}$  in  $\mathfrak{B}(G)$ .

**Proposition 1.** *Let  $K \supset J$  be an open compact subgroup of  $G$  such that  $\alpha := \mathrm{Ind}_J^K \tau$  is irreducible. Then  $(K, \alpha)$  is an  $\mathfrak{s}$ -type.*

*Proof.* Let  $\pi$  be an irreducible smooth representation of  $G$ . Using Frobenius reciprocity, we see that

$$\mathrm{Hom}_K(\alpha, \mathrm{Res}_K^G(\pi)) = \mathrm{Hom}_J(\tau, \mathrm{Res}_J^G(\pi)).$$

The result follows from the definition of  $\mathfrak{s}$ -types. □

Let  $J, J', K$  be subgroups of  $G$  with  $J, J'$  compact open and  $J \subset K, J' \subset K$ . Let  $\lambda, \lambda'$  be representations of  $J, J'$  on finite-dimensional vector spaces  $V, V'$ . Let  $g \in G$ . Then  $gJg^{-1} \cap J'$  is a subgroup of  $J'$ . We set  $\lambda^g(x) := \lambda(g^{-1}xg)$ . We define the  *$g$ -intertwining vector space* of  $(\lambda, \lambda')$  to be

$$\mathcal{I}_g(\lambda, \lambda') = \mathrm{Hom}_{gJg^{-1} \cap J'}(\lambda^g, \lambda').$$

**Definition 1.** (1) We say that  $g$  *intertwines*  $\lambda$  if  $\mathcal{I}_g(\lambda) \neq 0$ . The  *$K$ -intertwining set of  $\lambda$*  is

$$\mathcal{I}_K(\lambda) = \{g \in K : \mathcal{I}_g(\lambda) \neq 0\} \subset K.$$

(2) We say that  $g$  *intertwines*  $\lambda$  and  $\lambda'$  if  $\mathcal{I}_g(\lambda, \lambda') \neq 0$ . The  *$K$ -intertwining set of  $\lambda$  and  $\lambda'$*  is

$$\mathcal{I}_K(\lambda, \lambda') = \{g \in K : \mathcal{I}_g(\lambda, \lambda') \neq 0\} \subset K.$$

In [9, 10, 8], the results centre around identification of the  $G$ -intertwining set  $I_G(\lambda)$ . In our applications, we shall need only the  $K$ -intertwining set  $I_K(\lambda)$  where  $K$  is compact.

In order to study the induced representations and their decomposition into irreducible constituents, we need to use the Mackey formulas repeatedly.

We assume now that  $K$  is open compact. Then  $J, J'$  have finite index in  $K$ . We have the Mackey formula:

$$\mathrm{Hom}_K(\mathrm{Ind}_J^K(\lambda), \mathrm{Ind}_{J'}^K(\lambda')) \cong \bigoplus \mathcal{I}_x(\lambda, \lambda') \quad (3)$$

with  $x \in J \backslash K / J'$ . If  $\lambda = \lambda' \cong \lambda'$  then we set  $\mathcal{I}_g(\lambda) = \mathcal{I}_g(\lambda, \lambda')$  and we then have the isomorphism of  $\mathbb{C}$ -vector spaces

$$\mathrm{End}_K(\mathrm{Ind}_J^K(\lambda)) \cong \bigoplus \mathcal{I}_x(\lambda)$$

with  $x \in J \backslash K / J$ .

The following is an immediate consequence: we will use this result repeatedly.

**Proposition 2.** *If  $\mathcal{I}_K(\lambda) = J$  then  $\mathrm{Ind}_J^K(\lambda)$  is irreducible.*

We will use the following immediate result.

**Proposition 3.** *If  $\mathrm{Ind}_J^K(\lambda)$  and  $\mathrm{Ind}_{J'}^K(\lambda')$  are irreducible, then*

$$\mathrm{Ind}_J^K(\lambda) \cong \mathrm{Ind}_{J'}^K(\lambda') \iff \mathcal{I}_K(\lambda, \lambda') = JyJ'$$

for some element  $y$ .

**Proposition 4.** *Let  $(J^\mathfrak{s}, \tau^\mathfrak{s})$  be an  $\mathfrak{s}$ -type,  $(J^{\mathfrak{s}'}, \tau^{\mathfrak{s}'})$  be a  $\mathfrak{s}'$ -type with  $\mathfrak{s}, \mathfrak{s}'$  in  $\mathfrak{B}(G)$ ,  $\mathfrak{s} \neq \mathfrak{s}'$ . Let  $J$  be a compact open subgroup of  $G$  such that  $J^\mathfrak{s} \subset J$ ,  $J^{\mathfrak{s}'} \subset J$ . Then we have*

$$\dim_{\mathbb{C}} \mathrm{Hom}_J(\mathrm{Ind}_{J^\mathfrak{s}}^J \tau^\mathfrak{s}, \mathrm{Ind}_{J^{\mathfrak{s}'}}^J \tau^{\mathfrak{s}'}) = 0.$$

*Proof.* From the Mackey formula (3), it is equivalent to prove that  $\mathcal{I}_J(\tau^\mathfrak{s}, \tau^{\mathfrak{s}'}) = 0$ . The proof of the equivalence of (i) and (ii) of [9, Theorem 9.3.a] shows that  $\mathfrak{s} = \mathfrak{s}'$  if and only if  $\mathcal{I}_G(\tau^\mathfrak{s}, \tau^{\mathfrak{s}'}) \neq 0$ . The result follows.  $\square$

Let  $J$  be a compact open subgroup of  $G$  and  $(\tau, \mathcal{W})$  be an irreducible smooth representation of  $J$ . Let  $(\tau^\vee, \mathcal{W}^\vee)$  be the contragredient representation of  $(\tau, \mathcal{W})$ .

For any subgroup  $K$  of  $G$ , let  $\mathcal{H}(K, \tau)$  denote the space of compactly supported functions  $f: K \rightarrow \mathrm{End}_{\mathbb{C}}(\mathcal{W}^\vee)$  such that  $f(j_1 k j_2) = \tau^\vee(j_1) f(k) \tau^\vee(j_2)$ ,

for any  $j_i \in J$ ,  $k \in K$ . The standard convolution operation gives  $\mathcal{H}(K, \tau)$  the structure of an associative unital  $\mathbb{C}$ -algebra.

Let  $M$  be a Levi subgroup of  $G$ , and let  $(J_M, \tau_M)$  be a  $\mathfrak{t}$ -type, with  $\mathfrak{t} := [M, \sigma]_M$  a point of the Bernstein spectrum of  $M$ .

We recall from [9, Definition 8.1] that the pair  $(J, \tau)$  is a  $G$ -cover of  $(J_M, \tau_M)$  if  $J \cap M = J_M$  and  $\tau|_{J_M} \cong \tau_M$ , and if the following conditions hold for every parabolic subgroup  $P$  of  $G$  with Levi subgroup  $M$ :

- (1)  $(J, \tau)$  it is *decomposed with respect to  $(M, P)$* , that is,  $J$  admits the Iwahori decomposition:

$$J = J \cap U \cdot J_M \cdot J \cap \bar{U},$$

and the groups  $J \cap U$ ,  $J \cap \bar{U}$  are both contained in the kernel of  $\tau$  (here  $U$ ,  $\bar{U}$  denote the unipotent radicals of  $P$  and of its opposite parabolic subgroup, respectively),

- (2) there exists an invertible element of  $\mathcal{H}(G, \tau)$  supported on a double coset  $Jz_P J$ , where  $z_P$  is a central element in  $M$ , which is strongly  $(P, J)$ -positive in the sense of [9, Definition (6.16)].

The group  $\Psi(M)$  of unramified quasicharacters of  $M$  has the structure of a complex torus. The action (by conjugation) of  $N_G(M)$  on  $M$  induces an action of  $W(M) := N_G(M)/M$  on  $\mathfrak{B}(M)$ . Let  $W_{\mathfrak{t}}$  denote the stabilizer of  $\mathfrak{t} = [M, \sigma]_L$  in  $W(M)$ . Thus  $W_{\mathfrak{t}} = N_{\mathfrak{t}}/M$ , where

$$N_{\mathfrak{t}} = \{n \in N_G(M) : {}^n\sigma \cong \nu\sigma, \text{ for some } \nu \in \Psi(M)\} \quad (4)$$

denotes the  $N_G(M)$ -normalizer of  $\mathfrak{t}$ .

We will need the following Proposition which gives a bound for the compact intertwining.

**Proposition 5.** [11] *Let  $M$  be a Levi subgroup of  $G$ , let  $(J, \tau)$  be a  $G$ -cover of a  $\mathfrak{t}$ -type, with  $\mathfrak{t} = [M, \sigma]_M$  a point of the Bernstein spectrum of  $M$ , and let  $K$  be a compact subgroup of  $G$  which contains  $J$ . Let  $t$  denote the number of double classes  $J \backslash K / J$  which intertwine  $\tau$ . Then*

$$t \leq |W_{\mathfrak{t}}|.$$

*Proof.* It is a classical result that  $t$  is bounded by the dimension of  $\mathcal{H}(K, \tau)$ . The hypotheses listed in [11, §1.3] are satisfied, and so we can apply [11, Theorem 1.5(ii)]. We infer that

$$\dim_{\mathbb{C}} \mathcal{H}(K, \tau) \leq |W_{\mathfrak{t}}|.$$

□

### 3 Chamber homology groups

Let  $\mathfrak{o}_F$  denote the ring of integers of  $F$ , let  $\varpi = \varpi_F$  be a uniformizer in  $F$ , and  $\mathfrak{p}_F = \varpi_F \mathfrak{o}_F$  denote the maximal ideal of  $\mathfrak{o}_F$ . We set

$$\Pi = \Pi_N = \begin{pmatrix} 0 & \mathbf{I}_{N-1} \\ \varpi & 0 \end{pmatrix}.$$

Let  $s_0, s_1, \dots, s_{N-1}$  denote the standard involutions in  $G$ :  $s_i$  denote the matrix in  $G$  of the transposition  $i \leftrightarrow i+1$ , that is,

$$s_i = \begin{pmatrix} \mathbf{I}_{i-1} & & & \\ & 0 & 1 & \\ & 1 & 0 & \\ & & & \mathbf{I}_{N-i-1} \end{pmatrix},$$

for every  $i \in \{1, \dots, N-1\}$ , and  $s_0 = \Pi s_1 \Pi^{-1}$ .

The finite Weyl group is  $W_0 = \langle s_1, s_2, \dots, s_{N-1} \rangle$ , and the affine Weyl group is given by  $W = \langle s_0, s_1, \dots, s_{N-1} \rangle$ .

We set

$$\mathcal{R}(g) = \Pi^{-1} g \Pi$$

with  $g \in G$ , so that  $\mathcal{R}^N = 1$ .

We will use repeatedly, and without further comment, the fact that induction commutes with conjugation: in particular conjugation by  $\text{Ad } \Pi^i$ ,  $1 \leq i \leq N-1$ . We will use this in the following form:

$$\mathcal{R}^{-1}(\text{Ind}_{\mathcal{R}H}^{\mathcal{R}G}(\mathcal{R}\alpha)) \cong \text{Ind}_H^G(\alpha). \quad (5)$$

Note that

$$\mathcal{R}(s_i) = s_{i+1}, \quad \text{with } i = 0, 1, \dots, N-1 \pmod{N}.$$

The extended affine Weyl group is given by  $\widetilde{W} = W \rtimes \langle \Pi \rangle$ . We observe that

$$\widetilde{W} \cap \text{GL}(N, \mathfrak{o}_F) = W_0. \quad (6)$$

The standard Iwahori subgroup is

$$I = \begin{pmatrix} \mathfrak{o}_F^\times & \mathfrak{o}_F & \cdots & \mathfrak{o}_F \\ \mathfrak{p}_F & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \mathfrak{o}_F \\ \mathfrak{p}_F & \cdots & \mathfrak{p}_F & \mathfrak{o}_F^\times \end{pmatrix}.$$

Let  $A$  be the apartment attached to the diagonal torus and let  $\Delta$  denote the unique chamber of  $A$  which is stabilized by  $\langle \Pi \rangle I$ . We index the vertices  $L_0, L_1, \dots, L_{N-1}$  of  $\Delta$  in such a way that

- $s_i\Delta$  is the unique chamber of  $A$  which is adjacent to  $\Delta$  and such that  $s_i\Delta \cap \Delta$  is the  $(N-2)$ -simplex  $\{L_0, \dots, L_{N-1}\} \setminus \{L_i\}$ ;
- $\mathcal{R}(L_i) = L_{i+1}$  with  $i = 0, 1, \dots, N-1 \pmod N$ .

The  $L_i$  are the maximal standard parahoric subgroups of  $G$ ,

$$L_i = I \langle s_0, s_1, \dots, s_{i-1}, s_{i+1}, \dots, s_{N-1} \rangle I = \mathcal{R}^i(L_0),$$

and  $L_0 = \mathrm{GL}(N, \mathfrak{o}_F)$ .

The stabilizers of the facets of dimension  $N-1$  of  $\Delta$  are  $J_0, J_1, \dots, J_{N-1}$ , where

$$J_i = I \langle s_i \rangle I.$$

Each parahoric subgroup of  $G$  is defined by a facet of the building and the standard parahoric subgroups are the

$$J_S = I \langle s_j : j \in S \rangle I,$$

where  $S$  is any subset of  $\{0, 1, \dots, N-1\} \pmod N$ , [19, p. 118].

Hence,  $I = J_\emptyset$ ,  $J_i = J_{\{i\}}$ ,  $L_i = J_{\{0,1,\dots,i-1,i+1,\dots,N-1\}}$ .

The enlarged building  $\beta^1 G$  is labellable, that is, there exists a simplicial map  $\ell: \beta^1 G \rightarrow \Delta$ , which preserves the dimensions of the simplices. The labelling is unique, up to the automorphisms of  $\Delta$ . It allows us to fix an orientation of the simplices: one defines an incidence number  $\langle \eta : \sigma \rangle$  between an arbitrary facet  $\eta = (\eta_0, \dots, \eta_{i-1})$  of dimension  $i$  and any facet  $\sigma = (\sigma_0, \dots, \sigma_i)$  of dimension  $i+1$  which contains  $\eta$ , as follows

$$\langle \eta : \sigma \rangle = (-1)^i \text{ if } \{\ell(\eta_0), \dots, \ell(\eta_{i-1})\} \setminus \{\ell(\sigma_0), \dots, \ell(\sigma_i)\} = \emptyset.$$

The chamber homology groups are obtained by totalizing the bicomplex  $C_{**}$

$$\begin{array}{cccccccc} 0 & \longleftarrow & C_0 & \longleftarrow & \cdots & \longleftarrow & C_i & \longleftarrow & \cdots & \longleftarrow & C_{N-2} & \longleftarrow & C_{N-1} \\ & & \downarrow & & & & \downarrow & & & & \downarrow & & \downarrow \\ 0 & \longleftarrow & C_0 & \longleftarrow & \cdots & \longleftarrow & C_i & \longleftarrow & \cdots & \longleftarrow & C_{N-2} & \longleftarrow & C_{N-1} \end{array}$$

in which the chains are as follows:

$$C_i = \bigoplus_{\substack{S \subset \{0,1,\dots,N-1\} \\ |S|=N-1-i}} R(J_S) \quad (7)$$

and each vertical map is given by  $1 - \mathrm{Ad} \Pi$ . In particular, we have



- $C_0 = R(L_0) \oplus R(L_1) \oplus \cdots \oplus R(L_{N-1})$ ,
- $C_{N-2} = R(J_0) \oplus R(J_1) \oplus \cdots \oplus R(J_{N-1})$ ,
- $C_{N-1} = R(I)$ .

We will write an arbitrary element  $v$  in  $C_i$  as a  $\binom{N}{i}$ -uple  $[\eta]$ . Once an orientation has been chosen, the differentials are as follows: if  $v \in C_i$  then

$$\partial(v) = \sum_{\substack{\eta \subset \sigma \\ \dim \eta = i}} (-1)^{\langle \eta, \sigma \rangle} \text{Ind}_{G(\sigma)}^{G(\eta)}[\eta] \in C_{i-1}.$$

In particular:

- if  $v \in C_{N-1}$  then  $\partial(v) = (\text{Ind}_I^{J_0}(v), \text{Ind}_I^{J_1}(v), \dots, \text{Ind}_I^{J_{N-1}}(v))$ ,
- if  $v \in C_0$  then  $\partial(v) = 0$ .

When  $G = \text{GL}(3)$ , if  $v = (v_0, v_1, v_2) \in C_1$  then  $\partial(v)$  equals

$$(\text{Ind}_{J_2}^{L_0}(v_2) - \text{Ind}_{J_1}^{L_0}(v_1), \text{Ind}_{J_0}^{L_1}(v_0) - \text{Ind}_{J_2}^{L_1}(v_2), -\text{Ind}_{J_0}^{L_2}(v_0) + \text{Ind}_{J_1}^{L_2}(v_1)),$$

and, in the chain complex

$$0 \longleftarrow C_0 \longleftarrow C_1 \longleftarrow \cdots \longleftarrow C_2 \longleftarrow 0,$$

we have that  $v$  is a 1-cycle if and only if

$$\text{Ind}_{J_2}^{L_0}(v_2) = \text{Ind}_{J_1}^{L_0}(v_1), \text{Ind}_{J_0}^{L_1}(v_0) = \text{Ind}_{J_2}^{L_1}(v_2), \text{Ind}_{J_0}^{L_2}(v_0) = \text{Ind}_{J_1}^{L_2}(v_1),$$

*i.e.*, if and only if the 1-chain  $(v_0, v_1, v_2)$  is *vertex compatible*. Note that a true representation in  $R(I)$  can never be a 2-cycle; on the other hand, each 0-chain is a 0-cycle.

When we totalize the bicomplex we obtain the chain complex

$$0 \longleftarrow C_0 \longleftarrow C_0 \oplus C_1 \longleftarrow \cdots \longleftarrow C_{i-1} \oplus C_i \longleftarrow C_i \oplus C_{i+1} \longleftarrow \cdots \longleftarrow C_{N-1} \longleftarrow 0$$

**Definition 2.** The homology groups of this totalized complex are the chamber homology groups, as in [4].

To each point  $\mathfrak{s} \in \mathfrak{B}(G)$  we will associate a sub-bicomplex  $C_{**}(\mathfrak{s})$ :

$$\begin{array}{ccccccc} 0 & \longleftarrow & C_0(\mathfrak{s}) & \longleftarrow & \cdots & \longleftarrow & C_i(\mathfrak{s}) & \longleftarrow & \cdots & \longleftarrow & C_{N-1}(\mathfrak{s}) \\ & & \downarrow & & & & \downarrow & & & & \downarrow \\ 0 & \longleftarrow & C_0(\mathfrak{s}) & \longleftarrow & \cdots & \longleftarrow & C_i(\mathfrak{s}) & \longleftarrow & \cdots & \longleftarrow & C_{N-1}(\mathfrak{s}) \end{array}$$

in which each vertical differential is 0. By an *invariant chain* we shall mean a chain invariant under  $\text{Ad } \Pi$ .

Let  $\mathfrak{s}$  be a point in  $\mathfrak{B}(G)$  with  $\mathfrak{s} = [M, \sigma]_G$ . We recall that  $W(M)$  denotes the group  $N_G(M)/M$ . We take for  $M$  a standard Levi subgroup of  $G$ , isomorphic to  $\text{GL}(N_1) \times \cdots \times \text{GL}(N_r)$ , with  $(N_1 \geq N_2 \geq \cdots \geq N_r)$  a partition of  $N$ .

Given a point  $\mathfrak{s} \in \mathfrak{B}(G)$ , fix an  $\mathfrak{s}$ -type  $(J, \tau)$ . Such types exist [8, 9, 10]. There exists a parahoric subgroup  $J^{\mathfrak{s}}$  containing  $J$  such that  $(J^{\mathfrak{s}}, \alpha := \text{Ind}_J^{J^{\mathfrak{s}}} \tau)$  is also an  $\mathfrak{s}$ -type (see Theorems 5, 6, 7).

Then

- induce (if possible) each element in the orbit  $W(M) \cdot \alpha$  to the standard parahoric subgroups containing  $J^{\mathfrak{s}}$ , and rotate, *i.e.*, apply  $\mathcal{R}, \dots, \mathcal{R}^{N-1}$ ,
- take the free abelian groups generated by all the irreducible components which arise in this way.

Each of our sub-complexes  $C_{**}(\mathfrak{s})$  will come from some or all of this data. All the chain groups in  $C_{**}(\mathfrak{s})$  are finitely generated free abelian groups and comprise invariant chains. The homology groups of the chain complex

$$0 \longleftarrow C_0(\mathfrak{s}) \longleftarrow C_1(\mathfrak{s}) \longleftarrow \cdots \longleftarrow C_{N-1}(\mathfrak{s}) \longleftarrow 0$$

will be denoted  $h_*(\mathfrak{s})$ . We call this the *little complex*.

When we totalize the associated bicomplex  $C_{**}(\mathfrak{s})$  we obtain the chain complex

$$0 \longleftarrow C_0(\mathfrak{s}) \longleftarrow \cdots \longleftarrow C_{i-1}(\mathfrak{s}) \oplus C_i(\mathfrak{s}) \longleftarrow C_i(\mathfrak{s}) \oplus C_{i+1}(\mathfrak{s}) \longleftarrow \cdots \longleftarrow C_{N-1}(\mathfrak{s}) \longleftarrow 0$$

The following lemma will speed up our calculations.

**Lemma 1.** *The homology groups  $H_*(\mathfrak{s})$  of this complex are given by*

$$H_0(\mathfrak{s}) = h_0(\mathfrak{s}), \quad H_N(\mathfrak{s}) = h_{N-1}(\mathfrak{s})$$

$$H_{i+1}(\mathfrak{s}) = h_i(\mathfrak{s}) \oplus h_{i+1}(\mathfrak{s}), \quad 0 \leq i \leq N-2$$

$$H_{\text{ev}}(\mathfrak{s}) = h_0(\mathfrak{s}) \oplus h_1(\mathfrak{s}) \oplus \cdots \oplus h_{N-1}(\mathfrak{s}) = H_{\text{odd}}(\mathfrak{s})$$

*The even (resp. odd) chamber homology is precisely the total homology of the little complex.*

*Proof.* This is a direct consequence of the fact that each vertical differential in the bicomplex  $C_{**}(\mathfrak{s})$  is 0.  $\square$

## 4 Lattice chains and lattice sequences

Let  $V$  be an  $F$ -vector space of dimension  $N$ . We recall from [10, Def. 2.1] that a *lattice sequence* is a function  $\Lambda$  from  $\mathbb{Z}$  to the set of  $\mathfrak{o}_F$ -lattices in  $V$  such that

- $i \geq j$  implies  $\Lambda(i) \leq \Lambda(j)$ ;
- there exists  $e = e(\Lambda) \in \mathbb{Z}$ ,  $e \geq 1$ , such that  $\Lambda(i + e) = \mathfrak{p}_F \Lambda(i)$  for any  $i \in \mathbb{Z}$ .

The integer  $e$  is uniquely determined, and is called the *period* of  $\Lambda$ . We have  $e \leq N$ .

A lattice sequence which is injective as a function is called *strict*. We will put

$$\mathfrak{a}_n(\Lambda) := \{a \in A : a\Lambda(m) \subset \Lambda(m + n), m \in \mathbb{Z}\}, \quad n \in \mathbb{Z}. \quad (8)$$

The concept of lattice sequence generalizes the notion of lattice chain: as defined in [8, (1.11)], a *lattice chain* in  $V$  is a set  $\mathcal{L} = \{L_i : i \in \mathbb{Z}\}$  of  $\mathfrak{o}_F$ -lattices  $L_i$  in  $V$  such that

- $L_i \supset L_{i+1}$ ,  $L_i \neq L_{i+1}$ , for any  $i \in \mathbb{Z}$ ;
- there exists  $e = e(\mathcal{L}) \in \mathbb{Z}$  such that  $L_{i+e} = \mathfrak{p}_F L_i$ , for any  $i \in \mathbb{Z}$ .

The integer  $e$  is uniquely determined, and is called the *period* of  $\mathcal{L}$ .

Let  $k_F$  denote the residue field of  $F$ . For each  $i$ , the quotient  $L_i/L_{i+1}$  is a  $k_F$ -vector space. Write

$$d_i = d_i(\mathcal{L}) := \dim_{k_F}(L_i/L_{i+1}). \quad (9)$$

The function  $d(\mathcal{L}) : i \mapsto d_i$ ,  $i \in \mathbb{Z}$ , is periodic of period dividing  $e$ , and we have

$$\sum_{i=0}^{e-1} d_i = N. \quad (10)$$

To each lattice chain  $\mathcal{L}$  is attached a strict lattice sequence  $\Lambda_{\mathcal{L}}$  defined by  $\Lambda_{\mathcal{L}}(i) := L_i$ , for  $i \in \mathbb{Z}$ . In the opposite direction, to each lattice sequence  $\Lambda$  is attached a lattice chain  $\mathcal{L}_{\Lambda}$  defined by

$$\mathcal{L}_{\Lambda} := \{\Lambda(i) : i \in \mathbb{Z}\}. \quad (11)$$

As in [10, §2.6], we extend a lattice sequence  $\Lambda$  to a function on the real line  $\mathbb{R}$  by setting

$$\Lambda(x) := \Lambda(\lceil x \rceil), \quad x \in \mathbb{R}, \quad (12)$$

where  $\lceil x \rceil$  is the integer defined by the relation  $\lceil x \rceil - 1 < x \leq \lceil x \rceil$ .

Let  $\Lambda$  be a lattice sequence in  $V$  and let  $m$  be a positive integer. Then the function  $m\Lambda$  from  $\mathbb{Z}$  to the set of  $\mathfrak{o}_F$ -lattices in  $V$  defined by

$$(m\Lambda)(i) := \Lambda(i/m), \quad \text{for any } i \in \mathbb{Z},$$

is a lattice sequence in  $V$  with period  $m e(\Lambda)$ , and we have

$$(m\Lambda)(i) = \begin{cases} \Lambda(i/m) & \text{if } m \text{ divides } i, \\ \Lambda(1 + \lceil i/m \rceil) & \text{otherwise,} \end{cases} \quad (13)$$

and  $(m\Lambda)(x) = \Lambda(x/m)$ , for all  $x \in \mathbb{R}$  (see [10, Prop. 2.7]).

If we have a lattice sequence  $\Lambda$  in  $V$  and an integer  $t$ , we can define a lattice chain  $\Lambda + t$  by

$$(\Lambda + t)(i) := \Lambda(i + t), \quad \text{for any } i \in \mathbb{Z}. \quad (14)$$

Let  $m$  be a positive integer, and let  $V^1, V^2, \dots, V^m$  be  $m$  finite-dimensional  $F$ -vector spaces. Let  $\Lambda^1, \Lambda^2, \dots, \Lambda^m$  be  $m$  lattice sequences in  $V$ , with periods  $e_1, e_2, \dots, e_m$ , respectively. We denote by  $\Lambda = \Lambda^1 \oplus \dots \oplus \Lambda^m$  the *direct sum* of  $\Lambda^1, \dots, \Lambda^m$ : we recall from [10, §2.8] that  $\Lambda$  is defined by

$$\Lambda(ex) = \Lambda^1(e_1x) \oplus \dots \oplus \Lambda^m(e_mx), \quad \text{for each } x \in \mathbb{R}, \text{ where } e = \text{lcm}\{e_1, \dots, e_m\}. \quad (15)$$

The following example occurs in the construction of [10, §7.2]. See also [10, Example 2.8].

**Example 1.** We assume given  $m$  lattice chains  $\mathcal{L}^1, \mathcal{L}^2, \dots, \mathcal{L}^m$  in  $V^1, V^2, \dots, V^m$ , respectively, of same period  $e$ . We define a lattice chain

$$\mathcal{L} = \{L_i : i \in \mathbb{Z}\}$$

in  $V$  of period  $me$  by setting

$$L_{mj+k} := L_j^1 \oplus L_j^2 \oplus \dots \oplus L_j^{m-k} \oplus L_{j+1}^{m-k+1} \oplus \dots \oplus L_{j+1}^m,$$

any  $j \in \mathbb{Z}$  and  $0 \leq k \leq m-1$ . Using (13), (14), we obtain

$$\Lambda_{\mathcal{L}} = (m\Lambda^1 - m + 1) \oplus \dots \oplus (m\Lambda^{m-k} - k) \oplus \dots \oplus (m\Lambda^{m-1} - 1) \oplus m\Lambda^m.$$

## 4.1 Addition of lattice chains

Let  $A := \text{End}_F(V)$  and let  $E/F$  be a subfield of  $A$ . We denote by  $\mathfrak{o}_E$  the discrete valuation ring in  $E$ , by  $k_E$  its residue field, and by  $e(E|F)$  the ramification degree of  $E/F$ .

Let  $V^1, V^2, \dots, V^m$  be  $m$  finite-dimensional  $F$ -vector spaces of dimension  $N_1, N_2, \dots, N_m$ , respectively. We may consider each  $V^l$  as a  $E$ -vector space of dimension  $N_l/[E:F]$ .

Let  $\mathcal{L}^1, \mathcal{L}^2, \dots, \mathcal{L}^m$  be  $m$   $\mathfrak{o}_E$ -lattice chains in the  $E$ -vector spaces  $V^1, V^2, \dots, V^m$ , respectively, of period  $e'_1, e'_2, \dots, e'_m$ , respectively.

### 4.1.1 First addition procedure

We define first an  $\mathfrak{o}_E$ -lattice chain  $\mathcal{L}^1 + \mathcal{L}^2 = \{L_j^{[1,2]} : j \in \mathbb{Z}\}$  in  $V^1 \oplus V^2$  of period  $e'_1 + e'_2$  by

$$L_i^{[1,2]} := \begin{cases} L_0^1 \oplus L_i^2, & \text{if } 0 \leq i \leq e'_2 - 1 \\ L_{i-e'_2}^1 \oplus L_{e'_2}^2, & \text{if } e'_2 \leq i \leq e'_1 + e'_2 - 1. \end{cases} \quad (16)$$

Then let  $\mathcal{L}^1 + \mathcal{L}^2 + \mathcal{L}^3 = \{L_i^{[1,3]} : i \in \mathbb{Z}\}$  be the  $\mathfrak{o}_E$ -lattice chain  $(\mathcal{L}^1 + \mathcal{L}^2) + \mathcal{L}^3$  (which is the same as  $\mathcal{L}^1 + (\mathcal{L}^2 + \mathcal{L}^3)$ ). By applying (16) to the two  $\mathfrak{o}_E$ -lattice chains  $\mathcal{L}^1 + \mathcal{L}^2$  and  $\mathcal{L}^3$ , we get

$$L_i^{[1,3]} := \begin{cases} L_0^{[1,2]} \oplus L_i^3, & \text{if } 0 \leq i \leq e'_3 - 1 \\ L_{i-e'_3}^{[1,2]} \oplus L_{e'_3}^3, & \text{if } e'_3 \leq i \leq (e'_1 + e'_2) + e'_3 - 1, \end{cases}$$

that is, by using (16),

$$L_i^{[1,3]} := \begin{cases} L_0^1 \oplus L_0^2 \oplus L_i^3, & \text{if } 0 \leq i \leq e'_3 - 1 \\ L_0^1 \oplus L_{i-e'_3}^2 \oplus L_{e'_3}^3, & \text{if } e'_3 \leq i \leq e'_2 + e'_3 - 1 \\ L_{i-e'_3-e'_2}^1 \oplus L_{e'_2}^2 \oplus L_{e'_3}^3, & \text{if } e'_2 + e'_3 \leq i \leq e'_1 + e'_2 + e'_3 - 1. \end{cases} \quad (17)$$

Using this procedure, we finally obtain an  $\mathfrak{o}_E$ -lattice chain

$$\mathcal{L}^1 + \dots + \mathcal{L}^m = \mathcal{L}^{[1,m]} := \left\{ L_i^{[1,m]} : i \in \mathbb{Z} \right\} \quad (18)$$

of period  $e'_1 + e'_2 + \dots + e'_m$ . We have

$$L_i^{[1,m]} = L_0^1 \oplus \dots \oplus L_0^{j-1} \oplus L_k^j \oplus L_{e'_{j+1}}^{j+1} \oplus \dots \oplus L_{e'_m}^m, \quad (19)$$

for  $i = e'_{j+1} + \dots + e'_m + k$  with  $1 \leq j \leq m$  and  $0 \leq k \leq e'_j - 1$ .

We will need the  $\mathfrak{o}_E$ -lattice chain  $\mathcal{L}^1 + \cdots + \mathcal{L}^m$  in the special case when  $e'_1 = \cdots = e'_m = 1$ . In that case, the equation (19) becomes

$$L_i^{[1,m]} = L_0^1 \oplus \cdots \oplus L_0^{m-i-1} \oplus L_0^{m-i} \oplus L_1^{m-i+1} \oplus \cdots \oplus L_1^m, \quad (20)$$

for each  $i \in \{0, 1, \dots, m-1\}$ .

Since the  $\mathfrak{o}_E$ -lattice chains  $\mathcal{L}^1, \dots, \mathcal{L}^m$  all have period 1, we have  $\mathfrak{p}_E^j L_0^l = L_j^l$ , for each  $l \in \{1, \dots, m\}$  and each  $j \in \mathbb{Z}$ . Hence, since the  $\mathfrak{o}_E$ -lattice chain  $\mathcal{L}^{[1,m]}$  is of period  $m$ , we have

$$L_{mj+k}^{[1,m]} = \mathfrak{p}_E^j L_k^{[1,m]} = L_j^1 \oplus \cdots \oplus L_j^{m-k-1} \oplus L_j^{m-k} \oplus L_{j+1}^{m-k+1} \oplus \cdots \oplus L_{j+1}^m, \quad (21)$$

for each  $j \in \mathbb{Z}$  and each  $k \in \{0, 1, \dots, m-1\}$ .

Then we have

$$L_{mj+k+1}^{[1,m]} = L_j^1 \oplus \cdots \oplus L_j^{m-k-1} \oplus L_{j+1}^{m-k} \oplus L_{j+1}^{m-k+1} \oplus \cdots \oplus L_{j+1}^m.$$

It follows that

$$L_{mj+k}^{[1,m]} / L_{mj+k+1}^{[1,m]} \cong L_j^{m-k} / L_{j+1}^{m-k},$$

for each  $j \in \mathbb{Z}$  and each  $k \in \{0, 1, \dots, m-1\}$ .

Hence, setting  $d^l := d(\mathcal{L}^l)$  for any  $1 \leq l \leq m$ , we obtain

$$d_{mj+k}(\mathcal{L}^{[1,m]}) = d_j^{m-k} = d_0^{m-k} = \dim_E(V^{m-k}) = N_{m-k} / [E : F], \quad (22)$$

by (10), since  $e_{m-k} = 1$ .

We may consider each  $\mathcal{L}^l$ ,  $1 \leq l \leq m$ , as an  $\mathfrak{o}_F$ -lattice chain in the  $F$ -vector space  $V$ , of period  $e(E|F)$  (see [8, (1.2.4)]). Then  $\mathcal{L}^1 + \cdots + \mathcal{L}^m$ , viewed as an  $\mathfrak{o}_F$ -lattice chain, has period  $m e(E|F)$  (by [8, (1.2.4)]) and the equation (21) shows that it is the same as the chain  $\mathcal{L}$  considered in the Example 1.

#### 4.1.2 Second addition procedure

We keep assuming  $e'_1 = \cdots = e'_m = 1$ , and we will now consider the  $\mathfrak{o}_E$ -lattice chain

$$\mathcal{L}^m + \cdots + \mathcal{L}^1 = \left\{ L_i^{[m,1]} : i \in \mathbb{Z} \right\}.$$

We have

$$L_{mj+k}^{[m,1]} = L_{j+1}^1 \oplus \cdots \oplus L_{j+1}^k \oplus L_j^{k+1} \oplus \cdots \oplus L_j^m, \quad (23)$$

for each  $j \in \mathbb{Z}$  and each  $k \in \{0, 1, \dots, m-1\}$ .

It gives

$$L_{mj+k}^{[m,1]} / L_{mj+k+1}^{[m,1]} \cong L_j^{k+1} / L_{j+1}^{k+1}, \quad (24)$$

for each  $j \in \mathbb{Z}$  and each  $k \in \{0, 1, \dots, m-1\}$ . Hence we obtain

$$d_{mj+k}(\mathcal{L}^{[m,1]}) = d_j^{k+1} = d_0^{k+1} = \dim_E(V^{k+1}) = N_{k+1}/[E:F], \quad (25)$$

which in particular does not depend on  $m$ , in contrast with  $d_{mj+k}(\mathcal{L}^{[1,m]})$ .

As before, we may consider each  $\mathcal{L}^l$ ,  $1 \leq l \leq m$ , as an  $\mathfrak{o}_F$ -lattice chain in the  $F$ -vector space  $V$ , of period  $e(E|F)$ . Then  $\mathcal{L}^m + \dots + \mathcal{L}^1$ , viewed as an  $\mathfrak{o}_F$ -lattice chain, has period  $m e(E|F)$ .

## 5 Hereditary $\mathfrak{o}_F$ -orders

To any  $\mathfrak{o}_F$ -lattice chain  $\mathcal{L} = \{L_i\}$  in  $V$  is attached the following sequence of  $\mathfrak{o}_F$ -lattices in  $A$

$$\text{End}_{\mathfrak{o}_F}^n(\mathcal{L}) := \{x \in A : xL_i \subset L_{i+n}, i \in \mathbb{Z}\},$$

for each  $n \in \mathbb{Z}$ . In particular,  $\mathfrak{A} = \mathfrak{A}(\mathcal{L}) := \text{End}_{\mathfrak{o}_F}^0(\mathcal{L})$  is an hereditary  $\mathfrak{o}_F$ -order in  $A$ , and  $\mathfrak{P} := \text{End}_{\mathfrak{o}_F}^1(\mathcal{L})$  is the Jacobson radical of  $\mathfrak{A}$ . We will set

$$U(\mathfrak{A}) := \mathfrak{A}^\times \quad \text{and} \quad U^n(\mathfrak{A}) := 1 + \mathfrak{P}^n, \quad \text{for } n \geq 1. \quad (26)$$

We put

$$\mathfrak{K}(\mathfrak{A}) := \{g \in \text{Aut}_F(V) : g^{-1}\mathfrak{A}g = \mathfrak{A}\}. \quad (27)$$

**Definition 3.** For any partition  $(N_1, N_2, \dots, N_r)$  of  $N$ , we denote by

$$\mathfrak{A}(N_1, N_2, \dots, N_r)$$

the subset of  $M_N(F)$  consisting of the matrices of the following form: the  $(i, j)$ -block has dimension  $N_i \times N_j$ ,  $1 \leq i, j \leq r$ , and its entries lie in  $\mathfrak{o}_F$  if  $i \leq j$ , in  $\mathfrak{p}_F$  otherwise. Pictorially,

$$\mathfrak{A}(N_1, N_2, \dots, N_r) = \begin{pmatrix} \mathfrak{o}_F & \mathfrak{o}_F & \cdots & \mathfrak{o}_F \\ \mathfrak{p}_F & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \mathfrak{o}_F \\ \mathfrak{p}_F & \cdots & \mathfrak{p}_F & \mathfrak{o}_F \end{pmatrix}.$$

Let  $e := e(\mathcal{L})$  and  $d_i := d_i(\mathcal{L})$ . For each  $i \in \{0, 1, \dots, e-1\}$ , we choose elements  $v_{i,h} \in L_i$ ,  $1 \leq h \leq d_i$  such that the cosets  $v_{i,h} + L_{i+1}$  form a basis of the  $k_F$ -space  $L_i/L_{i+1}$ . Then

$$(v_{e-1,1}, v_{e-1,2}, \dots, v_{e-1,d_{e-1}}, v_{e-2,1}, v_{e-2,2}, \dots, v_{e-2,d_{e-2}}, \dots, v_{0,1}, v_{0,2}, \dots, v_{0,d_0})$$

is an  $F$ -basis of  $V$ . If we use this basis to identify  $A$  with the matrix algebra  $M_N(F)$ , then  $\mathfrak{A}$  becomes identified with  $\mathfrak{A}(d_0, d_2, \dots, d_{e-1})$ .

Now let  $V^1, V^2, \dots, V^m$  be  $m$  finite-dimensional  $F$ -vector spaces as in sections 4.1.1, 4.1.2, and let  $\mathcal{L}^1, \mathcal{L}^2, \dots, \mathcal{L}^m$  be  $m$   $\mathfrak{o}_E$ -lattice chains in  $V^1, V^2, \dots, V^m$ , all of period 1. We put

$$\mathfrak{A}^{[m,1]} := \mathfrak{A}(\mathcal{L}^{[m,1]}),$$

where  $\mathcal{L}^{[m,1]}$  is defined as in (23).

Let  $(m_1, \dots, m_r)$  be a partition of  $m$ . For each  $i \in \{1, \dots, r\}$ , we set  $\underline{m}_{i-1} := m_1 + \dots + m_{i-1}$ ,

$$\mathcal{L}^{[\underline{m}_i, \underline{m}_{i-1}+1]} := \mathcal{L}^{\underline{m}_i} + \mathcal{L}^{\underline{m}_i-1} + \dots + \mathcal{L}^{\underline{m}_{i-1}+2} + \mathcal{L}^{\underline{m}_{i-1}+1},$$

and

$$\mathfrak{A}^{[m_i, m_{i-1}+1]} := \mathfrak{A}(\mathcal{L}^{[\underline{m}_i, \underline{m}_{i-1}+1]}).$$

We set  $m_0 := 0$ . For each  $i \in \{1, \dots, r\}$ , we define  $V^{[m_{i-1}+1, m_i]}$  as

$$V^{[m_{i-1}+1, m_i]} := V^{\underline{m}_{i-1}+1} \oplus V^{\underline{m}_{i-1}+2} \oplus \dots \oplus V^{\underline{m}_i}.$$

Let  $M(m_1, \dots, m_r)$  denote the stabilizer of the decomposition

$$V = \bigoplus_{i=1}^r V^{[m_{i-1}+1, m_i]}.$$

**Lemma 2.** *We have*

$$M(m_1, \dots, m_r) \cap U(\mathfrak{A}^{[m,1]}) = \prod_{i=1}^r U(\mathfrak{A}^{[m_i, m_{i-1}]}).$$

*Proof.* We set  $e := e(E|F)$ . Let  $l \in \{1, 2, \dots, m\}$  and let  $j \in \{0, 1, \dots, e-1\}$ . Since the  $\mathfrak{o}_E$ -lattice chain  $\mathcal{L}^l$  has period 1, the equations (9) and (10) give

$$\dim_{k_E} L_j^l / L_{j+1}^l = \dim_{k_E} L_0^l / L_1^l = \frac{N_l}{[E:F]}.$$

It follows that

$$d_j^l = \dim_{k_F} L_j^l / L_{j+1}^l = [k_E : k_F] \dim_{k_E} L_j^l / L_{j+1}^l = [k_E : k_F] \frac{N_l}{[E:F]} = \frac{N_l}{e}.$$

Since  $d_j^l = d_0^l = N_l/e$ , we may and do fix an  $\mathfrak{o}_F$ -basis  $\mathcal{B}^l := (v_{0,1}^l, \dots, v_{0,N_l}^l)$  of  $\mathcal{L}^l$ , chosen to span  $L_0^l$  over  $\mathfrak{o}_F$ . We put

$$v_{j,h}^l := \begin{cases} v_{0,h}^l & \text{if } 1 \leq h \leq \delta_j^l, \\ \varpi_F v_{0,h}^l & \text{if } \delta_j^l + 1 \leq h \leq N_l, \end{cases}$$



where

$$\delta_j^l := \dim_{k_F} L_j^l / L_e^l = (e - j)d_j^l = \frac{e - j}{e} N_l.$$

The  $\mathfrak{o}_F$ -lattice  $L_j^l$  is then the  $\mathfrak{o}_F$ -linear span of the set  $\{v_{j,1}^l, \dots, v_{j,N_l}^l\}$ , the cosets  $v_{j,h}^l + L_{j+1}^l$  ( $1 \leq h \leq N_l/e$ ) form a basis of the  $k_F$ -space  $L_j^l / L_{j+1}^l$ , and

$$(v_{e-1,1}^l, \dots, v_{e-1,N_l/e}^l, \dots, v_{1,1}^l, \dots, v_{1,N_l/e}^l, v_{0,1}^l, \dots, v_{0,N_l/e}^l) = \mathcal{B}^l.$$

It follows that, for each  $i \in \{1, 2, \dots, r\}$ ,

$$\mathcal{B}^{[m_{i-1}+1, m_i]} := (\mathcal{B}^{m_{i-1}+1}, \mathcal{B}^{m_{i-1}+2}, \dots, \mathcal{B}^{m_i})$$

is an  $F$ -basis of the vector space  $V^{[m_{i-1}+1, m_i]}$  such that the cosets

$$v_{j,h}^{k+1} + L_{m_j+k+1}^{[m,1]}, \quad \text{for } 1 \leq h \leq N_{k+1}/e,$$

form a basis of the  $k_F$ -space

$$L_{m_j+k}^{[m,1]} / L_{m_j+k+1}^{[m,1]} \cong L_j^{k+1} / L_{j+1}^{k+1},$$

by (24).

Let  $\mathcal{B}$  denote the  $F$ -basis of  $V$  defined as

$$\mathcal{B} := (\mathcal{B}^{[1, m_1]}, \mathcal{B}^{[m_1+1, m_1+m_2]}, \dots, \mathcal{B}^{[m_{r-1}+1, m_r]}).$$

We observe that we have by construction

$$\mathcal{B} = \mathcal{B}^{[1, m]}, \tag{28}$$

where  $\mathcal{B}^{[1, m]}$  is the  $F$ -basis corresponding to the partition  $m$ .

We will now use the basis  $\mathcal{B}$  to identify  $A = \text{End}_F(V)$  with  $M_N(F)$  and use the basis  $\mathcal{B}^{[\underline{m}_{i-1}+1, \underline{m}_i]}$  to identify  $\text{End}_F(V^{[m_{i-1}+1, m_i]})$  with  $M_{N(i)}(F)$ , where

$$N(i) := N_{\underline{m}_{i-1}+1} + N_{\underline{m}_{i-1}+2} + \dots + N_{\underline{m}_i}.$$

Then  $\mathfrak{A}^{[m,1]}$  becomes identified with the matrices of the following form: the  $(h, h')$ -block has dimension

$$d_h(\mathcal{L}^{[m,1]}) \times d_{h'}(\mathcal{L}^{[m,1]}), \quad \text{if } 0 \leq h, h' \leq me - 1,$$

and its entries lie in  $\mathfrak{o}_F$  if  $i \leq i'$ , in  $\mathfrak{p}_F$  otherwise.

Now the product  $\prod_{i=1}^r \mathfrak{A}^{[\underline{m}_i, \underline{m}_{i-1}+1]}$  is viewed as diagonally embedded in  $M_N(F)$ , and  $\mathfrak{A}^{[\underline{m}_i, \underline{m}_{i-1}+1]}$  becomes then identified with the matrices of the following form: the  $(\underline{m}_{i-1}e + j, \underline{m}_{i-1}e + j')$ -block has dimension

$$d_j(\mathcal{L}^{[\underline{m}_i, \underline{m}_{i-1}+1]}) \times d_{j'}(\mathcal{L}^{[\underline{m}_i, \underline{m}_{i-1}+1]}), \quad \text{if } 0 \leq j, j' \leq m_i e - 1,$$

and its entries lie in  $\mathfrak{o}_F$  if  $j \leq j'$ , in  $\mathfrak{p}_F$  otherwise. Then the result follows from (25).  $\square$

## 6 Semisimple types

Let  $G = \mathrm{GL}(N, F) = \mathrm{GL}(V)$  and let  $\mathfrak{s} = [M, \sigma]_G$  be a point in the Bernstein spectrum  $\mathfrak{B}(G)$ . The Levi subgroup  $M$  is the stabilizer of a decomposition  $V = \bigoplus_{l=1}^m V^l$  of  $V$  as a direct sum of nonzero subspaces  $V^l$ . We set  $N_l := \dim_F V^l$ , and  $A_l := \mathrm{End}_F(V^l) \cong M_{N_l}(F)$ . Then  $N_1 + \cdots + N_m = N$ , and  $M$  is isomorphic to  $\mathrm{GL}(N_1, F) \times \cdots \times \mathrm{GL}(N_m, F)$ , and the supercuspidal representation  $\sigma$  of  $M$  is of the form  $\sigma = \pi_1 \otimes \cdots \otimes \pi_m$ , where  $\pi_l$  is an irreducible supercuspidal representation of the group  $\mathrm{GL}(N_l, F)$ , for  $l = 1, \dots, m$ . We set  $\mathfrak{t} := [M, \sigma]_M$ .

By [8, Theorem (8.4.1)], for each  $l$ , there is a maximal simple type  $(J^l, \lambda^l)$  occurring in  $\pi_l$ . The pair  $(J_M, \tau_M) := (J^1 \times \cdots \times J^m, \lambda^1 \otimes \cdots \otimes \lambda^m)$  is then an  $\mathfrak{t}$ -type in  $M$ .

By definition (see [8, (5.5.10)]), for each  $l$ , there exists an element  $\beta_l \in A^l$  for which the algebra  $E_l := F[\beta_l]$  is a field and a principal  $\mathfrak{o}_F$ -order  $\mathfrak{A}^l$  in  $A^l$ , of period  $e(E_l|F)$ , with Jacobson radical  $\mathfrak{P}_l$ , such that

$$J^l = \begin{cases} J(\beta_l, \mathfrak{A}^l) & \text{(as defined in [8, (3.1.14)]) if } \beta_l \notin F, \\ U(\mathfrak{A}^l) & \text{if } \beta_l \in F. \end{cases}$$

For each  $x \in A^l$ , we will write

$$\nu_{\mathfrak{A}^l}(x) := \max \{n \in \mathbb{Z} : x \in \mathfrak{P}_l^n\}. \quad (29)$$

Let  $\mathcal{L}^l$  denote the  $\mathfrak{o}_E$ -lattice chain defining the maximal  $\mathfrak{o}_E$ -order  $\mathfrak{B}^l := \mathfrak{A}^l \cap \mathrm{End}_E(V^l)$ . We have

$$J(\beta, \mathfrak{A}^l)/J^1(\beta, \mathfrak{A}^l) = U(\mathfrak{B}^l)/U^1(\mathfrak{B}^l) \cong \mathrm{GL}(f_l, k_E). \quad (30)$$

### 6.1 Simple types

We assume in this subsection that the  $N_l$  are all equal to  $N/m$  and that  $\pi_l \cong \pi_j \chi_j$ , with  $\chi_j$  an unramified character of  $\mathrm{GL}(N/m, F)$ , for each  $l, j \in \{1, \dots, m\}$ . In particular,  $M$  is then isomorphic to  $\mathrm{GL}(N/m, F)^m$ , and by [8, Theorem (8.4.2)], we can assume that all the  $\beta_l$ , all the  $\mathfrak{A}^l$ , all the  $\mathcal{L}^l$ , all the  $J^l$  and all the  $\lambda^l$  are equal. We will denote by  $E$  (resp.  $\beta$ ) the common value of the  $E_l$  (resp.  $\beta_l$ ).

Using the second addition procedure 4.1.2, we define: the  $\mathfrak{o}_E$ -lattice chain

$$\mathcal{L} := \mathcal{L}^m + \mathcal{L}^{m-1} + \cdots + \mathcal{L}^1, \quad \text{and} \quad \mathfrak{A} := \mathrm{End}_{\mathfrak{o}_F}^0(\mathcal{L}). \quad (31)$$

If  $\beta$  belongs to  $F$ , we set  $J := U(\mathfrak{A})$ . Otherwise, let  $n := -\nu_{\mathfrak{A}^1}(\beta)$ , then  $[\mathfrak{A}, mn, 0, \beta]$  is a simple stratum in the sense of [8, Definition (1.5.5)], let  $(J, \lambda) := (J(\beta, \mathfrak{A}), \lambda)$  be the corresponding simple type in  $G$ .

Let  $\mathfrak{B}$  denote the principal  $\mathfrak{o}_E$ -order in  $B := M_{N/[E:F]}(E)$  defined by  $\mathfrak{B} := B \cap \mathfrak{A}$ . We have  $m = e(\mathfrak{B}) = e(\mathfrak{B}|\mathfrak{o}_E)$ . In the case when  $\beta \in F$ , we have  $m = e(\mathfrak{A})$ .

**Definition 4.** We set

$$\mathfrak{A}^s := \mathfrak{A}(N/m, \dots, N/m) \quad \text{and} \quad J^s := U(\mathfrak{A}^s),$$

where  $\mathfrak{A}(N/m, \dots, N/m)$  is defined by Definition 3.

**Lemma 3.** *The  $\mathfrak{o}_F$ -order  $\mathfrak{A}$  is contained in the  $\mathfrak{o}_F$ -order  $\mathfrak{A}^s$ .*

*Proof.* We have  $\mathfrak{A} = oA(N/e(\mathfrak{A}), \dots, N/e(\mathfrak{A}))$ . In the case when  $J = U(\mathfrak{A})$ , we have  $\mathfrak{A}^s = \mathfrak{A}$ . Otherwise, the statement follows immediately from the above descriptions of the orders  $\mathfrak{A}$ ,  $\mathfrak{A}^s$ , and from the fact (see [8, Proposition (1.2.4)]) that

$$e(\mathfrak{A}) = m \cdot e(E|F).$$

Indeed, from the above descriptions of the orders  $\mathfrak{A}$ ,  $\mathfrak{A}^s$ , we have

$$\mathfrak{A}^s \cap U = \mathfrak{A} \cap U, \quad \mathfrak{A}^s \cap \overline{U} = \mathfrak{A} \cap \overline{U}, \quad (32)$$

$$M \cap \mathfrak{A}^s \cong (\mathrm{GL}(N/m, \mathfrak{o}_F))^m, \quad (33)$$

while  $M \cap \mathfrak{A}$  is isomorphic to the product of  $m$  copies of the order of  $e(E|F) \times e(E|F)$  blocks matrices of the following form: the  $(j, l)$ -block has dimension  $N/e(\mathfrak{A}) \times N/e(\mathfrak{A}) = (N/e(E|F)m) \times (N/e(E|F)m)$ ,  $0 \leq j, l \leq e(E|F) - 1$ , and its entries lie in  $\mathfrak{o}_F$  if  $j \leq l$ , in  $\varpi_F \mathfrak{o}_F$  otherwise, so that  $M \cap \mathfrak{A} \subset M \cap \mathfrak{A}^s$ .  $\square$

We set

$$f = \frac{N}{[E:F] \cdot m}. \quad (34)$$

Let  $K/E$  be an unramified field extension of degree  $f$  with

$$K^\times \subset \mathfrak{K}(\tilde{\mathfrak{B}}),$$

where  $\mathfrak{K}(\tilde{\mathfrak{B}})$  is defined by (27), and let  $C = \mathrm{End}_K(V) \cong M_m(K)$ . We view  $\varpi_E$  as a prime element of  $K$ . For  $i = 1, \dots, m-1$ , let  $s_{i,C}$  denote the matrix in  $C$  of the transposition  $i \leftrightarrow i+1$ , that is,

$$s_{i,C} = \begin{pmatrix} \mathbf{I}_{i-1} & & & \\ & 0 & 1 & \\ & 1 & 0 & \\ & & & \mathbf{I}_{m-i-1} \end{pmatrix},$$

and let  $s_{0,C} = \Pi_{m,C} s_{1,C} \Pi_{m,C}^{-1}$ , with

$$\Pi_{m,C} = \begin{pmatrix} 0 & \mathbf{I}_{m-1} \\ \varpi_E & 0 \end{pmatrix}.$$

We fix the embedding

$$\bigotimes \mathbf{I}_{N/m} : C \hookrightarrow \mathbf{M}_N(K) \quad c = (c_{ij}) \mapsto c \otimes \mathbf{I}_{N/m} = (c_{ij} \mathbf{I}_{N/m}),$$

$c \otimes \mathbf{I}_{N/m}$  being a block matrix with scalar blocks.

Let  $\widetilde{W}_C$  be the group generated by

$$S = \{s_{0,C} \otimes \mathbf{I}_{N/m}, s_{1,C} \otimes \mathbf{I}_{N/m}, \dots, s_{m-1,C} \otimes \mathbf{I}_{N/m}\}.$$

Then  $(\widetilde{W}_C, S)$  is a Coxeter group of type  $\widetilde{A}_{m-1}$ .

**Theorem 5.** *The representation  $\alpha = \text{Ind}_J^{J^s}(\lambda)$  is irreducible. Hence the pair  $(J^s, \alpha)$  is an  $\mathfrak{s}$ -type.*

*Proof.* In the case when  $J = U(\mathfrak{A})$ , we have  $J^s = J$ , so the result follows trivially in this case. We will assume from now on that  $J = J(\beta, \mathfrak{A})$ . For any  $i \in \{1, \dots, m-1\}$ ,

$$s_{i,C} \otimes \mathbf{I}_{N/m} = \begin{pmatrix} \mathbf{I}_{(i-1)N/m} & & & \\ & 0 & \mathbf{I}_{N/m} & \\ & \mathbf{I}_{N/m} & 0 & \\ & & & \mathbf{I}_{(m-i-1)N/m} \end{pmatrix} \notin J^s,$$

and

$$\Pi_{m,C} \otimes \mathbf{I}_{N/m} = \begin{pmatrix} 0 & \mathbf{I}_{(m-1)N/m} \\ \varpi_E \mathbf{I}_{N/m} & 0 \end{pmatrix} \notin J^s.$$

Hence  $\widetilde{W}_C \cap J^s = \{1\}$ , which gives

$$J^s \cap (J \cdot \widetilde{W}_C \cdot J) = J. \tag{35}$$

Then the result follows from the fact (see [8, Propositions (5.5.11) and (5.5.14) (iii)]) that

$$I_G(\lambda) \subset J \cdot \widetilde{W}_C \cdot J.$$

□

## 6.2 In the Levi subgroup $\widetilde{M}$

We will now consider the case of an arbitrary point  $\mathfrak{s} = [M, \sigma]_G$  in  $\mathfrak{B}(G)$ , with  $G = \mathrm{GL}(N, F)$ . Let  $\widetilde{M}$  denote the unique Levi subgroup of  $G$  which contains  $N_t$  (see 4) and is minimal for this property.

We write  $\sigma = \pi_1 \otimes \pi_2 \otimes \cdots \otimes \pi_m$  as

$$\sigma = (\sigma_1, \dots, \sigma_1, \sigma_2, \dots, \sigma_2, \dots, \sigma_t, \dots, \sigma_t),$$

where  $\sigma_j$ , a supercuspidal representation of  $\mathrm{GL}(N'_j, F)$ , is repeated  $\varepsilon_j$  times,  $1 \leq j \leq t$ , and  $\sigma_1, \dots, \sigma_t$  are pairwise distinct (after unramified twist). The integers  $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_t$  are called the *exponents* of  $\sigma$ . Then we have

$$M \cong \mathrm{GL}(N'_1, F)^{\varepsilon_1} \times \mathrm{GL}(N'_2, F)^{\varepsilon_2} \times \cdots \times \mathrm{GL}(N'_t, F)^{\varepsilon_t},$$

and

$$\widetilde{M} \cong \mathrm{GL}(\varepsilon_1 N'_1, F) \times \mathrm{GL}(\varepsilon_2 N'_2, F) \times \cdots \times \mathrm{GL}(\varepsilon_t N'_t, F).$$

For every  $j \in \{1, \dots, t\}$ , we set

$$\mathfrak{s}_j = [\mathrm{GL}(N'_j, F)^{\varepsilon_j}, \sigma_j^{\otimes \varepsilon_j}]_{\mathrm{GL}(\varepsilon_j N'_j, F)}.$$

Then let  $(K^j, \tau^j)$  be the  $\mathfrak{s}_j$ -type in  $\mathrm{GL}(\varepsilon_j N'_j, F)$  (a simple type) defined as in the previous section, and let  $(\widetilde{K}^j, \widetilde{\tau}^j)$  be the “modified simple type” attached to  $(K^j, \tau^j)$  as in [10, proof of Prop. 1.4].

**Lemma 4.** *We have  $\widetilde{K}^j \subset J^{\mathfrak{s}_j}$  and  $\alpha_i = \mathrm{Ind}_{\widetilde{K}^j}^{J^{\mathfrak{s}_j}}(\widetilde{\tau}^j)$  is irreducible.*

*Proof.* There is an isomorphism of Hecke algebras

$$\mathcal{H}(\mathrm{GL}(\varepsilon_j N'_j), \widetilde{\tau}^j) \cong \mathcal{H}(\mathrm{GL}(\varepsilon_j N_j), \tau^j)$$

such that, if  $\tilde{f} \in \mathcal{H}(\mathrm{GL}(\varepsilon_j N'_j), \widetilde{\tau}^j)$  has support  $\widetilde{K}^j g \widetilde{K}^j$ , for some element  $g \in \mathrm{GL}(\varepsilon_j N'_j, F)$ , then its image  $f$  in  $\mathcal{H}(\mathrm{GL}(\varepsilon_j N_j), \tau^j)$  has support  $K^j g K^j$  (see [8, (7.2.19)]). Then the result follows from Theorem 5.  $\square$

We set

$$\mathfrak{s}_{\widetilde{M}} = [M, \sigma]_{\widetilde{M}}, \quad \widetilde{\mathfrak{A}}^{\mathfrak{s}} = \mathfrak{A}^{\varepsilon_1} \times \cdots \times \mathfrak{A}^{\varepsilon_t}, \quad \widetilde{J}^{\mathfrak{s}} = U(\widetilde{\mathfrak{A}}^{\mathfrak{s}}),$$

$$\widetilde{K} = \widetilde{K}^1 \times \cdots \times \widetilde{K}^t \subset \widetilde{M}, \quad \widetilde{\tau} = \widetilde{\tau}^1 \otimes \cdots \otimes \widetilde{\tau}^t. \quad (36)$$

Note that

$$\widetilde{J}^{\mathfrak{s}} = \widetilde{M} \cap J^{\mathfrak{s}}. \quad (37)$$

It immediately follows from Lemma 4 that:

**Lemma 5.** *We have  $\widetilde{K} \subset \widetilde{J}^{\mathfrak{s}}$  and  $\tilde{\alpha} = \mathrm{Ind}_{\widetilde{K}}^{\widetilde{J}^{\mathfrak{s}}}(\widetilde{\tau})$  is irreducible.*

### 6.3 Recollection of the notion of endoclass [10]

We recall that a *simple pair*  $(k, \beta)$  over  $F$  consists of an integer  $k$  and a nonzero element  $\beta$  generating a field extension  $E$  of  $F$  such that

$$-k > \max \{k_0(\beta, \mathfrak{A}(E)), \nu_E(\beta)\},$$

where  $\nu_E$  is the standard additive valuation on  $E$  and  $k_0(\beta, \mathfrak{A}(E))$  is defined by [8, (1.4.5)], with  $\mathfrak{A}(E)$  denoting the unique hereditary  $\mathfrak{o}_F$ -order in  $\text{End}_F(E)$  such that  $\mathfrak{K}(\mathfrak{A}(E)) \supset E^\times$ .

Let  $(k, \beta)$  be a given simple pair in which  $k \geq 0$ . A *ps-character* (attached to the simple pair  $(k, \beta)$ ) is then a triple  $(\Theta, k, \beta)$ , where  $\Theta$  is a simple-character-valued function, such that to each triple  $(V, \mathfrak{B}, m)$ , where  $V$  is a finite-dimensional  $E$ -vector space,  $\mathfrak{B}$  is a hereditary  $\mathfrak{o}_E$ -order in  $\text{End}_E(V)$ , and  $m$  is an integer such that  $[m/e(\mathfrak{B}|\mathfrak{o}_E)] = k$ , the function  $\Theta$  attaches a simple character  $\Theta(\mathfrak{A}) \in \mathcal{C}(\mathfrak{A}, m, \beta)$ , called the *realization of  $\Theta$  on  $\mathfrak{A}$  of order  $m$* . (If we put  $n := -\nu_E(\beta) e(\mathfrak{B})$ , the stratum  $[\mathfrak{A}, n, m, \beta]$  is simple and the simple character set  $\mathcal{C}(\mathfrak{A}, m, \beta)$  of [8, (3.2)] is defined.)

These realizations are subject to the following coherence condition: if we have two realizations  $\Theta(\mathfrak{A}_1)$  and  $\Theta(\mathfrak{A}_2)$  of on orders  $\mathfrak{A}_1, \mathfrak{A}_2$ , they are related by  $\Theta(\mathfrak{A}_2) = \tau_{\mathfrak{A}_1, \mathfrak{A}_2, \beta}(\Theta(\mathfrak{A}_1))$ , where

$$\tau_{\mathfrak{A}_1, \mathfrak{A}_2, \beta}: \mathcal{C}(\mathfrak{A}_1, m, \beta) \rightarrow \mathcal{C}(\mathfrak{A}_2, m, \beta)$$

is the canonical bijection of [8, (3.6.14)].

Following [10, §4.3], we will say that two ps-characters  $(\Theta_1, k_1, \beta_1)$  and  $(\Theta_2, k_2, \beta_2)$  are *endo-equivalent* if there exists an  $F$ -vector space  $V$ , hereditary  $\mathfrak{o}_F$ -orders  $\mathfrak{A}_1, \mathfrak{A}_2$  in  $\text{End}_F(V)$ , and realizations  $\Theta_i(\mathfrak{A}_i)$  of the  $\Theta_i$  of same level, such that  $\mathfrak{A}_1 \cong \mathfrak{A}_2$  as  $\mathfrak{o}_F$ -orders, and such that the simple characters  $\Theta_i(\mathfrak{A}_i)$  intertwine in  $\text{Aut}_F(V)$ . Endo-equivalence is an equivalence relation on the set of ps-characters over  $F$ . One refers to the equivalence classes as *endo-classes* of simple characters.

If the supercuspidal representation  $\pi_l$  of  $\text{GL}(N_l, F)$  contains the trivial character of  $U^1(\mathfrak{A}^l) = 1 + \mathfrak{P}_l$ , then  $\pi_l$  is said to be of level-zero. Otherwise, there exists a simple stratum  $[\mathfrak{A}^l, n_l, 0, \beta_l]$  in  $A_l$  and a simple character  $\theta_l \in \mathcal{C}(\mathfrak{A}^l, 0, \beta_l)$  such that the restriction of  $\lambda^l$  to  $H^1(\beta_l, \mathfrak{A}^l)$  is a multiple of  $\theta_l$ . (Here  $H^1(\beta_l, \mathfrak{A}^l)$  is defined as in [8, (3.1.14)].) Since  $[\mathfrak{A}^l, n_l, 0, \beta_l]$  is simple, we have  $n_l = -\nu_{\mathfrak{A}^l}(\beta_l)$ . Then each representation  $\lambda^l$  is given as follows. There is a unique irreducible representation  $\eta_l$  of  $J^1(\beta, \mathfrak{A}^l)$  whose restriction to  $H^1(\beta, \mathfrak{A}^l)$  is a multiple of  $\theta_l$ . The representation  $\eta_l$  extends to a representation  $\kappa_l$  which is a  $\beta$ -extension of  $\eta_l$ , and we have  $\lambda^l = \kappa_l \otimes \rho_l$ , where  $\rho_l$  is the inflation of an irreducible representation of  $\text{GL}(f_l, k_E)$ , with  $f_l$  defined by (30).

If the representation  $\pi_l$  is of level zero, we set  $\Theta_{\pi_l} = \{\Theta^0\}$ , where  $\Theta^0$  is the trivial ps-character (that is, if  $\mathfrak{A}$  is a hereditary  $\mathfrak{o}_F$ -order in some  $\text{End}_F(V)$ , the realization of  $\Theta^0$  on  $\mathfrak{A}$  is the trivial character of  $U^1(\mathfrak{A})$ ). Otherwise, the simple character  $\theta_i$  determines a ps-character  $(\Theta_l, 0, \beta)$  and hence an endo-class  $\Theta_{\pi_l}$ .

We will denote by  $\Theta(1), \Theta(2), \dots, \Theta(q)$  the distinct endo-classes arising in the set  $\{\Theta_{\pi_1}, \dots, \Theta_{\pi_m}\}$ .

## 6.4 The homogeneous case

In this subsection, we assume that all the representations  $\pi_1, \pi_2, \dots, \pi_m$  admit the same endo-class. It follows that all the elements  $\beta_1, \dots, \beta_m$  may be assumed to be equal. We will denote by  $E$  (resp.  $\beta$ ) the common value of the  $E_l$  (resp.  $\beta_l$ ).

Let  $l \in \{1, \dots, m\}$ , and let  $(v_1^l, v_2^l, \dots, v_{N_l}^l)$  be an  $F$ -basis of  $V^l$ , with respect to which  $\mathfrak{A}^l = \mathfrak{A}(\mathcal{L}^l)$  is identified with  $\mathfrak{A}(N_l/e(E|F), \dots, N_l/e(E|F))$ . We have  $L_0^l = \mathfrak{o}_F v_1^l \oplus \dots \oplus \mathfrak{o}_F v_{N_l}^l$ . We set

$$L_{i,\max}^l := \mathfrak{p}^i L_0^l, \text{ for any } i \in \mathbb{Z}.$$

Then

$$\mathcal{L}_{\max}^l := \{L_{i,\max}^l : i \in \mathbb{Z}\} \quad (38)$$

is an  $\mathfrak{o}_F$ -lattice chain in  $V^l$  of period 1, and we have

$$\mathfrak{A}(\mathcal{L}_{\max}^l) := \text{End}_{\mathfrak{o}_F}^0(\mathcal{L}_{\max}^l) = \mathfrak{A}(N_l) = M_{N_l}(\mathfrak{o}_F) \supset \mathfrak{A}^l.$$

Following the second addition procedure defined in the subsection 4.1, we assemble the  $\mathfrak{o}_E$ -lattices chains  $\mathcal{L}^1, \dots, \mathcal{L}^m$  into the  $\mathfrak{o}_E$ -lattice chain

$$\bar{\mathcal{L}} := \mathcal{L}^m + \mathcal{L}^{m-1} + \dots + \mathcal{L}^1 \quad (39)$$

in  $V$ , of period  $m$ , and we assemble the  $\mathfrak{o}_F$ -lattices chains  $\mathcal{L}_{\max}^1, \dots, \mathcal{L}_{\max}^m$  into the  $\mathfrak{o}_F$ -lattice chain

$$\bar{\mathcal{L}}_{\max} := \mathcal{L}_{\max}^m + \mathcal{L}_{\max}^{m-1} + \dots + \mathcal{L}_{\max}^1 = \{\bar{L}_{\max,i} : i \in \mathbb{Z}\} \quad (40)$$

in  $V$ , of period  $m$ . Let  $j \in \mathbb{Z}$  and  $k \in \{0, 1, \dots, m-1\}$ . From (24), we have

$$\bar{L}_{\max,mj+k} / \bar{L}_{\max,mj+k+1} \cong L_{\max,j}^{k+1} / L_{\max,j+1}^{k+1}.$$

Hence:

$$d_{mj+k}(\bar{\mathcal{L}}_{\max}) = N_{k+1}. \quad (41)$$

It follows that

$$\mathfrak{A}(\bar{\mathcal{L}}_{\max}) = \mathfrak{A}(N_1, N_2, \dots, N_m).$$

We put

$$B := \text{End}_E(V) \quad \text{and} \quad \mathfrak{B} := \text{End}_{\mathfrak{o}_E}^0(\bar{\mathcal{L}}). \quad (42)$$

Considering  $\bar{\mathcal{L}}$  as an  $\mathfrak{o}_F$ -lattice chain, we put

$$\mathfrak{A} := \text{End}_{\mathfrak{o}_F}^0(\bar{\mathcal{L}}). \quad (43)$$

We have  $\mathfrak{B} = \mathfrak{A} \cap B$ .

The following definition, lemma and theorem generalize Definition 4, Lemma 3, and Theorem 5, respectively.

**Definition 5.** We set

$$\mathfrak{A}^s := \mathfrak{A}(N_1, N_2, \dots, N_m) \quad \text{and} \quad J^s := U(\mathfrak{A}^s).$$

**Lemma 6.** *The  $\mathfrak{o}_F$ -order  $\mathfrak{A}$  is contained in the  $\mathfrak{o}_F$ -order  $\mathfrak{A}^s$ .*

*Proof.* We have

$$\mathfrak{A}^s \cap U = \mathfrak{A} \cap U, \quad \mathfrak{A}^s \cap \bar{U} = \mathfrak{A} \cap \bar{U},$$

$$M \cap \mathfrak{A}^s \cong \prod_{l=1}^m \text{GL}(N_l, \mathfrak{o}_F).$$

In the notation of subsection 4.1.2, setting  $(m_1, \dots, m_r) = (1, \dots, 1)$ , we have  $r = m$ ,  $M = M(m_1, \dots, m_r)$ . Then  $\underline{m}_{l-1} = l - 1 = \underline{m}_l - 1$ , hence  $\bar{\mathcal{L}}^{[\underline{m}_l, \underline{m}_{l-1}+1]} = \mathcal{L}^l$ ,

$$\mathfrak{A}^l = \mathfrak{A}(\mathcal{L}^l) \cong \mathfrak{A}(N_l/e(E|F), \dots, N_l/e(E|F)),$$

and Lemma 2 gives

$$M \cap \mathfrak{A} \cong \prod_{l=1}^m U(\mathfrak{A}^l).$$

Since  $U(\mathfrak{A}^l) \subset \text{GL}(N_l, \mathfrak{o}_F)$ , the result follows.  $\square$

We set

$$n := \max(n_1, \dots, n_m). \quad (44)$$

**Lemma 7.** *With notation as above,  $[\mathfrak{A}, nm, 0, \beta]$  is a simple stratum.*

*Proof.* We have to check that the four conditions occurring in [8, Definition (1.5.5)] are satisfied.



- (i) We know that the algebra  $E = F[\beta]$  is a field, since the strata  $[\mathfrak{A}^l, n_l, 0, \beta]$  are simple.
- (ii) We defined  $\bar{\mathcal{L}} = \{\bar{L}_i : i \in \mathbb{Z}\}$  to be an  $\mathfrak{o}_E$ -lattice chain in the  $E$ -vector space  $V$ . Hence, by [8, Proposition (1.2.1)], we have  $E^\times \subset \mathfrak{K}(\mathfrak{A})$ .
- (iii) Let  $l \in \{1, \dots, m\}$ . We set  $\mathfrak{Q}_l := \mathfrak{B}_l \cap \mathfrak{P}_l$ . Since  $\nu_{\mathfrak{A}^l}(\beta) = -n_l$ , the definition (29) for  $\nu_{\mathfrak{A}^l}$  shows that

$$\beta \in \text{End}_E(V^l) \cap \mathfrak{P}_l^{-n_l} \quad \text{and} \quad \beta \notin \mathfrak{P}_l^{-n_l+1},$$

that is,

$$\beta \in \mathfrak{Q}_l^{-n_l} \quad \text{and} \quad \beta \notin \mathfrak{Q}_l^{-n_l+1}.$$

By [8, Proposition (1.2.4)], we know that  $\mathfrak{Q}_l$  is the Jacobson radical of the  $\mathfrak{o}_E$ -order  $\mathfrak{B}_l$ . Hence  $\mathfrak{Q}_l^i = \text{End}_{\mathfrak{o}_E}^i(\mathcal{L}^l)$  for each  $i \in \mathbb{Z}$ , and  $\beta(L_j^l)$  is contained in  $L_{j-n_l}^l$  and not in  $L_{j-n_l+1}^l$ . Now, it follows from (23) that

$$\beta(L_{mj+k}) = \beta(L_{j+1}^1) \oplus \dots \oplus \beta(L_{j+1}^k) \oplus \beta(L_j^{k+1}) \oplus \dots \oplus \beta(L_j^m),$$

for each  $j \in \mathbb{Z}$  and each  $k \in \{0, 1, \dots, m-1\}$ . Since  $n = \max(n_1, \dots, n_m)$ , we have  $L_{j-n_l}^l \subset L_{j-n}^l$ , for each  $l$ . It gives  $\beta(L_{mj+k}) \subset L_{m(j-n)+k}$ . On the other side there exists  $l_0 \in \{1, \dots, m\}$  such that  $n = n_{l_0}$ , and hence  $\beta(L_{j}^{l_0})$  is not contained in  $L_{j-n+1}^{l_0}$ . It follows that  $\beta(L_{mj+k})$  is not contained in  $L_{m(j-n)+k+1}$ , that is,

$$\beta \in \mathfrak{Q}^{-n} \quad \text{and} \quad \beta \notin \mathfrak{Q}^{-n+1},$$

where  $\mathfrak{Q}$  denotes the Jacobson radical of  $\mathfrak{B}$ . Since  $\mathfrak{Q}^i = B \cap \mathfrak{P}^i$  for each  $i \in \mathbb{Z}$  (by [8, Proposition (1.2.4)]), we get  $\nu_{\mathfrak{A}}(\beta) = -nm$ .

- (iv) Let  $A(E) := \text{End}_F(E)$ . The algebra  $A(E)$  contains the principal  $\mathfrak{o}_F$ -order

$$\mathfrak{A}(E) := \text{End}_{\mathfrak{o}_F}^0(\{\mathfrak{p}_E^i : i \in \mathbb{Z}\}).$$

We have  $E^\times \subset \mathfrak{K}(\mathfrak{A}(E))$  and [8, Proposition (1.4.13) (ii)] gives

$$k_0(\beta, \mathfrak{A}) = mk_0(\beta, \mathfrak{A}(E)) = k_0(\beta, \mathfrak{A}^l), \quad \text{for each } l \in \{1, \dots, m\}.$$

Since  $[\mathfrak{A}^l, n_l, 0, \beta]$  is a simple stratum, we have

$$0 < -k_0(\beta, \mathfrak{A}^l).$$

Hence  $0 < -k_0(\beta, \mathfrak{A})$  and  $[\mathfrak{A}, mn, 0, \beta]$  is simple.

□

Since  $[\mathfrak{A}, nm, 0, \beta]$  is a simple stratum, we can associate to it the compact open subgroups  $J(\beta, \mathfrak{A})$  and  $H^1(\beta, \mathfrak{A})$  of  $U(\mathfrak{A})$ , defined following [8, (3.1.14)].

As in [8, §7.1, 7.2], the set

$$K := H^1(\beta, \mathfrak{A}) \cap \overline{U} \cdot J(\beta, \mathfrak{A}) \cap P \quad (45)$$

is then a subgroup of  $U(\mathfrak{A})$  containing  $H^1(\beta, \mathfrak{A})$ .

Definition 5 and Lemma 6 imply

$$K \subset J^{\mathfrak{s}}. \quad (46)$$

As in [10, §7.2.1], it admits an irreducible representation  $\kappa$ , trivial on  $K \cap \overline{U}$ ,  $K \cap U$ , whose restriction to  $H^1(\beta, \mathfrak{A})$  is a multiple of  $\theta = \Theta(\mathfrak{A})$ , and such that  $\kappa|_{K \cap M}$  is of the form  $\kappa'_1 \otimes \cdots \otimes \kappa'_m$  for some  $\beta$ -extension  $\kappa'_l$  of  $\eta_l$ . As in [10, §7.2], we can choose the decomposition  $\lambda^l = \kappa_l \otimes \rho_l$  above so that  $\kappa_l = \kappa'_l$ ; we assume this has been done. We have canonically

$$K/K \cap J^1(\beta, \mathfrak{A}) \cong \prod_{l=1}^m J(\beta, \mathfrak{A}^l)/J^1(\beta, \mathfrak{A}^l) \cong \prod_{l=1}^m \mathrm{GL}(f_l, k_E),$$

and we can inflate the cuspidal representation  $\rho_1 \otimes \cdots \otimes \rho_m$  of  $\prod_{l=1}^m \mathrm{GL}(f_l, k_E)$  to a representation  $\rho$  of  $K$  and form

$$\tau = \kappa \otimes \rho. \quad (47)$$

Moreover similar proofs of those of [10, Theorem 7.2.1, Main Theorem 8.2] show that  $(K, \tau)$  is a  $G$ -cover of the pair  $(\tilde{K}, \tilde{\tau})$  defined in (36) and give the following formula for the intertwining:

$$\mathcal{I}_G(\tau) = K \cdot \mathcal{I}_{\tilde{M}}(\tilde{\tau}) \cdot K. \quad (48)$$

**Theorem 6.** *Let  $J^{\mathfrak{s}}$  defined as in Definition 5. Then the representation  $\alpha := \mathrm{Ind}_K^{J^{\mathfrak{s}}}(\tau)$  is irreducible. Hence the pair  $(J^{\mathfrak{s}}, \alpha)$  is an  $\mathfrak{s}$ -type.*

*Proof.* Using equations (46) and (48), we obtain

$$\mathcal{I}_{J^{\mathfrak{s}}}(\tau) = K \cdot \mathcal{I}_{\tilde{M} \cap J^{\mathfrak{s}}}(\tilde{\tau}) \cdot K.$$

On the other side, equation (37) and Lemma 5 imply that

$$\mathcal{I}_{\tilde{M} \cap J^{\mathfrak{s}}}(\tilde{\tau}) = \tilde{K} \subset K.$$

Hence  $\mathcal{I}_{J^{\mathfrak{s}}}(\tau) = K$ , and the result follows from Proposition 2.  $\square$

## 6.5 The general case

The Levi subgroup  $\widetilde{M}$  defined in the beginning of the subsection 6.2 is the  $G$ -stabilizer of a decomposition

$$V = \widetilde{V}^1 \oplus \widetilde{V}^2 \oplus \cdots \oplus \widetilde{V}^t,$$

of  $V$  as a direct sum of nonzero subspaces  $\widetilde{V}^j$ .

Since the endo-class of a supercuspidal representation only depends on the corresponding point in the Bernstein spectrum (see [10, Proposition 4.5]), we can associate to each  $\widetilde{V}^j$  an endo-class of simple characters, namely  $\Theta_{\pi_l}$  for any  $l$  such that  $V^l \subset \widetilde{V}^j$ .

Now let  $\bar{M} \supset M$  be the Levi subgroup in  $G$  defined as in [10, §8.1], that is, for each  $i$ , let  $\bar{V}^i$  be the sum of those  $\widetilde{V}^j$  whose associate endo-class  $\Theta_{\pi_j}$  is  $\Theta(i)$ , and write  $\bar{M}$  for the  $G$ -stabilizer of a decomposition

$$V = \bar{V}^1 \oplus \bar{V}^2 \oplus \cdots \oplus \bar{V}^q.$$

Setting  $\bar{N}_i := \dim_F \bar{V}^i$ , we get

$$\bar{M} \cong \mathrm{GL}(\bar{N}_1, F) \times \cdots \times \mathrm{GL}(\bar{N}_q, F).$$

We put

$$\bar{K} := K_1 \times K_2 \times \cdots \times K_q \quad \text{and} \quad \bar{\tau} := \tau_1 \times \tau_2 \times \cdots \times \tau_q, \quad (49)$$

where the pairs  $(K_i, \tau_i)$  are defined as in (45), (47). Then a similar proof as those of [10, §7.2] shows that the pair  $(\bar{K}, \bar{\tau})$  is a  $\bar{M}$ -cover of  $(J_M, \tau_M)$ .

For each  $i \in \{1, \dots, q\}$ , let  $\bar{\mathcal{L}}^i, \bar{\mathcal{L}}_{\max}^i$  respectively denote the  $\mathfrak{o}_E$ -lattice chain in the  $E$ -vector space  $\bar{V}^i$  defined by (39), and the  $\mathfrak{o}_F$ -lattice chain in the  $F$ -vector space  $\bar{V}^i$  defined by (40). Let  $m_i$  denote the number of representations  $\pi_l$  ( $1 \leq l \leq m$ ) with endo-class  $\theta_i$ . Then  $\bar{\mathcal{L}}^i$ , considered as an  $\mathfrak{o}_F$ -lattice chain, has period  $e_i := e(\bar{\mathcal{L}}^i) = e(E_i|F) m_i$ , and  $\bar{\mathcal{L}}_{\max}^i$  has period  $e(\bar{\mathcal{L}}_{\max}^i) = m_i$ .

Then let  $\Lambda^i$  (resp.  $\Lambda_{\max}^i$ ) denote the (strict) *lattice sequence* defined by the lattice chain  $\bar{\mathcal{L}}^i$  (resp.  $\bar{\mathcal{L}}_{\max}^i$ ), considered as  $\mathfrak{o}_F$ -lattice chains. Then, using the addition of lattice sequences recalled in (15), we define

$$\Lambda := \Lambda^1 \oplus \Lambda^2 \oplus \cdots \oplus \Lambda^q, \quad (50)$$

and

$$\Lambda_{\max} := e(E_1|F)\Lambda_{\max}^1 \oplus e(E_2|F)\Lambda_{\max}^2 \oplus \cdots \oplus e(E_q|F)\Lambda_{\max}^q. \quad (51)$$

Let  $\mathcal{L}_\Lambda, \mathcal{L}_{\Lambda_{\max}}$  denote the  $\mathfrak{o}_F$ -lattice chains attached to the lattice sequences  $\Lambda, \Lambda_{\max}$ , respectively, as in (11). Let  $\mathfrak{A}_\Lambda, \mathfrak{A}_{\Lambda_{\max}}$  denote the hereditary  $\mathfrak{o}_F$ -orders in  $A$  defined by the lattice chain  $\mathcal{L}_\Lambda, \mathcal{L}_{\Lambda_{\max}}$ , respectively. We have (see [10, Proposition 2.3. (i)]):

$$\mathfrak{A}_\Lambda = \mathfrak{A}(\mathcal{L}_\Lambda) = \mathfrak{a}_0(\Lambda) \quad \text{and} \quad \mathfrak{A}_{\Lambda_{\max}} = \mathfrak{A}(\mathcal{L}_{\Lambda_{\max}}) = \mathfrak{a}_0(\Lambda_{\max}), \quad (52)$$

where  $\mathfrak{a}_0(\Lambda), \mathfrak{a}_0(\Lambda_{\max})$  are defined as in (8).

**Lemma 8.** *The  $\mathfrak{o}_F$ -order  $\mathfrak{A}_\Lambda$  is contained in the  $\mathfrak{o}_F$ -order  $\mathfrak{A}_{\Lambda_{\max}}$ .*

*Proof.* Let  $e := \text{lcm}\{e_1, \dots, e_q\}$ . Both  $\Lambda$  and  $\Lambda_{\max}$  have period  $e$ . From (15), we have

$$\Lambda(ex) = \Lambda^1(e_1x) \oplus \cdots \oplus \Lambda^q(e_qx), \quad \Lambda_{\max}(ex) = \Lambda_{\max}^1(e_1x) \oplus \cdots \oplus \Lambda_{\max}^q(e_qx),$$

for each  $x \in \mathbb{R}$ . On the other side, (12) gives

$$\Lambda^i\left(\frac{e_i}{e}j\right) = \Lambda(l_i(j)) \quad \text{and} \quad \Lambda_{\max}^i\left(\frac{e_i}{e}j\right) = \Lambda_{\max}(l_i(j)),$$

for each  $j \in \mathbb{Z}$ , where  $l_i(j)$  is the integer defined by the relation

$$l_i(j) - 1 < \frac{e_i}{e}j \leq l_i(j).$$

Hence

$$\Lambda(j) = \Lambda^1(l_1(j)) \oplus \cdots \oplus \Lambda^q(l_q(j)), \quad \Lambda_{\max}(j) = \Lambda_{\max}^1(l_1(j)) \oplus \cdots \oplus \Lambda_{\max}^q(l_q(j)).$$

Then the result is consequence of Lemma 6.  $\square$

The following definition generalizes Definitions 4 and 5.

**Definition 6.** We set

$$\mathfrak{A}^s := \mathfrak{A}_{\Lambda_{\max}}, \quad \text{and} \quad J^s := U(\mathfrak{A}^s).$$

**Example 2.** We assume here that  $q = m$ , that is, the representations  $\pi_l$  have all distinct endo-classes. It implies that  $\bar{M} = \widetilde{M} = M$ . Then each lattice sequence  $\Lambda_{\max}^l$  has period 1, and so  $\Lambda_{\max}$  has also period 1. We get in this case  $J^s = \text{GL}(N, \mathfrak{o}_F)$ .

**Theorem 7.** *Then there exists a  $G$ -cover  $(J, \tau)$  of  $(J_M, \lambda_M)$  such that*

- $J \subset J^s$ ,

- $\alpha := \text{Ind}_J^{J^\mathfrak{s}}(\tau)$  is irreducible. Hence  $(J^\mathfrak{s}, \alpha)$  is an  $\mathfrak{s}$ -type.

*Proof.* Let  $(J, \tau)$  be the  $G$ -cover of  $(\bar{K}, \bar{\tau})$  constructed in the similar way as in [10, §8], in particular, we have

$$J \subset U(\mathfrak{A}_\Lambda).$$

Then the first assertion follows from Lemma 8.

On the other side the same proof as those of [10, §8.2, Main Theorem] gives the following formula for the intertwining:

$$\mathcal{I}_G(\tau) = J \cdot \mathcal{I}_{\widetilde{M}}(\tau_{\widetilde{M}}) \cdot J.$$

Since  $J \subset J^\mathfrak{s}$ , it implies:

$$\mathcal{I}_{J^\mathfrak{s}}(\tau) = J \cdot \mathcal{I}_{\widetilde{M} \cap J^\mathfrak{s}}(\tau_{\widetilde{M}}) \cdot J = J \cdot \mathcal{I}_{\widetilde{J}^\mathfrak{s}}(\widetilde{\tau}) \cdot J.$$

Now, by Lemma 5, we have

$$\mathcal{I}_{\widetilde{J}^\mathfrak{s}}(\tau_{\widetilde{M}}) = \widetilde{K}.$$

We get

$$\mathcal{I}_{J^\mathfrak{s}}(\tau) = J,$$

and the result follows from Proposition 2.  $\square$

## 7 Supercuspidal Bernstein components

Let  $\mathfrak{s} = [G, \pi]_G$ , where  $G = \text{GL}(N, F)$ . Here  $\pi$  is an irreducible unitary supercuspidal representation of  $G$ .

Let  $(J, \lambda)$  be a maximal simple type contained in  $\pi$ , as in [8]. We have  $e = 1$  and hence  $\mathfrak{A}_\mathfrak{s} = \text{M}(N, \mathfrak{o}_F)$ . It follows that  $J^\mathfrak{s} = \text{GL}(N, \mathfrak{o}_F) = L_0$ . By Proposition 5, the representation  $\alpha = \text{Ind}_J^{L_0}(\lambda)$  is irreducible. The pair  $(L_0, \alpha)$  is an  $\mathfrak{s}$ -type. The restriction to  $L_0$  of a smooth irreducible representation  $\pi'$  of  $G$  contains  $\alpha$  if and only if  $\pi'$  is isomorphic to  $\pi \otimes \chi \circ \det$ , where  $\chi$  is an unramified quasicharacter of  $F^\times$ . Moreover,  $\pi$  contains  $\alpha$  with multiplicity 1. In fact, the representation  $\alpha$  is the *unique* smooth irreducible representation  $\tau$  of  $L_0$  such that  $(L_0, \tau)$  is an  $\mathfrak{s}$ -type, see [16].

The little complex  $C_*(\mathfrak{s})$  determined by  $\alpha$  is

$$0 \longleftarrow C_0(\mathfrak{s}) \longleftarrow 0$$

where  $C_0(\mathfrak{s})$  is the free abelian group on the invariant 0-cycle

$$(\tau, \mathcal{R}(\tau), \mathcal{R}^2(\tau), \dots, \mathcal{R}^{n-1}(\tau))$$

The total homology of the little complex is given by  $h_0(\mathfrak{s}) = \mathbb{Z}$ . Therefore, by Lemma 1, we have

$$H_{\text{ev}}(\mathfrak{s}) = \mathbb{Z} = H_{\text{odd}}(\mathfrak{s}).$$

**Theorem 8.** *Let  $\pi$  be an irreducible unitary supercuspidal representation of  $\text{GL}(N)$ . Let  $\mathfrak{s} = [G, \pi]_G$ . Then we have*

$$H_{\text{ev}}(\mathfrak{s}) \cong K_0(\mathcal{A}^{\mathfrak{s}}), \quad H_{\text{odd}}(\mathfrak{s}) \cong K_1(\mathcal{A}^{\mathfrak{s}}).$$

*Proof.* The  $C^*$ -ideal  $\mathcal{A}^{\mathfrak{s}}$  is given by

$$\mathcal{A}^{\mathfrak{s}} = C(S^1, \mathfrak{K})$$

where  $\mathfrak{K}$  is the  $C^*$ -algebra of compact operators and

$$S^1 = \{\pi \otimes \chi \circ \det : \chi \in (F^\times)^\wedge\}.$$

The noncommutative  $C^*$ -algebra  $\mathcal{A}^{\mathfrak{s}}$  is strongly Morita equivalent to the commutative  $C^*$ -algebra  $C(S^1)$ . For this  $C^*$ -algebra we have

$$K_j(C(S^1)) \cong K^j(S^1) = \mathbb{Z}$$

where  $j = 0, 1$ . □

## 8 Generic Bernstein components attached to a maximal Levi subgroup

We assume in this section that  $\mathfrak{s} = [M, \sigma]_G$  with  $M \cong \text{GL}(N_1) \times \text{GL}(N_2)$  a 2-blocks Levi subgroup of  $G$  such that  $W_{\mathfrak{t}} = \{1\}$ . Note that the last conditions is always satisfied if  $N_1 \neq N_2$ .

Let  $(J_M, \lambda_M)$  be an  $\mathfrak{t}$ -type and let  $(J, \tau)$  be the  $G$ -cover of  $(J_M, \tau_M)$  considered in Theorem 7. We have shown there that  $\alpha := \text{Ind}_J^{J^{\mathfrak{s}}}(\tau)$  is irreducible. It then follows from Propositions 2 and 5 that  $\beta = \text{Ind}_J^{L_0}(\tau)$  is irreducible.

Let  $C_0(\tau), C_1(\tau)$  denote respectively the free abelian group on one generator  $(\beta, \mathcal{R}(\beta), \dots, \mathcal{R}^{N-1}(\beta))$ , and on  $(\alpha, \mathcal{R}(\alpha), \dots, \mathcal{R}^{N-1}\alpha)$ . The little complex is

$$0 \longleftarrow C_0(\mathfrak{s}) \xleftarrow{\partial} C_1(\mathfrak{s}) \longleftarrow 0$$

The map  $\partial$  is 0 by vertex compatibility of  $(\alpha, \mathcal{R}(\alpha), \dots, \mathcal{R}^{N-1}(\alpha))$ . Then  $h_0(\mathfrak{s}) = \mathbb{Z}, h_1(\mathfrak{s}) = \mathbb{Z}$  and so  $H_{\text{ev}}(\mathfrak{s}) = \mathbb{Z}^2 = H_{\text{odd}}(\mathfrak{s})$ .

The subset of the tempered dual of  $\text{GL}(N)$  which contains the  $\mathfrak{s}$ -type  $(J, \tau)$  has the structure of a compact 2-torus. But  $K^0(\mathbb{T}^2) = \mathbb{Z}^2 = K^1(\mathbb{T}^2)$  as required.

**Theorem 9.** *The  $\mathfrak{s}$ -type  $(J, \tau)$  generates a little complex  $C(\mathfrak{s})$ . For this complex we have*

$$H_{\text{ev}}(\mathfrak{s}) \cong K_0(\mathcal{A}^{\mathfrak{s}}) = \mathbb{Z}^2, \quad H_{\text{odd}}(\mathfrak{s}) \cong K_1(\mathcal{A}^{\mathfrak{s}}) = \mathbb{Z}^2$$

Note that the above Theorem applies to the intermediate principal series of  $\text{GL}(3)$ . In the next section, we will consider the principal series of  $\text{GL}(3)$ .

## 9 Principal series in $\text{GL}(3)$

Here  $s_0, s_1, s_2$  are the standard involutions

$$s_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad s_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \quad s_0 = \begin{pmatrix} 0 & 0 & \varpi^{-1} \\ 0 & 1 & 0 \\ \varpi & 0 & 0 \end{pmatrix}$$

where

$$\Pi = \Pi_3 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ \varpi & 0 & 0 \end{pmatrix}.$$

Note that  $\text{val}(\det(\Pi)) = 1$ . Restricted to the affine line  $\mathbb{R}$  in the enlarged building  $\beta^1\text{GL}(3) = \beta\text{SL}(3) \times \mathbb{R}$ ,  $\Pi$  sends  $t$  to  $t+1$ . We also have  $\Pi^3 = \varpi 1 \in \text{GL}(3)$ .

We have the double coset identities

$$0 \leq k \leq 2 \implies I \backslash J_k / I = \{1, s_k\} \tag{53}$$

$$r, s, t \text{ distinct} \implies J_r \backslash L_s / J_r = \{1, s_t\}. \tag{54}$$

Let  $\mathfrak{s} = [T, \sigma]_G$ , where  $T$  is the diagonal split torus in  $\text{GL}(3)$ :

$$T = \begin{pmatrix} F^\times & 0 & 0 \\ 0 & F^\times & 0 \\ 0 & 0 & F^\times \end{pmatrix},$$

and  $\sigma$  is an irreducible smooth character of  $T$ .

### 5.1. Construction of an $\mathfrak{s}$ -type, following Roche

For  $u \in F$ , we set

$$x_{1,2}(u) = \begin{pmatrix} 1 & u & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad x_{1,3}(u) = \begin{pmatrix} 1 & 0 & u \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad x_{2,3}(u) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & u \\ 0 & 0 & 1 \end{pmatrix},$$

$$x_{2,1}(u) = \begin{pmatrix} 1 & 0 & 0 \\ u & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad x_{3,1}(u) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ u & 0 & 1 \end{pmatrix}, \quad x_{3,2}(u) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & u & 1 \end{pmatrix},$$

and, for any  $k \in \mathbb{Z}$ ,

$$U_{i,j,k} = x_{i,j}(\mathfrak{p}_F^k).$$

Let  $\Phi = \{\alpha_{i,j} : 1 \leq i, j \leq 2\}$  be the set of roots of  $G$  with respect to  $T$ . For each root  $\alpha_{i,j}$ , let  $\alpha_{i,j}^\vee$  denotes the corresponding coroot. We have

$$\alpha_{1,2}^\vee(t) = \begin{pmatrix} t^{-1} & 0 & 0 \\ 0 & t & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \alpha_{2,1}^\vee(t) = \begin{pmatrix} t & 0 & 0 \\ 0 & t^{-1} & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

$$\alpha_{1,3}^\vee(t) = \begin{pmatrix} t^{-1} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & t \end{pmatrix}, \quad \alpha_{3,1}^\vee(t) = \begin{pmatrix} t & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & t^{-1} \end{pmatrix},$$

$$\alpha_{2,3}^\vee(t) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & t^{-1} & 0 \\ 0 & 0 & t \end{pmatrix}, \quad \alpha_{3,2}^\vee(t) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & t & 0 \\ 0 & 0 & t^{-1} \end{pmatrix}.$$

Define  $\sigma: T \rightarrow \mathbb{T}$  by

$$\sigma \begin{pmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{pmatrix} = \sigma_1(a)\sigma_2(b)\sigma_3(c),$$

where  $\sigma_i: F^\times \rightarrow \mathbb{T}$  is a character of  $F^\times$ , for  $i = 1, 2, 3$ .

Hence  $\sigma \circ \alpha_{i,j}^\vee: \mathfrak{o}_F^\times \rightarrow \mathbb{T}$  is the smooth character of  $\mathfrak{o}_F^\times$  defined by

$$\sigma \circ \alpha_{i,j}^\vee(t) = \sigma_j(t)\sigma_i(t^{-1}) = (\sigma_j\sigma_i^{-1})(t).$$

Now if  $\chi: \mathfrak{o}_F^\times \rightarrow \mathbb{T}$  is a smooth character, let  $c(\chi)$  be the conductor of  $\chi$ : the least integer  $n \geq 1$  such that  $1 + \mathfrak{p}_F^n \subset \ker(\chi)$ . We will write  $c_{i,j}$  for  $c(\sigma \circ \alpha_{i,j}^\vee)$ . We get

$$c_{i,j} = c(\sigma_j/\sigma_i) = c_{j,i}.$$

We can define a function  $f = f_\sigma: \Phi \rightarrow \mathbb{Z}$  (here  $\Phi$  is the set of roots) as follows:

$$f_\sigma(\alpha_{i,j}) = \begin{cases} [c_{i,j}/2] & \text{if } \alpha_{i,j} \in \Phi^+, \\ [(c_{i,j} + 1)/2] & \text{if } \alpha_{i,j} \in \Phi^-. \end{cases}$$

Here  $[x]$  denotes the largest integer  $\leq x$ .



Let

$$U_\sigma = \langle U_{i,j,f(\alpha_{i,j})} : \alpha_{i,j} \in \Phi \rangle,$$

and

$$J = \langle {}^\circ T, U_\sigma \rangle = {}^\circ T U_\sigma = U_\sigma {}^\circ T,$$

where  ${}^\circ T$  is the compact part of  $T$ ,

$${}^\circ T = \begin{pmatrix} \mathfrak{o}_F^\times & 0 & 0 \\ 0 & \mathfrak{o}_F^\times & 0 \\ 0 & 0 & \mathfrak{o}_F^\times \end{pmatrix}.$$

It follows that

$$J = \begin{pmatrix} \mathfrak{o}_F^\times & \mathfrak{p}_F^{[c_{1,2}/2]} & \mathfrak{p}_F^{[c_{1,3}/2]} \\ \mathfrak{p}_F^{[(c_{1,2}+1)/2]} & \mathfrak{o}_F^\times & \mathfrak{p}_F^{[c_{2,3}/2]} \\ \mathfrak{p}_F^{[(c_{1,3}+1)/2]} & \mathfrak{p}_F^{[(c_{2,3}+1)/2]} & \mathfrak{o}_F^\times \end{pmatrix}.$$

The group  $J$  will give the open compact group we are looking for.

Next, we need to figure out what is the correct character of  $J$ . In order to do that, we set

$$T_\sigma = \prod_{\alpha_{i,j} \in \Phi} \alpha_{i,j}^\vee (1 + \mathfrak{p}_F^{f(\alpha_{i,j}) + f(-\alpha_{i,j})}) \subset {}^\circ T.$$

Setting

$$U_\sigma^+ = U_\sigma \cap \begin{pmatrix} 1 & F & F \\ 0 & 1 & F \\ 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad U_\sigma^- = U_\sigma \cap \begin{pmatrix} 1 & 0 & 0 \\ F & 1 & 0 \\ F & F & 1 \end{pmatrix},$$

we obtain

$$U_\sigma = U_\sigma^- \cdot T_\sigma \cdot U_\sigma^+ \quad \text{and} \quad J = U_\sigma^- \cdot {}^\circ T \cdot U_\sigma^+.$$

It follows that

$$J/U_\sigma \cong {}^\circ T/T_\sigma.$$

By construction,  $T_\sigma \subset \ker(\sigma|_{{}^\circ T})$ . Hence  $\sigma|_{{}^\circ T}$  defines a character of  ${}^\circ T/T_\sigma$ , and so can be lifted to a character  $\tau$  of  $J$ . Then  $(J, \tau)$  an  $\mathfrak{s}$ -type by [18, Theorem 7.7].

## 5.2 Intertwining

We first recall that the following results ([18, Theorem 4.15])

$$I_G(\tau) = J \widetilde{W}(\sigma) J, \tag{55}$$

where

$$\widetilde{W}(\sigma) = \left\{ v \in \widetilde{W} : \sigma^v = \sigma \right\}.$$

More generally, it follows by the same proof as those of [18, Theorem 4.15], using [1, Prop. 9.3] instead of [18, Prop. 4.11], that, for each  $w \in W$ ,

$$I_G(\tau, \tau^w) = J \widetilde{W}(\sigma, \sigma^w) J^w, \quad (56)$$

where

$$\widetilde{W}(\sigma, \sigma^w) = \left\{ v \in \widetilde{W} : \sigma^v = \sigma^w \right\}.$$

Let

$$\Phi(\sigma) = \{ \alpha_{i,j} \in \Phi : (\sigma_i)|_{\mathfrak{o}_F^\times} = (\sigma_j)|_{\mathfrak{o}_F^\times} \} \subset \Phi.$$

The group  $W_0(\sigma)$  is equal to the group  $W_{\mathfrak{s}_T}$ , where  $\mathfrak{s}_T = [T, \lambda]_T$ . We observe that

$$I_{L_0}(\tau) = J W_0(\sigma) J. \quad (57)$$

**5.2.1 The case  $\Phi(\sigma) = \Phi$ .** Let  $\mathfrak{s} = [T, \sigma]_G$ , where  $\sigma = \psi \circ \det$  with  $\psi$  a smooth character of  $F^\times$ . In this case  $c_{i,j} = 1$  for any  $i, j$ . It follows that  $J = I$ .

The pair  $(I, \tau)$  is an  $\mathfrak{s}$ -type. We will construct cycles from this type. It follows from (57) that, as  $\mathbb{C}$ -algebras,

$$\text{End}_{L_0}(\text{Ind}_I^{L_0} \tau) \cong \mathcal{H}(\text{GL}(3, k_F) // B).$$

We also have, as  $\mathbb{C}$ -algebras,

$$\mathcal{H}(\text{GL}(3, k_F) // B) \cong \mathbb{C}[W_0] \cong \mathbb{C} \oplus \mathbb{C} \oplus M_2(\mathbb{C})$$

so that

$$\text{Ind}_I^{L_0} \tau = \lambda_{L_0} \oplus \mu_{L_0} \oplus \nu_{L_0} \oplus \nu_{L_0}$$

where  $\lambda_{L_0}, \mu_{L_0}, \nu_{L_0}$  are distinct.

We also have

$$\sigma|_{J_0} \hookrightarrow \text{Ind}_I^{J_0} \tau$$

by Frobenius reciprocity. The triple  $(\sigma|_{J_0}, \mathcal{R}(\sigma|_{J_0}), \mathcal{R}^2(\sigma|_{J_0}))$  is an invariant 1-cycle, and is not the boundary of  $1_I$ .

We now form the little complex:

- $C_0(\mathfrak{s})$  is the free abelian group on the three invariant 0-cycles

$$\lambda_L := (\lambda_{L_0}, \mathcal{R}(\lambda_{L_0}), \mathcal{R}^2(\lambda_{L_0}))$$

$$\mu_L := (\mu_{L_0}, \mathcal{R}(\mu_{L_0}), \mathcal{R}^2(\mu_{L_0}))$$

$$\nu_L := (\nu_{L_0}, \mathcal{R}(\nu_{L_0}), \mathcal{R}^2(\nu_{L_0}))$$

- $C_1(\mathfrak{s})$  is the free abelian group on the invariant 1-cycle

$$\lambda_J := (\sigma|_{J_0}, \mathcal{R}(\sigma|_{J_0}), \mathcal{R}^2(\sigma|_{J_0}))$$

In the little complex

$$0 \longleftarrow C_0(\mathfrak{s}) \xleftarrow{0} C_1(\mathfrak{s}) \longleftarrow 0$$

we have

$$h_0(\mathfrak{s}) = \mathbb{Z}^3, \quad h_1(\mathfrak{s}) = \mathbb{Z}.$$

The total homology of the little complex is  $\mathbb{Z}^4$ . As generating cycles we may take

$$\lambda_L, \mu_L, \nu_L, \lambda_J.$$

and so, by Lemma 1, the even (resp. odd) chamber homology groups are

$$H_{\text{ev}}(\mathfrak{s}) = \mathbb{Z}^4, \quad H_{\text{odd}}(\mathfrak{s}) = \mathbb{Z}^4.$$

Each irreducible representation  $\rho$  of a compact open subgroup  $J$  creates an *idempotent* in  $\mathcal{A}$  as follows. Let  $d$  denote the dimension of  $\rho$ , let  $\chi$  denote the character of  $\rho$ . Form the function  $d \cdot \chi : J \rightarrow \mathbb{C}$  and *extend by 0* to  $G$ . This function on  $G$  is a non-zero idempotent in  $\mathcal{A}$ , with the convolution product. We will denote this idempotent by  $e(\rho)$ :

$$e(\rho) * e(\rho) = e(\rho).$$

The inclusion

$$H_{\text{ev}}(\mathfrak{s}) \hookrightarrow K_0(\mathcal{A})$$

is given explicitly as follows:

$$\lambda_L \mapsto e(\lambda_{L_1}), \mu_L \mapsto e(\mu_{L_1}), \nu_L \mapsto e(\nu_{L_1}), \lambda_J \mapsto e(\lambda_{J_1}).$$

It follows from [17] that the  $C^*$ -ideal  $\mathcal{A}^{\mathfrak{s}}$  is given as follows:

$$\mathcal{A}^{\mathfrak{s}} \cong C(\text{Sym}^3 \mathbb{T}, \mathfrak{K}) \oplus C(\mathbb{T}^2, \mathfrak{K}) \oplus C(\mathbb{T}, \mathfrak{K}).$$

The symmetric cube  $\text{Sym}^3 \mathbb{T}$  is homotopy equivalent to  $\mathbb{T}$  via the product map

$$\text{Sym}^3 \mathbb{T} \sim \mathbb{T}, \quad (z_1, z_2, z_3) \mapsto z_1 z_2 z_3.$$

Hence  $K_0(\mathcal{A}^{\mathfrak{s}}) = \mathbb{Z}^4 = K_1(\mathcal{A}^{\mathfrak{s}})$  as required.

Note that

- $\text{Sym}^3 \mathbb{T}$  is in the minimal unitary principal series of  $\text{GL}(3)$

- $\mathbb{T}^2$  is in the intermediate unitary principal series of  $\mathrm{GL}(3)$
- $\mathbb{T}$  is in the discrete series of  $\mathrm{GL}(3)$ ; if  $\tau = 1$  then  $\mathbb{T}$  comprises the unramified unitary twists of the Steinberg representation of  $\mathrm{GL}(3)$

These are precisely the tempered representations of  $\mathrm{GL}(3)$  which contain the type  $(I, \tau)$ .

**Theorem 10.** *Let  $\mathfrak{s} = [T, \sigma]_G$  where  $\sigma = \psi \circ \det$  and  $\psi$  is a smooth (unitary) character of  $F^\times$ . Then we have*

$$H_{\mathrm{ev}}(\mathfrak{s}) \cong K_0(\mathcal{A}^{\mathfrak{s}}) = \mathbb{Z}^4, \quad H_{\mathrm{odd}}(\mathfrak{s}) \cong K_1(\mathcal{A}^{\mathfrak{s}}) = \mathbb{Z}^4.$$

### 5.2.2 The case $\emptyset \neq \Phi(\sigma) \neq \Phi$

Assume that  $(\sigma_1)_{|\mathfrak{o}_F^\times} = (\sigma_2)_{|\mathfrak{o}_F^\times} \neq (\sigma_3)_{|\mathfrak{o}_F^\times}$ . We have

$$J = \begin{pmatrix} \mathfrak{o}_F^\times & \mathfrak{o}_F & \mathfrak{p}_F^{[\ell/2]} \\ \mathfrak{p}_F & \mathfrak{o}_F^\times & \mathfrak{p}_F^{[\ell/2]} \\ \mathfrak{p}_F^{[(\ell+1)/2]} & \mathfrak{p}_F^{[(\ell+1)/2]} & \mathfrak{o}_F^\times \end{pmatrix},$$

where  $\ell = c_{1,3} = c_{2,3}$ , and

$$\tau \begin{pmatrix} a & * & * \\ * & b & * \\ * & * & c \end{pmatrix} = \sigma_1(a)\sigma_1(b)\sigma_3(c).$$

It is clear that  $s_1 \in I_{L_0}(\tau)$ . The Weyl group  $W_{s_T} = \mathbb{Z}/2\mathbb{Z}$  and so we have  $I_{L_0}(\tau) = J \cup J s_1 J$ . The complete list is as follows:

$$I_I(\tau) = J$$

$$I_{J_1}(\tau) = J \langle s_1 \rangle J, \quad I_{J_2}(\tau) = J, \quad I_{J_0}(\tau) = J$$

$$I_{L_1}(\tau) = J \langle s' \rangle J, \quad I_{L_2}(\tau) = J \langle s_1 \rangle J, \quad I_{L_0}(\tau) = J \langle s_1 \rangle J$$

where

$$s' = \begin{pmatrix} 0 & \varpi^{-1} & 0 \\ \varpi & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

**Lemma 9.** *Let  $\tau_1 = \mathrm{Ind}_J^I(\tau)$ . Then  $\tau_1$  is irreducible.*

*Proof.* This follows from proposition 2, since  $I_I(\tau) = J$ . It follows that  $(I, \tau_1)$  is an  $\mathfrak{s}$ -type.  $\square$

**Lemma 10.** *We have*

$$\mathrm{Ind}_I^{J_1} \tau_1 = \xi_1 \oplus \eta_1, \quad \mathrm{Ind}_I^{L_0} \tau_1 = \gamma_0 \oplus \delta_0.$$

*Proof.* We have  $I_{J_1}(\tau) = J \cup s_1 J$ . Hence

$$\mathrm{End}_{J_1}(\mathrm{Ind}_I^{J_1} \tau_1) = \mathcal{I}_1(\tau) \oplus \mathcal{I}_{s_1}(\tau) = \mathbb{C} \oplus \mathbb{C}.$$

This implies that  $\mathrm{Ind}_I^{J_1} \tau_1$  has two distinct irreducible constituents  $\xi_1, \eta_1$ . Now, we replace  $J_1$  by  $L_0$ , and infer that  $\mathrm{Ind}_I^{L_0} \tau_1$  has two distinct irreducible constituents  $\gamma_0, \delta_0$ .  $\square$

It follows that

$$\begin{aligned} \mathrm{Ind}_I^{J_2} \mathcal{R}(\tau_1) &= \mathcal{R}(\xi_1) \oplus \mathcal{R}(\eta_1), \\ \mathrm{Ind}_I^{J_0} \mathcal{R}^2(\tau_1) &= \mathcal{R}^2(\xi_1) \oplus \mathcal{R}^2(\eta_1). \end{aligned}$$

This creates two invariant 1-chains

$$\xi := (\xi_1, \mathcal{R}(\xi_1), \mathcal{R}^2(\xi_1)), \quad \eta := (\eta_1, \mathcal{R}(\eta_1), \mathcal{R}^2(\eta_1)).$$

It follows from (5) that

$$\begin{aligned} \mathrm{Ind}_I^{L_0} \tau_1 &\cong \mathrm{Ind}_I^{L_0} \mathcal{R}(\tau_1) \\ \zeta_1 &:= \mathrm{Ind}_I^{J_1} \mathcal{R}(\tau_1) \cong \mathrm{Ind}_I^{J_1} \mathcal{R}^2(\tau_1) \end{aligned}$$

By (17) we have

$$0 = \langle \mathrm{Ind}_I^{J_1} \tau, \mathrm{Ind}_I^{J_1} \mathcal{R}(\tau) \rangle.$$

Let  $C_0(\mathfrak{s})$  be the free abelian group generated by the two invariant 0-cycles

$$(\gamma_0, \mathcal{R}(\gamma_0), \mathcal{R}^2(\gamma_0)), \quad (\delta_0, \mathcal{R}(\delta_0), \mathcal{R}^2(\delta_0)).$$

Let  $C_1(\mathfrak{s})$  be the free abelian group generated by the two invariant 1-cycles  $\xi$  and  $\zeta$ .

The little complex is then

$$0 \longleftarrow C_0(\mathfrak{s}) \xleftarrow{0} C_1(\mathfrak{s}) \longleftarrow 0.$$

We have  $h_0(\mathfrak{s}) = \mathbb{Z}^2, h_1(\mathfrak{s}) = \mathbb{Z}^2$  and the total homology is  $\mathbb{Z}^4$  and so  $H_{\mathrm{ev}}(\mathfrak{s}) = \mathbb{Z}^4 = H_{\mathrm{odd}}(\mathfrak{s})$ .

The definition of  $\zeta := (\zeta_0, \mathcal{R}(\zeta_0), \mathcal{R}^2(\zeta_0))$  shows that

$$\partial(\tau_1 + \mathcal{R}(\tau_1) + \mathcal{R}^2(\tau_1)) = \xi + \eta + 2\zeta$$

so that  $\eta$  and  $-(\xi + 2\zeta)$  are homologous. Therefore the invariant 1-cycle  $\eta$  does not contribute a new homology class in  $H_1(G; \beta^1 G)$ .

The  $C^*$ -ideal  $\mathcal{A}^{\mathfrak{s}}$  is as follows:

$$C(\mathbb{T}^2, \mathfrak{K}) \oplus C(\mathrm{Sym}^2 \mathbb{T} \times \mathbb{T}, \mathfrak{K}).$$

To identify these ideals, we proceed as follows. First, let  $\Psi(F^\times)$  denote the group of unramified unitary characters of  $F^\times$ . The first summand is determined by the compact orbit

$$\mathcal{O}(\mathrm{St}(\sigma_1, 2) \otimes \sigma_3) = \{\chi_1 \mathrm{St}(\sigma_1, 2) \otimes \chi_2 \sigma_3 : \chi_j \in \Psi(F^\times)\}$$

where  $\mathrm{St}(\sigma_1, 2)$  is a generalized Steinberg representation; the second is determined by the compact orbit

$$\mathcal{O}(\sigma_1 \otimes \sigma_1 \otimes \sigma_3) = \{\chi_1 \sigma_1 \otimes \chi_2 \sigma_1 \otimes \chi_3 \sigma_3 : \chi_j \in \Psi(F^\times)\}.$$

The compact space  $\mathrm{Sym}^2 \mathbb{T} \times \mathbb{T}$  is homotopy equivalent to the 2-torus  $\mathbb{T}^2$ .

The space  $\mathrm{Sym}^2 \mathbb{T} \times \mathbb{T}$  is in the minimal unitary principal series of  $\mathrm{GL}(3)$  and the space  $\mathbb{T}^2$  is in the intermediate unitary principal series of  $\mathrm{GL}(3)$ . The union of these two compact spaces is precisely the set of tempered representations of  $\mathrm{GL}(3)$  which contain the  $\mathfrak{s}$ -type  $(J, \tau)$ .

The  $K$ -groups are now immediate:

$$K_j(\mathcal{A}^{\mathfrak{s}}) = \mathbb{Z}^4$$

with  $j = 0, 1$ .

**Theorem 11.** *Let  $\mathfrak{s} = [T, \sigma]_G$ . We have*

$$H_{\mathrm{ev}}(\mathfrak{s}) \cong K_0(\mathcal{A}^{\mathfrak{s}}) = \mathbb{Z}^4, \quad H_{\mathrm{odd}}(\mathfrak{s}) \cong K_1(\mathcal{A}^{\mathfrak{s}}) = \mathbb{Z}^4.$$

**5.2.3 The case  $\Phi(\sigma) = \emptyset$ .** The generic torus. The Bernstein component is  $[T, \sigma_1 \otimes \sigma_2 \otimes \sigma_3]$ . The Weyl group  $W(T) = W_0 = S_3$ , and the associated parahoric subgroup is the Iwahori subgroup  $I$ .

The restrictions of  $\sigma_1, \sigma_2$  and  $\sigma_3$  to  $\mathfrak{o}_F^\times$  are all distinct. We have  $\Phi(\sigma) = \emptyset$ . We have  $\widetilde{W}(\sigma) = D$ , where  $D$  is the subgroup of  $T$  whose eigenvalues are powers of  $\varpi$ . The subgroup  $D$  is free abelian of rank 3. The only compact element in  $D$  is  $1_G$ . The only double- $J$ -coset representative in  $L_0$  which  $G$ -intertwines  $\tau$  is  $1_G$ . This proves the following:

**Lemma 11.** *If  $r = 0, 1, 2$  then  $\mathrm{Ind}_J^{J_r}(\tau)$  is irreducible,  $\mathrm{Ind}_J^{L_r}(\tau)$  is irreducible.*

Let  $\alpha = \mathrm{Ind}_J^I(\tau)$ . Then  $\alpha$  is irreducible. Therefore  $(I, \alpha)$  is an  $\mathfrak{s}$ -type.

**Lemma 12.** *If  $w \in W_0$  then  $\text{Ind}_I^{L_0} \alpha = \text{Ind}_I^{L_0} \alpha^w$ .*

*Proof.* We have  $\text{Ind}_I^{L_0}(\alpha) = \text{Ind}_J^{L_0}(\tau)$  and  $\text{Ind}_I^{L_0}(\alpha^w) = \text{Ind}_J^{L_0}(\tau^w)$ . By Proposition 3, it is sufficient to prove that  $I_G(\tau, \tau^w) \neq \{0\}$ . But  $I_G(\tau, \tau^w) = J \widetilde{W}(\sigma, \sigma^w) J$ .  $\square$

**Lemma 13.** *If  $w \in W_0$  then*

$$\text{Ind}_I^{J_r}(\alpha) \cong \text{Ind}_I^{J_r}(w\alpha) \iff w \in \langle s_r \rangle$$

with  $0 \leq r \leq 2$ .

*Proof.* By Proposition 3,

$$\text{Ind}_I^{J_r}(\tau) \cong \text{Ind}_I^{J_r}(w\tau) \iff I_{J_r}(\tau, \tau^w) \neq \{0\}.$$

From (56), we have

$$I_{J_r}(\tau, \tau^w) = J_r \cap \widetilde{W}(\sigma, \sigma^w) = J_r \cap \widetilde{W}(\sigma) \cdot w = J_r \cap D \cdot w.$$

The result follows from the fact that  $J_r = I < 1, s_r > I$ .  $\square$

Inducing the orbit  $W_0 \cdot \alpha$  from  $J$  to  $J_1$  gives 3 distinct elements  $\rho_1, \phi_1, \psi_1$ , by Lemma 9. Inducing from  $J$  to  $L_0$  gives  $\gamma_0$ .

Set  $C_2(\mathfrak{s}) =$  free abelian group on the invariant 2-cycle

$$\epsilon := \sum_{w \in W_0} \text{sgn}(w) w \cdot \alpha.$$

Set  $C_1(\mathfrak{s}) =$  free abelian group on the three invariant 1-cycles

$$\rho := (\rho_1, \mathcal{R}(\rho_1), \mathcal{R}^2(\rho_1)),$$

$$\phi := (\phi_1, \mathcal{R}(\phi_1), \mathcal{R}^2(\phi_1)),$$

$$\psi := (\psi_1, \mathcal{R}(\psi_1), \mathcal{R}^2(\psi_1)).$$

Set  $C_0(\mathfrak{s}) =$  free vector abelian group on the invariant 0-cycle

$$\gamma := (\gamma_0, \mathcal{R}(\gamma_0), \mathcal{R}^2(\gamma_0)).$$

Note that

$$\partial \left( \sum_{w \in \text{Alt}(3)} w \cdot \alpha \right) = \rho + \phi + \psi$$

where  $Alt(3)$  is the alternating subgroup of  $W_0$ . Since  $s_1 s_2 \cdot \alpha = \mathcal{R}(\alpha)$ , we may also write this as

$$\partial(\alpha + \mathcal{R}(\alpha) + \mathcal{R}^2(\alpha)) = \rho + \phi + \psi.$$

It follows that  $\psi$  is homologous to  $-(\rho + \phi)$  in the top row of the double complex  $C_{**}$ . This implies that the *image* of  $C(\mathfrak{s})$  in  $C_{**}$  determines 4 homology classes. As representing cycles we may take the 2-cycle  $\epsilon$ , the two 1-cycles  $\rho, \phi$ , and the 0-cycle  $\gamma$ . Therefore

$$H_{\text{ev}}(G; \beta^1 G)^{\mathfrak{s}} = H_{\text{odd}}(G; \beta^1 G)^{\mathfrak{s}} = \mathbb{Z}^4.$$

**Theorem 12.** *The subspace of the tempered dual of  $GL(3)$  which contains the  $\mathfrak{s}$ -type  $(I, \alpha)$  has the structure of a compact 3-torus. This is a generic torus in the minimal unitary principal series of  $GL(3)$ . We have*

$$H_{\text{ev}}(G; \beta^1 G)^{\mathfrak{s}} \cong K_0(\mathcal{A}^{\mathfrak{s}}) = \mathbb{Z}^4, \quad H_{\text{odd}}(G; \beta^1 G)^{\mathfrak{s}} \cong K_1(\mathcal{A}^{\mathfrak{s}}) = \mathbb{Z}^4.$$

*Proof.* Let  $\Psi(F^\times)$  denote the group of unramified characters of  $F^\times$ . If  $\chi \in \Psi(F^\times)$  then  $\chi(x) = z^{\text{val}(x)}$  with  $z$  a complex number of modulus 1, so that

$$\Psi(F^\times) \cong \mathbb{T}.$$

Writing

$$\mathbb{T}^3 = \{\text{Ind}_T^G(\chi_1 \sigma_1 \otimes \chi_2 \sigma_2 \otimes \chi_3 \sigma_3) : \chi_j \in \Psi(F^\times)\}$$

we have

$$\mathcal{A}^{\mathfrak{s}} \cong C(\mathbb{T}^3, \mathfrak{K})$$

which is strongly Morita equivalent to  $C(\mathbb{T}^3)$ . The  $K$ -theory of the 3-torus is given by

$$K^j(\mathbb{T}^3) = \mathbb{Z}^4$$

where  $j = 0, 1$ . □

## A Chamber homology and K-theory

Let  $G = GL(N)$  and let  $\mathcal{A}$  denote the reduced  $C^*$ -algebra of  $G$ . Let  $\mathcal{H}(G)$  be the convolution algebra of uniformly locally constant, compactly supported, complex-valued functions on  $G$ , and let  $\mathcal{C}(G)$  be the Harish-Chandra Schwartz algebra of  $G$ . The following diagram serves as a framework for this



article:

$$\begin{array}{ccc}
K_j^{\text{top}}(G) & \xrightarrow{\mu} & K_j(\mathcal{A}) \\
\text{ch} \downarrow & & \downarrow \text{ch} \\
H_j(G; \beta^1 G) \otimes_{\mathbb{Z}} \mathbb{C} & \longrightarrow & \text{HP}_j(\mathcal{H}(G)) \xrightarrow{\iota_*} \text{HP}_j(\mathcal{C}(G))
\end{array}$$

with  $j = 0, 1$ . In this diagram,  $K_j^{\text{top}}(G)$  denotes the topological  $K$ -theory of  $G$ ,  $K_j(\mathcal{A})$  denotes  $K$ -theory for the  $C^*$ -algebra  $\mathcal{A}$ . In addition,  $\text{HP}_j(\mathcal{H}(G))$  denotes periodic cyclic homology of the algebra  $\mathcal{H}(G)$ , and  $\text{HP}_j(\mathcal{C}(G))$  denotes periodic cyclic homology of the topological algebra  $\mathcal{C}(G)$ . For periodic cyclic homology, see [12, 2.4].

The Baum-Connes assembly map  $\mu$  is an isomorphism [3, 15]. The map

$$H_*(G; \beta^1 G) \otimes_{\mathbb{Z}} \mathbb{C} \longrightarrow \text{HP}_*(\mathcal{H}(G))$$

is an isomorphism [14, 20]. The map  $\iota_*$  is an isomorphism by [3, 6]. The right hand Chern character is constructed in [7] and is an isomorphism after tensoring over  $\mathbb{Z}$  with  $\mathbb{C}$  [7, Theorem 3]. The left hand Chern character is the unique map for which the diagram is commutative.

## B The Bernstein spectrum

Let  $G$  be the group of  $F$ -points of a connected reductive algebraic group defined over  $F$ . We consider pairs  $(L, \sigma)$  where  $L$  is a Levi subgroup of a parabolic subgroup of  $G$ , and  $\sigma$  is an irreducible supercuspidal representation of  $L$ . We say two such pairs  $(L_1, \sigma_1)$ ,  $(L_2, \sigma_2)$  are *inertially equivalent* if there exist  $g \in G$  and an unramified quasicharacter  $\chi$  of  $L_2$  such that

$$L_2 = L_1^g \quad \text{and} \quad \sigma_1^g \cong \sigma_2 \otimes \chi.$$

Here,  $L_1^g := g^{-1}L_1g$  and  $\sigma_1^g(x) = \sigma_1(gxg^{-1})$  for all  $x \in L_1^g$ . We write  $[L, \sigma]_G$  for the inertial equivalence of the pair  $(L, \sigma)$  and  $\mathfrak{B}(G)$  for the set of all inertial equivalence classes. The set  $\mathfrak{B}(G)$  is the *Bernstein spectrum* of  $G$ . We will write  $\mathfrak{s} \in \mathfrak{B}(G)$ .

The Hecke algebra  $\mathcal{H}(G)$  is a unital  $\mathcal{H}(G)$ -module via left multiplication, and admits the canonical Bernstein decomposition as a purely algebraic direct

sum of two-sided ideals:

$$\mathcal{H}(G) = \bigoplus_{\mathfrak{s} \in \mathfrak{B}(G)} \mathcal{H}(G)^{\mathfrak{s}}.$$

This determines the canonical Bernstein decomposition of the reduced  $C^*$ -algebra as a  $C^*$ -direct-sum of two-sided  $C^*$ -ideals:

$$\mathcal{A} = \bigoplus_{\mathfrak{s} \in \mathfrak{B}(G)} \mathcal{A}^{\mathfrak{s}}.$$

Now  $C^*$ -direct sums are respected by the  $K$ -theory of  $C^*$ -algebras, and we have

$$K_j(\mathcal{A}) = \bigoplus_{\mathfrak{s} \in \mathfrak{B}(G)} K_j(\mathcal{A}^{\mathfrak{s}}) \quad (58)$$

with  $j = 0, 1$ . The abelian groups  $K_j(\mathcal{A}^{\mathfrak{s}})$  are finitely generated free abelian groups, see [17].

We will define  $H_{\text{ev/odd}}(G; \beta^1 G)^{\mathfrak{s}}$  as the pre-image of  $K_j(\mathcal{A}^{\mathfrak{s}})$  via the commutative diagram in Appendix A:

$$H_{\text{ev}}(G; \beta^1 G)^{\mathfrak{s}} \cong K_0(\mathcal{A}^{\mathfrak{s}}), \quad H_{\text{odd}}(G; \beta^1 G)^{\mathfrak{s}} \cong K_1(\mathcal{A}^{\mathfrak{s}}). \quad (59)$$

## C The formula for the rank

Let  $\mathfrak{s}$  be a point in the Bernstein spectrum  $\mathfrak{B}(G)$ , so that  $\mathfrak{s} = [L, \sigma]_G$ . We have

$$L = \text{GL}(m_1)^{e_1} \times \cdots \times \text{GL}(m_r)^{e_r}$$

with  $m_1 e_1 + \cdots + m_r e_r = N$ . The numbers  $e_1, \dots, e_r$  are called the *exponents* of  $\mathfrak{s}$ , as in [6]. According to [6, Lemma 3.2], we then have

$$\text{rank } K_j(\mathcal{A}^{\mathfrak{s}}) = 2^{r-1} \beta(e_1) \cdots \beta(e_r) \quad (60)$$

where

$$\beta(e) = \sum 2^{\kappa(\pi)-1}.$$

In this formula,  $\pi$  is a partition of  $e$ , the sum is over all partitions of  $e$ , and  $\kappa(\pi)$  is the number of unequal parts of  $\pi$ . For example, if  $\pi$  is the partition  $1 + 1 + 1 + 3 + 3 + 3 + 3 + 7 + 9$  of 31 then  $\kappa(\pi) = 4$ .

The ranks of the finitely generated abelian groups  $H_{\text{ev/odd}}(G; \beta^1 G)^{\mathfrak{s}}$  are given by

$$\text{rank } H_{\text{ev/odd}}(G; \beta^1 G)^{\mathfrak{s}} = 2^{r-1} \beta(e_1) \cdots \beta(e_r). \quad (61)$$

## D Invariants attached to $\mathfrak{s}$

We write the supercuspidal representation  $\sigma$  of the Levi subgroup

$$M \cong \prod_{i=1}^q \prod_{j=1}^{c_i} \mathrm{GL}(N_{i,j}, F)$$

as a vector  $\sigma = (\sigma_{1,1}, \dots, \sigma_{1,c_1}, \sigma_{2,1}, \dots, \sigma_{2,c_2}, \dots, \sigma_{q,1}, \dots, \sigma_{q,c_q})$  where  $\sigma_{i,j}$  is an irreducible supercuspidal representation of  $\mathrm{GL}(N_{i,j}, F)$ , and for each  $i \in \{1, \dots, q\}$ , the representations  $\sigma_{i,j}$  ( $1 \leq j \leq c_i$ ) admit the same endo-class. At the same time, for all  $1 \leq j \leq c_i$  and  $1 \leq j' \leq c_{i'}$ , the representations  $\sigma_{i,j}$  and  $\sigma_{i',j'}$  have distinct endo-classes if  $i' \neq i$ . This implies that, for a given  $i$ , in the construction of Bushnell-Kutzko, all the representations  $\sigma_{i,j}$  ( $1 \leq j \leq c_i$ ) may be assumed to correspond to the same field extension  $E_i$  of  $F$ . Let  $e(E_i|F)$  denote the ramification index of  $E_i$  over  $F$ . Then the parahoric subgroup  $J^{\mathfrak{s}}$  only depends on the integers  $N_{i,j}$ ,  $c_i$  and  $e(E_i|F)$  (see Definition 6).

For supercuspidal representations, the parahoric subgroup is always the same one, say  $\mathrm{GL}(N, \mathfrak{o}_F)$ ; when  $q = 1$  (that is, only one endo-class), the parahoric is given by the integers  $N_{1,1}, \dots, N_{1,c_1}$ , which are the sizes of the blocks of  $M$ . In the general case, the parahoric subgroup depends on the sizes of the blocks of  $M$ , of the block decomposition defined by the endo-classes (that is, those corresponding to the Levi subgroup  $\bar{M} \cong \prod_{i=1}^q \mathrm{GL}(\bar{N}_i)$ , with  $\bar{N}_i = \sum_{j=1}^{c_i} N_{i,j}$ ) and on the ramification indices.

## References

- [1] J. Adler, A. Roche, *An intertwining result for  $p$ -adic groups*, Canad. J. Math. **52** (2000), no. 3, 449–467.
- [2] P. Baum, A. Connes, N. Higson, *Classifying space for proper actions and  $K$ -theory of groups  $C^*$ -algebras*, Contemporary Math. **167** (1994) 241–291.
- [3] P. Baum, N. Higson, R. J. Plymen, *A proof of the Baum-Connes conjecture for  $p$ -adic  $\mathrm{GL}(n)$* , C. R. Acad. Sci. Paris **325** (1997) 171–176.
- [4] P. Baum, N. Higson, R. J. Plymen, *Representations of  $p$ -adic groups: a view from operator algebras*, Proc. Symp. Pure Math. **68** (2001) 111–149.
- [5] J. Bernstein (rédigé par P. Deligne), *Le “centre” de Bernstein. Représentations des groupes réductifs sur un corps local*. Hermann, Paris (1984) 1–32.

- [6] J. Brodzki, R.J. Plymen, *Complex structure on the smooth dual of  $GL(n)$*  Documenta Math. **7** (2002) 91–112.
- [7] J. Brodzki, R.J. Plymen, *Chern character for the Schwartz algebra of  $p$ -adic  $GL(n)$*  Bull. London Math.Soc. **34** (2002) 219–228.
- [8] C.J. Bushnell, P.C. Kutzko, *The admissible dual of  $GL(N)$  via compact open subgroups*, Annals of Math. Study **129** (1993) Princeton University Press.
- [9] C.J. Bushnell, P.C. Kutzko, *Smooth representations of reductive  $p$ -adic groups: structure theory via types*, Proc. London Math. Soc. **77** (1998) 582–634.
- [10] C.J. Bushnell, P.C. Kutzko, *Semisimple types in  $GL_n$* , Compos. Math. **119** (1999) 53–97.
- [11] C.J. Bushnell, P.C. Kutzko, *Types in reductive  $p$ -adic groups: the Hecke algebra of a cover*, Proc. AMS **129** (2001) 601–607.
- [12] J. Cuntz, G. Skandalis, B. Tsygan, *Cyclic homology in noncommutative geometry*, EMS 121, Springer-Verlag, Berlin 2004.
- [13] S.I. Gelfand, Yu.I. Manin, *Homological algebra*, Springer-Verlag, Berlin 1999.
- [14] N. Higson, V. Nistor, *Cyclic homology of totally disconnected groups acting on buildings*, J. Functional Analysis **141** (1996) 466–485.
- [15] V. Lafforgue,  *$K$ -théorie bivariante pour les algèbres de Banach et conjecture de Baum-Connes*. Invent. Math. 149 (2002), no. 1, 1–95.
- [16] V. Paskunas, *Unicity of types for supercuspidals*, math.NT/0306124.
- [17] R.J. Plymen, *Reduced  $C^*$ -algebra of the  $p$ -adic group  $GL(n)$* , J. Functional Analysis **72** (1987) 1–12.
- [18] A. Roche, *Types and Hecke algebras for principal series representations of split reductive  $p$ -adic groups*, Ann. scient. Éc. Norm. Sup. **31** (1998) 361–413.
- [19] M. Ronan, *Lectures on buildings*, Academic press (1989).
- [20] P. Schneider, *Equivariant homology for totally disconnected groups*, J.Algebra **203** (1998) 50–68.

- [21] J. Tits, *Reductive groups over local fields*, in *Automorphic forms, representations and L-functions*, Proc. Symp. Pure Math. **33** (1979), part 1, 29–69.

Anne-Marie Aubert, Institut de Mathématiques de Jussieu, U.M.R. 7586 du C.N.R.S., 175 rue du Chevaleret 75013 Paris, France.

Email: aubert@math.jussieu.fr

Samir Hasan, Mathematics Department, Damascus University, Damascus, Syria.

Email: Samir@damascus.every1.net

Roger Plymen, School of Mathematics, Manchester University, M13 9PL, England.

Email: plymen@manchester.ac.uk