

Telomeres shorten and then lengthen before fledging in Magellanic penguins (*Spheniscus magellanicus*)

Jack A. Cerchiara¹, Rosa Ana Risques², Donna Prunkard², Jeffrey R. Smith³, Olivia J. Kane¹, P. Dee Boersma^{1,4,5}

¹Department of Biology, University of Washington, Seattle, WA 98195, USA

²Department of Pathology, University of Washington, Seattle, WA 98195, USA

³School of Environmental and Forest Sciences, University of Washington, Seattle, WA 98195, USA

⁴Wildlife Conservation Society, Bronx, NY 10460, USA

⁵Global Penguin Society, University of Washington, Seattle, WA 98195, USA

Correspondence to: Jack Cerchiara; **email:** jackc44@u.washington.edu

Keywords: telomeres, growth, life history, aging

Received: December 8, 2016

Accepted: February 1, 2017

Published: February 8, 2017

ABSTRACT

For all species, finite metabolic resources must be allocated toward three competing systems: maintenance, reproduction, and growth. Telomeres, the nucleoprotein tips of chromosomes, which shorten with age in most species, are correlated with increased survival. Chick growth is energetically costly and is associated with telomere shortening in most species. To assess the change in telomeres in penguin chicks, we quantified change in telomere length of wild known-age Magellanic penguin (*Spheniscus magellanicus*) chicks every 15 days during the species' growth period, from hatching to 60 days-of-age. Magellanic penguins continue to grow after fledging so we also sampled a set of 1-year-old juvenile penguins, and adults aged 5 years. Telomeres were significantly shorter on day 15 than on hatch day but returned to their initial length by 30 days old and remained at that length through 60 days of age. The length of telomeres of newly hatched chicks, chicks aged 30, 45 and 60 days, juveniles, and adults aged 5 years were similar. Chicks that fledged and those that died had similar telomere lengths. We show that while telomeres shorten during growth, Magellanic penguins elongate telomeres to their length at hatch, which may increase adult life span and reproductive opportunities.

INTRODUCTION

In all species, finite metabolic resources must be allocated toward three competing systems: survival (or maintenance), reproduction, and growth [1, 2]. All species incur a resource bias towards growth during development [2-4]. Growth is energetically costly, and diverts resources away from maintenance systems in favor of growth [5, 6]. However, even with energetically demanding growth periods, species should evolve processes that maintain physiological systems important to survival.

All birds grow rapidly [5, 6]. Short-lived species with low survival rates develop and reach sexual maturity quickly [6-8]. With a decreased annual survival rate;

resource investment should favor growth and reproductive, rather than maintenance systems. In long-lived species, however, there is a fitness advantage to adult survival. While long-lived species will still incur the resource bias during growth, they should also allocate resources to systems that benefit increased longevity. Evidence of this tradeoff is known, as developmental conditions like brood size [9], growth rate [10] and being smaller than brood-mates [11] are shown to affect survival [12, 13]. We hypothesize that the continual investment cost in maintenance systems, like telomeres, increases the potential of adult survival and future reproductive events.

Telomeres are nucleoprotein complexes that protect the ends of chromosomes [14]. During each cycle of cell replication, telomeres are shortened because DNA

polymerase cannot fully replicate the 3' end of the DNA [14-16]. Telomere shortening is correlated negatively with adult survival in most species [17-20], and telomeres shorten more slowly in long-lived birds than in shorter-lived species [17]. In long-lived species, there is a fitness advantage to increased survival through more reproductive opportunities. Thus, long-lived species should allocate resources to increase adult survival, like telomeres.

In most species, growth is implicated in the shortening of telomeres. In King penguins (*Aptenodytes patagonicus*), chicks that had a faster growth rate showed higher oxidative damage and had accelerated telomere loss [10]. One explanation for the shortening is oxidative damage, which results in DNA damage and shorter telomeres [21-24]. Specifically, reactive oxygen species (ROS) produce DNA strand breaks, which are difficult to repair when they occur in telomeric DNA [24]. Reactive oxygen species are byproducts of mitochondrial metabolism, which is increased during growth [25, 26]. Thus, telomere shortening might be caused by the increase in mitochondrial metabolism during growth [24, 27]. While telomere length may shorten during the growth period, if telomere length and rate of shortening is important to survival, fledging with longer telomeres may maximize lifespan.

While telomeres shorten in most adult vertebrates, they can also be elongated by two primary mechanisms; (1) through the action of telomerase, a ribonucleic reverse transcriptase (TERT), or (2) homologous recombination-mediated DNA replication, termed the Alternative Lengthening of Telomeres (ALT) [28, 29]. Telomerase contains both an enzyme with reverse transcriptase activity to elongate linear DNA as well as carries its own RNA template [30-32]. Some vertebrates show increased telomerase activity during development [10, 31, 33, 34]. Additionally, increased telomerase activity in regions where cell lines develop could allow mature somatic cells to have longer telomeres as the individual ages. Telomeres in human sperm cells display this phenomenon. Sperm cell telomeres from older individuals are longer, as they are likely acted upon by enhanced telomerase activity in the gonads [35]. Additionally, telomerase activity is implicated in enhanced longevity observed in some taxa, including lobsters [36], *hydra* and *planaria* [37], of which the latter two are thought to be biologically immortal. A number of transcription factors regulate the activity of telomerase, so the expression is highly variable among cell conditions [38]. Oncogenes, however, can activate these transcription factors, and therefore telomerase is up-regulated in approximately 85-90% of cancer cases [38, 39].

Magellanic penguins (*Spheniscus magellanicus*) in the wild live more than 30 years [40], exceeding their maximum, mass-predicted lifespan by 26% [41]. We predict that, if growth is a stressor, and evolution favors adult penguins having the highest chance of survival, Magellanic penguins should finish growth with the longest possible telomeres. We quantified telomere length during growth of Magellanic penguins to characterize the impact of high cellular proliferation and energy demand on telomeres. We predicted that: (a) in young chicks, the lengths of telomeres would decrease but (b) telomere lengths would stop shortening by the time chicks reached fledging age.

RESULTS

Telomeres were significantly shorter when chicks were 15 days of age than when they hatched (Figure 1). By 30 days of age, telomere length was similar to their length at hatching, and they remained unchanged through 60 days of age ($p=0.002$, $n=65$, TukeyHSD, Figure 1). Telomeres of newly hatched chicks, 1-year-old juveniles, and adults aged 5 years were similar in length ($p=0.14$, $n=51$). There was no relationship between the change of bill depth, flipper length, foot length or weight and the change in telomere length between day 0 and day 15 (BD : $Rsq = 0.246$, $p=0.07$, $n = 11$; FL : $Rsq = 0.129$, $p = 0.15$, $n = 11$; FT : $Rsq = -0.076$, $p = 0.60$, $n = 11$; WT : $Rsq = 0.101$, $p = 0.18$, $n = 11$). However, there was a significant correlation in the change of bill length and telomere length change from day 0 to day 15 ($Rsq = 0.374$, $p = 0.026$, $n = 11$). One chick, that had very little change between 0 and 15 days, drove this trend, so it is unlikely to be biologically meaningful. Those individuals that survived to fledging and those that died while nestlings had similar telomere lengths ($t= 0.792$, $p=0.43$, $n=20$).

DISCUSSION

Telomeres shorten from hatching to day 15, but telomeres of newly hatched chicks, chicks ≥ 30 days, 1-year-old juveniles, and adults aged 5 years were similar in length. As was predicted, Magellanic penguin telomeres shorten immediately after hatch. Growth, in young chicks, requires high levels of energy and is characterized by high metabolism [5, 42, 43]. A high metabolic state can release reactive oxygen species (ROS), which shorten telomeres [24, 34]. This mechanism most likely explains what we observed between hatch day and day 15. King penguin chicks also showed a similar correlative link between growth and telomere shortening [10]. After day 15, however, Magellanic penguin telomeres elongated back to hatch day length by fledging and penguins entered adult size with telomeres of similar length to when they hatched.




Figure 1. Telomere length shortens and then elongates in Magellanic penguins. Telomere length shortens in early life, but returns to hatch day length by age 30 days. Groups (A/B) that do not share the same heading letter are significantly different. Dark bars within boxes represent group means, sample sizes presented under means, and bars are standard error.

This phenomena was not observed in King penguins [10]. King penguins live only 73% of their mass-predicted lifespan, where lifespan = $17.6 (mass \text{ in kg})^{0.20}$ [41], shorter than Magellanic penguins [40, 44]. Since telomeres are linked to adult survival in most species, elongation of telomeres to hatch length by the time Magellanic penguins are adult size could contribute to this longevity [17-20]. In fact, we showed previously that Magellanic penguins conserve their telomere length from age 5 years to older than 27 years, even if they reproduce more (Cerchiara et al, in review).

Telomerase elongates telomeres [14, 45]. After birth in most mammals, telomeres in somatic cells begin shortening [46], and, in adults, telomerase is only found up-regulated in germ and stem cell lines and remains at low levels in somatic cell lines of kidneys, lymphocytes and epithelial cells, which is likely part of a tumor suppression mechanism [30, 32, 47, 48]. There is some evidence, however, that telomerase may remain active in some adult long-lived birds (*Sterna hirundo*; *Oceanodroma leucorhoa*) compared to short lived ones [45], which could explain the telomere elongation we observed as it is the only known reverse-transcriptase to elongate telomeres [30, 32, 47-53]. To our knowledge, late-growth elongation, like we observed, is only known for one species, the fish *Oryzias latipes* [54]. Increased telomerase activity is the proposed mechanism for developmental telomere elongation observed in *O. latipes* [54], so this is likely the same mechanism

contributing to our results. Although we did not measure telomerase directly, we measured an increase in telomere length from shortest length on day 15 back to hatch day length by day 30, similar to what was shown in *O. latipes*.

Magellanic penguins elongated telomeres by the end of their growth period, making them one of the few species to evolve a physiological system that allows telomeres to recover to their original length at hatching. This mechanism allows Magellanic penguins to enter reproductive age with telomere lengths comparable to their length at hatching.

MATERIALS AND METHODS

Samples

In 1982, we began studying Magellanic penguins at Punta Tombo, Argentina (44°02'S, 65°11'W), banding chicks, juveniles, and breeding adults [55]. We know the lay date of individual eggs, and because we checked the nests daily before hatching, we know the exact hatch day of individual chicks [56].

We collected blood from wild Magellanic penguin chicks from hatching (November 2010) to fledging or death (January 2011). Chicks were marked on the day they hatched and bled on hatch day (day 0) and then captured and bled every 15 days until they were 60 days

of age. Chick mortality reduced sample sizes during the nestling period: hatching day (“day 0”, n=20 chicks), 15 days old (n=12), 30 days old (n=11), 45 days old (n=11), and 60 days old (n=11). We also collected blood from 20 juveniles (1 year, January 2011). We collected blood from 15 known-age adult Magellanic penguins, aged 5 years from September to December 2007 at Punta Tombo, Argentina. We measured morphological traits (bill, flipper, foot and weight) of all individuals on the day of sample collection [55].

Sample collection and processing

We collected whole blood into a heparinized capillary tube (Thermo Fisher Scientific Inc.), and immediately placed the blood into anti-lyses buffer (10% DMSO/90% Newborn Bovine Serum), and placed the samples on ice. We froze the samples at -18°C within 1hr of collection. We stored the samples at -80°C at the University of Washington, until processing.

We extracted DNA from a lightly centrifuged cell pack, consisting primarily of erythrocytes (Qiagen DNeasy Mini-kit), and then quantified DNA via nanodrop spectrophotometer (*mean 260/280 ratio*±SE = 1.85±0.02). We measured telomere length by Quantitative Polymerase Chain Reaction (qPCR), a method used to measure telomere length in other penguins [10, 21].

qPCR – telomere length

We ran two PCR assays for each sample. The first PCR amplified the telomeric DNA and the second amplified a single-copy control gene (36B4, acidic ribosomal phosphoprotein PO). The control gene PCR is used to normalize the starting amount of DNA. We included a four-point standard curve (2-fold serial dilutions from 10 to 1.25ng of DNA) in all PCRs to allow the transformation of Ct (cycle threshold) into nanograms of DNA. All samples were run in triplicates and for analyses we used the median value.

We ran all PCR reactions in a Rotor-Gene 3000 (Corbett Research, Sydney, Australia), with a final volume of 20ul including: 1X PCR buffer (Invitrogen, Carlsbad, CA), 0.2mM dNTPs, 0.4X SybrGreen (Molecular Probes, Eugene, OR), 2.5mM DTT, 1% DMSO and 5ng of DNA. The telomere PCR used 0.8ul of Platinum Taq (Invitrogen), 1.5mM MgCl₂, and 300nM of each primer (tel1b: CGGTTTGGTTTGGGTTTGGGTTTGGGTTTGGGTTTGGTT; tel2b: GGCTTGCCTTACCCTTACCCTTACCCTTACCCTTACCCTTACCCT). PCR conditions were: 95°C for 15min and 30 cycles of amplification at 95°C for

15sec and at 56°C for 60sec. For the control gene, we used 36B4 primers designed with Primer3 with the 36B4 gene sequence in the common chicken (*Gallus gallus*; NW_001471461.2) and zebra finch (*Taeniopygia guttata*; NW_002197395.1) from the nucleotide database, GenBank. The control-gene PCR used 0.5ul of Platinum Taq, 3.5mM MgCl₂, 300nM of forward primer (MAPE1: AGGGAGAAGAGGGACTGGAC) and 500nM of reverse primer (MAPE2-CAATCCCACACACACCTCAG). Control gene PCR conditions were: 95°C for 15min and 35 cycles of amplification at 95°C for 15sec and at 56°C for 20sec and 72°C for 20sec. A melting (dissociation) curve, run at the end of every control-gene PCR, confirmed the presence of a single amplification product [57]. We verified the primers showed the same product (at ~75bp) for all samples when ran on a 2% gel, with no secondary products. We targeted the 75bp oligomer of the 36B4 reference gene, so we are confident in the accuracy of our control gene [58].

We imported the raw data from the Rotor Gene software into Excel (v14.4.4) and aligned all amplification plots to a baseline height of 2% in the first 5 cycles of amplification. The fluorescence threshold for determination of the Ct was set at 20% of maximum products at the beginning of the exponential phase of the plot. We used a four point standard curve to calculate the corresponding nanograms for telomere analysis (mean telomere efficiency ± SE = 0.74±0.01, mean 36B4 efficiency ± SE = 0.88±0.02; all R_{sq} > 0.99). Due to lower efficiency values, we checked the absorbance curves for every sample and, if replicates were not overlapping (visual evaluation), we removed these data from our analyses. We re-ran removed samples again on additional trials, and evaluated the absorbance curves again.

The amount of telomeric DNA was divided by the amount of control-gene DNA, producing a relative measurement of the telomere length of the sample. For every trial, we ran two control samples for normalization between trials and to assess reproducibility of accurate measurements. The intra- and inter-trial variability (coefficient of variation) for the qPCR was 7% and 8%, respectively [59].

Statistics

Previous work showed that, in 60 days, Magellanic penguins develop from newly hatched chicks (~76g) to fledging weight (~1800g) [55]. Since we used repeated samples of the same individuals during growth all analyses used a mixed-effects model approach with age (as a factor) as a fixed effect and chick-ID as a random effect to test if telomere length was predicted by chick

age. We used a TukeyHSD to test for difference between age groups for chicks and for adults.

We used a linear model to determine if the growth rate (percent change in body part measurements between sample dates) of body parts predicted telomere length (percent change in telomere length). We also used a binomial linear model to determine in the telomere length at hatching predicted those chicks that fledged and those that died during growth. For statistical tests, we used R Statistical software (R Foundation for Statistical Computing: Development Core Team (v3.1.3)).

AUTHOR CONTRIBUTIONS

JAC and PDB designed the research, JAC, RAR, DP, JRS and OJK performed the research, PDB, RAR and DP contributed reagents and analytical tools, RAR and JAC analyzed data, JAC wrote the paper, and JAC, RAR, DP, JRS, OJK and PDB assisted in editing the paper.

ACKNOWLEDGEMENTS

The Penguin Project is affiliated with the Wildlife Conservation Society (WCS), the Global Penguin Society (GPS), the University of Washington, National Scientific and Technical Research Council (CONICET) and Centro Nacional Patagonico (CENPAT), Argentina. The research was carried out under a joint agreement between the WCS and the Office of Tourism, Province of Chubut, Argentina. The La Regina family gave permission to work on their land, and we had permits from Chubut to work in the Provincial reserve of Punta Tombo from provincial authorities in the departments of Turismo and Flora and Fauna. We thank the many students and field volunteers who collected data over the years. Garcia Borboroglu, William Conway, Graham Harris, and Patricia Harris facilitated the research and helped with permits. We also thank Calvin Ngo for his help with the laboratory analysis, and Courtney Wenneborg for help in the field. We thank Gordon Orians, Ginger Rebstock and anonymous reviewers for improving the manuscript.

CONFLICTS OF INTEREST

The authors have no conflict of interests to declare.

FUNDING

Funds for research have included the WCS, Exxonmobil Foundation, the Pew Fellows Program in Marine Conservation, the Disney Worldwide Conservation Fund, the Chase, Cunningham, CGMK, Offield, Peach,

Thorne, Tortuga, and Kellogg Foundations, the Wadsworth Endowed Chair in Conservation Science, and Friends of the Penguins. The NASA Space Grant Fellowship and the Wadsworth Student Fellowship supported JAC. In addition, the Offield Fellowship and Riddiford-Truman Fellowship supported JAC in field and laboratory work.

REFERENCES

1. Stearns SC. Life-history tactics: a review of the ideas. *Q Rev Biol.* 1976; 51:3–47. doi: 10.1086/409052
2. Wikelski M, Ricklefs RE. The physiology of life histories. *Trends Ecol Evol.* 2001; 16:479–81. doi: 10.1016/S0169-5347(01)02279-0
3. Ricklefs RE, Wikelski M. The physiology / life- history nexus. *Trends Ecol Evol.* 2002; 17:462–68. doi: 10.1016/S0169-5347(02)02578-8
4. Austad SN. An experimental paradigm for the study of slowly aging organisms. *Exp Gerontol.* 2001; 36:599–605. doi: 10.1016/S0531-5565(00)00229-1
5. Ricklefs RE. Preliminary models for growth rates in altricial birds. *Ecology.* 1969; 506:1031–39. doi: 10.2307/1936894
6. Ricklefs RE. Density Dependence, Evolutionary Optimization, and the Diversification of Avian Life Histories. *Condor.* 2000; 102:9. doi: 10.1650/0010-5422(2000)102[0009:DDEOAT]2.0.CO;2
7. Saether BE. Pattern of covariation between life-history traits of European birds. *Nature.* 1988; 331:616–17. doi: 10.1038/331616a0
8. Promislow D, Harvey P. Living fast and dying young: A comparative analysis of life-history variation among mammals. *J Zool (Lond).* 1990; 220:417–37. doi: 10.1111/j.1469-7998.1990.tb04316.
9. Voillemot M, Hine K, Zahn S, Criscuolo F, Gustafsson L, Doligez B, Bize P. Effects of brood size manipulation and common origin on phenotype and telomere length in nestling collared flycatchers. *BMC Ecol.* 2012; 12:17. doi: 10.1186/1472-6785-12-17
10. Geiger S, Le Vaillant M, Lebard T, Reichert S, Stier A, LE Maho Y, Criscuolo F. Catching-up but telomere loss: half-opening the black box of growth and ageing trade-off in wild king penguin chicks. *Mol Ecol.* 2012; 21:1500–10. doi: 10.1111/j.1365-294X.2011.05331.x
11. Nettle D, Monaghan P, Gillespie R, Brilot B, Bedford T, Bateson M. An experimental demonstration that early-life competitive disadvantage accelerates telomere loss. *Proc Biol Sci.* 2015; 282:20141610. doi: 10.1098/rspb.2014.1610
12. Pauliny A, Wagner RH, Augustin J, Szép T, Blomqvist D.

- Age-independent telomere length predicts fitness in two bird species. *Mol Ecol.* 2006; 15:1681–87. doi: 10.1111/j.1365-294X.2006.02862.x
13. Boonekamp JJ, Mulder GA, Salomons HM, Dijkstra C, Verhulst S. Nestling telomere shortening, but not telomere length, reflects developmental stress and predicts survival in wild birds. *Proc Biol Sci.* 2014; 281:20133287. doi: 10.1098/rspb.2013.3287
 14. Blackburn EH. Structure and function of telomeres. *Nature.* 1991; 350:569–73. doi: 10.1038/350569a0
 15. Watson JD. Origin of concatemeric T7 DNA. *Nat New Biol.* 1972; 239:197–201. doi: 10.1038/newbio239197a0
 16. McClintock B. The Stability of Broken Ends of Chromosomes in Zea Mays. *Genetics.* 1941; 26:234–82.
 17. Haussmann MF, Winkler DW, O'Reilly KM, Huntington CE, Nisbet IC, Vleck CM. Telomeres shorten more slowly in long-lived birds and mammals than in short-lived ones. *Proc Biol Sci.* 2003; 270:1387–92. doi: 10.1098/rspb.2003.2385
 18. Haussmann MF, Mauck RA. Telomeres and longevity: testing an evolutionary hypothesis. *Mol Biol Evol.* 2008; 25:220–28. doi: 10.1093/molbev/msm244
 19. Salomons HM, Mulder GA, van de Zande L, Haussmann MF, Linskens MH, Verhulst S. Telomere shortening and survival in free-living corvids. *Proc Biol Sci.* 2009; 276:3157–65. doi: 10.1098/rspb.2009.0517
 20. Bize P, Criscuolo F, Metcalfe NB, Nasir L, Monaghan P. Telomere dynamics rather than age predict life expectancy in the wild. *Proc Biol Sci.* 2009; 276:1679–83. doi: 10.1098/rspb.2008.1817
 21. Beaulieu M, Reichert S, Maho YL, Ancel A, Criscuolo F. Oxidative status and telomere length in a long-lived bird facing a costly reproductive event. *Funct Ecol.* 2011; 25:577–85. doi: 10.1111/j.1365-2435.2010.01825.x
 22. Kotrschal A, Ilmonen P, Penn DJ. Stress impacts telomere dynamics. *Biol Lett.* 2007; 3:128–30. doi: 10.1098/rsbl.2006.0594
 23. Kirkwood TB, Austad SN. Why do we age? *Nature.* 2000; 408:233–38. doi: 10.1038/35041682
 24. von Zglinicki T. Oxidative stress shortens telomeres. *Trends Biochem Sci.* 2002; 27:339–44. doi: 10.1016/S0968-0004(02)02110-2
 25. Miquel J, Economos AC, Fleming J, Johnson JE Jr. Mitochondrial role in cell aging. *Exp Gerontol.* 1980; 15:575–91. doi: 10.1016/0531-5565(80)90010-8
 26. Harman D. Aging: a theory based on free radical and radiation chemistry. *J Gerontol.* 1956; 11:298–300. doi: 10.1093/geronj/11.3.298
 27. Passos JF, Saretzki G, von Zglinicki T. DNA damage in telomeres and mitochondria during cellular senescence: is there a connection? *Nucleic Acids Res.* 2007; 35:7505–13. doi: 10.1093/nar/gkm893
 28. Bryan TM, Englezou A, Gupta J, Bacchetti S, Reddel RR. Telomere elongation in immortal human cells without detectable telomerase activity. *EMBO J.* 1995; 14:4240–48.
 29. Counter CM, Avilion AA, LeFeuvre CE, Stewart NG, Greider CW, Harley CB, Bacchetti S. Telomere shortening associated with chromosome instability is arrested in immortal cells which express telomerase activity. *EMBO J.* 1992; 11:1921–29.
 30. Tollefsbol TO, Andrews LG. Mechanisms for telomerase gene control in aging cells and tumorigenesis. *Med Hypotheses.* 2001; 56:630–37. doi: 10.1054/mehy.2000.1241
 31. Chan SR, Blackburn EH. Telomeres and telomerase. *Philos Trans R Soc Lond B Biol Sci.* 2004; 359:109–21. doi: 10.1098/rstb.2003.1370
 32. Dong CK, Masutomi K, Hahn WC. Telomerase: regulation, function and transformation. *Crit Rev Oncol Hematol.* 2005; 54:85–93. doi: 10.1016/j.critrevonc.2004.12.005
 33. Youngren K, Jeanclos E, Aviv H, Kimura M, Stock J, Hanna M, Skurnick J, Bardeguet A, Aviv A. Synchrony in telomere length of the human fetus. *Hum Genet.* 1998; 102:640–43. doi: 10.1007/s004390050755
 34. Monaghan P, Haussmann MF. Do telomere dynamics link lifestyle and lifespan? *Trends Ecol Evol.* 2006; 21:47–53. doi: 10.1016/j.tree.2005.11.007
 35. Eisenberg DT, Hayes MG, Kuzawa CW. Delayed paternal age of reproduction in humans is associated with longer telomeres across two generations of descendants. *Proc Natl Acad Sci USA.* 2012; 109:10251–56. doi: 10.1073/pnas.1202092109
 36. Klapper W, Kühne K, Singh KK, Heidorn K, Parwaresch R, Krupp G. Longevity of lobsters is linked to ubiquitous telomerase expression. *FEBS Lett.* 1998; 439:143–46. doi: 10.1016/S0014-5793(98)01357-X
 37. Tan TC, Rahman R, Jaber-Hijazi F, Felix DA, Chen C, Louis EJ, Aboobaker A. Telomere maintenance and telomerase activity are differentially regulated in asexual and sexual worms. *Proc Natl Acad Sci USA.* 2012; 109:4209–14. doi: 10.1073/pnas.1118885109
 38. Cong YS, Wen J, Bacchetti S. The human telomerase catalytic subunit hTERT: organization of the gene and characterization of the promoter. *Hum Mol Genet.* 1999; 8:137–42. doi: 10.1093/hmg/8.1.137

39. Kim NW, Piatyszek MA, Prowse KR, Harley CB, West MD, Ho PL, Coviello GM, Wright WE, Weinrich SL, Shay JW. Specific association of human telomerase activity with immortal cells and cancer. *Science*. 1994; 266:2011–15. doi: 10.1126/science.7605428
40. Boersma PD, Esteban F, Kane O, Pozzi L, Putz K, Rey A, Rebstock R, Simeone A, Smith J, Van Buren A, Yorio P, Borboroglu P. (2013). Magellanic Penguins. In: Borboroglu P and Boersma PD, eds. *Penguins: Natural History and Conservation*. (Seattle: University of Washington), pp. 233–263.
41. Lindstedt S, Calder W. Body size and longevity in birds. *The Condor*. 1976; 78:91–94.
42. Breuner CW. (2010) Stress and Reproduction in Birds. In: D. Norris and K. Lopez, eds. *Hormones and Reproduction in Vertebrates*. (Academic Press), pp. 129–151.
43. Ricklefs RE. Natural selection and the development of mortality rates in young birds. *Nature*. 1969; 223:922–25. doi: 10.1038/223922a0
44. Halsey LG, Le Coadic M, Handrich Y. Estimating the critical body mass of king penguins. *Polar Biol*. 2008; 31:395–98. doi: 10.1007/s00300-007-0395-7
45. Haussmann MF, Winkler DW, Huntington CE, Nisbet IC, Vleck CM. Telomerase activity is maintained throughout the lifespan of long-lived birds. *Exp Gerontol*. 2007; 42:610–18. doi: 10.1016/j.exger.2007.03.004
46. Bekaert S, Derradji H, Baatout S. Telomere biology in mammalian germ cells and during development. *Dev Biol*. 2004; 274:15–30. doi: 10.1016/j.ydbio.2004.06.023
47. Stewart SA, Hahn WC, O'Connor BF, Banner EN, Lundberg AS, Modha P, Mizuno H, Brooks MW, Fleming M, Zimonjic DB, Popescu NC, Weinberg RA. Telomerase contributes to tumorigenesis by a telomere length-independent mechanism. *Proc Natl Acad Sci USA*. 2002; 99:12606–11. doi: 10.1073/pnas.182407599
48. Hornsby PJ. Telomerase and the aging process. *Exp Gerontol*. 2007; 42:575–81. doi: 10.1016/j.exger.2007.03.007
49. Bernardes de Jesus B, Vera E, Schneeberger K, Tejera AM, Ayuso E, Bosch F, Blasco MA. Telomerase gene therapy in adult and old mice delays aging and increases longevity without increasing cancer. *EMBO Mol Med*. 2012; 4:691–704. doi: 10.1002/emmm.201200245
50. Flores I, Blasco MA. The role of telomeres and telomerase in stem cell aging. *FEBS Lett*. 2010; 584:3826–30. doi: 10.1016/j.febslet.2010.07.042
51. Forsyth NR, Wright WE, Shay JW. Telomerase and differentiation in multicellular organisms: turn it off, turn it on, and turn it off again. *Differentiation*. 2002; 69:188–97. doi: 10.1046/j.1432-0436.2002.690412.x
52. Jaskelioff M, Muller FL, Paik JH, Thomas E, Jiang S, Adams AC, Sahin E, Kost-Alimova M, Protopopov A, Cadiñanos J, Horner JW, Maratos-Flier E, Depinho RA. Telomerase reactivation reverses tissue degeneration in aged telomerase-deficient mice. *Nature*. 2011; 469:102–06. doi: 10.1038/nature09603
53. Prowse KR, Greider CW. Developmental and tissue-specific regulation of mouse telomerase and telomere length. *Proc Natl Acad Sci USA*. 1995; 92:4818–22. doi: 10.1073/pnas.92.11.4818
54. Hatakeyama H, Yamazaki H, Nakamura K, Izumiya-Shimomura N, Aida J, Suzuki H, Tsuchida S, Matsuura M, Takubo K, Ishikawa N. Telomere attrition and restoration in the normal teleost *Oryzias latipes* are linked to growth rate and telomerase activity at each life stage. *Aging (Albany NY)*. 2016; 8:62–76. doi: 10.18632/aging.100873
55. Boersma PD, Stokes D, Yorio P. (1990). Reproductive variability and historical change of Magellanic Penguins (*Spheniscus magellanicus*) at Punta Tombo, Argentina. In: Davis LS and Darby JT, eds. *Penguin Biology*. (San Diego: Academic Press), pp. 15–43.
56. Boersma PD, Rebstock GA. Climate change increases reproductive failure in Magellanic penguins. *PLoS One*. 2014; 9:e85602. doi: 10.1371/journal.pone.0085602
57. Ringsby TH, Jensen H, Pärn H, Kvalnes T, Boner W, Gillespie R, Holand H, Hagen IJ, Rønning B, Sæther BE, Monaghan P. On being the right size: increased body size is associated with reduced telomere length under natural conditions. *Proc Biol Sci*. 2015; 282:20152331. doi: 10.1098/rspb.2015.2331
58. O'Callaghan NJ, Fenech M. A quantitative PCR method for measuring absolute telomere length. *Biol Proced Online*. 2011; 13:3. doi: 10.1186/1480-9222-13-3
59. Martin-Ruiz CM, Baird D, Roger L, Boukamp P, Krunich D, Cawthon R, Dokter MM, van der Harst P, Bekaert S, de Meyer T, Roos G, Svenson U, Codd V, et al. Reproducibility of telomere length assessment: an international collaborative study. *Int J Epidemiol*. 2014.