

Distinct and Shared Roles of β -Arrestin-1 and β -Arrestin-2 on the Regulation of C3a Receptor Signaling in Human Mast Cells

Arpana Vibhuti¹, Kshitij Gupta¹, Hariharan Subramanian, Qiang Guo, Hydar Ali*

Department of Pathology, School of Dental Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, United States of America

Abstract

Background: The complement component C3a induces degranulation in human mast cells via the activation of cell surface G protein coupled receptors (GPCR; C3aR). For most GPCRs, agonist-induced receptor phosphorylation leads to the recruitment of β -arrestin-1/ β -arrestin-2; resulting in receptor desensitization and internalization. Activation of GPCRs also leads to ERK1/2 phosphorylation via two temporally distinct pathways; an early response that reflects G protein activation and a delayed response that is G protein independent but requires β -arrestins. The role of β -arrestins on C3aR activation/regulation in human mast cells, however, remains unknown.

Methodology/Principal Findings: We utilized lentivirus short hairpin (sh)RNA to stably knockdown the expression of β -arrestin-1 and β -arrestin-2 in human mast cell lines, HMC-1 and LAD2 that endogenously expresses C3aR. Silencing β -arrestin-2 attenuated C3aR desensitization, blocked agonist-induced receptor internalization and rendered the cells responsive to C3a for enhanced NF- κ B activity as well as chemokine generation. By contrast, silencing β -arrestin-1 had no effect on these responses but resulted in a significant decrease in C3a-induced mast cell degranulation. In shRNA control cells, C3a caused a transient ERK1/2 phosphorylation, which peaked at 5 min but disappeared by 10 min. Knockdown of β -arrestin-1, β -arrestin-2 or both enhanced the early response to C3a and rendered the cells responsive for ERK1/2 phosphorylation at later time points (10–30 min). Treatment of cells with pertussis toxin almost completely blocked both early and delayed C3a-induced ERK1/2 phosphorylation in β -arrestin1/2 knockdown cells.

Conclusion/Significance: This study demonstrates distinct roles for β -arrestins-1 and β -arrestins-2 on C3aR desensitization, internalization, degranulation, NF- κ B activation and chemokine generation in human mast cells. It also shows that both β -arrestin-1 and β -arrestin-2 play a novel and shared role in inhibiting G protein-dependent ERK1/2 phosphorylation. These findings reveal a new level of complexity for C3aR regulation by β -arrestins in human mast cells.

Citation: Vibhuti A, Gupta K, Subramanian H, Guo Q, Ali H (2011) Distinct and Shared Roles of β -Arrestin-1 and β -Arrestin-2 on the Regulation of C3a Receptor Signaling in Human Mast Cells. PLoS ONE 6(5): e19585. doi:10.1371/journal.pone.0019585

Editor: Karl-Wilhelm Koch, University of Oldenburg, Germany

Received: March 5, 2011; **Accepted:** April 6, 2011; **Published:** May 12, 2011

Copyright: © 2011 Vibhuti et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by National Institutes of Health Grants HL085774 and AI080852. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: alih@upenn.edu

These authors contributed equally to this work.

Introduction

The anaphylatoxin C3a is generated following bacterial infection and from IgE/Fc ϵ RI stimulated human mast cells [1]. Accordingly, C3a has been proposed to play critical roles in innate immunity and allergic diseases such as asthma [2,3,4]. C3a activates its cell surface G protein coupled receptor (GPCR; C3aR) to induce chemotaxis in human mast cell line (HMC-1) and degranulation in human skin mast cells, peripheral blood CD34⁺ cell-derived mast cells and a differentiated mast cell line, LAD2 [1,5,6,7,8]. C3a induces mast cell degranulation via the activation of phospholipase C β and mobilization of intracellular Ca²⁺ [7,9]. However, the mechanism(s) involved in regulation of C3aR signaling in mast cells remain poorly understood.

It is well established that for most GPCRs, receptor phosphorylation by G protein coupled receptor kinases (GRKs)

and the subsequent recruitment of β -arrestin provides an important mechanism for their desensitization and internalization [10]. Two isoforms of β -arrestins, (β -arrestin-1 and β -arrestin-2) are known and each can differentially regulate GPCR desensitization and internalization. Thus, for protease activated receptor-1 (PAR-1) only β -arrestin-1 is capable for receptor desensitization but receptor internalization is independent of either β -arrestins [11]. By contrast, both isoforms of β -arrestins can promote desensitization of β 2-adrenergic receptors (β AR2) and angiotensin II type 1A receptor (AT₁AR) [12]. Although, only β -arrestin-2 promotes internalization of β AR2 both isoforms are required for the internalization of AT₁AR. We have previously shown that in transfected rat basophilic leukemia (RBL-2H3) cells, C3aR associates with β -arrestin-2 following agonist stimulation [9]. However, the roles of β -arrestin-1 and β -arrestin-2 on C3aR desensitization and internalization have not been determined.

Previous studies with transfected RBL-2H3 cells showed that phosphorylation-deficient chemoattractant/chemokine receptors that do not associate with β -arrestins respond to ligands for more sustained Ca^{2+} mobilization and degranulation when compared with cells expressing wild-type receptors [9,13,14,15,16]. These findings are consistent with the view that β -arrestins play an important role in desensitization. By contrast, activation of the chemokine receptor CXCR1 in human neutrophils leads to receptor internalization and complex formation between β -arrestin-2 and Src kinases, (Hck and c-Fgr) which translocate to secretory granules to promote degranulation [17]. β -arrestin-2 also forms a complex with Ral-GDP dissociation stimulator (Ral-GDS) in the cytoplasm of human neutrophils [18]. Furthermore, activation of fMLP receptor results in the translocation of the complex to the plasma membrane. This is followed by the release of Ral-GDS from β -arrestin and the activation of Ral resulting in actin cytoskeleton rearrangement presumably leading to degranulation. The roles of β -arrestins on C3a-induced mast cell degranulation, however, remain unknown.

In addition to receptor desensitization, internalization and degranulation, β -arrestins modulate the activity of the transcription factor, NF- κ B. Witherow et al., [19], using a yeast two-hybrid screen, first demonstrated that the inhibitor of NF- κ B, I κ B α binds to β -arrestin-1. Furthermore, both β -arrestin-1 and 2 interact with I κ B α in transfected cells. However, siRNA-mediated knockdown studies indicated that β -arrestin-1 but not β -arrestin-2 inhibits TNF- α -induced NF- κ B activation. By contrast, Gao et al., [20] showed that β -arrestin-2, but not β -arrestin-1 interacts with I κ B α to inhibit NF- κ B activation. Studies with primary leukocytes from β -arrestin-2 knockout mice showed that this adapter molecule is involved in the internalization of the chemokine receptor, CXCR2 [21]. Furthermore, *in vivo* studies showed that β -arrestin-2 deletion promotes tumor growth and angiogenesis and these responses are associated with enhanced chemokine generation [22]. Other studies have shown that both β -arrestins promote NF- κ B activation following the activation GPCRs [23,24]. The roles of β -arrestins on C3a-induced NF- κ B activation and chemokine production in mast cells have not been determined.

The mitogen-activated protein kinases (MAPKs), extracellular signal-regulated kinases 1 and 2 (ERK1/2) play important roles in a variety of cellular responses and have been studied for a variety

of GPCRs. Activation of GPCRs in transfected cell lines induce ERK1/2 phosphorylation via two temporally distinct pathways; an early response that reflects G protein activation and a delayed response that is G protein independent but requires the formation of signaling complexes involving Src/ERK with β -arrestin [25,26]. Although C3a induces ERK1/2 phosphorylation in mast cells [7,9,27], whether or not β -arrestin-1 and β -arrestin-2 regulate this response remains unknown.

The goal of the present study was to determine the roles of β -arrestins on the regulation of C3aR signaling in human mast cells. To achieve this goal, we utilized lentivirus shRNA to stably knockdown the expression of β -arrestin-1 and β -arrestin-2 in human mast cells (HMC-1 and LAD2). Using these systems, we report unexpected findings regarding distinct roles of these adapter molecules on C3aR desensitization, internalization, degranulation, NF- κ B activation and chemokine generation. Furthermore, we provide first demonstration that both β -arrestins acts as novel inhibitors of C3a-induced G-protein-mediated ERK1/2 phosphorylation in human mast cells.

Results

Stable knockdown of β -arrestin-1 and β -arrestin-2 in a human mast cell, HMC-1

To determine the role of β -arrestins on the regulation of C3aR signaling in mast cells, we used the Mission shRNA lentivirus system to stably knockdown the expression of β -arrestin-1 and β -arrestin-2 in a human mast cell line; HMC-1 cells. Cells were transduced with 5 different shRNA constructs targeting different regions of β -arrestin-1 and β -arrestin-2. For control, we used a scrambled shRNA construct. After transduction and selection with puromycin, quantitative real-time PCR was performed to determine the extent of β -arrestin knockdown. As shown in Fig. 1A and B, all five β -arrestin shRNA constructs decreased the expression of β -arrestin-1 and β -arrestin-2 to variable levels. Clone 3 (TRCN0000230149) for β -arrestin-1 and clone 5 (TRCN0000159482) for β -arrestin-2 showed >80% decrease in mRNA. We therefore, used these clones to generate double knockdown (both β -arrestin-1 and β -arrestin-2) in HMC-1 cells. As shown in Fig. 1C, we were able to generate




Figure 1. Stable knockdown of β -arrestin-1 and β -arrestin-2 in human mast cells. HMC-1 cells were stably transduced with scrambled shRNA control lentivirus or different clones of shRNA lentivirus targeted against β -arrestin-1 and β -arrestin-2 (Panels A and B). For double knockdown HMC-1 cell were transduced with shRNA lentivirus, Clone 3 of β -arrestin-1 and Clone 5 of β -arrestin-2 (C). Quantitative PCR was employed to assess β -arrestin-1 or -2 mRNA levels. Results are expressed as a ratio of β -arrestin to GAPDH mRNA levels. Data represent the mean \pm SEM from three independent experiments. Statistical significance was determined by one way ANOVA. ** indicates $p < 0.001$. doi:10.1371/journal.pone.0019585.g001

HMC-1 cells with $\sim 80\%$ knockdown of both genes. These cells were used in subsequent studies described below.

β -arrestin-2, but not β -arrestin-1, is required for C3aR desensitization and internalization


Intracellular Ca^{2+} mobilization provides a rapid, sensitive and real-time assay to measure receptor desensitization [28]. We have previously shown that receptors that undergo desensitization respond to agonists with an initial Ca^{2+} spike, which decays rapidly and reaches baseline within $\sim 2\text{--}3$ min [28]. By contrast, phosphorylation-deficient receptors that do not associate with β -arrestin respond to agonist for a similar initial Ca^{2+} spike, followed by a sustained response that remains elevated for an extended period of time [29,30]. We therefore, used Ca^{2+} mobilization as an assay to determine the effects of β -arrestin-1 and β -arrestin-2 knockdowns on C3aR desensitization. As shown in Fig. 2A and B, C3a caused a rapid increase in Ca^{2+} mobilization in shRNA control and β -arrestin-1 knockdown cells. By contrast, in β -arrestin-2 knockdown

C3a caused a similar initial spike but subsequent response was sustained (Fig. 2C). Furthermore, deletion of both β -arrestins resulted in a Ca^{2+} response similar to that observed in β -arrestin-2 knockdown cells (Fig. 2C and D). These findings suggest that β -arrestin-2, but not β -arrestin-1, mediates desensitization of C3aR.


GPCRs that undergo desensitization display reduced responsiveness to a second stimulation with the same agonist [28]. To test further the effects of β -arrestins on desensitization, shRNA control or knockdown cells were exposed to C3a and washed twice before re-exposure to the same concentration of C3a. In shRNA control and β -arrestin-1 knockdown cells, there was little or no response to second C3a stimulation. Interestingly, β -arrestin-2 knockdown cells responded to re-exposure to C3a for Ca^{2+} mobilization (Fig. 2C). Notably, for initial Ca^{2+} mobilization and desensitization, double knockdown cells responded similarly to β -arrestin-2 knockdown cells (Fig. 2D).

To investigate the role of β -arrestins on agonist-induced C3aR internalization, shRNA control, β -arrestin-1 and β -arrestin-2 knockdown cells were exposed to buffer or C3a and receptor


A: shRNA



B: β -arrestin-1 KD



C: β -arrestin-2 KD



D: β -arrestin-1 and -2 KD




Figure 2. Knockdown of β -arrestin-2, but not β -arrestin-1, attenuates C3aR desensitization. (A), shRNA control, (B) β -arrestin-1 KD (knockdown), (C) β -arrestin-2 KD and (D) double β -arrestin-1 and 2 KD cells were loaded with Indo-1(1 μM), stimulated with C3a (100 nM) for 5 min and intracellular Ca^{2+} mobilization was determined (black solid lines). The cells were immediately washed three times with ice-cold buffer, resuspended in warm buffer and exposed to a second stimulation of C3a (100 nM) and intracellular Ca^{2+} mobilization was again determined (red broken lines). Data shown are representative of three similar experiments. doi:10.1371/journal.pone.0019585.g002

internalization was determined by flow cytometry. In shRNA control cells, C3a caused a robust internalization of its receptors (Fig. 3A). In β -arrestin-1 knockdown cells, there was no marked difference in the extent of receptor internalization (Fig. 3A, B). Interestingly, internalization of C3aR was substantially reduced in β -arrestin-2 knockdown cells (Fig. 3C and D). These findings clearly demonstrate that β -arrestin-2, but not β -arrestin-1, is involved in C3aR desensitization and internalization in human mast cell line, HMC-1.

β -arrestin-1, but not β -arrestin-2, promotes C3a-induced mast cell degranulation

Our next goal was to determine the roles of β -arrestin-1 and β -arrestin-2 on C3a-induced mast cell degranulation. We could not use HMC-1 cells for these studies because this immature mast cell line has little or no capacity to degranulate. LAD2 mast cells express C3aR and responds to ligand for Ca^{2+} mobilization and degranulation [7]. We therefore knocked down the expression of β -arrestin-1 and β -arrestin-2 in LAD2 cells. As in HMC-1 cells,

lentiviral shRNA induced $\sim 80\%$ knockdown of the β -arrestin-1 and β -arrestin-2 in LAD2 mast cells (Fig. 4A). Furthermore, consistent with the findings in HMC-1 cells, β -arrestin-1 knockdown in LAD2 cells had little or no effect on C3a-induced Ca^{2+} mobilization (Fig. 4B and C) while β -arrestin-2 silencing resulted in a more sustained Ca^{2+} mobilization and loss of desensitization (Fig. 4D). Surprisingly, however, knockdown of β -arrestin-2 had no effect on C3a-induced mast cell degranulation but the absence of β -arrestin-1 resulted in a significantly decreased degranulation response (Fig. 4E).

β -arrestin-2, but not β -arrestin-1, inhibits C3a-induced NF- κ B activation and chemokine CCL4 generation

β -arrestin-1 and β -arrestin-2 bind to $\text{I}\kappa\text{B}\alpha$ to inhibit GPCR-induced NF- κ B activity in transfected cell lines [19,20]. We therefore sought to determine the roles of these adapter molecules on C3a-induced NF- κ B luciferase activity in human mast cells. We used HMC-1 cells for these studies because they are more amenable to transfection than LAD2 cells. C3a did not induce NF-




Figure 3. Knockdown of β -arrestin-2, but not β -arrestin-1, inhibits agonist-induced C3aR internalization. (A) shRNA control HMC-1 cells (B) β -arrestin-1 KD and (C) β -arrestin-2 KD cells were exposed to buffer ($-$ C3a) or C3a (100 nM) for 5 min. Cells were washed with ice-cold FACS buffer, incubated with a mouse anti-C3aR antibody or an isotype control antibody followed by PE-labeled donkey anti-mouse IgG antibody and analyzed by flow cytometry. Representative histograms showing cell surface C3aR expression in (A) shRNA control, (B) β -arrestin-1 KD and (C) β -arrestin-2 KD cells are shown. (D) shRNA control, β -arrestin-1 KD and β -arrestin-2 KD cells were exposed to C3a for different time periods and receptor internalization was determined as described above. Internalization is expressed as the percentage loss of C3aR following exposure to C3a. Data represent the mean \pm SEM from three experiments. Statistical significance was determined by two way ANOVA with Bonferroni's post test. * indicates $p < 0.05$.

doi:10.1371/journal.pone.0019585.g003




Figure 4. Knockdown of β -arrestin-1, but not β -arrestin-2, inhibits C3a-induced degranulation in LAD2 mast cells. (A) LAD2 cells were transfected with shRNA control, β -arrestin-1, and β -arrestin-2 lentivirus. After puromycin selection, quantitative PCR was employed to assess the β -arrestin-1 or β -arrestin-2 mRNA level and results are expressed as a ratio of β -arrestin to GAPDH mRNA levels. (B, C, D) A representative desensitization experiment in shRNA control, β -arrestin-1 and β -arrestin-2 knockdown LAD2 cells is shown. (E) shRNA control, β -arrestin-1 and -2 KD LAD2 mast cells were stimulated with different concentrations of C3a and percent degranulation (β -hexosaminidase release) was determined. Data represent the mean \pm SEM from three independent experiments. Statistical significance was determined by two way ANOVA with Bonferroni's post test. * indicates $p < 0.05$ and ** indicates $p < 0.001$. doi:10.1371/journal.pone.0019585.g004

κ B luciferase activity in shRNA control or β -arrestin-1 silenced HMC-1 cells (Fig. 5A). By contrast, β -arrestin-2 knockdown cells showed a significant enhancement in C3a-induced NF- κ B luciferase activity as compared to shRNA control cells. Given that NF- κ B plays an important role in the generation of proinflammatory cytokines, we tested the effects of β -arrestin-1 and β -arrestin-2 knockdown in C3a-induced chemokine CCL4 production. Consistent with NF- κ B activation, C3a induced CCL4 only in β -arrestin-2 silenced cells (Fig. 5B).

β -arrestin-1 and β -arrestin-2 inhibit C3a-induced ERK1/2 phosphorylation

Activation of GPCRs leads to ERK1/2 phosphorylation via two temporally distinct pathways; an early response that reflects G protein activation and a delayed response that is G protein independent but requires β -arrestins [31]. We therefore investigated the effects of silencing the expression of β -arrestin-1, β -arrestin-2 or both on the time course of C3a-induced ERK1/2 phosphorylation in HMC-1 cells. In shRNA control cells, C3a caused a transient ERK1/2 phosphorylation that peaked between

1–5 min and returned to basal thereafter (Fig. 6A). Surprisingly, silencing β -arrestin-1 or β -arrestin-2 expression enhanced the magnitude of this early response and rendered the cells responsive to C3a for ERK1/2 phosphorylation even at later time points (10–30 min). Furthermore, in double knockdown cells, C3a-induced ERK1/2 phosphorylation was greater in magnitude than single β -arrestin knockdown cells (Fig 6A and 6B).

To determine if the delayed C3a-induced ERK1/2 phosphorylation in β -arrestin knockdown cells is mediated via a G protein-dependent pathway, shRNA control and β -arrestin-1 and β -arrestin-2 double knockdown cells were exposed to pertussis toxin and the effects of C3a on ERK1/2 phosphorylation was determined. As shown in Fig. 7A and 7B both early and delayed responses were almost completely inhibited in pertussis toxin treated cells.

Discussion

β -arrestins are well known for their roles in GPCR desensitization and internalization. They also modulate downstream signaling pathways such as those for ERK and NF- κ B indepen-




Figure 5. Knockdown of β -arrestin-2, but not β -arrestin-1, enhanced C3a-induced NF- κ B activation and chemokine CCL4 generation. shRNA control, β -arrestin-1 KD or β -arrestin-2 KD HMC-1 cells were transiently transfected with NF- κ B luciferase reporter gene construct. (A) Cells were stimulated with C3a (100 nM for 6 hr) and NF- κ B-dependent transcriptional activity was determined by luciferase activity assay. (B) Control or β -arrestin KD cells were stimulated with C3a (100 nM for 6 hr) and CCL4 production was determined from the supernatant by ELISA. Data shown are mean \pm SEM of three experiments performed in triplicate. Statistical significance was determined by two way ANOVA with Bonferroni's post test. ** indicates $p < 0.001$. doi:10.1371/journal.pone.0019585.g005

dent of receptor desensitization. Most previous studies on GPCR regulation have been performed using mouse embryonic fibroblasts (MEFs) derived from β -arrestin null mice [11], transfected cell lines overexpressing β -arrestins or siRNA-mediated β -arrestin knockdown in HEK293 cells [32,33,34,35]. For the present study, we utilized lentivirus shRNA to stably knockdown the expression of β -arrestin-1 and β -arrestin-2 in human mast cell lines, HMC-1 cells and LAD2 cells that endogenously express C3aR. Using this approach, we have uncovered distinct roles of β -arrestin-1 and β -arrestin-2 on C3aR desensitization, internalization, degranulation, NF- κ B activation and chemokine generation. Furthermore, we provided the first demonstration that β -arrestin-1 and β -arrestin-2 act as novel inhibitors of C3a-induced G protein-dependent ERK1/2 phosphorylation in human mast cells.


Previous studies indicated that β -arrestin-2 either inhibits or promotes chemoattractant/chemokine induced degranulation. Thus, in response to agonist stimulation wild-type chemoattractant receptors associate with β -arrestin-2 in transfected RBL-2H3 cells but phosphorylation-deficient mutants do not [9,14,16,30]. Furthermore, agonist-induced Ca^{2+} mobilization and degranulation are enhanced in cells expressing phosphorylation-deficient receptors when compared to wild-type receptors. These findings are consistent with the notion that receptor phosphorylation and β -arrestin-2 participate in receptor desensitization. By contrast, Barlic et al., [17] showed that agonist induced phosphorylation of the chemokine receptor CXCR1 leads to β -arrestin-2-mediated receptor internalization and the formation of β -arrestin-2-Hck complex, which migrates to secretory granules initiating the process of degranulation. The finding in the present study that enhanced Ca^{2+} response in the absence of β -arrestin-2 did not promote greater degranulation provides a possible explanation for the previously published conflicting data for the role of β -arrestin-2 on degranulation. It suggests that β -arrestin-2 plays a dual role on GPCR-induced degranulation; inhibition via desensitization and activation via its association with Hck. Thus, the inability of

enhanced Ca^{2+} response to promote greater degranulation in β -arrestin-2 knockdown cells probably reflects the loss of β -arrestin-2-mediated Hck signaling (see Model in Fig. 8A).

An interesting finding of the present study was that while knockdown of β -arrestin-1 had no effect on C3aR desensitization (as measured by Ca^{2+} mobilization) or receptor internalization its absence resulted in a substantial inhibition of C3a-induced mast cell degranulation. Because β -arrestin-1 does not participate in C3aR internalization, C3a is unlikely to promote Hck- β -arrestin-1 interaction. Our studies with confocal microscopy in live cells indicated that C3a causes translocation of β -arrestin-1 to the plasma membrane (data not shown). Furthermore, β -arrestin-1 forms a complex with Ral-GDS in the cytoplasm of human neutrophils [18]. This raises the interesting possibility that upon C3aR activation, β -arrestin-1/Ral GDS complex translocates to the plasma membrane to promote degranulation and that knockdown of β -arrestin-1 leads to attenuated response due to the absence of this complex (see Model in Fig. 8A). Whether this or other mechanism(s) participate on the effect of β -arrestin-1 on C3a-induced mast cell degranulation remains to be determined.

β -arrestins have been shown to promote or inhibit NF- κ B activity depending on the cell type and receptors utilized [14,19,20,23,24]. Our results clearly demonstrate that β -arrestin-2, but not β -arrestin-1, inhibits C3a-induced NF- κ B activation and chemokine generation. Gao et al., [20] recently observed similar differences between β -arrestin-1 and β -arrestin-2 in cytokine production in HeLa cells and THP-1 monocytes. This difference was thought to reflect a reduced ability of β -arrestin-1 to form a complex with the inhibitory $\text{I}\kappa\text{B}\alpha$ when compared to β -arrestin-2. In the present study, we showed that while β -arrestin-1 does not participate in agonist-induced C3aR internalization, β -arrestin-2 is essential for this response. It is therefore possible that internalized C3aR- β -arrestin-2 complex interacts with $\text{I}\kappa\text{B}\alpha$ to keep NF- κ B inactive and that depletion of β -arrestin-2 removes this inhibitory constraint to allow NF- κ B activation and chemokine generation (Fig. 8B). It is also

A



B




Figure 6. C3a-induced ERK1/2 phosphorylation is enhanced in β -arrestin-1, β -arrestin-2 and double KD cells. shRNA control or β -arrestin KD HMC-1 cells (1×10^6 /ml) were exposed to C3a (100 nM) for 1, 5 and 10, 15 and 30 min. Cell lysates were separated on SDS-PAGE and blots were probed with anti-phospho-ERK1/2 antibody followed by anti-rabbit IgG-HRP. The blots were then stripped and reprobbed with anti-ERK1/2 antibody followed by anti-rabbit IgG-HRP. Immunoreactive band were visualized by SuperSignal West Femto maximum sensitivity substrate. (A) Representative immunoblots from three similar experiments are shown. (B) ERK1/2 phosphorylation was quantified using Image J as shown in the line graph. Data represent the mean \pm SEM from three independent experiments. Statistical significance was determined by two way ANOVA with Bonferroni's post test. ** indicates $p < 0.001$. doi:10.1371/journal.pone.0019585.g006

possible that enhanced signaling as manifested by a more sustained Ca^{2+} mobilization in β -arrestin-2 knockdown cells results in greater NF- κ B activation and chemokine generation.

An interesting finding of the present study was that silencing β -arrestin-1 and β -arrestin-2 enhanced early ERK1/2 phosphorylation in response to C3a and also rendered the cells responsive to C3a at later time points (15–30 min). This finding is in direct contrast to situations with many other GPCRs, where β -arrestins are required for delayed ERK phosphorylation [31]. One possible interpretation of our finding is that knockdown β -arrestins attenuate C3aR desensitization leading to more sustained ERK1/2 phosphorylation. This explanation is, however, unlikely as β -arrestin-1 knockdown, which had no impact on C3aR desensitization, rendered the cells responsive to C3a for ERK1/2 phosphorylation. Ahn et al. [33] recently demonstrated that β -arrestin-2 inhibits angiotensin II type 1A-mediated ERK1/2 activation in HEK293 cells and silencing β -arrestin-1 expression enhances this response by blocking β -arrestin-2 mediated inhibition. This type of reciprocal regulation is unlikely for C3a-induced response in mast cells as knockdown of both β -arrestins either individually or together enhanced C3a-induced ERK1/2 phosphorylation.

It is noteworthy that *Drosophila* genome encodes a single β -arrestin, Kurtz (Krz), which controls olfaction, behavior, sensitivity to osmotic stress, and is essential for survival of the fly [36,37,38]. Tipping et al., [39] recently showed that Krz directly binds to and sequesters an inactive form of ERK, thus preventing its activation by the upstream kinase, MEK. We have shown that pertussis toxin completely blocks the transient C3a-induced ERK1/2 phosphorylation in shRNA control cells as well as the sustained response in β -arrestin-1 and β -arrestin-2 double knockdown cells. This finding suggests that C3a causes ERK1/2 phosphorylation via a GPCR-mediated pathway and that arrestins inhibit this response by forming a direct complex with ERK and preventing its activation by MEK (Fig. 8B).

In summary, we demonstrated distinct roles for β -arrestins-1 and β -arrestins-2 on C3aR desensitization, internalization, degranulation, NF- κ B activation and chemokine generation in human mast cells. Most importantly, we provided the first demonstration that both β -arrestin-1 and β -arrestin-2 act as novel inhibitors of C3a-induced G-protein-mediated ERK1/2 phosphorylation in human mast cells. In addition to C3aR, human mast cells express a large number of other GPCRs, Fc ϵ RI, toll-like




Figure 7. Enhanced C3a-induced ERK1/2 phosphorylation in β -arrestin KD cells mast is mediated via a G protein-dependent pathway. shRNA control and double β -arrestin KD cells were pretreated with vehicle or Pertussis toxin (PTx; 100 ng/ml, 16 hr). Cell were then washed twice in serum free medium and stimulated with C3a (100 nM) for 1, 5 and 10, 15 and 30 min and ERK1/2 phosphorylation were determined. (A) Representative immunoblots from three similar experiments are shown. (B). ERK1/2 phosphorylation was quantified using Image J as shown in the bar graph. Data represent the mean \pm SEM from three independent experiments. Statistical significance was determined by two way ANOVA with Bonferroni's post test. ** indicates $p < 0.001$. doi:10.1371/journal.pone.0019585.g007

receptors and IL-33 receptor T1/ST2 [40,41,42,43,44]. Given that β -arrestins regulate GPCR and non-GPCR signaling [45,46], it is likely that they regulate other receptor/signaling pathways in human mast cells. Our future studies will focus on the receptor specificity of human mast cell regulation by β -arrestins.

Materials and Methods

Materials

Mission shRNA bacterial glycerol stocks for β -arrestins were purchased from Sigma Life Sciences (St. Louis, MO). Indo-1 AM was from Molecular Probes (Eugene, OR). All tissue culture reagents were purchased from Invitrogen (Gaithersburg, MD). Anti-human C3aR was obtained from Santa Cruz Biotechnology (Santa Cruz, CA), PE-labeled donkey anti-mouse IgG was purchased from eBioscience (San Diego, CA). All recombinant human cytokines were purchased from Peprotech (Rocky Hill, NJ). Rabbit anti-ERK1 and anti-phospho-ERK1/2 antibodies were purchased from Cell Signaling (Beverly, MA). SuperSignal[®] West Femto Maximum

Sensitivity Substrate and HRP labeled Goat anti-rabbit IgG were from Thermo Scientific (Rockford, IL). Purified C3a was obtained from Advanced Research Technologies (San Diego, CA). CCL4 ELISA kit was purchased from R&D Systems (Minneapolis, MN).

Mast cell culture

HMC-1 cells were cultured in Iscove's modified Dulbecco's medium (IMDM) supplemented with 10% FCS, glutamine (2 mM), penicillin (100 IU/mL) and streptomycin (100 μ g/mL) [47]. LAD2 cells were maintained in complete StemPro-34 medium supplemented with 100 ng/mL rhSCF [48].

Lentivirus and stable transduction of shRNAs in mast cells

The following β -arrestin-1 and -2 targeted shRNAs in Lentiviral construct plasmid were purchased from Sigma-Aldrich (St. Louis, MO): β -arrestin-1 (NM_004041) Clone 1 TRCN000023-0148, Clone 2 TRCN0000230147, Clone 3 TRCN0000230149,



(A): Desensitization, Internalization and Degranulation**(B): ERK/NF- κ B**

Figure 8. Model for the Regulation of C3aR signaling in human mast cells by β -arrestin-1 and β -arrestin-2. (A): β -arrestin-2 causes desensitization and internalization of C3aR but both β -arrestins promote G protein-independent signaling for degranulation via the activation of Hck and/or Ral-GDS-mediated signaling pathways [17,18]. (B) β -arrestin-2 mediates inhibition of C3a-induced NF- κ B activity but both β -arrestins block C3a-induced G-protein-mediated ERK1/2 phosphorylation. Red dotted lines denote inhibition. doi:10.1371/journal.pone.0019585.g008

Clone 4 TRCN0000230150, Clone 5 TRCN0000219075; β -arrestin-2 (NM_004313) Clone 1 TRCN0000165387, Clone 2 TRCN0000164794, Clone 3 TRCN0000159332, Clone 4 TRCN0000161834, Clone 5 TRCN0000159482 and control non-target vector SHC002. Cell transduction was conducted by mixing 1.5 ml of virus with 3.5 ml of HMC-1 or LAD-2 cells (5×10^6). For the double knockdown of β -arrestin-1 and -2, 1.5 ml of each virus of specific clones were transduced in 2 ml of HMC-1 or LAD-2 cells (5×10^6). Eight hr post-infection, medium was changed to virus-free complete medium, and antibiotic (puromycin; 2 μ g/ml Sigma-Aldrich) selection was initiated 16 h later. Cells were analyzed for β -arrestin knockdown one week after initiation of puromycin selection.

Real-Time PCR

Total RNA was extracted from 4×10^6 of cells using TRIZOL, treated with DNase I and subsequently purified for genomic DNA contamination with RNeasy mini Kit (Qiagen) according to the manufacture's instruction. cDNA was synthesized from genomic DNA-free RNA using the cDNA synthesis kit from GE Healthcare. Gene expression was analyzed using real time PCR with Taqman[®] Fast Universal PCR Master Mix on a Taqman 7500 Fast Real-Time PCR System (Applied Biosystems, Foster City, CA). Taqman hGAPDH, β -arrestin-1 and β -arrestin-2 primers were used for real time PCR to analyze the knockdown efficiency. The amplification conditions were as follows: initial denaturation at 95°C for 20 sec, followed by 40 cycles of amplification: 95°C for 3 sec, 60°C for 30 sec. Analysis was performed according to $\Delta\Delta$ -Ct method. The results were expressed as β -arrestin-1 or -2/*GAPDH* ratio.

C3a Receptor desensitization

Receptor desensitization assay based on Ca²⁺ mobilization was determined as described previously [49]. Briefly, 1×10^6 HMC-1

or 0.25×10^6 LAD-2 cells were washed twice with buffer (119 mM NaCl, 5 mM KCl, 25 mM HEPES, 5.6 mM Glucose, 0.4 mM MgCl₂, 1 mM CaCl₂) containing 1 mg/ml BSA and incubated with 1 μ M of Indo-1 for 30 min in dark. Cells were then washed and resuspended in 1.5 ml of the same buffer and time course of Ca²⁺ mobilization (0–5 min) was determined using Hitachi F-2500 Fluoro spectrophotometer (San Jose, CA) with an excitation wavelength of 355 nm and an emission wavelength of 410 nm [9]. For desensitization assay, cells were removed from the cuvette, washed twice and Ca²⁺ mobilization to a subsequent exposure of C3a (100 nM) was determined.

Degranulation Assay

LAD-2 cells (1.2×10^4) were seeded into 96-well plates in a total volume of 50 μ l of buffer containing 1 mg/ml BSA and exposed to different concentrations of C3a (1, 10 and 100 nM). For total β -hexosaminidase release, control cells were lysed in 50 μ l of 0.1% Triton X-100. Aliquots (20 μ l) of supernatants or cell lysates were incubated with 20 μ l of 1 mM p-nitrophenyl-N-acetyl- β -D-glucosamine for 1.5 hour at 37°C. The reaction was stopped by adding 250 μ l of a 0.1 M Na₂CO₃/0.1 M NaHCO₃ buffer and absorbance measured at 405 nm [49].

Receptor Internalization

ShRNA control and β -arrestin knockdown HMC-1 cells (2.5×10^5) were stimulated with or without C3a (100 nM) at 37°C. Cells were washed twice and resuspended in 50 μ l of ice-cold FACS buffer (PBS containing 2% FBS). C3aR antibody or isotype control (2 μ l) was added and the cells were incubated on ice for 1 h. Cells were washed twice and re-suspended in 48.5 μ l of ice-cold FACS buffer. Phycoerythrin (PE)-labeled donkey anti-mouse (1.5 μ l) was added and incubated on ice for 1 h. Cells were washed twice with cold FACS buffer and fixed in 250 μ l of 2% formaldehyde. Receptor internalization was quantified as the loss

of cell-surface receptors, as analyzed on a BD LSR II flow cytometer (BD Biosciences).

ERK1/2 Phosphorylation

ShRNA control and β -arrestin knockdown HMC-1 cells were serum starved overnight. The following day, cells were washed twice and resuspended in serum free IMDM medium at a concentration of 1×10^6 /ml and stimulated C3a (100 nM) for different time points. Three-fold volume of ice-cold PBS containing 1 mM sodium orthovanadate was added to stop the reaction. Total cell lysate was prepared with RIPA buffer (150 mM NaCl, 1.0% NP-40, 0.5% Sodium-deoxycholate, 0.10% SDS, 50 mM Tris [pH 8.0], 5 mM EDTA, 10 mM NaF, 10 mM Na-pyrophosphate and protease inhibitor cocktail) and subsequently analyzed by Western blot using rabbit monoclonal antibodies for phospho-p44/42 MAPK (pERK1/2) and p44/42 MAPK (ERK1/2).

NF- κ B luciferase reporter activity

ShRNA control and β -arrestin knockdown HMC-1 cells (3×10^6) were seeded in 12-well plates. The following day, cells were co-transfected with NF- κ B luciferase reporter gene construct (pNF- κ B-LUC and p-Renilla Stratagene, Santaclara, CA) (in a 10:1 ratio) using Lipofectamine 2000 reagent (Invitrogen, Carlsbad, CA) in serum-free IMDM medium as per manufacturer protocol. Six hour post-transfection, medium was replaced with IMDM containing 10% FBS. After 18 hr of incubation in complete medium, cells were re-plated and stimulated in the presence or absence of 100 nM C3a for 6 hr. Cells were then harvested, washed in ice-cold PBS and finally lysed in Promega passive lysis buffer (Dual Luciferase assay kit; Promega, Madison, WI). NF- κ B luciferase activity was measured using Turner

biosystem 20/20 Luminometer (Promega, Madison, WI). Results expressed have been normalized to Renilla.

CCL4 chemokine release assay

Chemokine release assay was performed as previously described [27]. HMC-1 shRNA control, β -arrestin-1 and β -arrestin-2 knockdown cells (0.2×10^6 cells) were stimulated with 100 nM C3a for 6 hours. CCL4 chemokine levels were quantified by sandwich ELISA according to the manufacturer's protocol.

Data analysis

The results are expressed as \pm S.E.M for the values obtained from experiment. GraphPad Prism software (Graph Pad, Version 5.0 San Diego, CA) was used to analyze data for statistical significance. The statistical significance was determined by one-way analysis of variance (ANOVA) with Dunnett's multiple comparison post hoc test, and two way ANOVA with Bonferroni's post test.

Acknowledgments

We are grateful to Dr. Joseph Butterfield (Mayo Clinic, Rochester, MN) for supplying us with HMC-1 cells. We also thank Drs. Arnold Kirshenbaum and Dean Metcalfe (NIAID/NIH) for providing LAD2 mast cells and the FACS core facilities of the School of Dental Medicine, University of Pennsylvania.

Author Contributions

Conceived and designed the experiments: HA. Performed the experiments: AV KG HS QG. Analyzed the data: AV KG HS QG. Contributed reagents/materials/analysis tools: AV KG HS QG HA. Wrote the paper: HA KG HS.

References

- Fukuoka Y, Xia HZ, Sanchez-Munoz LB, Dellinger AL, Escribano L, et al. (2008) Generation of anaphylatoxins by human β -tryptase from C3, C4, and C5. *J Immunol* 180: 6307–6316.
- Drouin SM, Corry DB, Hollman TJ, Kildsgaard J, Wetsel RA (2002) Absence of the complement anaphylatoxin C3a receptor suppresses Th2 effector functions in a murine model of pulmonary allergy. *J Immunol* 169: 5926–5933.
- McNeil HP, Adachi R, Stevens RL (2007) Mast Cell-restricted Trypsases: Structure and Function in Inflammation and Pathogen Defense. *J Biol Chem* 282: 20785–20789.
- Humbles AA, Lu B, Nilsson CA, Lilly C, Israel E, et al. (2000) A role for the C3a anaphylatoxin receptor in the effector phase of asthma. *Nature* 406: 998–1001.
- Hartmann K, Henz BM, Kruger-Krasagakes S, Kohl J, Burger R, et al. (1997) C3a and C5a stimulate chemotaxis of human mast cells. *Blood* 89: 2863–2870.
- Lappalainen J, Lindstedt KA, Kovanen PT (2007) A protocol for generating high numbers of mature and functional human mast cells from peripheral blood. *Clin Exp Allergy* 37: 1404–1414.
- Venkatesha RT, Berla Thangam E, Zaidi AK, Ali H (2005) Distinct regulation of C3a-induced MCP-1/CCL2 and RANTES/CCL5 production in human mast cells by extracellular signal regulated kinase and PI3 kinase. *Mol Immunol* 42: 581–587.
- Woolhiser MR, Brockow K, Metcalfe DD (2004) Activation of human mast cells by aggregated IgG through Fc γ RI: additive effects of C3a. *Clin Immunol* 110: 172–180.
- Ahamed J, Haribabu B, Ali H (2001) Cutting edge: Differential regulation of chemoattractant receptor-induced degranulation and chemokine production by receptor phosphorylation. *J Immunol* 167: 3559–3563.
- DeWire SM, Ahn S, Lefkowitz RJ, Shenoy SK (2007) β -arrestins and cell signaling. *Annu Rev Physiol* 69: 483–510.
- Paing MM, Stutts AB, Kohout TA, Lefkowitz RJ, Trejo J (2002) β -Arrestins regulate protease-activated receptor-1 desensitization but not internalization or Down-regulation. *J Biol Chem* 277: 1292–1300.
- Kohout TA, Lin FS, Perry SJ, Conner DA, Lefkowitz RJ (2001) β -Arrestin 1 and 2 differentially regulate heptahelical receptor signaling and trafficking. *Proc Natl Acad Sci U S A* 98: 1601–1606.
- Ahamed J, Ali H (2002) Distinct roles of receptor phosphorylation, G protein usage, and mitogen-activated protein kinase activation on platelet activating factor-induced leukotriene C(4) generation and chemokine production. *J Biol Chem* 277: 22685–22691.
- Venkatesha RT, Ahamed J, Nuesch C, Zaidi AK, Ali H (2004) Platelet-activating Factor-induced Chemokine Gene Expression Requires NF- κ B Activation and Ca²⁺/Calcineurin Signaling Pathways: inhibition by receptor phosphorylation and β -arrestin recruitment. *J Biol Chem* 279: 44606–44612.
- Richardson RM, DuBose RA, Ali H, Tomhave ED, Haribabu B, et al. (1995) Regulation of human interleukin-8 receptor A: identification of a phosphorylation site involved in modulating receptor functions. *Biochemistry* 34: 14193–14201.
- Vines CM, Xue M, Maestas DC, Cimino DF, Prossnitz ER (2002) Regulation of N-formyl peptide-mediated degranulation by receptor phosphorylation. *J Immunol* 169: 6760–6766.
- Barlic J, Andrews JD, Kelvin AA, Bosinger SE, DeVries ME, et al. (2000) Regulation of tyrosine kinase activation and granule release through β -arrestin by CXCR1. *Nat Immunol* 1: 227–233.
- Bhattacharya M, Anborgh PH, Babwah AV, Dale LB, Dobransky T, et al. (2002) β -arrestins regulate a Ral-GDS Ral effector pathway that mediates cytoskeletal reorganization. *Nat Cell Biol* 4: 547–555.
- Wetherow DS, Garrison TR, Miller WE, Lefkowitz RJ (2004) β -Arrestin inhibits NF- κ B activity by means of its interaction with the NF- κ B inhibitor I κ B α . *Proc Natl Acad Sci U S A*.
- Gao H, Sun Y, Wu Y, Luan B, Wang Y, et al. (2004) Identification of β -arrestin2 as a G protein-coupled receptor-stimulated regulator of NF- κ B pathways. *Mol Cell* 14: 303–317.
- Su Y, Raghuvanshi SK, Yu Y, Nanney LB, Richardson RM, et al. (2005) Altered CXCR2 signaling in β -arrestin-2-deficient mouse models. *J Immunol* 175: 5396–5402.
- Raghuvanshi SK, Nasser MW, Chen X, Strieter RM, Richardson RM (2008) Depletion of β -arrestin-2 promotes tumor growth and angiogenesis in a murine model of lung cancer. *J Immunol* 180: 5699–5706.
- Yang M, He RL, Benovic JL, Ye RD (2009) β -Arrestin1 interacts with the G-protein subunits β 1 γ 2 and promotes β 1 γ 2-dependent Akt signalling for NF- κ B activation. *Biochem J* 417: 287–296.
- Sun J, Lin X (2008) β -arrestin 2 is required for lysophosphatidic acid-induced NF- κ B activation. *Proc Natl Acad Sci U S A* 105: 17085–17090.
- DeFea KA, Vaughn ZD, O'Bryan EM, Nishijima D, Dery O, et al. (2000) The proliferative and antiapoptotic effects of substance P are facilitated by formation of a β -arrestin-dependent scaffolding complex. *Proc Natl Acad Sci U S A* 97: 11086–11091.

26. Luttrell LM, Ferguson SS, Daaka Y, Miller WE, Maudsley S, et al. (1999) β -arrestin-dependent formation of β 2 adrenergic receptor-Src protein kinase complexes [see comments]. *Science* 283: 655–661.
27. Ali H, Ahamed J, Hernandez-Munain C, Baron JL, Krangel MS, et al. (2000) Chemokine production by G protein-coupled receptor activation in a human mast cell line: roles of extracellular signal-regulated kinase and NFAT. *J Immunol* 165: 7215–7223.
28. Tomhave ED, Richardson RM, Didsbury JR, Menard L, Snyderman R, et al. (1994) Cross-desensitization of receptors for peptide chemoattractants. Characterization of a new form of leukocyte regulation. *J Immunol* 153: 3267–3275.
29. Haribabu B, Richardson RM, Fisher I, Sozzani S, Peiper SC, et al. (1997) Regulation of human chemokine receptors CXCR4. Role of phosphorylation in desensitization and internalization. *J Biol Chem* 272: 28726–28731.
30. Richardson RM, Haribabu B, Ali H, Snyderman R (1996) Cross-desensitization among receptors for platelet activating factor and peptide chemoattractants. Evidence for independent regulatory pathways. *J Biol Chem* 271: 28717–28724.
31. Defea K (2008) β -arrestins and heterotrimeric G-proteins: collaborators and competitors in signal transduction. *Br J Pharmacol* 153 Suppl 1: S298–309.
32. Wei H, Ahn S, Shenoy SK, Karmik SS, Hunyady L, et al. (2003) Independent β -arrestin 2 and G protein-mediated pathways for angiotensin II activation of extracellular signal-regulated kinases 1 and 2. *Proc Natl Acad Sci U S A*.
33. Ahn S, Wei H, Garrison TR, Lefkowitz RJ (2004) Reciprocal regulation of angiotensin receptor-activated extracellular signal-regulated kinases by β -arrestins 1 and 2. *J Biol Chem* 279: 7807–7811.
34. Gesty-Palmer D, Chen M, Reiter E, Ahn S, Nelson CD, et al. (2006) Distinct β -arrestin- and G protein-dependent pathways for parathyroid hormone receptor-stimulated ERK1/2 activation. *J Biol Chem* 281: 10856–10864.
35. Shenoy SK, Drake MT, Nelson CD, Houtz DA, Xiao K, et al. (2006) β -arrestin-dependent, G protein-independent ERK1/2 activation by the β 2 adrenergic receptor. *J Biol Chem* 281: 1261–1273.
36. Ge H, Krishnan P, Liu L, Krishnan B, Davis RL, et al. (2006) A Drosophila nonvisual arrestin is required for the maintenance of olfactory sensitivity. *Chem Senses* 31: 49–62.
37. Johnson EC, Tift FW, McCauley A, Liu L, Roman G (2008) Functional characterization of kurtz, a Drosophila non-visual arrestin, reveals conservation of GPCR desensitization mechanisms. *Insect Biochem Mol Biol* 38: 1016–1022.
38. Liu L, Davis RL, Roman G (2007) Exploratory activity in Drosophila requires the kurtz nonvisual arrestin. *Genetics* 175: 1197–1212.
39. Tipping M, Kim Y, Kyriakakis P, Tong M, Shvartsman SY, et al. β -arrestin Kurtz inhibits MAPK and Toll signalling in Drosophila development. *EMBO J*.
40. Iikura M, Suto H, Kajiwara N, Oboki K, Ohno T, et al. (2007) IL-33 can promote survival, adhesion and cytokine production in human mast cells. *Lab Invest* 87: 971–978.
41. Kubo Y, Fukuishi N, Yoshioka M, Kawasoe Y, Iriguchi S, et al. (2007) Bacterial components regulate the expression of Toll-like receptor 4 on human mast cells. *Inflamm Res* 56: 70–75.
42. Oskeritzian CA, Zhao W, Min HK, Xia HZ, Pozez A, et al. (2005) Surface CD88 functionally distinguishes the MC_{1TC} from the MC_T type of human lung mast cell. *J Allergy Clin Immunol* 115: 1162–1168.
43. Oskeritzian CA, Price MM, Hait NC, Kapitonov D, Falanga YT, et al. Essential roles of sphingosine-1-phosphate receptor 2 in human mast cell activation, anaphylaxis, and pulmonary edema. *J Exp Med* 207: 465–474.
44. Kuehn HS, Gilfillan AM (2007) G protein-coupled receptors and the modification of Fc ϵ RI-mediated mast cell activation. *Immunol Lett* 113: 59–69.
45. Lefkowitz RJ, Whalen EJ (2004) β -arrestins: traffic cops of cell signaling. *Curr Opin Cell Biol* 16: 162–168.
46. Kovacs JJ, Hara MR, Davenport CL, Kim J, Lefkowitz RJ (2009) Arrestin development: emerging roles for β -arrestins in developmental signaling pathways. *Dev Cell* 17: 443–458.
47. Butterfield JH, Weiler DA (1989) In vitro sensitivity of immature human mast cells to chemotherapeutic agents. *Int Arch Allergy Appl Immunol* 89: 297–300.
48. Kirshenbaum AS, Akin C, Wu Y, Rottem M, Goff JP, et al. (2003) Characterization of novel stem cell factor responsive human mast cell lines LAD 1 and 2 established from a patient with mast cell sarcoma/leukemia; activation following aggregation of Fc ϵ RI or Fc γ RI. *Leuk Res* 27: 677–682.
49. Ali H, Richardson RM, Tomhave ED, DuBose RA, Haribabu B, et al. (1994) Regulation of stably transfected platelet activating factor receptor in RBL-2H3 cells. Role of multiple G proteins and receptor phosphorylation. *J Biol Chem* 269: 24557–24563.