

Survival rates of a reintroduced population of the Crested Ibis *Nipponia nippon* in Ningshan County (Shaanxi, China)

YANFANG LI, XINPING YE, MIN WANG, XIA LI, RONG DONG,
ZHIPING HUO and XIAOPING YU

Summary

A reintroduction project for the endangered Crested Ibis *Nipponia nippon* was conducted in Ningshan County (Shaanxi Province) on May 31, 2007. Post-release monitoring of the survival of the reintroduced population was carried out extensively from 2008 to 2015. Data collected over eight years after release were used to estimate the annual survival rate for different cohorts using a Cormack-Jolly-Seber model with capture-recapture data. The mean annual survival rates for all individuals were estimated to be 0.738 (95% CI: 0.547–0.801) and 0.752 (95% CI: 0.478–0.887) for released birds. For different age classes in the recipient population, the survival rates were estimated to be 0.384 (95% CI: 0.277–0.504), 0.853 (95% CI: 0.406–0.978), and 0.812 (95% CI: 0.389–0.950) for yearlings, juveniles and adults, respectively. The higher mortality for yearlings has greatly decreased the survival rate and our focal population was indeed sensitive to changes in yearling survival. Therefore, effective protection of yearlings was crucial to population persistence, as well as to juveniles and adults. The large proportion of mature individuals in our focal population indicated a gradually growing population. There was a slight bias towards males in the adult sex ratio with the increase of wild-born offspring, but it was not statistically significant. Therefore, we conclude that the primary goal of establishing a self-sustaining population of the Crested Ibis in part of their historical range has been achieved. Finally, we discuss factors affecting the survival of the reintroduced population and we propose some changes for future management of endangered species.

Introduction

Reintroduction of a species aims to re-establish a viable population within the indigenous range of the focal species (IUCN 1998, 2013). The measure of success of a reintroduction programme is the establishment of a self-sustaining population (Scott and Carpenter 1987), which ultimately relies on the survival ability of the released animals at the release site. A population is considered self-sustaining if it shows fidelity to the release area, long-term survival and breeding success (Armstrong *et al.* 1999). The success of reintroduction programmes depends on the quality of the habitat in which the reintroduction takes place, the adaptive ability of the birds to new conditions, the total number and the age structure of the released birds (Griffith *et al.* 1989, Wolf *et al.* 1996, Armstrong *et al.* 2002, Tweed *et al.* 2006). To establish a new population, it is essential for the released birds to survive during the establishment phase, to settle in the new reintroduced area, to pair and nest successfully in the release habitat and to produce a sufficient number of young individuals to balance mortality (Yu *et al.* 2015). In addition, the age structure of a reintroduced population should be considered to see whether the population will increase continuously because a population with more adults has a potential to grow faster (Komers and Curman 2000). Among all

reintroduction programmes for Ciconiiformes species, the reintroduction project for the White Stork *Ciconia ciconia* in Switzerland was relatively successful (Schaub *et al.* 2004), and the results provided useful insight for the protection of the Crested Ibis *Nipponia nippon*.

The Crested Ibis is now extirpated in almost all of its former range except in China (BirdLife International 2001). The reasons for the rapid disappearance of the species in most of its historical range remains unclear, but deforestation, illegal hunting, and the excessive use of pesticides were generally believed to be largely contributable to the successive extirpation during the 20th century (Shi *et al.* 1991). The rediscovery of the last seven individuals (two pairs and three nestlings) in a remote mountainous village in Yangxian gave new hope for the survival of the species (Liu 1981).

Over the past three decades, a great deal of conservation effort, both *in situ* and *ex situ*, has been undertaken by researchers and the local government. The wild ibis population has increased to approximately 1,100 (Wang *et al.* 2014) and more than 1,000 captive-bred individuals were raised in 12 conservation breeding centres or zoos in China, Japan and South Korea (Yu *et al.* 2015). The great success in multiple conservation projects (BirdLife International 2001) enabled the reintroduction of the Crested Ibis within its former ranges (Yu *et al.* 2009, Nagata and Yamagishi 2013). By October 2011, 56 captive-bred individuals (29 males and 27 females) were reintroduced into the wild in four subsequent releases (Huo *et al.* 2014, Yu *et al.* 2015). All released individuals and wild-born fledglings were monitored over an eight-year period (Wang *et al.* 2016).

In this article, we aim (i) to quantify age structure and adult sex ratio (ASR) of the reintroduced population, (ii) to employ a modern capture-recapture analysis method to estimate the survival rates of different cohorts in the reintroduced population, and (iii) to propose suggestions for subsequent releases of the Crested Ibis and other endangered species.

Methods

Study area

Our study area is located in Ningshan County (33°07′–33°50′ N, 108°02′–108°56′ E), (Shaanxi Province) on the southern slope of the Qinling Mountains (Figure 1). The area is used by the ibis for breeding and foraging and the landscape mosaic is characterised by intermontane basins, often developed for rice fields, interspersed with hills. Numerous shallow streams flow into the Changan River, where most individuals of the population bred and foraged. The specific geographical location and natural conditions have been reported in details by Yu *et al.* (2015).

Study birds and field observations

Up to the present study, a total of 56 captive-reared individuals have been reintroduced into the study area. From May to October in 2007 (13 females, 13 males); 2008 (four females, two males); 2009 (six females, eight males), and 2011 (three females, seven males), were released, respectively. Two breeding pairs formed and fledged three nestlings successfully immediately after release in 2008 (Yu *et al.* 2009). New breeding pairs were formed continuously, and the number of fledglings increased during the following years (Yu *et al.* 2015). By June 2015, 26 breeding pairs have produced 88 fledglings over eight breeding seasons (Yu *et al.* unpubl. data). In addition, three females immigrating into the released area from the wild population in Yangxian County mated with the released birds or their offspring.

For post-release monitoring, all released birds were marked individually by numeric metal rings and coloured alphanumeric plastic bands (provided by the National Bird Banding Centre of China) on their legs to allow resighting after release. Since 2008, all wild-born fledglings were marked at 23–25 days of age when the wings were still inadequate for flying freely but their tarsi were developed enough for ringing. Eleven captive-bred males and five of the strongest nestlings (typically the eldest) of 23–25 days old were fitted each with a radio transmitter weighing 15g, approximately 1% of the body weight (model RI-2D; frequency 216.368–216.691 MHz; battery life ~18 months; Holohil Systems, Carp, Ontario, Canada).




Figure 1. Map of released region showing the location of the release site (pentagram), breeding sites (solid circle), foraging sites (grass) and roost sites (tree) for the reintroduced population. An adult male of the reintroduced population travelled south-east (to Hanyin County) and paired with another female from the wild population of Yangxian County.

The annual cycle of the resident ibis can be divided into three stages: breeding period (March to June), wandering period (July to October) and wintering period (November to February) (Shi and Cao 2001). The annual resighting and recovery data was obtained using banding and radio tracking from 2007 to 2014. At 1-week intervals in the breeding period, all breeding pairs in the study area were checked for their unique leg bands by spotting scopes and binoculars. We monitored females throughout the breeding season and measured reproduction in terms of the number of fledglings produced per female per year (Armstrong *et al.* 2002). All ibises, as well as the radio-marked individuals generally formed wandering flocks in the non-breeding period (autumn and winter) for approximately five months. We observed the different groups at dusk (17h40–18h40) when all members of a wandering flock perched in a night-roosting site and the individuals observed in a fledgling group between 07h00 and 12h00 hours on a rotational basis at 2-week intervals.

Age structure and adult sex ratio of the reintroduced Crested Ibis population

Age structure, defined as the number of individuals in different age groups, is valuable in reflecting the quantity configuration of different age groups, in order to understand the population's history, and to forecast population dynamics (Sun *et al.* 2005). The dynamics of the wild Crested Ibis population was illustrated by an age pyramid (Lu *et al.* 1997). In the present study, age pyramids of the wild-born ibises were used to determine the overall age distribution, indicating the fecundity and probability of survival of our focal population.

The proportion of the sexes during the period of production was calculated as the female/male ratio (Komers and Curman 2000) and defined as adult sex ratio (ASR) (Donald 2007). We estimated

ASR of the population using a method of direct field observation (Donald 2007) to roughly estimate the population sex ratio. Although all birds were individually marked, the estimation of ASR was difficult due to the lack of external sexual differences (Baumgardt *et al.* 2013) in the Crested Ibis. Therefore, the sexes of all wild-born individuals were determined only by observation of copulation behaviour when they became sexually mature at least ≥ 2 years old (Yu *et al.* 2015). The statistical significance of departure from an expected sex ratio of 0.5 was assessed using Pearson's χ^2 goodness-of-fit statistic. Results were considered statistically significant at $P = 0.05$ and values were reported as mean \pm SE.

Estimation of the survival probabilities of the reintroduced population

We used capture-recapture data collected from 2008 to 2015 to estimate the correlative annual survival probabilities of the reintroduced population using Cormack-Jolly-Seber (CJS) model in the program MARK 7.1, which was especially developed for analysing capture-recapture data (Lebreton *et al.* 1992). The basic CJS model estimate survival and recapture (resighting) probability for each interval and assumes no differences among animals. The CJS can be modified by dividing animals into groups, introducing age structure, and creating relationships among intervals (Armstrong and Ewen 2002). We conducted survival analyses for released individuals, all individuals combined, and individuals of different age classes of recruitment from 2008 to 2015, respectively. Thus, we prepared three detailed encounter histories for different cohorts at the beginning of the year of release. The encounter history files are matrices of ones and zeros showing which individuals were encountered in each survey. Crested Ibis generally forage and roost in pairs and/or flocks and radio-tagged individuals can be the resighting cue for the others. Hence we assumed that all individuals had the same probabilities of resighting and survival.

We developed a set of candidate models: a model with full-time dependence in both survival and recapture ($\varphi(t)p(t)$); a model with constant in both survival and probability of recapture ($\varphi(.)p(.)$); and a model with constant survival ($\varphi(.)p(t)$) and constant probability of recapture ($\varphi(t)p(.)$). We selected the most parsimonious one based on Akaike's Information Criterion corrected for small sample sizes (AIC_c). The model with the lowest AIC_c was considered as the best supported model (Anderson *et al.* 2000, Burnham and Anderson 2002). We considered models with differences of AIC_c (ΔAIC_c) values ≤ 2 from the best-fit model equally parsimonious (Burnham and Anderson 2002), and in such cases included variables from all models with $\Delta AIC_c \leq 2$ in the analysis. We also calculated AIC_c weights for each candidate model (Buckland *et al.* 1997), which represent the weight of evidence in support of each model (Alisauskas and Lindberg 2002). The most general model was tested with the bootstrap goodness-of-fit test in MARK. The variance inflation factor (\hat{c}) was also estimated for detecting multicollinearity (Burnham and Anderson 1998).

Results

Reproduction status, age structure and adult sex ratio

Since 2008, a total of 26 pairs were recorded during an eight-year period (two in 2008, five in 2009, seven in 2010, seven in 2011, six in 2011, nine in 2013, ten in 2014 and seven in 2015). By 2015, a total of 88 birds had fledged successfully, and two second generation birds formed a breeding pair and fledged one young bird (the third generation) successfully in the wild. Additionally, annual counts of the individuals indicated that the reintroduced population grew each year over the course of our research period.

By the end of the breeding season in 2015, the minimal estimated population size was 108 individuals (Figure 2), which included 11 yearlings (0–1 year old), 22 juveniles (1–2 years old), 59 adults (≥ 2 years old) and 16 old individuals (> 10 years old). The percentage of yearlings and juveniles, breeding adults and old individuals were 30.6%, 54.6% and 14.8%, respectively (Figure 3a). For wild-born birds, the percentage of mature individuals accounted for 66.8% of our focal population.




Figure 2. The population size of the reintroduced Crested Ibis between 2007 and 2015.

The large proportion of mature individuals in our focal population indicated a gradually growing population (Figure 3b). The released individuals were mainly composed of adults and juveniles whose mean age at release was 5.2 ± 2.5 years (range 2–10, $n = 27$) for females and 6.4 ± 2.9 years (range 2–11, $n = 29$) for males, indicating higher fecundity of the founder population (Figure 3c).

The ASR of the released ibises (29 males and 27 females) was 0.536 in order to reintroduce an initial population with non-skewed ASR. There was a slight bias towards males in the ASR with the increase of wild-born offspring. By 2015, 57 mature individuals including 28 females and 29 males were observed in the reintroduced population (Table 1). The bias of ASRs (0.33–0.51) was not statistically significant, indicating a population with a balanced ASR at the present stage.

Survival rates

During our study, 56 captive-bred ibises and 88 wild-born fledglings were included in the survival analysis. The model with temporal variation in the survival rate and no temporal variation in the recapture ($\varphi(t)p(\cdot)$) was more parsimonious overall given the lowest AIC_c (Table 2). Estimate of survival rates for the released individuals ranged from 0.472 (95% CI: 0.289–0.663) in 2008 to 0.832 (95% CI: 0.414–0.972) in 2015 (Table 3). The annual survival rates of released individuals increased without further release since 2011, indicating a progress in acclimatisation of the individuals to the release site.

The survival rates for all individuals with the best model $\varphi(t)p(\cdot)$ averaged 0.738 (95% CI: 0.547–0.801), fluctuating between 0.482 and 0.857 (Tables 3, 5). The model $\varphi(a_3 - ./ ./)p(\cdot)$ was more parsimonious overall for different age classes (Table 4). As shown in Table 5, the survival rates for adults (0.757; 95% CI: 0.389–0.950) was much higher than that for yearlings (0.384; 95% CI: 0.277–0.504) but slightly lower than that of juveniles (0.830; 95% CI: 0.406–0.978).

Discussion

Since the subsequent release of 56 captive-bred individuals from 2007 to 2011, reproduction (Yu *et al.* 2015), post-fledging dispersal (Huo *et al.* 2014), mortality (Li *et al.* 2013) and survival rate (present study) of the reintroduced population have been studied over an eight-year period. Observations indicate that the number and the range of the reintroduced population increased gradually without further releases (Wang *et al.* 2016). Although there is no general agreement on




Figure 3. The age structure of different cohorts for the reintroduced populations of the Crested Ibis. The age of all individuals (a) and wild-born individuals (b) show the values for 2015 and released birds for years at release.

what constitutes a successful reintroduction, there are several criteria of success that can be referred to (Sarrazin and Barbault 1996, Seddon 1999, Robert *et al.* 2015). Additionally, the largest proportion of mature individuals and a balanced ASR in our focal population may be a positive indicator of population trends. Based on these criteria, we conclude that the primary goal of establishing a

Table 1. Adult sex ratio (ASR) estimates from wild-born individuals of the focal Crested Ibis population.

Year	Observed Number		Sample size (<i>n</i>)	Adult sex ratios (ASR)	Pearson's Chi-square test
	Females	Males			
2010	2	1	3	0.33	$\chi_1^2=0.67, P=0.41$
2011	6	5	11	0.45	$\chi_2^2=0.18, P=0.67$
2012	11	9	20	0.45	$\chi_3^2=0.40, P=0.53$
2013	17	16	33	0.48	$\chi_4^2=0.06, P=0.81$
2014	24	23	47	0.49	$\chi_5^2=0.04, P=0.84$
2015	28	29	57	0.51	$\chi_6^2=0.04, P=0.85$

self-sustaining population in one of the species' former ranges has been achieved. Our reintroduction programme could be a good example for future conservation of other endangered species.

The establishment probability of a released population can be affected by the size and composition of the release group (Armstrong and Seddon 2007). The age of introduced animals appeared to be important for the success of reintroductions owing to the higher fecundity of mature females (Bronson 1989, Komers and Curman 2000). As shown in Figure 3c, the released birds were mainly composed of adults and juveniles. Compared with the wild population (Lu *et al.* 1997), wild-born offspring were exclusively young birds including yearlings, juveniles and breeders no more than eight-years old (Figure 3b). These may be contributable to the rapid growth of our focal population (Wang *et al.* 2016).

The population-level offspring sex ratio (OSR) generally does not differ significantly from equality (Yamaguchi *et al.* 2004, Eraud *et al.* 2006), but skewed ASRs with a bias towards males are common in wild bird populations, especially in globally threatened species (Donald 2007). There is currently no quantitative evidence that a male-biased ASR skew occurs either in the wild (Lu *et al.* 1997) or in a released population (present study; Table 1). It is generally believed that a skewed ASR can be explained by higher mortality of females (Githiru *et al.* 2006, Székely *et al.* 2006). For the Crested Ibis, the sexual differences in survival rates and offspring sex ratio remain poorly studied. Therefore we cannot explain the current balanced ASRs of the ibis population. There is evidence that a general correlation exists between the ASR and the population trend, or the habitat quality (Zanette 2001, Wilkinson *et al.* 2002, Johnson *et al.* 2006). The ASR might be a useful indicator of a species' population trajectory or conservation status (Donald 2007). In monogamous mating systems (e.g. Crested Ibis), the extinction probability of small populations is likely to be lowest when the sex ratio is balanced (Bessa-Gomes *et al.* 2004). This might be one of the causes resulting in the rapid growth of wild and reintroduced ibis populations.

Since the release, the reintroduced population has been monitored extensively with particular attention being paid to the survival of wild-born birds over an eight-year period (Yu *et al.* 2015).

Table 2. Model selection for the survival (φ) and resighting probability (p) of the released and all individuals in the reintroduced Crested Ibis population in Ningshan County (2007-2015), based on a variance inflation factor (\hat{c}) of 0.96 ($P=0.51$) and 0.99 ($P=0.47$), respectively.

Cohorts	Model	AIC_c	ΔAIC_c	AIC_c Weight	NO. P	Deviance
Released individuals	φ (t) p (.)	245.79	0.00	0.582	9	93.84
	φ (.) p (.)	246.51	0.71	0.407	2	109.76
	φ (t) p (t)	254.64	8.84	0.007	15	88.41
	φ (.) p (t)	256.00	10.21	0.003	9	104.05
All individuals	φ (t) p (.)	511.06	0.00	0.770	9	183.39
	φ (t) p (t)	513.93	2.88	0.183	15	173.19
	φ (.) p (.)	517.09	6.03	0.038	2	204.00
	φ (.) p (t)	519.82	8.76	0.009	9	192.15

Survival rates (φ) and the probability of recapture (p) were modeled as constant over time (.) or as an annual function of time (t). The number of estimated parameters is given by (NO. P).

Table 3. Annual survival estimates (ϕ), SE, 95% confidence intervals and relative important values for the best model of the released and all individuals.

Years	Released individuals			All individuals		
	ϕ	SE	LCI-UCI (with 95% CI)	ϕ	SE	LCI-UCI (with 95% CI)
2008	0.472	0.100	0.289-0.663	0.482	0.102	0.293-0.675
2009	0.800	0.107	0.519-0.937	0.809	0.107	0.519-0.943
2010	0.566	0.096	0.378-0.737	0.523	0.084	0.362-0.680
2011	0.886	0.091	0.566-0.978	0.845	0.077	0.634-0.945
2012	0.748	0.092	0.531-0.885	0.831	0.064	0.667-0.924
2013	0.896	0.082	0.603-0.980	0.753	0.067	0.601-0.861
2014	0.816	0.107	0.523-0.947	0.810	0.066	0.647-0.908
2015	0.832	0.138	0.414-0.972	0.857	0.076	0.641-0.953

With exception of yearlings, the average survival estimates for each cohort such as released individuals, juveniles and adults were relatively high (Table 5). The survival rate of adults (0.812) was slightly higher than that of the wild population (71%; Ding 2004). The survival rate of yearlings may be underestimated because some fledglings emigrated outside our study area (Yu *et al.* 2015).

Most reintroduction projects aimed at saving endangered animals in the 1970s and 1980s failed to establish viable populations, partly due to poor planning and subsequent monitoring (Griffith *et al.* 1989, Wolf *et al.* 1996). The success of reintroduction programmes is influenced by the quality of the habitat where the reintroductions took place, the adaptive ability of the birds to new conditions, as well as the number and age structure of the released birds (Griffith *et al.* 1989, Wolf *et al.* 1996, Armstrong *et al.* 2002, Tweed *et al.* 2006). Small release groups could fail to establish a self-sustaining population due to random factors (demographic stochasticity), low survival rates at low densities (Allee effects) (Caughley 1994). Our modelling results indicated that the initial population showed lower survival rate (0.472; Table 3) due to poor acclimatisation and high level of post-release dispersal immediately after release. Of 20 mortalities from 56 released individuals, eight (40%) died from food shortage, especially in winter, six (30%) from flight collision, five (25%) from electrocution on power lines, and one (5%) from unknown cause (Li *et al.* 2013). It is often believed that dispersal and/or mortality can be reduced by holding animals at the release site for a specified period. For example, housing animals in an enclosure could reduce dispersal and mortality for mammals (Tuberville *et al.* 2005, Shier and Owings 2006, Hamilton *et al.* 2010), yet it was useless for bird species. Therefore, a method was put forward to broadcast animal calls in the vicinity of the release sites (Molles *et al.* 2008). For the reintroduced Crested Ibis, subsequent releases were conducted in Ningshan County in the southern Qinling Mountains (Yu *et al.* 2009, 2015), Tongchuan City (Wang 2016) and Qianhu Wetland Park in the northern Qinling Mountains (Yu *et al.* unpubl. data). These showed that soft release and food supplementation are useful to

Table 4. Model selection for the survival rate (ϕ) and resighting probability (p) for the different age classes of the reintroduced Crested Ibis in Ningshan County (2008-2015), based on a variance inflation factor (\hat{c}) of 0.97 ($P=0.47$).

Model	AIC_c	ΔAIC_c	AIC_c Weight	NO. P	Deviance
ϕ (a3-././.) p (.)	216.12	0.00	0.5533	4	65.29
ϕ (a3-././t) p (t)	219.18	3.06	0.1197	14	45.11
ϕ (a3-././.) p (t)	219.47	3.35	0.1037	10	55.15
ϕ (a3-t././.) p (.)	219.67	3.55	0.0938	10	55.35
ϕ (a3-t././t) p (.)	220.25	4.13	0.0701	14	46.18
ϕ (a3-t././.) p (t)	221.77	5.65	0.0328	16	42.57

Six relatively good models describing survival rate (ϕ) and resighting probability (p) were selected. The number of estimated parameters is given by (NO. P).

Table 5. Survival estimates (ϕ), SE, 95% confidence intervals and relative important values for the best model of the different age classes, and mean survival probabilities for released and all individuals.

Cohorts	ϕ	SE	LCI-UCI (with 95% CI)	<i>n</i>
Yearlings	0.384	0.050	0.277-0.504	11
Juveniles	0.853	0.058	0.406-0.978	30
Adults	0.812	0.054	0.389-0.950	47
Released individuals	0.752	0.153	0.478-0.887	56
All individuals	0.738	0.149	0.547-0.801	144

hold the individuals around the release site. For example, most of the breeders nested near the release site in Tongchuan (Wang 2016) and Qianyang (Yu *et al.* unpubl. data) ibis populations.

The success of a reintroduction can be affected by several factors including population size, post-release management, habitat conditions and genetic makeup at population level (Armstrong and Seddon 2007). It is generally believed that there is a positive relationship between release group size and establishment success (Griffith *et al.* 1989, Wolf *et al.* 1996). A study on even-toed ungulate (Artiodactyla) species showed that the variation in the rate of increase of reintroduced animals was larger when < 20 animals were released (Komers and Curman 2000). Where post-release dispersal and mortality are low, populations can potentially establish successfully from < 10 released individuals (Taylor *et al.* 2005). However, high post-release dispersal and mortality would create a large disparity between the release group size and the effective initial population size. For example, in spite of a larger release group size (107 captive-bred individuals), the average breeding success of the Sado ibis population in Japan is much lower than that of our focal population (Wang *et al.* 2016). Therefore, we conclude that the effective initial population size may vary with habitat conditions in different release sites.

In addition to factors mentioned above, habitat conditions such as food availability and native predators are crucial to population survival (Huo *et al.* 2014). In the wild and our focal populations, Crested Ibises mainly forage in rice fields, shallow rivers, arid croplands and grasslands during different seasons (Shi and Cao 2001, Huo *et al.* 2014). However, in the northern Qinling Mountains, the rice fields in the vast areas along the Wei River almost disappeared over the past three decades, so we speculated that the other two reintroduced populations (Tongchuan and Qianhu) will be increasingly difficult to sustain because they would have to forage exclusively in shallow rivers. Another factor that has significantly decreased the survival rate is nest predation, such as by snakes (especially the king rat snake *Elaphe carinata*) and birds of prey (e.g. the Goshawk *Accipiter gentilis*). They usually attack eggs and chicks; in contrast, weasels *Mustela* spp. may also prey on adults as well as on eggs and nestlings (Cao *et al.* 1995, Xi *et al.* 1997, Lu *et al.* 1997, Zhang *et al.* 2000, Li *et al.* 2013). The king rat snake was a significant nest predator that usually preyed on nidicolous nestlings of the Crested Ibis. By 2014, four nestlings from the reintroduced population (Li *et al.* 2013) and 15 nestlings from the wild population by 2006 (Yu *et al.* 2006) had died from predation by this dangerous reptile. The predation of the Goshawk on chicks resulted in complete failure of two breeding pairs in 2013 (Li *et al.* 2013).

Although the reintroduction project in Ningshan County, as well as in other places, has great promise with some necessary interventions, threats to the population remain. As a consequence, the following proposals are made. First, more efforts are needed to reduce post-release mortality and dispersal with a focus on pre-release anti-predator training and habitat restoration. Another approach is usually to model the relationship based on the available data on survival, reproduction and dispersal rates for the species (Komers and Curman 2000, Towns and Ferreira 2001). Second, in the case of highly endangered flagship species, high intensity post-release intervention (e.g. food supplementation, veterinary care or predator control) is clearly warranted (Seddon 1999). Food supplementation has been proved to be more effective for enhancing reproductive success both in wild and reintroduced populations of the Crested Ibis (Yu *et al.* 2006, 2015). Some approaches to prevent the birds from dying of predation (e.g. the king rat snake) have not been very effective. Therefore, there is an urgent need for a

cost-effective management technique that can be applied at an ecosystem level (Yu *et al.* 2015). Third, although habitat conditions will be the main drivers of population growth, this will also be affected by the genetic diversity of the organism. Great care should be taken in the selection of reintroduction sites, as species with low genetic diversity are thought to be more limited in their ability to tolerate a wide range of environmental factors (Zhang *et al.* 2004). Hence, research is urgently needed, not only to predict how management will affect genetic diversity of reintroduced populations, but also to predict the effects on population growth and persistence. Finally, a long-term and effective international cooperation on ibis conservation among China, Japan, Korea, and Russia should be established.

Acknowledgements

We are very grateful to the reintroduction base of Crested Ibis in Ningshan County for providing logistical help and research aid. We also thank Cao Hai-xin, Guo Jun-feng and Ma Xiao-chun for obtaining the survival data and Liu Chao for drawing the maps. This work was completely supported by the National Nature Science Foundation of China under grant number 31172103 and 31572282.

References

- Alisauskas, R. T. and Lindberg, M. S. (2002) Effects of neckbands on survival and fidelity of white-fronted and Canada geese captured as non-breeding adults. *J. Appl. Stat.* 29: 521–537.
- Armstrong, D. P., Castro, I., Alley, J. C., Feenstra, B. and Perrott, J. K. (1999) Mortality and behaviour of HiHi, an endangered New Zealand honeyeater, in the establishment phase following translocation. *Biol. Conserv.* 89: 329–339.
- Armstrong, D. P., Davidson, R. S., Dimond, W. J., Perrott, J. K., Castro, I., Ewen, J. G., Griffiths, R. and Taylor, J. (2002) Population dynamics of reintroduced forest birds on New Zealand islands. *J. Biogeogr.* 29: 609–621.
- Armstrong, D. P. and Ewen, J. G. (2002) Dynamics and viability of a New Zealand Robin population reintroduced to regenerating fragmented habitat. *Conserv. Biol.* 16: 1074–1085.
- Armstrong, D. P. and Seddon, P. J. (2007) Directions in reintroduction biology. *Trends Ecol. Evol.* 23: 20–25.
- Baumgardt, J. A., Goldberg, C. S., Reese, K. P., Connelly, J. W., Musil, D. D., Garton, E. O. and Waits, L. P. (2013) A method for estimating population sex ratio for sage-grouse using noninvasive genetic samples. *Mol. Ecol. Resour.* 13: 393–402.
- Bessa-Gomes, C., Legendre, S. and Clobert, J. (2004) Allee effects, mating systems and the extinction risk in populations with two sexes. *Ecol. Lett.* 7: 802–812.
- BirdLife International (2001) *Threatened birds of Asia: The BirdLife International Red Data Book*. Cambridge, UK: BirdLife International.
- Bronson, F. H. (1989) *Mammalian reproductive biology*. Chicago, USA: Chicago University Press.
- Buckland, S. T., Burnham, K. P. and Augustin, N. H. (1997) Model selection: An integral part of inference. *Biometrics* 53: 603–618.
- Burnham, K. P. and Anderson, D. R. (1998) *Model selection and inference - a practical information-theoretic approach*. New York, USA: Springer.
- Burnham, K. P. and Anderson, D. R. (2002) *Model selection and multimodel inference*. New York, USA: Springer-Verlag.
- Cao, Y. H., Lu, X. R. and Zhai, T. Q. (1995) The harm made by snakes to the Crested Ibis *Nipponia nippon*, in breeding season. *J. Northwest Univ.* 25 (SL): 722–724. (In Chinese).
- Caughley, G. (1994) Directions in conservation biology. *J. Anim. Ecol.* 63: 215–244.
- Ding, C. Q. (2004) Research on the crested ibis. Shanghai, China: Shanghai Press of Science and Education. (In Chinese).
- Donald, P. F. (2007) Adult sex ratios in wild bird populations. *Ibis* 149: 671–692.
- Eraud, C., Lallemand, J. and Lormée, H. (2006) Sex-ratio of Skylark *Alauda arvensis* broods in relation to timing of breeding. *Bird Study* 53: 319–322.
- Griffith, B., Scott, J. M., Carpenter, J. W. and Reed, C. (1989) Translocation as a species

- conservation tool: status and strategy. *Science* 245: 477–480.
- Githiru, M., Lens, L., Bennun, L. A. and Perrins, C. (2006) Experimental evidence of ‘floaters’ in two isolated populations of an Afrotropical forest bird. *Ostrich* 77: 28–35.
- Hamilton, L. P., Kelley, P. A., William, D. E., Kelt, D. A. and Wittmer, H. U. (2010) Factors associated with survival of reintroduced riparian brush rabbits in California. *Biol. Conserv.* 143: 999–1007.
- Huo, Z. P., Guo, J. F., Li, X. and Yu, X. P. (2014) Post-fledging dispersal and habitat use of a reintroduced population of the Crested Ibis (*Nipponia nippon*). *Avian Res.* 5: 7.
- IUCN (1998) *Guidelines for reintroductions*. Gland, Switzerland and Cambridge, UK: International Union for Conservation of Nature (World Conservation Union/SSC) SSC Re-introduction Specialist Group.
- IUCN (2013) *Guidelines for reintroductions and other conservation translocations. Version 1.0*. Gland, Switzerland: IUCN Species Survival Commission.
- Johnson, M. D., Sherry, T. W., Holmes, R. T. and Marra, P. P. (2006) Assessing habitat quality for a migratory songbird wintering in natural and agricultural habitats. *Conserv. Biol.* 20: 1433–1444.
- Komers, P. E. and Curman, G. P. (2000) The effect of demographic characteristics on the success of ungulate re-introductions. *Biol. Conserv.* 93: 187–194.
- Lebreton, J. D., Burnham, K. P., Clobert, J. and Anderson, D. R. (1992) Modeling survival and testing biological hypotheses using marked animals: a unified approach with case studies. *Ecol. Monogr.* 62: 67–118.
- Li, X., Huo, Z. P. and Yu, X. P. (2013) Cause of death of a reintroduced population of the Crested Ibis of in Ningshan, Shaanxi. *Chi. J. Zool.* 48: 701–706. (In Chinese).
- Liu, Y. Z. (1981) The re-discovery of the Crested Ibis in Qinling Mountain. *Acta. Zool. Sinica* 27: 273. (In Chinese).
- Lu, B. Z., Zhai, T. Q., Zhang, Y. M. and Geng, Z. Z. (1997) Wild population structure and relationship of oriental crested ibis *Nipponia nippon*. *J. Shaanxi Normal Univ.* 25 (SL): 98–104. (In Chinese).
- Molles, L. E., Calcott, A., Peters, D., Delamare, G., Hudson, J., Innes, J., Flux, I. and Waas, J. (2008) “Acoustic anchoring” and the successful translocation of North Island kokako *Callaeas cinerea wilsoni* to a New Zealand mainland management site within continuous forest. *Notornis* 55: 57–68.
- Nagata, H. and Yamagishi, S. (2013) Re-introduction of crested ibis on Sado Island, Japan. *Global Re-introduction Perspectives* 2013: 58–62.
- Robert, A., Colas, B., Guigon, I., Kerbiriou, C., Mihou, J. B., Saint-Jalme, M. and Sarrazin, F. (2015) Defining reintroduction success using IUCN criteria for threatened species: a demographic assessment. *Anim. Conserv.* 18: 397–406.
- Sarrazin, F. and Barbault, R. (1996) Reintroduction: challenges and lessons for basic ecology. *Trends Ecol. Evol.* 11: 474–478.
- Schaub, M., Pradel, R. and Lebreton, J. D. (2004) Is the Reintroduced white stock *Ciconia ciconia* population in Switzerland self-sustainable? *Biol. Conserv.* 119: 105–114.
- Scott, J. M. and Carperter, J. W. (1987) Release of captive-reared or translated endangered birds: what do we need to know? *Auk* 104: 544–545.
- Seddon, P. J. (1999) Persistence without intervention: assessing success in wildlife re-introductions. *Trends Ecol. Evol.* 14: 503.
- Shi, D. C. and Cao, Y. H. (2001) *The Chinese Crested Ibis*. Beijing: China Forestry Publishing House. (In Chinese).
- Shier, D. M. and Owings, D. H. (2006) Effects of predator training on behaviour and post-release survival of captive prairie dogs *Cynomys ludovicianus*. *Biol. Conserv.* 132: 126–135.
- Sun, R. Y., Li, Q. F., Niu, C. J. and Lou, A. R. (2005) *Basic ecology*. Beijing: Advanced Education Press, 67. (In Chinese).
- Székely, T., Thomas, G. H. and Cuthill, I. C. (2006) Sexual conflict, ecology and breeding systems in birds. *Bioscience* 56: 801–808.
- Taylor, S. S., Jamieson, I. G. and Armstrong, D. P. (2005) Successful island reintroductions of New Zealand robins and saddlebacks with small numbers of founders. *Anim. Conserv.* 8: 415–420.
- Towns, D. R. and Ferreira, S. M. (2001) Conservation of New Zealand lizards

- Lacertilia: Scincidae* by translocation of small populations. *Biol. Conserv.* 98: 211–222.
- Tuberville, T., Buhlmann, K. A., Bjorkland, R. K. and Booher, D. (2005) Ecology of the Jamaican slider turtle *Trachemys terrapen*, with implications for conservation and management. *Chelonian Conserv. Biol.* 4: 908–915.
- Tweed, E. J., Foster, J. T., Woodworth, B. L., Monahan, W. B., Kellerman, J. L. and Lieberman, A. (2006) Breeding biology and success of a reintroduced population of the critically endangered Puaiohi *Myadestes palmeri*. *Auk* 123: 753–763.
- Wang, H. Q. (2016) Reproduction of reintroduced *Nipponia nippon* in Tongchuan of Shaanxi Province. *Sichuan J. Zool.* 35: 471–474. (In Chinese).
- Wang, C., Liu, D. P., Qing, B. P., Ding, H. H., Cui, Y. Y., Ye, Y. X., Lu, J., Yan, L., Ke, L. and Ding, C. Q. (2014) The current population and distribution of wild Crested Ibis *Nipponia nippon*. *Chinese J. Zool.* 49: 666–671. (In Chinese).
- Wang, M., Ye, X. P., Li, Y. F., Huo, Z. P., Li, X. and Yu, X. P. (2016) On the sustainability of a reintroduced Crested Ibis population in Qinling Mountains, Shaanxi, Central China. *Restoration Ecol.* (doi: 10.1111/rec.12409).
- Wilkinson, N. I., Langston, R. H. W., Gregory, R. D., Gibbons, D. W. and Marquiss, M. (2002) Capercaillie *Tetrao urogallus* abundance and habitat use in Scotland, in winter 1998–99. *Bird Study* 49: 177–185.
- Wolf, C. M., Griffith, B., Reed, C. and Temple, S. A. (1996) Avian and mammalian translocations: update and reanalysis of 1987 survey data. *Conserv. Biol.* 10: 1142–1154.
- Xi, Y. M., Lu, B. Z., Geng, Z. Z., Zhai, T. Q. and Zhang, Y. M. (1997) Cure of the Crested Ibis. *Chinese Wildl.* 18: 28–30. (In Chinese).
- Yamaguchi, N., Kawano, K. K., Eguchi, K. and Yahara, T. (2004) Facultative sex ratio adjustment in response to male tarsus length in the Varied Tit *Parus varius*. *Ibis* 146: 108–113.
- Yu, X. P., Liu, N. F., Xi, Y. M. and Lu, B. Z. (2006) Reproductive success of the Crested Ibis (*Nipponia nippon*). *Bird Conserv. Internatn.* 16: 325–343.
- Yu, X. P., Chang, X. Y., Li, X., Chen, W. G. and Shi, L. (2009) Return of the Crested Ibis *Nipponia nippon*: a reintroduction programme in Shaanxi province, China. *Birding ASIA* 11: 80–82.
- Yu, X. P., Li, X. and Huo, Z. P. (2015) Breeding ecology and success of a reintroduced population of the endangered Crested Ibis *Nipponia nippon*. *Bird Conserv. Internatn.* available on CJO2014. <http://dx.doi.org/10.1017/S0959270914000136>.
- Zanette, L. (2001) Indicators of habitat quality and the reproductive output of a forest songbird in small and large fragments. *J. Avian Biol.* 32: 38–46.
- Zhang, Y. M., Lu, B. Z., Zhai, T. Q., Xi, Y. M. and Wang, Y. J. (2000) Analysis on the death reason and conservation measures of Crested Ibis. Pp. 117–122 in *Proceedings of International Workshop on the Crested Ibis Conservation' 99*. Beijing: China Forestry Publishing House. (In Chinese).
- Zhang, B., Fang, S. G. and Xi, Y. M. (2004) Low genetic diversity in the Endangered Crested Ibis *Nipponia nippon* and implications for conservation. *Bird Conserv. Internatn.* 14: 183–190.

YANFANG LI, XINPING YE, MIN WANG, RONG DONG, ZHIPING HUO, XIAOPING YU*
College of Life Sciences, Shaanxi Normal University, 710119, Xi'an City, China.

XIA LI

Crested Ibis Reintroduction Centre, Shaanxi Ningshan Forestry Administrations, Ankang City,
725000, China.

*Author for correspondence; e-mail: yuxp64@163.com

Received 10 November 2015; revision accepted 19 December 2016;

Published online 9 May 2017