

Epidemiological of and risk factors for Alzheimer's disease: A review

Jana Povova^a, Petr Ambroz^a, Michal Bar^b, Veronika Pavukova^a, Omar Sery^c, Hana Tomaskova^a, Vladimir Janout^a

Background. Alzheimer's disease (AD) is the most common form of dementia. It is a degenerative, incurable and terminal disease. The increasing prevalence of AD is, among other reasons, due to population aging, which is, to a certain extent, seen worldwide. Continuous advances in health care keep increasing life expectancy. Official statistics are likely to significantly underestimate the actual prevalence of AD. Alzheimer's disease represents an important public health problem. Its aetiology is still unknown and for this reason, it is necessary to study all potential risk factors which may contribute to the development of this disease.

Methods. We searched original and review articles addressing Alzheimer's disease using key words Alzheimer's disease, epidemiology, risk factors and prevention. We found and used one hundred and four references.

Conclusions. Based on epidemiological studies, genetic studies, neuroimaging methods and neuropathology research, three basic etiological hypotheses of the development of AD have been formulated: genetic, vascular and psychosocial. At present, the level of evidence is insufficient for the etiological role of other factors, such as nutrition, occupational exposure to various substances and inflammation. From the point of view of early diagnosis and application of primary or secondary prevention principles, genetic factors are the most important.

Key words: Alzheimer's disease, epidemiology, risk factors, prevention

Received: September 9, 2011; Accepted with revision: May 17, 2012; Available online: June 14, 2012

<http://dx.doi.org/10.5507/bp.2012.055>

^aDepartment of Epidemiology and Public Health, Faculty of Medicine, University of Ostrava, Czech Republic

^bDepartment of Clinical Studies, Faculty of Medicine, University of Ostrava

^cDepartment of Biochemistry, Faculty of Science, Masaryk University Brno

Corresponding author: Jana Povova, e-mail: Jana.povova@osu.cz

INTRODUCTION

Alzheimer's disease (AD) is the most common form of dementia. It is a degenerative, incurable and terminal disease. Although it is mostly diagnosed in people over 65 years, a less prevalent early-onset form may develop much earlier.

Alzheimer's disease affects most (up to 75%) of the more than 35 million people suffering from dementia worldwide. The prevalence is believed to double every 20 years. Thus, it is estimated that approximately 115 million people may be affected by AD in 2050. The disease has a major impact not only on the sufferers but also on persons caring for them as well as the entire society.

The etiological factors are mostly unknown. There is increasing evidence of the role that certain risk factors play in the development of the disease, such as genetic and vascular factors or disorders. Also known are psychosocial factors which may positively influence the pathogenesis and clinical manifestation of the disease. Therefore, intervention in these areas may lower the risk of its development or at least delay the clinical manifestation.

The increasing prevalence of AD is, among other reasons, due to population aging, which is, to a certain extent, seen worldwide.

OCCURRENCE

Prevalence

Aggregate data from studies on the prevalence of AD in Europe reported the age-standardized prevalence of AD in people aged 65 or more to be 4.4% (ref.¹). A US study performed in a representative sample of persons over 70 years of age reported AD prevalence of 9.7% (ref.²). The global prevalence of AD is estimated at 3.9% in people older than 60 years, with regional variations in individual continents³.

In developed countries, approximately 1 in 10 persons over 65 years of age suffers from a form of dementia compared with more than one third of those older than 85 years^{4,5}.

Incidence

Two US studies of persons aged 65 or more reported an AD incidence of 15.0 per 1,000 person-years. The incidence rates for males and females were 13.0 and 16.9 per 1,000 person-years, respectively^{6,7}.

In a 3 year follow-up, a Swedish prospective study involving 987 persons aged 75 years or older found the incidence rates of dementia in the age group of 75-79 years to be 19.6 and 12.4 per 1,000 person-years in females and males, respectively. In those aged 90 or more, the incidence rates of dementia were 86.7 per 1,000 person-years in females and 15.0 per 1,000 person-years in males.

A similar distribution by age and gender was seen in AD. The risk of the development of any form of dementia is reported to be approximately twice as high in females as in males and the risk of Alzheimer's disease is as much as three times higher in females⁸.

Numerous other studies report the incidence rates of dementia and AD as the number of persons per year.

For example, in a British prospective cohort study of 1,070 persons aged 65 or more and followed for 3 years, the incidence of all types of dementia was 9.2 per 1,000 population and year, of which the AD incidence was 6.3 per 1,000 population and year⁹.

Koukolik reported approximately 35,000 AD patients aged 65 years or more in former Czechoslovakia in 1983 (ref.¹⁰).

According to the Czech Alzheimer Society, approximately 130,000 people in the Czech Republic currently suffer from dementia, two thirds of which have AD.

RISK FACTORS

Hypotheses

Based on epidemiological studies, neuroimaging methods and neuropathology research, three etiological hypotheses of the development have been reported, with moderate or strong evidence¹¹:

- genetic,
- vascular, and
- psychosocial.

At present, the level of evidence is insufficient for the etiological role of other factors, such as nutrition, occupational exposure to various substances and inflammation.

Genetic hypothesis

Early-onset familial AD is usually caused by autosomal dominant mutations in the genes for amyloid precursor protein (APP), presenilin 1 and presenilin 2. This form of AD accounts for approximately 2–5% of all AD cases¹².

First-degree relatives of patients with AD are at higher lifetime risk of developing the disease than the rest of the population¹³. Both genetic and other factors contribute to the increased risk and familial aggregation of AD cases. As far as genetic factors are concerned, this risk may be partly due to the presence of the apolipoprotein E (APOE) apoE4 allele. Other genes also contribute to the pathogenesis of this mental disorder¹⁴. The apoE4 allele is the only proven genetic factor so far identified in the development of both the early- and late-onset forms of AD. This factor increases susceptibility to AD but it is neither necessary nor sufficient for the development of this disease. The higher the number of the apoE4 alleles, the higher the risk of AD and the lower the age of onset. The risk effect of the presence of the apoE4 allele decreases with age. Generally, approximately 15–20% of AD cases may be attributed to this risk¹⁵.

Tens of other candidate genes with polymorphisms affecting the risk of the development of AD have been studied. Predominantly, these are secretase genes – e.g.

BACE1 (beta-site amyloid precursor protein-cleaving enzyme 1), presenilin 1, peptidase genes – e.g. ECE1 (endothelin-converting enzyme), IDE (insulin-degrading enzyme), microtubule and cytoskeletal genes – e.g. MAPT (microtubule-associated protein tau), synaptic genes – e.g. ABCA1 (ATP-binding cassette A1 transporter), anti-apoptotic genes – e.g. IL-1 (interleukin 1), protease genes -e.g. ACE (angiotensin-converting enzyme) and other genes such as the gene for APP.

Vascular hypothesis

Vascular risk factors, such as smoking, obesity and high total cholesterol levels, together with vascular morbidity, such as hypertension, diabetes mellitus and asymptomatic cerebral infarction, are associated with a higher risk of dementia including Alzheimer's disease.

Smoking

Earlier cross-sectional studies often reported a lower prevalence of AD among smokers compared with non-smokers¹⁶. This seemingly protective effect was probably due to survivor bias since the proportion of smokers among the prevalent cases was smaller.

When incident cases of AD were studied, however, the situation was completely reversed^{17–19}. That is, numerous analytical studies found a significantly increased risk of AD associated with cigarette smoking, especially in apoE4 allele non-carriers^{20–22}. Meta-analyses of these analytical studies concluded that current smoking was associated with an increased risk of the development of AD, with RR=1.79; 95% CI 1.43–2.23 (ref.^{23,24}).

Alcohol

It is well recognized that alcohol abuse causes alcohol dementia. Moreover, middle-aged heavy drinkers, especially apoE4 allele carriers, were found to have a more than 3-fold higher risk of dementia and AD later in their lives²⁵. On the other hand, the risk of developing dementia and AD was reduced in light and moderate alcohol consumers^{26,27}.

In heavy consumers, alcohol clearly damages the brain. Even light to moderate alcohol consumption was found to be related to brain atrophy and volume loss^{28,29}.

Overweight and obesity

Higher BMI in middle age is a risk factor for AD and other dementias. Higher BMI or obesity (in particular abdominal obesity) at around the age of 50 years means an increased risk of dementia 20–25 years later^{30–33}. Several analytical studies in the elderly suggested that a significant decrease in BMI was associated with a higher risk of AD in the following 5–6 years^{37,38}.

Blood pressure and management of hypertension

Increased blood pressure in middle age, especially if uncontrolled, was associated with a higher risk of the later development of AD in several observational studies^{39,40}.

Findings from analytical studies of the association between blood pressure and the risk of dementia are not

consistent. Studies with a relatively short follow-up (less than 3 years) found no or even an inverse association between blood pressure values and the risk of dementia and AD (ref.⁴¹). Since dementia has a long latent period and blood pressure may drop in its pre-clinical stage, an absent or inverse association may be interpreted as a result of the process^{42,43}.

However, analytical studies with a longer follow-up (e.g. more than 6 years) reported an inverse association^{44,46}, suggesting that low blood pressure later in the life may contribute to the development or clinical manifestation of dementia including AD (ref.^{41,47}).

Longitudinal studies have repeatedly shown a protective effect of antihypertensive drugs against the development of dementia and AD (ref.^{41,48,49}). Recent analytical studies have suggested that the protective effect of antihypertensive therapy may depend on its duration and the patient's age. Greater protective effects were found in the younger elderly (e.g. under 75 years of age) and in those with long-term treatment^{50,51}. Antihypertensive therapy may protect against dementia and AD by delaying the atherosclerotic process, decreasing the number of atherosclerotic lesions and improving cerebral perfusion⁴¹.

Hypercholesterolemia and statin therapy

High total serum cholesterol levels in middle age were found to be a risk factor for the development of AD at a later age^{52,53}. High total cholesterol in middle age is a risk factor for the development of AD and other dementias 20 years later but decreasing serum cholesterol levels in late middle age may be due to ongoing disease processes and may represent a marker for later AD and other dementias⁵⁴.

According to several cross-sectional and case-control studies, the use of statins significantly decreases the prevalence of AD (ref.^{55,56}). Whereas an analytical study (the Rotterdam Study) showed that the use of statins was associated with a lower risk of AD (ref.⁵⁷), other prospective studies found either no beneficial effect or only a slightly decreased risk of AD related to the use of statins^{58,59}. Experimental studies have suggested that statins may decrease the production of beta amyloid both *in vitro* and *in vivo*. Statins also have various other effects that may be beneficial for the CNS and thus may lower the risk of AD.

Nutritional factors

Several analytical studies showed a decreased risk of AD associated with higher intake of antioxidants such as vitamins E and C (ref.^{60,61}), either in the diet or in dietary supplements. However, some studies found a negative effect⁶². Yet other studies showed that the Mediterranean diet (high intake of fish, fruit and vegetables rich in antioxidants) was associated with a lower risk of AD, independent of vascular risk factors^{63,64}. A Cochrane systematic review concluded that supplementation with folic acid and vitamin B12 had no beneficial effect on cognitive functions despite the fact that folic acid and vitamin B12 effectively decrease serum homocysteine levels⁶⁵.

It was also reported that a diet rich in saturated fats and cholesterol increased the risk of AD (ref.⁶⁶) whereas

polyunsaturated fatty acids and fish may be protective against dementia^{67,68}.

Diabetes mellitus

Many longitudinal studies found an increased risk of not only neurodegenerative but also vascular dementia in people with diabetes⁶⁹⁻⁷¹. This risk was confirmed by a systematic review⁷². The presence of diabetes in middle age or a longer duration of diabetes may play a key role in the development of AD and other dementias^{73,74}. Moreover, borderline conditions, prediabetes or impaired glucose tolerance are also related to an increased risk of AD and other dementias in very old people⁷⁵. This relationship may be partly explained by diabetic comorbidities such as hypertension and dyslipidaemia⁷⁶⁻⁷⁸.

A relationship was also found between a higher prevalence of AD in an elderly population in Finland⁷⁹, although an analytical study of a multiethnic cohort of the elderly in the USA found no association between the metabolic syndrome and either the prevalence or incidence of AD, despite the fact that two components of the syndrome, i.e. diabetes and hyperinsulinaemia, were associated with increased risk of incident AD (ref.⁸⁰).

Cardiovascular and cerebrovascular diseases

A significantly increased risk of dementia and AD was found in patients who had suffered a stroke as well as clinically silent cerebral infarction confirmed by magnetic resonance imaging^{81,82}.

Also cardiovascular diseases are associated with an increased incidence of dementia and AD, with the highest risk of dementia in persons with peripheral artery disease, suggesting that extensive peripheral atherosclerosis is a risk factor for AD (ref.^{83,84}).

Neuropathological studies have suggested that cerebrovascular lesions, atherosclerosis and neurodegenerative changes often coexist with each other in the brain, producing a clinical manifestation of the dementia syndrome^{85,86}.

Psychosocial hypothesis

A systematic review found that psychosocial factors and an active lifestyle throughout life may decrease the risk of dementias including AD (ref.⁸⁷).

Education and socioeconomic status

Lower education is tied to increased risk of dementia and AD. This link has been confirmed by many cross-sectional and longitudinal studies^{88,89}. Although education and socioeconomic status are highly correlated with each other, when studied separately, an independent association was only found for education⁹⁰.

Social network and social engagement

Longitudinal observational studies suggest that a poor social network or a lack of social engagement are associated with decreased cognitive functions and dementia^{87,91}. The risk of dementia was also higher in elderly persons with increased social isolation and less frequent and unsatisfactory contacts with relatives and friends. Persons with low neuroticism combined with high extraversion

had a lower risk of dementia⁹². Low levels of social engagement in late life and decreased social engagement from middle to late life were associated with a two-fold increase in the risk of the development of dementia and AD later in life^{93,94}.

Physical activity

Regular physical exercise is associated with a delay in onset of dementia and AD in the elderly without cognitive impairment⁹⁵. Physical activity in the form of various leisure activities rather than sports or specific physical exercise led to a decrease in the risk of dementia⁹⁶. Even low-intensity physical activity such as walking may lower the risk of dementia and cognitive impairment⁹⁷. A significant protective effect of regular physical activity in middle age, with respect to the development of dementia and AD later in the life, was found especially in persons with the apoE4 allele⁹⁸.

Mental activity

Various activities requiring a mental effort, such as reading, social and cultural activities, knitting, gardening, dancing, tabletop games, playing musical instruments, watching specific TV programmes, showed a protective effect against dementia and AD (ref.^{99,72}). A study of Swedish twins showed that complexity of work, in particular more complex work with people, may reduce the risk of AD (ref.¹⁰⁰). A recent neuroimaging study suggested that a high level of complex mental activity across the lifespan was correlated with reduced hippocampal atrophy¹⁰¹.

PREVENTION

Primary prevention

Although the mechanisms of action of vascular and psychosocial factors participating in the pathogenesis and clinical manifestation of AD have not been fully understood, primary prevention is possible since most vascular and psychosocial factors including lifestyle factors are modifiable^{11,102}.

Primary prevention strategies addressing the vascular pathway in the development of AD include:

- measures against hypertension, obesity, increased glucose levels and diabetes mellitus in middle age,
- measures against cardiac failure and preventing very low blood pressure and thus maintaining sufficient cerebral perfusion.

Primary prevention strategies focused on maintaining an active and socially integrated lifestyle:

- ensuring an extensive social network,
- frequent participation in social physical and intellectual stimulating activities^{11,87}.

Secondary prevention

AD is characterized by a pre-clinical stage lasting several years during which progressive neurodegenerative changes occur in the brain, even before the onset of

typical clinical signs¹⁰³. However, this pre-clinical stage is difficult to identify although some clinical markers, neuroimaging biomarkers and biochemical markers are available¹⁰⁴:

- mild cognitive impairment (isolated memory loss),
- biochemical markers in the serum and cerebrospinal fluid (beta amyloid and tau protein),
- neuroimaging methods are considered an effective tool for diagnosing AD in the pre-clinical stage (amyloid positron emission tomography is able to measure beta amyloid in the brain *in vivo*),
- volumetric magnetic resonance imaging detecting medial temporal lobe atrophy.

Tertiary prevention

The goal of tertiary prevention of AD is to rule out functional disability and, if possible, improve the quality of life:

- cognitive training,
- treatment with cholinesterase inhibitors (donepezil, rivastigmine, galantamine),
- treatment with N-methyl D-aspartate receptor antagonists (memantine).

CONCLUSION

Alzheimer's disease represents an important public health problem. Its aetiology is still unknown and for this reason, it is necessary to study all potential risk factors which may contribute to its development.

From the point of view of early diagnosis and potential application of primary or secondary prevention, genetic factors are the most important.

ABBREVIATIONS

AD, Alzheimer's disease.

ACKNOWLEDGMENT

Supported by the Czech Ministry of Health grant project no. NT11152-6/2010.

CONFLICT OF INTEREST STATEMENT

Author's conflict of interest disclosure: The authors stated that there are no conflicts of interest regarding the publication of this article.

REFERENCES

1. Lobo A, Launer LJ, Fratiglioni L, Andersen K, Di Carlo A, Breteler MM, Copeland JR, Dartigues JF, Jagger C, Martinez-Lage J, Soininen H, Hofman A. Prevalence of dementia and major subtypes in Europe: A collaborative study of population-based cohorts. *Neurology* 2000;54:54-9.

2. Plassman BL, Langa KM, Fischer GG, Heeringa SG, DR Weir, Ofstedal MB, Burke JR, Hurd MD, Potter GG, Rodgers WL, Steffens DC, Willis RJ, Wallace RB. Prevalence of dementia in the United States: the aging, demographics, and memory study. *Neuroepidemiology* 2007;29:125-32.
3. Ferri CP, Prince M, Brayne C, Brodaty H, Fratiglioni L, Ganguli M, Hall K, Hasegawa K, Hendrie H, Huang Y, Jorm A, Mathers C, Menezes PR, Rimmer E, Sczufca M. Global prevalence of dementia: a Delphi consensus study. *Lancet* 2005;366:2112-7.
4. von Strauss E, Viitanen M, De Ronchi D, Winblad B, Fratiglioni L. Aging and the occurrence of dementia: findings from a population-based cohort with a large sample of nonagenarians. *Arch Neurol* 1999;56:587-92.
5. Corrada MM, Brookmeyer R, Berlau D, Paganini-Hill A, Kawas CH. Prevalence of dementia after age 90: results from The 90+ Study. *Neurology* 2008;71:337-43.
6. Kawas C, Gray S, Brookmeyer R, Fozard J, Zonderman A. Age-specific incidence rates of Alzheimer's disease: the Baltimore Longitudinal Study of Aging. *Neurology* 2000;54:2072-7.
7. Kukull WA, Higdon R, Bowen JD, McCormick WC, Teri L, Schellenberg GD, van Belle G, Jolley L, Larson EB. Dementia and Alzheimer disease incidence: a prospective cohort study. *Arch Neurol* 2002;59:1737-46.
8. Fratiglioni L, Viitanen M, von Strauss E, Tontodonati V, Herlitz A, Winblad B. Very old women at highest risk of dementia and Alzheimer's disease: Incidence data from the Kungsholmen Project, Stockholm. *Neurology* 1997;48,132-8.
9. Copeland JRM, Davidson IA, Dewey ME, Gilmore C, Larkin BA, McWilliam C, Saunders PA, Scott, Sharma V, Sullivan C. Alzheimer's disease other dementias, depression and pseudodementia: Prevalence, Incidence and Three Year Outcome in Liverpool. *Br J Psychiatry* 1992;161,230-9.
10. Koukolík F., Jiráček R. Alzheimerova nemoc a další demence, Galén,1998.
11. Fratiglioni L, von Strauss E, Qiu CX. Epidemiology of the dementias of old age. In Dening T, Jacoby R, Oppenheimer C, Thomas A, eds. *The Oxford Textbook of Old Age Psychiatry*. 4th ed. New York, NY: Oxford University Press 2008;391-406.
12. Blennow K, de Leon MJ, Zetterberg H. Alzheimer's disease. *Lancet* 2006;368:387-403.
13. Green RC, Cupples LA, Go R, Benke KS, Edeki T, Griffith PA, Williams M, Hipps Y, Graff-Radford N, Bachman D, Farrer LA. Risk of dementia among white and African American relatives of patients with Alzheimer disease. *JAMA* 2002;287:329-36.
14. Huang W, Qiu C, von Strauss E, Winblad B, Fratiglioni L. APOE genotype, family history of dementia, and Alzheimer disease risk: a 6-year follow-up study. *Arch Neurol* 2004;61:1930-4.
15. Qiu C, Kivipelto M, Agüero-Torres H, Winblad B, Fratiglioni L. Risk and protective effects of the APOE gene towards Alzheimer's disease in the Kungsholmen project: variation by age and sex. *J Neurol Neurosurg Psychiatry* 2004;75:828-33.
16. Fratiglioni L, Wang HX. Smoking and Parkinson's and Alzheimer's disease: review of the epidemiological studies. *Behav Brain Res* 2000;113:117-20.
17. Wang HX, Fratiglioni L, Frisoni GB, Viitanen M, Winblad B. Smoking and the occurrence of Alzheimer's disease: cross-sectional and longitudinal data in a population-based study. *Am J Epidemiol* 1999;149:640-4.
18. Hill G, Connelly J, Hébert R, Lindsay J, Millar W. Neyman's bias revisited. *J Clin Epidemiol* 2003;56:293-6.
19. Tyas SL, White LR, Petrovitch H, Webster Ross G, Foley DJ, Heimovitz HK, Launer LJ. Mid-life smoking and late-life dementia: the Honolulu-Asia Aging Study. *Neurobiol Aging* 2003;24:589-96.
20. Ott A, Slooter AJ, Hofman A, van Harskamp F, Witteman JC, Van Broeckhoven C, van Duijn CM, Breteler MM. Smoking and risk of dementia and Alzheimer's disease in a population-based cohort study: the Rotterdam Study. *Lancet* 1998;351:1840-3.
21. Merchant C, Tang MX, Albert S, Manly J, Stern Y, Mayeux R. The influence of smoking on the risk of Alzheimer's disease. *Neurology* 1999;52:1408-12.
22. Aggarwal NT, Bienias JL, Bennett DA, Wilson RS, Morris MC, Schneider JA, Shah RC, Evans DA. The relation of cigarette smoking to incident Alzheimer's disease in a biracial urban community population. *Neuroepidemiology* 2006;26:140-6.
23. Anstey KJ, von Sanden C, Salim A, O'Kearney R. Smoking as a risk factor for dementia and cognitive decline: a meta-analysis of prospective studies. *Am J Epidemiol* 2007;166:367-78.
24. Peters R, Poulter R, Warner J, Beckett N, Burch L, Bulpitt C. Smoking, dementia and cognitive decline in the elderly, a systematic review. *BMC Geriatr* 2008;8:36.
25. Anttila T, Helkala EL, Viitanen M, Kåreholt I, Fratiglioni L, Winblad B, Soininen H, Tuomilehto J, Nissinen A, Kivipelto M. Alcohol drinking in middle age and subsequent risk of mild cognitive impairment and dementia in old age: a prospective population based study. *BMJ* 2004;329:539.
26. Huang W, Qiu C, Winblad B, Fratiglioni L. Alcohol consumption and incidence of dementia in a community sample aged 75 years and older. *J Clin Epidemiol* 2002;55:959-64.
27. Ruitenberg A, van Swieten JC, Witteman JC, Mehta KM, van Duijn CM, Hofman A, Breteler MM. Alcohol consumption and risk of dementia: the Rotterdam Study. *Lancet* 2002;359:281-6.
28. Ding J, Eigenbrodt ML, Mosley TH Jr, Hutchinson RG, Folsom AR, Harris TB, Nieto FJ. Alcohol intake and cerebral abnormalities on magnetic resonance imaging in a community-based population of middle-aged adults: the Atherosclerosis Risk in Communities (ARIC) study. *Stroke* 2004;35:16-21.
29. Paul CA, Au R, Fredman L, Massaro JM, Seshadri S, Decarli C, Wolf PA. Association of alcohol consumption with brain volume in the Framingham study. *Arch Neurol* 2008;65:1363-7.
30. Kivipelto M, Ngandu T, Fratiglioni L, M Viitanen, Kåreholt I, Winblad B, Helkala EL, Tuomilehto J, H Soininen, Nissinen A. Obesity and vascular risk factors at midlife and the risk of dementia and Alzheimer disease. *Arch Neurol* 2005;62:1556-60.
31. Rosengren A, Skoog I, Gustafson D, Wilhelmsen L. Body mass index, other cardiovascular risk factors, and hospitalization for dementia. *Arch Intern Med* 2005;165:321-6.
32. Whitmer RA, Gunderson EP, Barrett-Connor E, Quesenberry CP Jr, Yaffe K. Obesity in middle age and future risk of dementia: a 27 year longitudinal population based study. *BMJ* 2005;330:1360.
33. Fitzpatrick AL, Kuller LH, Lopez OL, Diehr P, O'Meara ES, Longstreth WT Jr, Luchsinger JA. Midlife and late-life obesity and the risk of dementia: cardiovascular health study. *Arch Neurol* 2009;66:336-42.
34. Buchman AS, Wilson RS, Bienias JL, Shah RC, Evans DA, Bennett DA. Change in body mass index and risk of incident Alzheimer disease. *Neurology* 2005;65:892-7.
35. Johnson DK, Wilkins CH, Morris JC. Accelerated weight loss may precede diagnosis in Alzheimer disease. *Arch Neurol* 2006;63:1312-7.
36. Dahl AK, Löppönen M, Isoaho R, Berg S, Kivelä SL. Overweight and obesity in old age are not associated with greater dementia risk. *J Am Geriatr Soc* 2008;56:2261-66.
37. Nourhashemi F, Deschamps V, Larrieu S, Letenneur L, Dartigues JF, Barberger-Gateau P. Body mass index and incidence of dementia: the PAQUID study. *Neurology* 2003;60:117-9.
38. Atti AR, Palmer K, Volpato S, Winblad B, De Ronchi D, Fratiglioni L. Late-life body mass index and dementia incidence: nine-year follow-up data from the Kungsholmen Project. *J Am Geriatr Soc* 2008;56:111-6.
39. Launer LJ, Ross GW, Petrovitch H, Masaki K, Foley D, White LR, Havlik RJ. Midlife blood pressure and dementia: the Honolulu-Asia aging study. *Neurobiol Aging* 2000;21:49-55.
40. Kivipelto M, Helkala EL, Laakso MP, Hänninen T, Hallikainen M, Alhainen K, Soininen H, Tuomilehto J, Nissinen A. Midlife vascular risk factors and Alzheimer's disease in later life: longitudinal, population based study. *BMJ* 2001;322:1447-51.
41. Qiu C, Winblad B, Fratiglioni L. The age-dependent relation of blood pressure to cognitive function and dementia. *Lancet Neurol* 2005;4:487-99.
42. Skoog I, Lernfelt B, Landahl S, Palmertz B, Andreasson LA, Nilsson L, Persson G, Odén A, Svanborg A. 15-year longitudinal study of blood pressure and dementia. *Lancet* 1996;347:1141-5.
43. Petitti DB, Crooks VC, Buckwalter JG, Chiu V. Blood pressure levels before dementia. *Arch Neurol* 2005;62:112-6.
44. Morris MC, Scherr PA, Hebert LE, Glynn RJ, Bennett DA, Evans DA. Association of incident Alzheimer disease and blood pressure measured from 13 years before to 2 years after diagnosis in a large community study. *Arch Neurol* 2001;58:1640-6.
45. Qiu C, von Strauss E, Fastbom J, Winblad B, Fratiglioni L. Low blood pressure and risk of dementia in the Kungsholmen project: a 6-year follow-up study. *Arch Neurol* 2003;60:223-8.

46. Verghese J, Lipton RB, Hall CB, Kuslansky G, Katz MJ. Low blood pressure and the risk of dementia in very old individuals. *Neurology* 2003;61:1667-72.
47. Ruitenberg A, den Heijer T, Bakker SL, van Swieten JC, Koudstaal PJ, Hofman A, Breteler MM. Cerebral hypoperfusion and clinical onset of dementia: the Rotterdam Study. *Ann Neurol* 2005;57:789-94.
48. Yasar S, Corrada M, Brookmeyer R, Kawas C. Calcium channel blockers and risk of AD: the Baltimore Longitudinal Study of Aging. *Neurobiol Aging* 2005;26:157-63.
49. Khachaturian AS, Zandi PP, Lyketsos CG, Hayden KM, Skoog I, Norton MC, Tschanz JT, Mayer LS, Welsh-Bohmer KA, Breitner JC. Antihypertensive medication use and incident Alzheimer disease: the Cache County Study. *Arch Neurol* 2006;63:686-92.
50. Peila R, White LR, Masaki K, Petrovitch H, Launer LJ. Reducing the risk of dementia: efficacy of long-term treatment of hypertension. *Stroke* 2006;37:1165-70.
51. Haag MD, Hofman A, Koudstaal PJ, Breteler MM, Stricker BH. Duration of antihypertensive drug use and risk of dementia. A prospective cohort study. *Neurology* 2009;72:1727-34.
52. Kivipelto M, Helkala EL, Laakso MP, Hänninen T, M Hallikainen, Alhainen K, Iivonen S, Mannermaa, Tuomilehto J, Nissinen, Soininen H. Apolipoprotein E ϵ 4 allele, elevated midlife total cholesterol level, and high midlife systolic blood pressure are independent risk factors for late-life Alzheimer disease. *Ann Intern Med* 2002;137:149-55.
53. Whitmer RA, Sidney S, Selby J, Johnston SC, Yaffe K. Midlife cardiovascular risk factors and risk of dementia in late life. *Neurology* 2005;64:277-81.
54. Solomon A, Kåreholt I, Ngandu T, Winblad B, Nissinen, Tuomilehto J, H Soininen, Kivipelto M. Serum cholesterol changes after midlife and late-life cognition: twenty-one-year follow-up study. *Neurology* 2007;68:751-6.
55. Jick H, Zornberg GL, Jick SS, Seshadri S, Drachman DA. Statins and the risk of dementia. *Lancet* 2000;356:1627-31.
56. Rockwood K, Kirkland S, Hogan DB, MacKnight C, Merry H, Verreault R, Wolfson C, McDowell I. Use of lipid-lowering agents, indication bias, and the risk of dementia in community-dwelling elderly people. *Arch Neurol* 2002;59:223-7.
57. Haag MD, Hofman A, Koudstaal PJ, Stricker BH, Breteler MM. Statins are associated with a reduced risk of Alzheimer disease regardless of lipophilicity: the Rotterdam Study. *J Neurol Neurosurg Psychiatry* 2009;80:13-7.
58. Rea TD, Breitner JC, Psaty BM, Fitzpatrick AL, Lopez OL, Newman AB, Hazzard WR, Zandi PP, Burke GL, Lyketsos CG, Bernick C, Kuller LH. Statin use and the risk of incident dementia: the Cardiovascular Health Study. *Arch Neurol* 2005;62:1047-51.
59. Zandi PP, Sparks DL, Khachaturian AS, Khachaturian AS, Tschanz J, Norton M, Steinberg M, Welsh-Bohmer KA, Breitner JC. Do statins reduce risk of incident dementia and Alzheimer disease? The Cache County Study. *Arch Gen Psychiatry* 2005;62:217-24.
60. Zandi PP, Anthony JC, Khachaturian AS, Stone SV, Gustafson D, Tschanz JT, Norton MC, Welsh-Bohmer KA, Breitner JC. Reduced risk of Alzheimer disease in users of antioxidant vitamin supplements: the Cache County Study. *Arch Neurol* 2004;61:82-8.
61. Barberger-Gateau P, Raffaitin C, Letenneur L, C Berr, Tzourio C, Dartigues JF, Alépérovitch A. et al. Dietary patterns and risk of dementia: the Three-City cohort study. *Neurology* 2007;69:1921-30.
62. Gray SL, Anderson ML, Crane PK, Breitner JC, McCormick W, Bowen JD, Teri L, Larson E. Antioxidant vitamin supplement use and risk of dementia or Alzheimer's disease in older adults. *J Am Geriatr Soc* 2008;56:291-5.
63. Scarmeas N, Stern Y, Mayeux R, Luchsinger JA. Mediterranean diet, Alzheimer disease, and vascular mediation. *Arch Neurol* 2006;63:1709-17.
64. Scarmeas N, Stern Y, Tang MX, Mayeux R, Luchsinger JA. Mediterranean diet and risk for Alzheimer's disease. *Ann Neurol* 2006;59:912-21.
65. Malouf M, Grimley EJ, Areosa SA. Folic acid with or without vitamin B12 for cognition and dementia. *Cochrane Database Syst Rev* 2003; CD004514.
66. Laitinen MH, Ngandu T, Rovio S, Helkala EL, Uusitalo U, Viitaniemi M, Nissinen A, Tuomilehto J, Soininen H, Kivipelto M. Fat intake at midlife and risk of dementia and Alzheimer's disease: a population-based study. *Dement Geriatr Cogn Disord* 2006;22:99-107.
67. Huang TL, Zandi PP, Tucker KL, Fitzpatrick AL, Kuller LH, Fried LP, Burke GL, Carlson MC. Benefits of fatty fish on dementia risk are stronger for those without APOE ϵ 4. *Neurology* 2005;65:1409-14.
68. Schaefer EJ, Bongard V, Beiser AS, Lamon-Fava S, Robins SJ, Au R, Tucker KL, Kyle DJ, Wilson PW, Wolf PA. Plasma phosphatidylcholine docosahexaenoic acid content and risk of dementia and Alzheimer disease: the Framingham Heart Study. *Arch Neurol* 2006;63:1545-50.
69. Arvanitakis Z, Wilson RS, Bienias JL, Evans DA, Bennett DA. Diabetes mellitus and risk of Alzheimer disease and decline in cognitive function. *Arch Neurol* 2004;61:661-6.
70. Akomolafe A, Beiser A, Meigs JB, Au R, Green RC, Farrer LA, Wolf PA, Seshadri S. Diabetes mellitus and risk of developing Alzheimer disease: results from the Framingham Study. *Arch Neurol* 2006;63:1551-5.
71. Irie F, Fitzpatrick AL, Lopez OL, et al. Enhanced risk for Alzheimer disease in persons with type 2 diabetes and APOE ϵ 4: the Cardiovascular Health Study Cognition Study. *Arch Neurol* 2008;65:89-93.
72. Biessels GJ, Staekenborg S, Brunner E, Brayne C, Scheltens P. Risk of Dementia in diabetes mellitus: a systematic review. *Lancet Neurol* 2006;5:64-74.
73. Roberts RO, Geda YE, Knopman DS, Christianson TJ, Pankratz VS, Boeve BF, Vella A, Rocca WA, Petersen RC. Association of duration and severity of diabetes mellitus with mild cognitive impairment. *Arch Neurol* 2008;65:1066-73.
74. Xu W, Qiu C, Gatz M, Pedersen NL, Johansson B, Fratiglioni L. Mid- and late-life diabetes in relation to the risk of dementia: a population-based twin study. *Diabetes* 2009;58:71-7.
75. Xu WL, Qiu CX, Winblad B, Fratiglioni L. The effect of borderline diabetes mellitus on the risk of dementia and Alzheimer disease. *Diabetes* 2007;56:211-6.
76. Korf ES, White LR, Scheltens P, Launer LJ. Brain aging in very old men with type 2 diabetes: the Honolulu-Asia Aging Study. *Diabetes Care* 2006;29:2268-74.
77. Rönnemaa E, Zethelius B, Sundelöf J, Sundström J, Degerman-Gunnarsson M, Berne C, Lannfelt L, Kilander L. Impaired insulin secretion increases the risk of Alzheimer disease. *Neurology* 2008;71:1065-71.
78. Xu WL, von Strauss E, Qiu CX, Winblad B, Fratiglioni L. Uncontrolled diabetes increases the risk of Alzheimer's disease: a population-based cohort study. *Diabetologia* 2009;52:1031-9.
79. Vanhanen M, Koivisto K, Moilanen L, Helkala EL, Hänninen T, Soininen H, Kervinen, Kesäniemi YA, Laakso M, Kuusisto J. Association of metabolic syndrome with Alzheimer disease: a population-based study. *Neurology* 2006;67:843-7.
80. Muller M, Tang MX, Schupf N, Manly JJ, Mayeux R, Luchsinger JA. Metabolic syndrome and dementia risk in a multiethnic elderly cohort. *Dement Geriatr Cogn Disord* 2007;24:185-92.
81. Vermeer SE, Prins ND, den Heijer T, Hofman A, Koudstaal PJ, Breteler MM. Silent brain infarcts and the risk of dementia and cognitive decline. *N Engl J Med* 2003;348:1215-22.
82. Honig LS, Tang MX, Albert S, R Costa, Luchsinger J, J Manly, Stern Y, Mayeux R. Stroke and the risk of Alzheimer disease. *Arch Neurol* 2003;60:1707-12.
83. Newman AB, Fitzpatrick AL, Lopez O, Jackson S, Lyketsos C, Jagust W, Ives D, Dekosky ST, Kuller LH. Dementia and Alzheimer's disease incidence in relationship to cardiovascular disease in the Cardiovascular Health Study cohort. *J Am Geriatr Soc* 2005;53:1101-7.
84. Beerli MS, Rapp M, Silverman JM, Schmeidler J, Grossman HT, Fallon JT, Purohit DP, Perl DP, Siddiqui A, Lesser G, Rosendorff C, Haroutunian V. Coronary artery disease is associated with Alzheimer disease neuropathology in APOE4 carriers. *Neurology* 2006;66:1399-1404.
85. Snowdon DA, Greiner LH, Mortimer JA, Riley KP, Greiner PA, Marksbery WR. Brain infarction and the clinical expression of Alzheimer disease: the Nun Study. *JAMA* 1997;277:813-7.
86. Esiri MM, Nagy Z, Smith MZ, Barnetson L, Smith AD. Cerebrovascular disease and threshold for dementia in the early stages of Alzheimer's disease. *Lancet* 1999;354:919-20.
87. Fratiglioni L, Paillard-Borg S, Winblad B. An active and socially integrated lifestyle in late life might protect against dementia. *Lancet Neurol* 2004;3:343-53.
88. Qiu C, Bäckman L, Winblad B, Agüero-Torres H, Fratiglioni L. The influence of education on clinically diagnosed dementia: incidence and mortality data from the Kungsholmen Project. *Arch Neurol* 2001;58:2034-9.

89. Ngandu T, von Strauss E, Helkala EL, Winblad B, Nissinen A, Tuomilehto J, Soininen H, Kivipelto M. Education and dementia: what lies behind the association? *Neurology* 2007;69:1442-50.
90. Karp A, Kåreholt I, Qiu C, Bellander T, Winblad B, Fratiglioni L. Relation of education and occupation-based socioeconomic status to incident Alzheimer's disease. *Am J Epidemiol* 2004;159:175-83.
91. Fratiglioni L, Wang HX, Ericsson K, Maytan M, Winblad B. Influence of social network on occurrence of dementia: a community-based longitudinal study. *Lancet* 2000;355:1315-9.
92. Wang HX, Karp A, Herlitz A, Crowe M, Kåreholt I, Winblad B, Fratiglioni L. Personality and lifestyle in relation to dementia incidence. *Neurology* 2009;72:253-9.
93. Wang HX, Karp A, Winblad B, Fratiglioni L. Late-life engagement in social and leisure activities is associated with a decreased risk of dementia: a longitudinal study from the Kungsholmen project. *Am J Epidemiol* 2002;155:1081-7.
94. Saczynski JS, Pfeifer LA, Masaki K, Korf ES, Laurin D, White L, Launer LJ. The effect of social engagement on incident dementia: the Honolulu-Asia Aging Study. *Am J Epidemiol* 2006;163:433-40.
95. Larson EB, Wang L, Bowen JD, McCormick WC, Teri L, Crane P, Kukull W. Exercise is associated with reduced risk for incident dementia among persons 65 years of age and older. *Ann Intern Med* 2006;144:73-81.
96. Karp A, Paillard-Borg S, Wang HX, Silverstein M, Winblad B, Fratiglioni L. Mental, physical and social components in leisure activities equally contribute to decrease dementia risk. *Dement Geriatr Cogn Disord* 2006;21:65-73.
97. Abbott RD, White LR, Ross GW, Masaki KH, Curb JD, Petrovitch H. Walking and dementia in physically capable elderly men. *JAMA* 2004;292:1447-53.
98. Rovio S, Kåreholt I, Helkala EL, Viitanen M, Winblad B, Tuomilehto J, Soininen H, Nissinen A, Kivipelto M. Leisure-time physical activity at midlife and the risk of dementia and Alzheimer's disease. *Lancet Neurol* 2005;4:705-11.
99. Crowe M, Andel R, Pedersen NL, Johansson B, Gatz M. Does participation in leisure activities lead to reduced risk of Alzheimer's disease? A prospective study of Swedish twins. *J Gerontol B Psychol Sci Soc Sci* 2003;58:249-55.
100. Andel R, Crowe M, Pedersen NL, Mortimer J, Crimmins E, Johansson B, Gatz M. Complexity of work and risk of Alzheimer's disease: a population-based study of Swedish twins. *J Gerontol B Psychol Sci Soc Sci* 2005;60:251-8.
101. Valenzuela MJ, Sachdev P, Wen W, Chen X, Brodaty H. Lifespan mental activity predicts diminished rate of hippocampal atrophy. *PLoS ONE* 2008;3:e2598.
102. Qiu CX, De Ronchi D, Fratiglioni L. The epidemiology of the dementias: an update. *Curr Opin Psychiatry* 2007;20:380-5.
103. Bäckman L, Small BJ, Fratiglioni L. Stability of the preclinical episodic memory deficit in Alzheimer's disease. *Brain* 2001;124:96-102.
104. DeKosky ST, Marek K. Looking backward to move forward: early detection of neurodegenerative disorders. *Science* 2003;302:830-4.

Alzheimer's Disease International. World Alzheimer Report 2014 Dementia and Risk Reduction. An analysis of protective and modifiable factors. The Global Observatory for Ageing and Dementia Care. The Global Observatory for Ageing and Dementia Care, hosted at the Health Service and Population Research Department, King's College London, was founded in 2013. The background concepts and methodological challenges of epidemiological studies of dementia are here briefly presented. Epidemiological concepts. Epidemiology is the discipline that studies the distribution, the causes and impact of health and ill health in populations. Alzheimer's Disease Risk Factors. Medically reviewed by Timothy J. Legg, PhD, CRNP on September 23, 2016. Written by Wendy Leonard, MPH. Alzheimer's disease is a condition that affects the way your brain functions. In the early stages, people with Alzheimer's often experience memory loss, such as: forgetting conversations. Alzheimer's disease is a progressive disease, which means it gets worse over time. People with late-stage Alzheimer's often need help with most of their everyday activities, such as eating, dressing, and bathing. Researchers still aren't sure what causes Alzheimer's disease. But certain factors increase your likelihood of developing this incurable disease. You can control some factors by making different lifestyle choices. The greatest known risk factor for Alzheimer's is increasing age, but Alzheimer's is not a normal part of aging. While age increases risk, it is not a direct cause of Alzheimer's. Most individuals with the disease are 65 and older. After age 65, the risk of Alzheimer's doubles every five years. It is estimated that less than 1 percent of Alzheimer's cases are caused by deterministic genes (genes that cause a disease, rather than increase the risk of developing a disease). Learn more: Genetics and Alzheimer's. Other risk factors you may be able to influence.