

## Livestock grazing management and biodiversity conservation in Australian temperate grassy landscapes

J. Dorrough<sup>A,F</sup>, A. Yen<sup>A,E</sup>, V. Turner<sup>A</sup>, S. G. Clark<sup>B</sup>, J. Crosthwaite<sup>C</sup>, and J. R. Hirth<sup>D</sup>

<sup>A</sup>Department of Sustainability and Environment, Arthur Rylah Institute for Environmental Research, PO Box 137, Heidelberg, Vic. 3084, Australia.

<sup>B</sup>Department of Primary Industry, Primary Industries Research Victoria—Hamilton, Private Bag 105, Hamilton, Vic. 3300, Australia.

<sup>C</sup>Department of Sustainability and Environment, 8 Nicholson Street, East Melbourne, Vic. 3002, Australia.

<sup>D</sup>Department of Primary Industry, Primary Industries Research Victoria—Rutherglen, RMB 1145, Rutherglen, Vic. 3685, Australia.

<sup>E</sup>Department of Primary Industry, Primary Industries Research Victoria—Knoxfield, Private Mail Bag 15, Ferntree Gully Delivery Centre, Knoxfield, Vic. 3156, Australia.

<sup>F</sup>Corresponding author; email: josh.dorrough@nre.vic.gov.au

*Abstract.* There is an increasing interest in the development of livestock grazing management strategies that achieve environmental sustainability and maintain or improve the long-term production capacity of commercial grazing systems. In temperate Australia, these strategies are generally focussed on reducing perennial pasture decline, soil loss, acidity, and salinity. An additional challenge facing land managers and researchers is developing grazing strategies that also maintain and enhance local and regional biodiversity. However, few studies have assessed the compatibility of management practices for maintaining long-term productivity and biodiversity conservation. We still have only a very basic understanding of the effects of different grazing strategies and pasture management on biodiversity and this is a major impediment to the development of appropriate and compatible best management practice. We argue that although there is an increasing desire to find management strategies that protect and enhance biodiversity without hindering long-term agricultural production, in many cases this may not be possible. Current knowledge suggests that compatibility is most likely to be achieved using low-input systems in low productivity (fragile) landscapes, whereas in highly productive (robust) landscapes there is less opportunity for integration of productive land-use and biodiversity conservation. There is an urgent need for improved communication and collaboration between agronomic and ecological researchers and research agencies to ensure that future programs consider sustainability in terms of biodiversity as well as pasture and livestock productivity and soil and water health.

*Additional keywords:* sustainability, ecosystem function, vegetation heterogeneity, grazing strategies, perennial grasses, agriculture.

### Introduction

In temperate Australia, conservation of native biodiversity is rarely viewed as an integral component of economically and socially viable livestock production. In this paper, biodiversity is defined as the composition of native biota and the interactions among biota and between biota and their environment. The structural and functional components of an ecosystem are also considered as part of this definition as they determine the arrangement of species and habitats in space and time. Research examining the long-term sustainability of livestock and pasture production has also often excluded consideration of biodiversity conservation. Even so it is increasingly believed that the level of internal

regulation of ecosystem function in agro-ecosystems is partly dependent on plant and animal diversity (Altieri 1999). It has been argued that biodiversity may enhance production of certain ecological services including nutrient recycling, regulation of microclimate and local hydrological processes, suppression of undesirable organisms, and detoxification of noxious chemicals (Altieri 1999).

In this paper we consider whether livestock grazing systems can be enhanced to improve biodiversity outcomes in the temperate grasslands and grassy woodlands of Australia. To achieve this we firstly examined the effects of past management strategies on biodiversity and demonstrate that the loss of native biodiversity is linked to other aspects



**Fig. 1.** The temperate climate zone of Australia estimated using the Köppen classification scheme. This approximates the distribution of temperate grassy vegetation types in southern Australia prior to European settlement (source: Bureau of Meteorology).

of landscape degradation. Consideration of the processes responsible for change may also help to identify grazing systems that can cater for biodiversity and sustainable management of soils, water, vegetation, and livestock. Following this we examine whether some landscapes are more susceptible to biodiversity loss or landscape degradation than others and thus whether strategies for maintaining biodiversity should vary within and among landscapes. We suggest a framework that places landscapes on a continuum from fragile to robust (as per Tongway and Ludwig 2002) and argue that different strategies will be required along this continuum. The final section provides a critique of specific livestock management strategies that are either currently practiced or have been recommended from research. We consider whether these strategies may cater for biodiversity conservation in grazed landscapes.

#### *The temperate grassland and grassy woodland zone of southern Australia*

The temperate zone of southern Australia covers much of Victoria, Tasmania, and the eastern portion of NSW and

includes isolated areas in southern South Australia and the south-west of Western Australia (Fig. 1). At the time of European settlement, much of this area was dominated by perennial grassland or grassy woodlands (Groves and Williams 1981; Orchard and Thompson 1999) and it was these areas that became the initial focus for livestock and associated agricultural land-use. Although other vegetation types occur in this region, they are not the subject of this paper.

Since European settlement, the extent of native-dominated grassland and grassy woodland vegetation has significantly declined (McDougall and Kirkpatrick 1994; Yates and Hobbs 1997). In some cases, more than 95% of the native vegetation has been cleared and sown to exotic pastures or crops (Table 1). In remaining native grasslands and grassy woodlands, grazing, fertilisers, and exotic plant species invasion have led to further biodiversity loss and habitat modification (Moore 1970; Whalley *et al.* 1978; McIntyre and Lavorel 1994a, 1994b; Prober and Thiele 1995; Hamilton 2001). These modifications have resulted in improved livestock production but have also had considerable effect on soil structure, nutrient cycling, water,

**Table 1. The estimated extant percentage cover of three vegetation types on private land in temperate Australia**  
Disturbed vegetation has been sown to exotic pastures or crops. Woodland is all areas not cleared of native tree cover.  
Areas of native pasture include vegetation that has been fertilised and/or invaded by exotic plant species and is not necessarily dominated by native plants

Region	%Disturbed	%Woodland	%Native pasture	Reference
Victorian Basalt Plain, Vic	>95	<1?	0.1–5	P. Quigley (unpublished data); McDougall and Kirkpatrick 1994
Ararat Region, SW Vic.	61	2	37	Dorrrough and Moxham (unpublished data)
Monaro Tablelands, NSW	31.5	22.3	46.3	Garden <i>et al.</i> 2000a
Southern Tablelands, NSW	49.7	19.4	30.9	Garden <i>et al.</i> 2000a
Central Tablelands, NSW	41.6	10.8	47.6	Garden <i>et al.</i> 2000a

pasture productivity and palatability, and thus on the long-term sustainability and productivity of the livestock industries.

The development of local and regional management strategies aimed at mitigating the environmental effects of the livestock industries, while maintaining or improving their productivity, has become an increasing focus in these landscapes. On-ground strategies and research have focussed on: (1) increasing the drought tolerance and persistence of pastures (Kemp and Culvenor 1994; Kemp and Dowling 2000); (2) reducing soil and nutrient loss (Greenwood and McKenzie 2001), acidification (Kemp and Dowling 2000), and the effects and extent of salinity (Mitchell *et al.* 2001); (3) preventing or minimising invasion by low palatability or toxic plants (Grigulis *et al.* 2001); and (4) controlling invertebrate pest outbreaks (Delfosse 1993).

A result of recent pastoral research has been an increasing recognition of the contribution of native perennial plant species to livestock production in temperate Australia (Dowling *et al.* 1993; Lodge 1994; Johnston 1996; Garden *et al.* 2000b, 2001; Kemp and Dowling 2000), which contrasts significantly with attitudes of previous decades (Donald 1970). In some regions within the temperate zone, 60–70% of land supporting livestock enterprises is unsown native pasture or woodland (Table 1) and further areas previously sown are likely to have been re-invaded by native plants (Whalley *et al.* 1978; Garden *et al.* 2001).

Another branch of research has focussed on understanding the loss of native fauna and flora (McIntyre and Lavorel 1994b; Prober and Thiele 1995; Prober 1996; Bennett and Ford 1997) and developing strategies to maintain or restore biodiversity within the pastoral landscape (McIntyre *et al.* 2002). This research has clearly demonstrated that although many native perennial grasses may be widespread in the temperate zone, other biota have significantly declined. However, most research has focussed almost entirely on vascular plants and compared sites that have been intermittently grazed (densities unknown), continuously grazed (densities estimated), and ungrazed.

Experimental research considering effects of alternative grazing strategies on native biodiversity is currently lacking.

#### **Biodiversity loss and land degradation in temperate grazing lands**

Livestock grazing management practices have resulted in considerable changes to the structure and composition of both the understorey and overstorey vegetation of temperate grassy vegetation. Three general changes to the understorey vegetation have been observed: (1) a decline in the diversity of native perennial forb species (McIntyre and Lavorel 1994a; Tremont and McIntyre 1994; Pettit *et al.* 1995), (2) a shift towards species with cool-season growth (Moore and Biddiscombe 1964; Moore 1970), and (3) an increasing dominance by annual plant species (Moore and Biddiscombe 1964; Moore 1970; Pettit *et al.* 1995). The overstorey has been modified by clearing while grazing by livestock, and other factors such as increased soil nutrient availability, altered insect–plant dynamics, and waterlogging have led to further declines in remnant woodland tree cover, via a condition known as tree die-back (Landsberg *et al.* 1990; Reid 2000). As we shall discuss below, these considerable changes in vegetation have also affected native fauna, soil microbial activity, nutrient cycling, soil hydrology, and the sustainability of grazing systems.

Prior to European settlement, perennial grass and forb species dominated the ground layer of temperate grassy ecosystems, whereas annuals were minor components (Costin 1954; Willis 1964; Moore 1970; Tremont and McIntyre 1994). Provision of watering points, supplemental feeding during droughts, and addition of fertilisers and nitrogen-fixing legumes enabled land managers to maintain herbivore densities at levels much higher than those before European settlement. In addition to increased herbivore densities, the foot pressure exerted by most introduced ungulates is greater than of native macropods (Bennett 1999). As a result of high herbivore densities and high rates of biomass removal, the rate and extent of bare ground creation is increased, particularly in late summer when soil

moisture is low. This disturbance regime has favoured the establishment of invasive annual plant species, most of which are exotic, at the expense of native perennial species (Moore 1970). This pattern is particularly pronounced in the south of the temperate zone where rainfall is primarily winter-dominant. The typical grazing management in much of temperate Australia, set-stocking in relatively large paddocks (>40 ha), also results in uneven grazing pressure, increasing herbivore selectivity and loss of palatable species (Leigh and Holgate 1978).

The shift from diverse native perennial grasslands to exotic cool-season pastures dominated by annuals underlies significant environmental changes that have negative effects on ecosystem function and livestock productivity. The loss of perennial plant cover is a major factor in increasing the susceptibility of grasslands to weed invasion (Grigulis *et al.* 2001), and continual selection of palatable plant species by grazing animals favours increased dominance by less palatable plant species (Moretto and Distel 1999). These vegetation changes gradually reduce the productivity of grasslands for grazing by livestock. Dominance by annual species increases drought susceptibility and variability of pasture production, with peaks of production in spring and feed shortages in late summer and early autumn. Where exotic annual plants have replaced diverse native perennial grasslands or woodland, water capture and use is reduced, particularly in summer and autumn months when annual species are not actively growing (Johnston 1996, 2003). Consequently, rates of soil erosion can be high and increased rates of deep drainage can lead to dryland salinisation (Johnston 2001).

The sowing of annual nitrogen-fixing legumes and addition of phosphate fertilisers is considered to have been one of the major advances in Australian agriculture because it has greatly promoted grass growth without the need for costly nitrogen fertilisers (Smith 2000). However, these practices are generally more detrimental to the persistence of native biodiversity than grazing alone (Dorrough and Ash 1999; Fairfax and Fensham 2000; Dorrough 2001; Hamilton 2001; McIntyre and Martin 2001). In addition, in pastures dominated by cool-season exotic annuals and perennials, there can be incomplete utilisation of available nitrates following summer rain, hastening acidification of soils (Helyar *et al.* 1990; Johnston 2001). Acidification has now been observed throughout temperate Australia and, although lime can ameliorate acidity, it is only effective and economical if acidification occurs within the top 0–15 cm of soil, not when it occurs at depth (Simpson 1999). Some of the sown exotic species (e.g. phalaris and lucerne) are sensitive to increasing acidity and this can result in even further annual-plant dominance (Simpson and Langford 1996).

High livestock densities can also lead to loss of litter cover and soil microtopography, degradation of surface soil structure (Yates *et al.* 2000), and reduced richness and cover

of microphytic soil crust flora (Rogers and Lange 1971). Soil crust flora play important roles in reducing erosion (Eldridge 1998), increasing water infiltration and nutrient cycling (Rogers *et al.* 1966), and may also have positive benefits for seedling establishment (Scarlett 1994). Declines in the abundance of microphytic soil crusts may also have hastened invasion by exotic annual flora, degraded invertebrate habitat, and led to general declines in ecosystem function, particularly in areas with low resource availability.

In conjunction with changes in plant species composition and richness, modifications to vegetation structure through livestock grazing has affected fauna habitat. High herbivore densities reduce litter cover and vegetation biomass and create a more even sward structure. The loss of perennial tussocks and increased bare ground may have effectively eliminated the habitat of many grassland fauna, particularly those requiring closed tussock vegetation. Some of the remaining species, such as the Earless Dragon (*Tympanocryptis pinguicolla*) (Smith 1994) and the Plains Wanderer (*Pedionomus torquatus*) (Baker-Gabb *et al.* 1990), are apparently favoured by an open tussock structure, but other species, such as the Striped Legless Lizard (*Delma impar*), are dependent on relatively closed perennial swards (Coulson 1990).

Most studies on invertebrate communities in grazing systems indicate that more intensively managed grazing systems have lower invertebrate richness and abundance (Rushton *et al.* 1989; Siepel *et al.* 1989; Dennis *et al.* 1997; Downie *et al.* 1999; Di Giulo *et al.* 2001). In particular, the abundance of most micro-, meso-, and macro-invertebrate groups, predatory invertebrates, and detritivorous and herbivorous invertebrates declines as grazing intensity increases (Hutchinson and King 1970, 1980; King and Hutchinson 1976, 1983; Abensperg-Traun *et al.* 1995, 1996). Insects such as the ptunarra brown butterfly and many species of xanthorhine geometrid moths, whose larval stages feed on native grasses, are adversely affected by grazing, fire, fertiliser application, and the sowing of exotic plant species (Neyland 1993; McQuillan 1999). Changes in the composition and richness of invertebrate communities are apparently due to many factors including declines in plant species richness, loss of tussock-forming plant species, changes in ground surface microtopography and microclimate, increased compaction of soil, and fertiliser inputs (King *et al.* 1976; Hutchinson and King 1980; King and Hutchinson 1983; Di Giulo *et al.* 2001).

Not all research has found livestock management systems to have negative effects on invertebrates. Although ungrazed woodlands and native pastures are associated with a greater diversity of invertebrate orders, some studies have found total invertebrate abundance and biomass to be greater in exotic pastures (Hutchinson and King 1980; Bromham *et al.* 1999). Ants also tend to be more abundant in heavily grazed areas (Hutchinson and King 1980), most likely because

increased bare ground, caused by grazing, favours opportunist species.

### Value of native biodiversity to livestock production systems

Despite the history of negative effects of pastoral grazing systems on Australian biodiversity, it is increasingly understood that grazing, or at least some form of biomass removal, is essential for the maintenance of plant biodiversity in most grasslands (Lunt 1991; Gilfedder and Kirkpatrick 1994; Freidel and James 1995; Fensham 1998). Herbivore grazing can create gaps and reduce litter cover and the stature of dominant plants, leading to increased rates of plant invasion, thus increasing diversity (Armesto and Pickett 1985; O'Connor 1991; Reader and Buck 1991; Lavorel *et al.* 1994; Morgan 1998a). In the absence of grazing, short-lived species dependent on establishment from seed and low-statured, shade-intolerant species decline, leading to overall loss of plant species richness (Belsky 1992; Tremont 1994) and a gradual reduction in structural complexity. These results suggest that, in areas historically managed for livestock production, opportunities for maintaining native biodiversity do exist.

It is evident that many of the same processes that lead to the loss of biodiversity also contribute to declining productivity. Although grazing can be compatible with persistence of some native species, biodiversity is likely to continue to decline in high-input grazing systems (grazing systems that include high rates of resource additions and pasture replacement) that aim to maximise production and profit. Low intensity and low-input grazing systems (utilising minimal input of resources, except those required to replace nutrients lost from the system) based on perennial native plant species can be implemented with lower costs and risks, but short-term financial benefits may be less (Crosthwaite and Malcolm 2000). High-input systems often become annual-dominated and, although annual pastures use less water and less available solar energy than perennial pastures, because little of their production goes into roots or long-term structural components, their above-ground production is generally greater. However, by shifting to a more diverse perennial system, there can be an increase in the long-term persistence and yield stability of the pasture (Tilman *et al.* 1996, 2001), with long-term production benefits (Lefroy 2001).

There is anecdotal evidence that livestock utilising native grassland have fewer health problems than those grazed on improved and fertilised pastures (Rankin 1993; Crosthwaite 1996; Crosthwaite and Macleod 2000). Low-input native grasslands are perceived as 'clean' pastures, providing stock with a diverse diet and reducing disease and intestinal worms (Crosthwaite and Macleod 2000). Observations and research suggest that pest outbreaks are most common in high-input introduced and degraded pastures (Reid 2000), whereas

small remnants of native grasslands with a tussock structure in western Victoria were found to have no exotic slugs and very few red-legged earth mites (Horne 1999). The wingless grasshopper does not thrive in perennial native grasslands and has become a pest of pastures as summer-growing native grasses have been progressively replaced by introduced, winter- and spring-growing annual grasses and legumes (Farrow and Baker 1993). Colonisation rates by dispersing winged adults are greater in overgrazed paddocks (Baker 1993) and in pastures disturbed by ploughing and sowing; overgrazing of annual pastures and droughts can lead to outbreaks.

In certain landscapes, particularly on low fertility and acidic soils, native pastures can be equally or more productive than sown exotic perennial pastures (Jones 1996; Simpson and Langford 1996). Regardless of soil fertility, the productive capacity of pasture types varies between and within particular years and, at certain times, native-dominated pastures can be as or more productive than sown pastures (Jones 1996; Simpson and Langford 1996). As a result, native plant-dominated areas are used by some graziers to fill the late summer–early autumn feed gap when active growth by exotic cool-season perennial and annual grass pastures has ceased.


Significant production value can also arise from non-grass components of the native grasslands. In terms of species richness, most vascular plant species in temperate grasslands are inter-tussock forbs (Tremont and McIntyre 1994). Although they constitute a small percentage of the actual vegetative cover, these species are often highly palatable and can form a large part of the diet of domestic livestock (Leigh and Holgate 1978; Wimbush and Costin 1979). However, many perennial inter-tussock forbs are susceptible to grazing effects, and although they may contribute considerably to livestock production when common, their value decreases dramatically as they become rare.

### Should strategies vary across the landscape? Resource availability and ecosystem resilience

The year-round availability of resources to plants is likely to play a large role in determining the resilience of ecosystems to disturbances or stress, such as that imposed by livestock grazing. On this basis we consider ecosystems to fall along a continuum from fragile (low resource availability and supply) to robust (high resource availability and supply) as per Tongway and Ludwig (2002) (Table 2). The resilience of ecosystems is likely to be lower in 'fragile' landscapes where water and/or nutrients limit plant growth and persistence (Fig. 2a). In fragile landscapes, reductions in plant cover due to grazing are more likely to result in soil degradation and extreme micro-climates, limiting the potential for plant growth (Ludwig and Tongway 1996; Yates *et al.* 2000) and re-establishment (Anderson *et al.* 1996; Ludwig and Tongway 1996; Yates *et al.* 2000). In contrast, in robust

**Table 2. Abiotic characteristics of robust and fragile landscapes**

Variable	Robust	Fragile
Soil nutrient availability	High	Low
Soil structure	Heavy	Light
Erodability	Low	High
Temporal variability in available soil moisture	Low	High
Mean monthly soil moisture availability	High	Low
Length of plant growing season	Long	Short



**Fig. 2.** Relationship between ecosystem function, stress, and biodiversity in fragile and robust landscapes. (a) As stress increases, ecosystem function declines as a result of loss in perennial grass cover, soil nutrients, and organic matter, and deterioration of soil structure. This is likely to be most rapid in fragile landscapes. (Derived from Tongway and Ludwig 2002.) (b) If the relationship between stress and native biodiversity is similar in most landscapes, then the dependency of ecosystem function on native biodiversity is likely to be greater in fragile landscapes.

landscapes, grazing-sensitive native plant species might be replaced with productive or invasive species at high grazing intensity, thus maintaining ecosystem functions (Fig. 2a).

There has been no detailed comparative work undertaken within Australia examining whether the relationship between native biodiversity and grazing differs between fragile or robust parts of the landscape. However, if the relationship is similar in both fragile and robust environments (assumed to be a non-linear negative function), in fragile landscapes the threshold beyond which there is long-term loss in production and functioning is likely to be very closely linked to native biodiversity (Fig. 2b). In robust landscapes, native biodiversity may contribute little to function or production (Fig. 2b).

These patterns suggest that alternative strategies need to be applied, depending on the fragility or robustness of environments. In those landscapes dominated by robust environments it is unlikely that successful biodiversity conservation can be achieved through direct integration with grazing systems without significant loss in productivity. In these cases, strategies solely targetted at biodiversity conservation should have greatest success. In these landscapes, many native fauna and flora are already restricted to small isolated remnants. Their persistence may be dependent upon enhancing biodiversity values in parts of the intervening landscape where the current focus is on production of agricultural products. In contrast, in less arable fragile landscapes, although grazing and associated management do lead to significant modification of vegetation, many native species persist under these regimes (Fensham 1998; Landsberg *et al.* 1999; McIntyre and Martin 2001). In this latter case, incorporating biodiversity conservation into production systems is also likely to have benefits for livestock productivity and the development of sustainable low-input systems.

Within landscapes, resource availability is both spatially and temporally patchy. Thus, within any one region, property or paddock, resilience and rates of ecosystem recovery following disturbance will vary through both space and time. The size and relationship of fragile and robust patches in a landscape will have considerable bearing on how an area could be managed to optimise productivity and biodiversity conservation. Where localised patches with high resource availability occur in a predominantly low resource landscape, these areas can be subject to intense grazing pressure in contrast to the surrounding landscape (McNaughton 1984). An alternative state can also be reached where high resource patches become dominated by large

plant species when grazing pressure is light (van de Koppel *et al.* 1996). In the first case, palatable plant species and native fauna dependent on high resource patches are likely to decline as a result of severe grazing (Morton 1990). In the second case the lack of grazing will result in the loss of shade-intolerant plants (Tremont 1994). Production outcomes are also likely to vary, with maximum utilisation in the first scenario and under-utilisation in the second.

The above observations indicate that subdivision of the landscape into manageable units based on inherent landscape productivity and physical features (environmentally defined management 'cells') should be a first priority. This is in line with current trends in grazing management based on land capability (Simpson and Langford 1996). Across the landscape the timing and intensity of grazing should be varied to provide a wide range of grazing regime defined niches for grazing-susceptible and -tolerant species to persist and disperse. Management in some cells may focus primarily on productivity, whereas in other cells it may focus on particular products, weed control outcomes, or rare species and communities. Management should also aim to account for temporal variation in climate, disturbances, plant growth rates, and habitat quality.

### Best management practices and thresholds

Factors such as loss of vegetative cover, soil erosion, weed invasion, soil structure decline, acidification, and salinity reduce the capacity of graziers to increase productivity in the long term (Crosthwaite and Malcolm 2000). As a result, there is a growing body of research that investigates how current grazing management practices can be modified to improve sustainability and long-term productive potential of grazing systems (for a review, see Kemp and Dowling 2000). Although native biodiversity may be integral for the development of low-input grazing systems and many of the processes influencing pastoral sustainability have contributed to biodiversity loss (see above), little research has been undertaken that compares the effects of different grazing strategies for maintaining or enhancing biodiversity in these ecosystems. In addition, experimental grazing trials undertaken in diverse native vegetation often fail to collect and/or present the data that would be necessary for assessment of their effects on native biodiversity. As a result, those management strategies that have been proposed tend to be based on anecdotal evidence and limited experimental results.

In this section we discuss current management strategies and more recently recommended strategies in relation to their potential effects on biodiversity and long-term livestock productivity. We consider the strategies at both the paddock and landscape scale as it is recognised that strategies that have negative effects on biodiversity, ecosystem function, or production within a paddock may be offset across a property or region.

### Grazing management

Under the typical continuous grazing systems (otherwise known as set stocking), animals are run in medium- to large-sized paddocks (>40 ha) and stock are either moved when vegetation or livestock condition begins to deteriorate or are fed conserved fodder or grain. Livestock production from continuous grazing can be high, as animals are allowed to selectively graze. However, continuous selection of the most palatable plant species can lead to their loss from the grassland and a gradual deterioration in perennial grass content and productivity. For these reasons, strict continuous grazing is no longer regarded as a sustainable grazing regime (Kemp *et al.* 2000). Continuous grazing with sheep commonly results in the transport of nutrients from the broader paddock to localised areas in sheep camps (Hilder and Mottershead 1963), increasing the need for fertiliser additions and increasing the risk of soil acidification in the broader paddock. As discussed above, it is generally accepted that continuous grazing is a major cause of native biodiversity loss. However, if grazing densities are low, in large paddocks where there is considerable spatial distribution of vegetation, soils, and topography, grazing effect can be patchy. Although this can lead to over-utilisation in some patches with subsequent bare ground and weed invasion (Johnston 2001), under-utilisation can also occur (Willms *et al.* 1988), enabling localised persistence of grazing-intolerant species (Huntly 1991). This is less likely to occur in grazing systems where grazing densities are high and pressure is even (e.g. rotational grazing systems, see below).

Under many circumstances, livestock grazing has been shown to be compatible with the conservation of a high diversity of native grassland species (McIntyre and Lavorel 1994b; Fensham 1998). Thus, it has been argued that in diverse native grasslands, if past grazing management regimes are known, then these should be continued (Diez and Foreman 1997; Foreman 1999; Milne *et al.* 1999). This approach, known as 'status quo management' (Diez and Foreman 1997), is based on the premise that past grazing management has led to the current composition of the grassy vegetation and maintaining that management strategy, if it is known, should maintain the current biota. Therefore, the underlying assumption is that the current composition of vegetation and fauna is stable and current levels of stress are not leading to irreversible changes in ecosystem function. However, concerns have been raised that such a management strategy fails to acknowledge that although many species may persist in grazed landscapes, their population growth rates may be negative (McIntyre and Lavorel 1994b; Kirkpatrick *et al.* 1995; Dorrrough and Ash 2004). Indeed there is little reason to assume that these grazed ecosystems should be stable or that under current grazing management they are not likely to shift to alternate, less diverse states as a result of interactions with climatic conditions or other

environmental factors (e.g. drought or fire). Indeed, much ecological research has emphasised the instability of grazing systems, particularly in semi-arid rangelands (van de Koppel *et al.* 1997).

Terrick Terrick National Park in the Victorian Riverina is often cited as an example of successful 'status quo' management (Foreman 1999). Continuing past grazing pressure has been argued to be essential for maintaining the diversity of native annual forbs (Foreman 1997; Milne *et al.* 1999). Despite this, exclosures established for 7 years indicated that the abundance and diversity of native annual forbs were more closely correlated to climatic patterns than to grazing (Conway 2000), and the diversity of native perennial plants increased and bare ground declined under grazing exclusion (Foreman 1997; Conway 2000). These results suggest that reductions in grazing intensity and frequency could improve landscape function and with little or no negative biodiversity effect.

Variations in continuous grazing management, based on strategic resting of pastures, could benefit biodiversity and the persistence of dominant native perennial grasses. Strategic resting can enhance flowering, growth, and survival of plant species (Lodge and Whalley 1985; FitzGerald and Lodge 1997; Ash and McIvor 1998). It can also influence invertebrate biodiversity, with potential flow-on effects through to litter and soil structure and upon some of the food web processes concerning fungi and bacteria.

Decisions regarding the timing of strategic rest from grazing can be based on: (1) livestock condition, (2) vegetation biomass, and (3) plant phenology. Approaches based on livestock condition are considered to be the least sustainable, leading to long-term adverse effects on vegetation and soil structure and function (Jones 1996; Lodge *et al.* 1998). Quantitative approaches based on perennial grass biomass are now widely recommended and some general rules have been developed (Kemp *et al.* 2000). Biomass-based strategies can provide a more flexible framework than those based on phenology and do not require knowledge about individual plant growth and reproductive cycles. However, phenologically based strategies provide more accurate predictions about changes in plant community composition over time.

The basis of the phenological approach is to identify the critical periods of a plant's lifecycle, including flowering, rapid growth, or recruitment, when rest from grazing may be beneficial. For example, it has been suggested that spring and summer rest from grazing may promote and maintain species richness within diverse grasslands by allowing spring and early summer flowering species to set seed and replenish root reserves (Diez and Foreman 1997; Milne *et al.* 1999; Eddy 2002). Similar recommendations have been made to increase the cover of exotic perennial pasture species, particularly following drought events (Dowling *et al.* 1996; FitzGerald and Lodge 1997; Lodge *et al.* 1998). Although

such a strategy should increase the flowering and seed production of native perennial plants, there has been no experimental work undertaken to determine whether this results in increased local plant diversity or higher probabilities of persistence for grazing susceptible plant species. No information is available on the benefits or otherwise for fauna, although theory would suggest that an increase in the density of flowering plants should promote beneficial invertebrates such as predators, parasitoids, and pollinators (Wratten *et al.* 1998).

Regardless of the lack of experimental research, some private landholders and government agencies are using spring resting to manage diverse native-dominated grasslands (J. Bellchambers, pers. comm.; P. Foreman, pers. comm.). Landholders utilising this strategy have remarked on an increase in flowering perennial species, but it is unknown whether this represents an increase in flowering of existing plants, an increase in species richness, or increased establishment of pre-existing species.

Spring resting could result in short-term reductions in livestock production and economic returns (Crosthwaite and Malcolm 2001). The feasibility of spring resting is also dependent on access to pastures that would not suffer from grazing pressure at this time (e.g. exotic perennial pastures). However, since perennial plant persistence and available summer and autumn forage should be increased by spring resting, it is possible that resting may have little long-term effect on the economic output of a farm, and may have positive outcomes during drought.

Spring resting could have some undesired ecological consequences. High rainfall and warm weather in spring can result in rapid plant growth, leading to rank pastures by the time the stock are introduced in summer (K. Calvert, pers. comm.). It is also possible that increased competitive effects in spring may reduce the abundance of some native plant species and, in sites with a high cover of exotic cool-season annual species, resting in spring may promote their abundance in the soil seed bank.

Gap formation in late summer and early autumn is a key precursor to invasion by exotic annuals and declines in native perennial species (Grigulis 1999; Dorrough 2001; Grigulis *et al.* 2001). Therefore, controlling grazing in late summer, during times of high soil moisture deficit, is also critical. One solution to enable utilisation of available summer forage and reduce competitive effects of grasses without enhancing weed invasion, is short-duration crash grazing (very high density of grazing animals over a short period of time) in mid-summer (T. Barlow, pers. comm.). This reduces competitive effects of dominant grasses and provides summer-growing grasses with enough time to regrow and fill large gaps prior to autumn germination of annual plants. Successful implementation of crash grazing strategies may be dependent on establishment of infrastructure similar to that necessary for cell or rotational grazing systems (see below).




Alternative grazing strategies requiring intensive management and infrastructure such as intensive rotational grazing, have been argued to provide increased benefits for biodiversity, productivity, and sustainability (Norton 1998). Rotational grazing generally describes a controlled grazing system using high stocking rates of livestock moved among many small paddocks with the intention that grazing time, and therefore grazing pressure, in any one area is minimised and rest is maximised (McCosker 2000). Rotational grazing has been shown to lead to an improvement in the utilisation of pasture, a reduction in selective grazing, and a maintenance of pasture cover (Earl and Jones 1996; Sanford *et al.* 2003). Observations of high plant species diversity in intermittently grazed reserves (McIntyre and Lavorel 1994b) have also been the basis of suggestions that rotational grazing strategies might be more beneficial for biodiversity than modified set stocking (Barlow 1998). However, grazing frequency in stock reserves is typically lower than in commercial rotational grazing systems and therefore it is unknown whether similar outcomes would be observed.

On the Monaro Tablelands of NSW, rotational grazing after 3–5 years was found to have no effect on either small- or large-scale plant species richness, although it did reduce spatial heterogeneity in vegetation structure (Bruce 1998). Spatial structural variation is important for plant species co-existence in grassland communities, and grazing-susceptible and twining or climbing plant species that co-occur with tussocks may be less likely to persist if structural heterogeneity declines. Low structural heterogeneity may also have negative consequences for fauna utilising fine-scale vegetation mosaics such as that created by patchy shrubs and tussocks and inter-tussock spaces (e.g. small- to medium-sized strictly terrestrial vertebrate fauna and terrestrial invertebrates). However, rotational grazing could increase between-paddock heterogeneity and this may favour some of the more mobile fauna species.

Other than limited research on dominant grass species, as far as we are aware no research has yet been published examining the effects of rotational grazing systems on native biodiversity in diverse temperate grasslands or grassy woodlands. Because of the potential for these grazing strategies to provide for both production and conservation goals, there is an urgent need for such research to be undertaken.

#### Vegetation structure

In Australian grasslands and rangelands, palatable perennial grasses are the main functional group driving ecosystem function (Hodgkinson 1992; Freudenberger *et al.* 1997; Kemp and Dowling 2000). These species form the vegetation matrix, structuring fauna habitat and the availability of resources in space and time for inter-tussock plant species. Retaining the dominant palatable perennial grasses should therefore be a key management goal for both sustaining




**Fig. 3.** The relationship between the richness of native plant species and the percentage of eighty-one 0.0011-m<sup>2</sup> quadrats occupied by perennial grasses in thirty-two 390-m<sup>2</sup> sites ( $R^2 = 0.46$ ,  $F_{1,31} = 26.8$ ,  $P < 0.001$ ). All sites were located within grazed grasslands on the Monaro Tablelands, NSW, and excluded areas that had been cultivated and sown to exotic perennial grass species. (J. Dorrrough, J. Ash, and S. Bruce, unpublished data.)

productivity (Kemp *et al.* 2000) and biodiversity (McIntyre *et al.* 2000) in temperate grazing lands.

Maintaining perennial grass cover and increasing standing biomass of dominant perennials can result in increased microbial biomass and reductions in runoff, groundwater discharge, and nitrate leaching (White *et al.* 2000). The implications of an increased cover of dominant perennials for native biodiversity are little known, although should this involve an increase in the cover of native perennial plant species, some positive gains are likely (Fig. 3). Alternatively, an increase in the cover of exotic perennial grasses is more likely to have negative outcomes (Morgan 1998b).

From a biodiversity perspective, it has been recommended that native perennial tussock-forming grasses should be no less than 60–70% of the ground cover (McIntyre *et al.* 2000). From a production point of view, perennial grass biomass generally needs to be maintained above 0.5–1.5 t dry matter/ha to prevent death of grasses in drought and minimise weed invasion (Kemp *et al.* 2000). Optimal production value will be obtained if the sward is even and kept at active stages of growth. In contrast, for biodiversity, the structural heterogeneity of vegetation should be maximised (Fig. 4) and tall and medium tussock-forming species should dominate the perennial grass layer. Although there may be minor production advantages of a variable vegetation structure dominated by tall tussocks (e.g. protection of lambs), diversifying the vertical structure of the pasture can result in a reduction in the utilisable forage, particularly in sheep enterprises.

Long-ungrazed vegetation has low vegetation heterogeneity, resulting in low habitat value and high competition for space and light for subordinate forb species (see above). Light to moderate grazing intensity reduces total



**Fig. 4.** Hypothesised relationship between vegetation biomass (dashed line), structural heterogeneity (solid line), and grazing pressure under a set-stocked grazing regime. Optimal conditions for biodiversity are likely to occur when biomass and heterogeneity are high, and in contrast, optimal livestock production is likely to be achieved when heterogeneity is low and biomass is moderate.

biomass, but increases structural heterogeneity (Fig. 4). With further increases in grazing pressure, vegetation biomass declines and heterogeneity reaches a peak and then begins to decline. Under rotational grazing regimes, heterogeneity may decline at lower grazing pressure. Fig. 4 indicates that production may need to be compromised if total biomass and vegetation structural heterogeneity are optimised. Research is required to parameterise these relationships.

#### Fertilisers

Sustainability in agroecosystems implies that there is limited use of non-renewable resources. Even so, the addition of non-renewable fertilisers is an important strategy in increasing native and exotic pasture production and replacing nutrients lost through leaching, stock movement, and the export of commodities (Cayley *et al.* 1987, 1999). At present, little is known about the effects of interactions between nutrient additions and grazing for native biodiversity, although negative effects are expected. In Australia, increased soil fertility in conjunction with disturbance or grazing can lead to high rates of invasion by annual plant species and declines in dominance by native perennial grasses (Trumble and Fraser 1932; Hobbs and Atkins 1988; Hobbs and Huenneke 1992; Robinson *et al.* 1993). Similar patterns have been observed along natural fertility gradients (Bridgewater and Backshall 1981; McIntyre and Lavorel 1994a). Fertiliser addition can also facilitate higher livestock grazing densities (Cayley *et al.* 1999), with subsequent negative effects on the persistence of grazing-susceptible species. Increased fertiliser can also decrease the number of soil mesofauna, ants and spiders

(Siepel *et al.* 1989; Di Giulo *et al.* 2001). The threatened Golden Sun Moth (*Synemon plana*) inhabits grasslands in south-eastern Australia that have a cover greater than 40% of *Austrodanthonia* spp. and low levels of phosphorus (O'Dwyer and Attiwill 1999). The addition of fertiliser to these grasslands increases weed cover, especially the introduced perennial ryegrass (*Lolium perenne*), and reduces the cover of *Austrodanthonia* spp., rendering the sites unsuitable for *S. plana* (O'Dwyer and Attiwill 1999).

Increased nutrient loss through surface run-off and deep drainage can also occur as a result of fertiliser application (Ridley *et al.* 2003). This can have effects on in-stream biodiversity (ANZECC 2000) and alter plant community composition within adjacent remnant vegetation (Hobbs and Huenneke 1992). Ridley *et al.* (2003) state that 'planting wide buffer strips to protect waterways is an essential component of more environmentally sustainable farming systems'. Similar consideration should also be given to isolated remnant vegetation adjoining fertilised pastures.

#### Grassland replacement

Replacement of native grasslands with a mix of exotic grasses and legumes in conjunction with high rates of fertiliser is a recommended practice for improving animal production in much of temperate Australia (Malcom *et al.* 1996; Cayley *et al.* 1999). However, in many cases, sown pastures have degraded over time, becoming increasingly dominated by annual grasses and broad-leafed weeds (Kemp and Dowling 1991; Garden *et al.* 2001). Pasture replacement strategies also have negative effects on the persistence of many species of native fauna and flora (Dorrough and Ash 1999; Fairfax and Fensham 2000; Dorrough 2001; Hamilton 2001; McIntyre and Martin 2001). Nitrogen fixation by introduced legumes is also instrumental in increasing annual grass and broad-leafed weed biomass and soil acidification in pasture. Despite this, annual legumes are still considered to play an essential role in sustainable grazing systems (Kemp *et al.* 2000), although perhaps this role should be examined.

There has been recent interest in the development of native grass pasture species for use on less fertile, saline, or acidic soils (Johnston *et al.* 2001; Mitchell *et al.* 2001). However, any replacement strategy, whether it uses native or exotic grass species, is likely to lead to biodiversity and sustainability effects (Jones 1996). A better approach is to develop grazing strategies that maximise the benefit of retaining and recruiting persistent native species.

It has been argued that intensively utilised improved pastures (i.e. high input, exotic species) can play a role in conserving biodiversity by off-setting production losses from less intensively utilised areas of a property (Crosthwaite and Malcolm 2000). In addition, run-down introduced pastures have both low biodiversity and production values (Crosthwaite and Malcolm 2000) and may

have additional adverse environmental effects (e.g. nitrate leaching, weeds, and soil loss) that can be reduced through appropriate intensification (Cayley *et al.* 1999; White *et al.* 2000). Maximising production in certain areas of a property can be further justified by considering the cost of restoring biodiversity in fertile productive areas compared with enhancing or maintaining biodiversity in areas with lower fertility and/or a less intense history of land use.

### *Fire*

Prior to European settlement, fire is thought to have been a major factor structuring grassland and grassy woodland communities (Lunt 1997; Lunt and Morgan 2002). Many of the more significant species-rich grassland remnants in south-eastern Australia have a history of regular burning and little or no livestock grazing (Stuwe and Parsons 1977). Fire plays an important role in maintaining the tussock and inter-tussock structure in native grasslands, enabling co-existence of a diversity of small native forb species (Lunt 1991; Morgan 1998a; Morgan and Lunt 1999). Since European settlement, fires have generally been suppressed, although burning has been used in conjunction with grazing to reduce dominance by large tussocks, particularly in drainage lines. The effect of fire on invertebrates is dependent upon timing, intensity, and extent of the fire, and on the length of the vegetation recovery period. The recovery response depends upon feeding habit (herbivore, predator, detritivore, fungivore), distance to source and mode of dispersal, and whether they survive the fire by going underground (Warren *et al.* 1987). The activity of some invertebrates, such as ants, increases dramatically after fire (Andersen and Yen 1985).

Because many species-rich grasslands have had a long history of regular burning, fire has often been used in grazed grasslands resumed for conservation. However, in sites with a long history of grazing, imposing a management regime of regular burning can quickly result in invasion by exotic plant species (Lunt 1990; Lunt and Morgan 1999). Contrary to this, recent trials in the Victorian Riverina have shown that a combination of fire and grazing can be used successfully to enhance native plant abundance and reduce weed cover in previously cropped and grazed grasslands (I. Lunt, pers. comm.). Further research is required to investigate the role that fire may play in maintaining biodiversity in grazed grasslands and grassy woodlands. If beneficial effects can be shown, trials are then required to determine how fire could be used within a commercial grazing enterprise without detrimental effects on production.

### *Invertebrate pest control*

Invertebrate pests consume foliage, roots, and sap, transmit plant pathogens, and predispose plants to pathogens. This leads to loss in pasture quality, which affects animal production. There is a lack of information to evaluate pest

effects on productivity because of a lack of records on pest occurrence (time and space), density of pests, level of damage, and a lack of methods to relate losses in pasture quality and quantity to livestock production (Madin 1993).

Invertebrate pests can be native species that have responded to better quality food associated with managed pastures (e.g. armyworms, cutworms, wingless grasshopper, redheaded pasture cockchafer, blackheaded pasture cockchafer, and the black field cricket) or can be introduced species without their natural enemies (aphids, sitona weevil, red legged earthmite, lucerne flea, and white snails) (Cullen 1993; Panetta *et al.* 1993). Different species are favoured by different factors: height of pasture, botanical composition, occurrence of bare patches of ground, amount of plant cover, proximity of alternative food plants, and habitat such as trees (Panetta *et al.* 1993).

Chemical applications are commonly used to control pests, but farmers are increasingly preferring to minimise or avoid the use of insecticides, and allow control by natural enemies or through manipulation of grazing regimes, pasture composition, and cropping frequency (Panetta *et al.* 1993). The effects of such management activities upon pest populations and non-target organisms are still largely unknown (Panetta *et al.* 1993). Chemical insecticides can be expensive, cause adverse damage to non-target animals, and may leave chemical residues in the soil and produce. The need to apply insecticides has to be ascertained first, and any application has to be properly timed. Some farmers apply fertilisers to compensate for pasture losses to pests. Although this may be more cost-effective than applying insecticides, this strategy may exacerbate existing problems or promote other insect pests (McQuillan 1993).

Tillage is also sometimes used to reduce numbers of soil insect pests, but zero tillage can increase populations of detritivorous and predaceous invertebrates (Robertson *et al.* 1994; Horne and Edward 1998) and generally does not lead to increased numbers of soil insect pests (Wilson-Rummenie *et al.* 1999). As a result, better long-term pest management outcomes might be achieved in zero-tillage systems.

### *Landscape-scale thresholds*

The direct effects of grazing and other land management practices clearly play an important role in the persistence of native biodiversity within temperate grasslands and grassy woodlands (see above). Landscape processes, primarily the extent and spatial pattern of remnant and modified vegetation, also influence patterns of native biodiversity, ecosystem processes, and pasture production within agricultural landscapes (Loyn 1987; Young *et al.* 1996; Davies and Margules 1998; Walpole 1999; McAlpine *et al.* 2002). Thus, it is necessary for grazing management to not just consider the requirements of biodiversity within individual paddocks, but also the context of the paddocks within the property or the region.

Theory indicates that the effect of landscape-scale habitat loss on species persistence is non-linear (Koopowitz *et al.* 1994). Recent research has focussed on attempting to quantify these theoretical models to determine whether there are clear landscape-scale thresholds of vegetation cover below which there is an irreversible and disproportionate loss of biodiversity (Bennett and Ford 1997). Results of a study on birds in the Northern Plains region of Victoria, an intensive agricultural region, showed that below 10% tree cover, bird species richness dramatically declined (Bennett and Ford 1997). Many other species appear to have also declined in abundance, suggesting that future extinctions may occur even in the absence of further clearing (Loyn 1987). In some areas that were once heavily wooded, tree cover is now well below 10%, emphasising that revegetation and protection of existing vegetation will be essential for the persistence of many species (Bennett and Ford 1997).

Based on both empirical and theoretical research on habitat clearance, it has been recommended that native vegetation cover should be retained over 50% of the landscape (McAlpine *et al.* 2002), with lower targets (approx. 30%) suggested at the property scale (Walpole 1999; McIntyre *et al.* 2000). Below these thresholds there is likely to be significant loss of biodiversity, disruption of ecosystem processes, and loss of pasture production. In most landscapes in south-eastern Australia, native grassland and grassy woodland vegetation cover has already been reduced well below these thresholds (Table 1) and broad-scale restoration of native vegetation appears necessary. The most likely focus for restoration will be pastures where native grasses still persist but other components of their natural system have been lost through grazing, nutrient application, and disturbance. However, there has been no attempt to restore these vegetation types at broad-scales and success at small-scales has been poor (Morgan 1999). Research is required to determine: (1) the composition, extent, and spatial arrangement of restored native vegetation desirable for broad-scale nature conservation objectives; (2) the technical aspects of broad-scale grassy ecosystem reconstruction; (3) the economic costs of broad-scale revegetation at farm and regional scales.

## Conclusions

It has been argued that the conservation of biodiversity in agricultural regions is hampered by mismatches between the scales of biological processes and units of management as well as regional and local policy and planning structures (Briggs 2001). This review has shown that insufficient knowledge of key ecological processes is also a major factor in preventing the integration of biodiversity conservation with agriculture. At present there are few empirical data examining the effect of different grazing strategies (i.e. season of rest, rotational systems) on biodiversity in remnant grassy vegetation in south-eastern Australia. The limited

information indicates that year-round set-stock grazing, fertiliser application, and pasture replacement have negative effects on the persistence of native grassland species. Intermittent grazing strategies are more likely to have long-term biodiversity and production benefits, particularly if they can be manipulated such that vegetation heterogeneity can be maintained through space and time.

Not only do we lack essential information on management effects for biodiversity but, in many cases, there is a lack of knowledge of biodiversity itself. Current knowledge of biodiversity in grasslands and grassy woodlands of south-eastern Australia is primarily restricted to vascular plants, threatened species of vertebrates, and pest invertebrates, although this too is limited. Much of our current knowledge is primarily descriptive. There is a dearth of knowledge regarding certain key elements of biodiversity such as invertebrates (Yen 1999) and non-vascular plants, their ecological functions, and the interactions between all these elements. A certain level of understanding of the roles of these biodiversity elements is necessary before we can determine whether proposed best management practices will lead to long-term sustainability.

Increasing the cover and abundance of perennial grasses is one of the most important strategies for ensuring sustainability. The majority of research in temperate Australia has only focussed on the role of dominant perennial grasses; however, grazing strategies to favour these species may also have benefits for other aspects of biodiversity, particularly if the dominant grasses are native. Conflict is most likely to arise with regard to vegetation structure. Cover and structure optimal for the persistence of perennial native grasses and maintenance of maximum livestock production will not always coincide with optimal co-existence of native fauna and flora.

Considerable collaborative research, extension, and policy development are required before a sustainable and biodiverse grazing system can be achieved in Australian temperate grassy ecosystems. It is essential that agronomists and ecologists develop joint projects to tackle many of the issues raised in this paper. In particular, effort is required to identify the links between long-term sustainability in production and biodiversity. It will also be necessary to re-train extension staff to ensure that they are aware of both biodiversity conservation and sustainable grazing management strategies. Government, industry, and universities should actively foster joint research, development, and application and this may require new approaches to both the structure and funding allocation of our agricultural research institutions.

## Acknowledgments

The authors thank Jann Williams, Matt White, and Russell Fairfax for their extremely helpful comments on drafts of this manuscript. Earlier discussions with Sue McIntyre, Julian

Ash, Tim Barlow, Ian Lunt, Denys Garden, and David Kemp were instrumental. The manuscript was considerably improved as a result of thorough comments made by two anonymous referees. Preparation of the manuscript was financially supported by the Ecologically Sustainable Agriculture Initiative of the Victorian Government.

## References

- Abensperg-Traun M, Smith GT, Arnold GW, Steven DE (1996) The effects of habitat fragmentation and livestock grazing on animal communities in remnants of gimlet *Eucalyptus salubris* woodland in the Western Australian wheatbelt. I. Arthropods. *Journal of Applied Ecology* **33**, 1281–1301.
- Abensperg-Traun M, Smith GT, Steven DE, Atkins L (1995) Nature conservation in the Western Australian wheatbelt. *Western Australian Journal of Agriculture* **36**, 88–93.
- Altieri MA (1999) The ecological role of biodiversity in agroecosystems. *Agriculture Ecosystems and Environment* **74**, 19–31. doi:10.1016/S0167-8809(99)00028-6
- Andersen AN, Yen AL (1985) Immediate effects of fire on ants in the semi-arid region of northwestern Victoria. *Australian Journal of Ecology* **10**, 25–30.
- Anderson VJ, Hodgkinson KC, Grice AC (1996) The influence of recent grazing pressure and landscape position on grass recruitment in a semi-arid woodland of eastern Australia. *Rangeland Journal* **18**, 3–9.
- ANZECC (2000) 'Australian and New Zealand guidelines for fresh and marine water quality. Vol. 2. Aquatic ecosystems.' (Australian and New Zealand Environment and Conservation Council, Agriculture and Resource Management Council of Australia and New Zealand: Canberra)
- Armesto JJ, Pickett STA (1985) Experiments on disturbance in old-field plant communities: impact on species richness and abundance. *Ecology* **66**, 230–240.
- Ash AJ, McIvor JG (1998) How season of grazing and herbivore selectivity influence monsoon tall-grass communities of northern Australia. *Journal of Vegetation Science* **9**, 123–132.
- Baker GH (1993) Current outbreak and pest-status of wingless grasshopper, *Phaulacridium vittatum*, in south-eastern Australia. In 'Pests of pastures: weed, invertebrate and disease pests of Australian sheep pastures'. (Ed. ES Delfosse) pp. 121–127. (CSIRO Information Services: Melbourne, Vic.)
- Baker-Gabb DJ, Benshemesh J, Maher PN (1990) A revision of the distribution, status and management of the Plains-wanderer *Pedionomus torquatus*. *Emu* **90**, 161–168.
- Barlow T (1998) 'Grassy guidelines: how to manage native grasslands and grassy woodlands on your property.' (Trust for Nature: Melbourne, Vic.)
- Belsky AJ (1992) Effects of grazing, competition, disturbance and fire on species composition and diversity in grassland communities. *Journal of Vegetation Science* **3**, 187–200.
- Bennett AF, Ford LA (1997) Land use, habitat change and the conservation of birds in fragmented rural environments: a landscape perspective from the Northern Plains, Victoria, Australia. *Pacific Conservation Biology* **3**, 244–261.
- Bennett MB (1999) Foot areas, ground reaction forces and pressures beneath the feet of kangaroos, wallabies and rat-kangaroos (Marsupialia: Macropodoidea). *Journal of Zoology* **247**, 365–369. doi:10.1017/S0952836999003088
- Bridgewater PB, Backshall DJ (1981) Dynamics of some Western Australian ligneous formations with special reference to the invasion of exotic species. *Vegetatio* **46**, 141–148.
- Briggs SV (2001) Linking ecological scales and institutional frameworks for landscape rehabilitation. *Ecological Management and Restoration* **2**, 28–35. doi:10.1046/J.1442-8903.2001.00065.X
- Bromham L, Cardillo M, Bennett AF, Elgar MA (1999) Effects of stock grazing on the ground invertebrate fauna of woodland remnants. *Australian Journal of Ecology* **24**, 199–207. doi:10.1046/J.1442-9993.1999.00963.X
- Bruce S (1998) The role of resources and grazing in determining pasture community structure on the Monaro. Honours thesis, Australian National University, Canberra, ACT.
- Cayley JWD, Cumming KN, Flinn PC (1987) The annual requirement of pasture for phosphate. In 'Proceedings of the 4th Australia Agronomy Conference'. La Trobe University, Melbourne. (Ed. TG Reeves) p. 250. (The Australian Society of Agronomy: Parkville, Vic.)
- Cayley JWD, Kearney GA, Saul GR, Lescun CL (1999) The long-term influence of superphosphate and stocking rate on the production of spring-lambing Merino sheep in the high rainfall zone of southern Australia. *Australian Journal of Agricultural Research* **50**, 1179–1190.
- Conway MG (2000) The effects of grazing exclusion on a long-grazed species-rich Riverina grassland. Honours thesis, Charles Sturt University, Wagga Wagga, NSW.
- Costin AB (1954) 'A study of the ecosystems of the Monaro region of New South Wales with special reference to soil erosion.' (Soil Conservation Service of NSW: Sydney, NSW)
- Coulson G (1990) Conservation biology of the striped legless lizard (*Delma impar*). An initial investigation. Series No. 106, Arthur Rylah Institute for Environmental Research, Department of Conservation and Environment, Victoria.
- Crosthwaite J (1996) Economic benefits of native grassland on farms. Grassland Ecology Program Project No. GEP 017, Australian Nature Conservation Agency, Canberra, ACT.
- Crosthwaite J, Macleod ND (2000) Retaining native vegetation on farms: understanding its private value. In 'Nature conservation 5: conservation in production environments; managing the matrix'. (Eds JL Craig, N Mitchell, DA Saunders). (Surrey Beatty & Sons: Chipping Norton, NSW)
- Crosthwaite J, Malcolm B (2000) Looking to the farm business: approaches to managing native grassland in south-eastern Australia. Land and Water Resources Research and Development Corporation Report, Canberra, ACT.
- Crosthwaite J, Malcolm B (2001) Farm scale economics of native grasses. In 'Proceedings of the 2nd National Conference of the Native Grasses Association'. Dookie. pp. 13–23. (Stipa Native Grass Association: Gulgong, NSW)
- Cullen JM (1993) Biological control of weed, invertebrate and disease pests of Australian sheep pastures. In 'Pests of pastures: weed, invertebrate and disease pests of Australian sheep pastures'. (Ed. ES Delfosse) pp. 293–312. (CSIRO Information Services: Melbourne, Vic.)
- Davies KF, Margules CR (1998) Effects of habitat fragmentation on carabid beetles: experimental evidence. *Journal of Animal Ecology* **67**, 460–471. doi:10.1046/J.1365-2656.1998.00210.X
- Delfosse ES (Ed.) (1993) 'Pests of pastures: weed, invertebrate and disease pests of Australian sheep pastures.' (CSIRO Information Services: Melbourne, Vic.)
- Dennis P, Young MR, Howard CL, Gordon IJ (1997) The response of epigeal beetles (Col.: Carabidae, Staphylinidae) to varied grazing regimes on upland *Nardus stricta* grasslands. *Journal of Applied Ecology* **34**, 433–443.
- Diez S, Foreman P (1997) 'The management of grasslands on the riverine plain.' (Department of Natural Resources and Environment: Melbourne, Vic.)

- Di Giulio M, Edwards PJ, Meister E (2001) Enhancing insect diversity in agricultural grasslands: the roles of management and landscape structure. *Journal of Applied Ecology* **38**, 310–319. doi:10.1046/J.1365-2664.2001.00605.X
- Donald CM (1970) Temperate pasture species. In 'Australian grasslands'. pp. 303–320. (Ed. RM Moore) (Australian National University Press: Canberra, ACT)
- Dorrrough J (2001) The impact of grazing and exotic invasion on the persistence of native grassland. PhD thesis, Australian National University, Canberra, ACT.
- Dorrrough J, Ash JE (1999) Using past and present habitat to predict the current distribution and abundance of a rare cryptic lizard, *Delma impar* (Pygopodidae). *Australian Journal of Ecology* **24**, 614–624. doi:10.1046/J.1442-9993.1999.00995.X
- Dorrrough J, Ash JE (2004) The impact of livestock grazing on the persistence of a perennial forb in a temperate Australian grassland. *Pacific Conservation Biology* **9** (in press).
- Dowling PM, Garden DL, Simpson PC, Eddy D, Kemp DR (1993) Native grasses: their distribution and management in south-eastern Australia. In 'Proceedings of the XVII International Grassland Congress'. Palmerston North, NZ. pp. 327–328. (New Zealand Grassland Association, Tropical Grasslands Society of Australia, New Zealand Society of Animal Production, Australian Society of Animal Production – Queensland Branch, New Zealand Institute of Agricultural Science: Palmerston North, NZ)
- Dowling PM, Kemp DR, Michalk DL, Klein TA, Millar GD (1996) Perennial grass response to seasonal rests in naturalised pastures of central New South Wales. *Rangeland Journal* **18**, 309–326.
- Downie IS, Wilson WL, Abernethy VJ, McCracken DI, Foster GN, Ribera I, Murphy KJ, Waterhouse A (1999) The impact of different agricultural land-uses on epigeal spider diversity in Scotland. *Journal of Insect Conservation* **3**, 273–286. doi:10.1023/A:1009649222102
- Earl JM, Jones CE (1996) The need for a new approach to grazing management: is cell grazing the answer? *Rangeland Journal* **18**, 327–350.
- Eddy DA (2002) 'Managing native grassland: a guide to management for conservation, production and land protection.' (WWF Australia: Sydney, NSW)
- Eldridge DJ (1998) Trampling of microphytic crusts on calcareous soils, and its impact on erosion under rain-impacted flow. *CATENA* **33**, 221–239. doi:10.1016/S0341-8162(98)00075-7
- Fairfax RJ, Fensham RJ (2000) The effect of exotic pasture development on floristic diversity in central Queensland, Australia. *Biological Conservation* **94**, 11–21. doi:10.1016/S0006-3207(99)00169-X
- Farrow RA, Baker G (1993) The reduction of wingless grasshopper infestations by sound farm management practices. In 'Pests of pastures: weed, invertebrate and disease pests of Australian sheep pastures'. (Ed. ES Delfosse) pp. 202–205. (CSIRO Information Services: Melbourne, Vic.)
- Fensham RJ (1998) The grassy vegetation of the Darling Downs, south-eastern Queensland, Australia. Floristic and grazing effects. *Biological Conservation* **84**, 301–310. doi:10.1016/S0006-3207(97)00105-5
- FitzGerald RD, Lodge GM (1997) Grazing management of temperate pastures: literature reviews and grazing guidelines for major species. Technical Bulletin No. 47. NSW Agriculture, Orange, NSW.
- Foreman P (1997) 'A floristic classification and description of the indigenous vegetation of the Victorian Riverina.' (Department of Natural Resources and Environment: Bendigo, Vic.)
- Foreman P (1999) Grazing as a conservation management tool at Terrick Terrick National Park: a new grassland reserve in the Victorian Riverina. In 'Proceedings of a Conference on the Management of Grassy Ecosystems'. (Eds V Craigie, C Hocking) pp. 75–82. (Victoria University: St Albans, Vic.)
- Freidel MH, James CD (1995) How does grazing of native pastures affect their biodiversity. In 'Conserving biodiversity: threats and solutions'. (Eds RA Bradstock, TD Auld, DA Keith, RT Kingsford, D Lunney, DP Sivertsen) pp. 249–259. (Surrey Beatty and Sons and NSW National Parks and Wildlife Service: Sydney, NSW)
- Freudenberger D, Noble JC, Hodgkinson K (1997) Management for production and conservation goals in rangelands. In 'Landscape ecology, function and management: principles from Australia's rangelands'. (Eds JA Ludwig, DJ Tongway, D Freudenberger, JC Noble, K Hodgkinson) pp. 93–106. (CSIRO Publishing: Melbourne, Vic.)
- Garden DL, Dowling PM, Eddy DA, Nicol HI (2000a) A survey of farms on the central, southern and Monaro tablelands of NSW: management practices, farmer knowledge of native grasses, and extent of native grass areas. *Australian Journal of Experimental Agriculture* **40**, 1081–1088. doi:10.1071/EA98157
- Garden DL, Dowling PM, Eddy DA, Nicol HI (2001) The influence of climate, soil, and management on the composition of native grass pastures on the central, southern, and Monaro tablelands of New South Wales. *Australian Journal of Agricultural Research* **52**, 925–936. doi:10.1071/AR98184
- Garden DL, Lodge GM, Friend DA, Dowling PM, Orchard BA (2000b) Effects of grazing management on botanical composition of native grass-based pastures in temperate south-east Australia. *Australian Journal of Experimental Agriculture* **40**, 225–245. doi:10.1071/EA98010
- Gilfedder L, Kirkpatrick JB (1994) Climate, grazing and disturbance, and the population dynamics of *Leucochrysum albicans* at Ross, Tasmania. *Australian Journal of Botany* **42**, 417–430.
- Greenwood KL, McKenzie BM (2001) Grazing effects on soil physical properties and the consequences for pastures: a review. *Australian Journal of Experimental Agriculture* **41**, 1231–1250. doi:10.1071/EA00102
- Grigulis K (1999) The comparative population dynamics of *Echium plantagineum* L. between its native and invaded ranges. PhD thesis, Australian National University, Canberra, ACT.
- Grigulis K, Sheppard AW, Ash JE, Groves RH (2001) The comparative demography of the pasture weed *Echium plantagineum* between its native and invaded ranges. *Journal of Applied Ecology* **38**, 281–290. doi:10.1046/J.1365-2664.2001.00587.X
- Groves RH, Williams OB (1981) Natural grasslands. In 'Australian vegetation'. (Ed. RH Groves) pp. 293–316. (Cambridge University Press: Cambridge, UK)
- Hamilton SD (2001) Impacts of agricultural land use on the floristic composition and cover of a box woodland in northern Victoria. *Pacific Conservation Biology* **7**, 169–184.
- Helyar KR, Cregan PP, Godyn DL (1990) Soil acidity in New South Wales—current pH values and estimates of acidification rates. *Australian Journal of Agricultural Research* **28**, 523–537.
- Hilder EJ, Mottershead BE (1963) The redistribution of plant nutrients through free-grazing sheep. *Australian Journal of Science* **26**, 88–89.
- Hobbs RJ, Atkins L (1988) Effect of disturbance and nutrient addition on native and introduced annuals in plant communities in the Western Australian wheatbelt. *Australian Journal of Ecology* **13**, 171–179.
- Hobbs RJ, Huenneke LF (1992) Disturbance, diversity and invasion: implications for conservation. *Conservation Biology* **6**, 324–337. doi:10.1046/J.1523-1739.1992.06030324.X



- Hodgkinson K (1992) Elements of grazing strategies for perennial grass management in rangelands. In 'Desertified grasslands: their biology and management'. (Ed. GP Chapman) pp. 77–94. (The Linnean Society of London: London)
- Horne PA (1999) Pest and beneficial invertebrates. In 'Down to Grass Roots. Proceedings of a Conference on the Management of Grassy Ecosystems'. (Eds V Craigie, C Hocking) pp. 84–86. (Victoria University: St Albans, Vic.)
- Horne PA, Edward CL (1998) Effects of tillage on pest and beneficial beetles in the Wimmera region of Victoria, Australia. *Australian Journal of Entomology* **37**, 60–63.
- Huntly N (1991) Herbivores and the dynamics of communities and ecosystems. *Annual Review of Ecology and Systematics* **22**, 477–503. doi:10.1146/ANNUREV.E.S.22.110191.002401
- Hutchinson KJ, King KL (1970) Sheep numbers and soil arthropods. *Search* **1**, 41–42.
- Hutchinson KJ, King KL (1980) The effects of sheep stocking levels on invertebrate abundance, biomass and energy utilisation in a temperate, sown grassland. *Journal of Applied Ecology* **17**, 369–387.
- Johnston WH (1996) The place of C-4 grasses in temperate pastures in Australia. *New Zealand Journal of Agricultural Research* **39**, 527–540.
- Johnston WH (2001) Approaches to managing the water balance of grasslands in southeastern Australia. In 'Proceedings of the 2nd National Conference of the Native Grasses Association'. Dookie. pp. 106–115. (Stipa Native Grass Association: Gulgong, NSW)
- Johnston WH (2003) The impact of pasture development and grazing on water-yielding catchments in the Murray–Darling Basin in south-eastern Australia. *Australian Journal of Experimental Agriculture* **43**, 817–841.
- Johnston WH, Mitchell ML, Mulham WE, Waterhouse DB (2001) LIGULE: An evaluation of indigenous perennial grasses for dryland salinity management in southeastern Australia. Part 1. A base germplasm collection. *Australian Journal of Agricultural Research* **52**, 343–350. doi:10.1071/AR99140
- Jones CE (1996) Pastoral value and production from native pastures. *New Zealand Journal of Agricultural Research* **39**, 449–456.
- Kemp DR, Culvenor RA (1994) Improving the grazing and drought tolerance of temperate perennial grasses. *New Zealand Journal of Agricultural Research* **37**, 365–378.
- Kemp DR, Dowling PM (1991) Species distribution within improved pastures over central NSW in relation to rainfall and altitude. *Australian Journal of Agricultural Research* **42**, 647–659.
- Kemp DR, Dowling PM (2000) Towards sustainable perennial pastures. *Australian Journal of Experimental Agriculture* **40**, 125–132. doi:10.1071/EA98003
- Kemp DR, Michalk D, Virgona JM (2000) Towards more sustainable pastures: lessons learnt. *Australian Journal of Experimental Agriculture* **40**, 343–356. doi:10.1071/EA99001
- King KL, Hutchinson KJ (1976) The effects of sheep stocking on the abundance and distribution of mesofauna in pastures. *Journal of Applied Ecology* **13**, 41–55.
- King KL, Hutchinson KJ (1983) The effects of sheep grazing on invertebrate numbers and biomass in unfertilized natural pastures of the New England Tableland (NSW). *Australian Journal of Ecology* **8**, 245–255.
- King KL, Hutchinson KJ, Greenslade P (1976) The effects of sheep numbers on associations of Collembola in sown pastures. *Journal of Applied Ecology* **13**, 731–739.
- Kirkpatrick JB, McDougall K, Hyde M (1995) 'Australia's most threatened ecosystem: the southeastern lowland native grasslands.' (Surrey Beatty & Sons: NSW)
- Koopowitz H, Thornhill AD, Andersen M (1994) A general stochastic model for the prediction of biodiversity losses based on habitat conversion. *Conservation Biology* **8**, 425–438. doi:10.1046/J.1523-1739.1994.08020425.X
- van de Koppel J, Huisman J, van derWal R, Olff H (1996) Patterns of herbivory along a productivity gradient: an empirical and theoretical investigation. *Ecology* **77**, 736–745.
- van de Koppel J, Rieterk M, Weissing FJ (1997) Catastrophic vegetation shifts and soil degradation in terrestrial grazing systems. *Trends in Ecology, Evolution and System* **12**, 352–356. doi:10.1016/S0169-5347(97)01133-6
- Landsberg J, Lavorel S, Stol J (1999) Grazing response groups among understorey plants in arid rangelands. *Journal of Vegetation Science* **10**, 683–696.
- Landsberg J, Morse J, Khanna P (1990) Tree dieback and insect dynamics in remnants of native woodlands on farms. *Proceedings of the Ecological Society of Australia* **16**, 149–165.
- Lavorel S, Lepart J, Debussche M, Lebreton J, Beffy J (1994) Small scale disturbances and the maintenance of species diversity in Mediterranean old fields. *Oikos* **70**, 455–473.
- Lefroy EC (2001) Applying ecological principles to the re-design of agricultural landscapes. In 'Proceedings of the 10th Australian Agronomy Conference'. Hobart, Tas. (Australian Society of Agronomy) [http://www.regional.org.au/au/asa/2001/plenary/4/lefroy.htm#P2\\_74](http://www.regional.org.au/au/asa/2001/plenary/4/lefroy.htm#P2_74)
- Leigh JH, Holgate MD (1978) Effects of pasture availability on the composition of the diet selected by sheep grazing native, degenerate and improved pastures in the Upper Shoalhaven Valley, New South Wales. *Australian Journal of Experimental Agriculture* **18**, 381–390.
- Lodge GM (1994) The role and future use of perennial native grasses for pastures in Australia. *New Zealand Journal of Agricultural Research* **37**, 419–426.
- Lodge GM, Scott JM, King KL, Hutchinson KJ (1998) A review of sustainable pasture production issues in temperate native and improved pastures. *Animal Production in Australia* **22**, 79–89.
- Lodge GM, Whalley RDB (1985) The manipulation of species composition of natural pastures by grazing management on the Northern Slopes of NSW. *Australian Rangeland Journal* **7**, 6–16.
- Loyn RH (1987) Effects of patch area and habitat on bird abundances, species numbers and tree health in fragmented Victorian forests. In 'Nature conservation: the role of remnants of native vegetation'. (Eds DA Saunders, GW Arnold, AA Burbidge, AJM Hopkins) pp. 65–77. (Surrey Beatty: Sydney, NSW)
- Ludwig JA, Tongway DJ (1996) Rehabilitation of semiarid landscapes in Australia. 2. Restoring vegetation patches. *Restoration Ecology* **4**, 398–406.
- Lunt I, Morgan JW (2002) The role of fire in temperate lowland grasslands of southeastern Australia. In 'Flammable Australia: the fire regimes and biodiversity of a continent'. (Eds RA Bradstock, AM Gill, JE Williams) pp. 177–196. (Cambridge University Press: Cambridge, UK)
- Lunt ID (1990) Impact of an autumn fire on a long-grazed *Themeda triandra* (kangaroo grass) grassland: implications for management of invaded, remnant vegetation. *Victorian Naturalist* **107**, 45–51.
- Lunt ID (1991) Management of remnant lowland grasslands and grassy woodlands for nature conservation: a review. *Victorian Naturalist* **108**, 56–66.
- Lunt ID (1997) Effects of long-term vegetation management on remnant grassy forests and anthropogenic native grasslands in south-eastern Australia. *Biological Conservation* **81**, 287–297. doi:10.1016/S0006-3207(96)00159-0
- Lunt ID, Morgan JW (1999) Vegetation changes after 10 years of grazing exclusion and intermittent burning in a *Themeda triandra* (Poaceae) grassland reserve in south-eastern Australia. *Australian Journal of Botany* **47**, 537–552. doi:10.1071/BT98011

- Madin RW (1993) Weed, invertebrate and disease pests of Australian sheep pastures—an overview. In 'Pests of pastures: weed, invertebrate and disease pests of Australian sheep pastures'. (Ed. ES Delfosse) pp. 3–20. (CSIRO Information Services: Melbourne, Vic.)
- Malcom B, Sale PWG, Egan A (1996) 'Agriculture in Australia; an introduction.' (Oxford University Press: South Melbourne, Vic.)
- McAlpine CA, Fensham RJ, Temple-Smith DE (2002) Biodiversity conservation and vegetation clearing in Queensland: principles and thresholds. *Rangeland Journal* **24**, 36–55.
- McCosker T (2000) Cell grazing—the first 10 years in Australia. *Tropical Grasslands* **34**, 207–218.
- McDougall K, Kirkpatrick JB (1994) 'Conservation of lowland native grasslands in south-eastern Australia.' (World Wildlife Fund: Sydney, NSW)
- McIntyre S, Lavorel S (1994a) How environmental and disturbance factors influence species composition in temperate Australian grasslands. *Journal of Vegetation Science* **5**, 373–384.
- McIntyre S, Lavorel S (1994b) Predicting richness of native, rare, and exotic plants in response to habitat and disturbance variables across a variegated landscape. *Conservation Biology* **8**, 521–531. doi:10.1046/J.1523-1739.1994.08020521.X
- McIntyre S, Martin TG (2001) Biophysical and human influences on plant species richness in grasslands: comparing variegated landscapes in subtropical and temperate regions. *Austral Ecology* **26**, 233–245. doi:10.1046/J.1442-9993.2001.01108.X
- McIntyre S, McIvor JG, Heard KM (2002) 'Managing and conserving grassy woodlands.' (CSIRO Publishing: Melbourne, Vic.)
- McIntyre S, McIvor JG, Macleod ND (2000) Principles for sustainable grazing in eucalypt woodlands: landscape scale indicators and the search for thresholds. In 'Management for sustainable ecosystems'. (Eds P Hale, A Petrie, D Moloney, P Sattler) pp. 92–100. (The University of Queensland: Brisbane, Qld)
- McNaughton SJ (1984) Grazing lawns: animals in herds, plant form, and coevolution. *American Naturalist* **124**, 863–886. doi:10.1086/284321
- McQuillan PB (1993) Cultural control of weed, invertebrate and disease pests of Australian sheep pastures. In 'Pests of pastures: weed, invertebrate and disease pests of Australian sheep pastures'. (Ed. ES Delfosse) pp. 183–192. (CSIRO Information Services: Melbourne, Vic.)
- McQuillan PB (1999) The effect of changes in Tasmanian grasslands on the geometrid moth tribe Xanthorhoini (Geometridae: Larentiinae). In 'The other 99%. The conservation and biodiversity of invertebrates'. (Eds W Ponder, D Lunney) pp. 121–128. (Royal Zoological Society of New South Wales: Mosman, NSW)
- Milne R, Westbrooke ME, Norman L, Jackson J (1999) 'Terrick Terrick National Park: land use history and grazing management guidelines.' (Centre for Environmental Management, University of Ballarat: Ballarat, Vic.)
- Mitchell ML, Koen TB, Johnston WH, Waterhouse DB (2001) LIGULE: An evaluation of indigenous perennial grasses for dryland salinity management in south-eastern Australia 2. Field performance and the selection of promising ecotypes. *Australian Journal of Agricultural Research* **52**, 351–365. doi:10.1071/AR99141
- Moore RM (1970) South-eastern temperate woodlands and grasslands. In 'Australian grasslands'. (Ed. RM Moore) pp. 169–190. (Halstead Press: Canberra, ACT)
- Moore RM, Biddiscombe EF (1964) The effects of grazing on grasslands. In 'Grasses and grasslands'. (Ed. C Barnard) pp. 221–235. (Macmillan and Company Ltd: London)
- Moretto AS, Distel RA (1999) Effects of selective defoliation on the competitive interaction between palatable and unpalatable grasses native to a temperate semi-arid grassland of Argentina. *Journal of Arid Environments* **42**, 167–175. doi:10.1006/JARE.1999.0510
- Morgan JW (1998a) Importance of canopy gaps for recruitment of some forbs in *Themeda triandra* dominated grasslands in south-eastern Australia. *Australian Journal of Botany* **46**, 609–627.
- Morgan JW (1998b) Patterns of invasion of an urban remnant of a species-rich grassland in southeastern Australia by non-native plant species. *Journal of Vegetation Science* **9**, 181–190.
- Morgan JW (1999) Have tubestock plantings successfully established populations of rare grassland species into reintroduction sites in western Victoria? *Biological Conservation* **89**, 235–243. doi:10.1016/S0006-3207(99)00014-2
- Morgan JW, Lunt ID (1999) Effects of time-since-fire on the tussock dynamics of a dominant grass (*Themeda triandra*) in a temperate Australian grassland. *Biological Conservation* **88**, 379–386. doi:10.1016/S0006-3207(98)00112-8
- Morton SR (1990) The impact of European settlement on the vertebrate animals of arid Australia: a conceptual model. *Proceedings of the Ecological Society of Australia* **16**, 201–213.
- Neyland MG (1993) The ecology and conservation management of the ptunarra brown butterfly *Oreixenica ptunarra* (Lepidoptera; Nymphalidae; Satyrinae) in Tasmania, Australia. *Papers and Proceedings of the Royal Society of Tasmania* **127**, 43–48.
- Norton BE (1998) The application of grazing management to increase sustainable livestock production. *Proceedings of the Australian Society for Animal Production* **22**, 15–26.
- O'Connor TG (1991) Patch colonisation in a savanna grassland. *Journal of Vegetation Science* **2**, 245–254.
- O'Dwyer C, Attiwill PM (1999) A comparative study of habitats of the Golden Sun Moth *Synemon plana* Walker (Lepidoptera: Castniidae): implications for restoration. *Biological Conservation* **89**, 131–141. doi:10.1016/S0006-3207(98)00157-8
- Orchard AE, Thompson HS (Eds) (1999) 'Flora of Australia.' (ABRS/CSIRO: Melbourne, Vic.)
- Panetta FD, Ridsdill-Smith TJ, Barbetti MJ, Jones RAC (1993) The ecology of weeds, invertebrate pests and diseases of Australian sheep pastures. In 'Pests of pastures: weed, invertebrate and disease pests of Australian sheep pastures'. (Ed. ES Delfosse) pp. 87–114. (CSIRO Information Services: Melbourne, Vic.)
- Pettit NE, Freund RM, Ladd PG (1995) Grazing in remnant woodland vegetation: changes in species composition and life form groups. *Journal of Vegetation Science* **6**, 121–130.
- Prober SM (1996) Conservation of the grassy white box woodlands: rangewide floristic variation and implications for reserve design. *Australian Journal of Botany* **44**, 57–77.
- Prober SM, Thiele KR (1995) Conservation of the grassy whitebox woodlands: relative contributions of size and disturbance to floristic composition and diversity of remnants. *Australian Journal of Botany* **43**, 349–366.
- Rankin G (1993) A practical look at native pastures. In 'Proceedings of the 34th Annual Conference of the Grassland Society of Victoria'. Melbourne. (Ed. P Sale) pp. 73–80. (Grassland Society of Victoria: Parkville, Vic.)
- Reader RJ, Buck J (1991) Community response to experimental soil disturbance in a midsuccessional, abandoned pasture. *Vegetatio* **92**, 151–159.
- Reid N (2000) Maintaining tree cover health in temperate pastoral landscapes. In 'Balancing conservation and production in grassy landscapes'. (Eds T Barlow, R Thorburn) pp. 177–187. (Environment Australia: Canberra, ACT)
- Ridley AM, Christy BP, White RE, McLean T, Green R (2003) North-east Victoria SGS national experiment site: water and nutrient losses from grazing systems on contrasting soil types and levels of input. *Australian Journal of Experimental Agriculture* **43**, 799–815. doi:10.1071/EA00171



- Robertson LN, Kettle BA, Simpson GB (1994) The influence of tillage practices on soil macrofauna in a semi-arid agroecosystem in northeastern Australia. *Agriculture Ecosystems and Environment* **48**, 149–156. doi:10.1016/0167-8809(94)90085-X
- Robinson JB, Munnich DJ, Simpson PC, Orchard PW (1993) Pasture associations and their relation to environment and agronomy in the Goulburn district. *Australian Journal of Botany* **41**, 627–636.
- Rogers RW, Lange RT (1971) Lichen populations on arid soil crusts around sheep watering places in South Australia. *Oikos* **22**, 93–100.
- Rogers RW, Lange RT, Nicholas DJ (1966) Nitrogen fixation by lichens of arid zone crusts. *Nature* **209**, 96–97.
- Rushton SP, Luff ML, Eyre MD (1989) Effects of pasture improvement and management on the ground beetle and spider communities of upland grasslands. *Journal of Applied Ecology* **26**, 489–503.
- Sanford P, Cullen BR, Dowling PM, Chapman DF, Garden DL, Lodge GM, Andrew MH, Quigley PE, Murphy SR, King WMcG, Johnston WH, Kemp DR (2003) SGS pasture theme: effect of climate, soil factors and management on pasture production and stability across the high rainfall zone of southern Australia. *Australian Journal of Experimental Agriculture* **43**, 945–959.
- Scarlett N (1994) Soil crusts, germination and weeds—issues to consider. *The Victorian Naturalist* **3**, 125–130.
- Siepel H, Meijer J, Mabelis AA, Den Boer MH (1989) A tool to assess the influence of management practices on grassland surface macrofaunas. *Journal of Applied Entomology* **108**, 271–290.
- Simpson P (1999) Recognising and managing landscape and pasture diversity. In 'Getting the best from your farm: Proceeding 14th Annual Conference of the Grassland Society of NSW'. Queanbeyan, NSW. (Eds D Garden, HL Davies, D Michalk, H Dove) p. 136. (The Grassland Society of NSW: Orange, NSW)
- Simpson P, Langford C (1996) Whole-farm management of grazing systems based on native and introduced species. *New Zealand Journal of Agricultural Research* **39**, 601–609.
- Smith DF (2000) 'Natural gain: in the grazing lands of southern Australia.' (UNSW Press: Sydney, NSW)
- Smith W (1994) The ecology and taxonomy of the southern lined earless dragon (*Tympanocryptis lineata pinguicollis*). Honours thesis, Australian National University, Canberra, ACT.
- Stuwe J, Parsons RF (1977) *Themeda australis* grasslands on the Basalt Plains, Victoria: floristics and management effects. *Australian Journal of Ecology* **2**, 467–476.
- Tilman D, Reich PB, Knops JMH, Wedin D, Mielke T, Lehman C (2001) Diversity and productivity in a long-term grassland experiment. *Science* **294**, 843–845. doi:10.1126/SCIENCE.1060391
- Tilman D, Wedin D, Knops J (1996) Productivity and sustainability influenced by biodiversity in grassland ecosystems. *Nature* **379**, 718–720. doi:10.1038/379718A0
- Tongway DJ, Ludwig JA (2002) Desertification, reversing. In 'Encyclopedia of soil science'. (Ed. R Lal) pp. 343–345. (Marcel Dekker: New York)
- Tremont RM (1994) Life-history attributes of plants in grazed and ungrazed grasslands on the Northern Tablelands of New South Wales. *Australian Journal of Botany* **42**, 511–530.
- Tremont RM, McIntyre S (1994) Natural grassy vegetation and native forbs in temperate Australia: structure, dynamics and life histories. *Australian Journal of Botany* **42**, 641–658.
- Trumble HC, Fraser KM (1932) The effect of top-dressing with artificial fertilisers on the annual yield, botanical composition, and carrying capacity of a natural pasture over a period of seven years. *The Journal of Agriculture* **35**, 1342–1353.
- Walpole SC (1999) Assessment of the economic and ecological impacts of remnant vegetation on pasture productivity. *Pacific Conservation Biology* **5**, 28–35.
- Warren SD, Scifres CJ, Teel PD (1987) Response of grassland arthropods to burning: a review. *Agriculture Ecosystems and Environment* **19**, 105–130. doi:10.1016/0167-8809(87)90012-0
- Whalley RBD, Robinson GG, Taylor JA (1978) General effects of management and grazing by domestic livestock in the rangelands of the northern tablelands of NSW. *Australian Rangeland Journal* **1**, 174–190.
- White RE, Helyar KR, Ridley AM, Chen D, Heng LK, *et al.* (2000) Soil factors affecting the sustainability and productivity of perennial and annual pastures in the high rainfall zone of southeastern Australia. *Australian Journal of Experimental Agriculture* **40**, 267–283. doi:10.1071/EA98013
- Willis JH (1964) Vegetation of the basalt plains in western Victoria. *Proceedings of the Royal Society of Victoria* **77**, 397–418.
- Willms WD, Dormaar JF, Schaalje GB (1988) Stability of grazed patches on rough fescue grasslands. *Journal of Range Management* **41**, 503–508.
- Wilson-Rummenie AC, Badford BJ, Robertson LN, Simpson GB, Bell KL (1999) Reduced tillage increases population density of soil macrofauna in a semiarid environment in central Queensland. *Environmental Entomology* **28**, 163–172.
- Wimbush DJ, Costin AB (1979) Trends in vegetation at Kosciusko. I Grazing trials in the subalpine zone, 1957–1971. *Australian Journal of Botany* **27**, 741–787.
- Wratten SD, van Emden HF, Thomas MB (1998) Within-field and border refugia for the enhancement of natural enemies. In 'Enhancing biological control'. (Eds CH Pickett, RL Bugg) pp. 375–403. (University of California Press: Berkeley, CA)
- Yates CJ, Hobbs R (1997) Temperate eucalypt woodlands: a review of their status, processes threatening their persistence and techniques for restoration. *Australian Journal of Botany* **45**, 949–973.
- Yates CJ, Norton DA, Hobbs RJ (2000) Grazing effects on plant cover, soil and microclimate in fragmented woodlands in south-western Australia: implications for restoration. *Austral Ecology* **25**, 36–47. doi:10.1046/J.1442-9993.2000.01030.X
- Yen AL (1999) Grassland invertebrates of the western Victorian basalt plains: plant crunchers or forgotten lunches? In 'The great plains crash. Proceedings of a Conference on the Grasslands and Grassy Woodlands of Victoria'. Victorian Institute of Technology. (Ed. RN Jones) pp. 57–68. (Indigenous Flora and Fauna Association/Victorian National Parks Association: East Melbourne, Vic.)
- Young A, Boyle T, Brown T (1996) The population and genetic consequences of habitat fragmentation for plants. *Trends in Ecology Evolution and Systematics* **11**, 413–418. doi:10.1016/0169-5347(96)10045-8

Manuscript received 29 January 2003, accepted 15 December 2003.