

RESEARCH ARTICLE

Shade Avoidance Components and Pathways in Adult Plants Revealed by Phenotypic Profiling

Kazunari Nozue, An V. Tat, Upendra Kumar Devisetty, Matthew Robinson, Maxwell R. Mumbach, Yasunori Ichihashi, Saradadevi Lekkala, Julin N. Maloof*

Department of Plant Biology, University of California, Davis, Davis, California, United States of America

* jnmaloof@ucdavis.edu



 OPEN ACCESS

Citation: Nozue K, Tat AV, Kumar Devisetty U, Robinson M, Mumbach MR, Ichihashi Y, et al. (2015) Shade Avoidance Components and Pathways in Adult Plants Revealed by Phenotypic Profiling. *PLoS Genet* 11(4): e1004953. doi:10.1371/journal.pgen.1004953

Editor: Nathan M. Springer, University of Minnesota, United States of America

Received: April 1, 2014

Accepted: December 11, 2014

Published: April 15, 2015

Copyright: © 2015 Nozue et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All R scripts for this paper and raw data are available at <https://bitbucket.org/knozue/sasphenotyping>. RNA-seq data in this study have been deposited in the NCBI SRA (Study ID PRJNA214254) and the NCBI GEO database (accession GSE66967).

Funding: We acknowledge financial support of National Science Foundation Integrative Organismal Systems (<http://www.nsf.gov/div/index.jsp?div=IOS>; IOS-0923752) and United States Department of Agriculture USDA NIFA project (<http://www.csrees.usda.gov/>; CA-D-PLB-7226-H) to JNM. The funders

Abstract

Shade from neighboring plants limits light for photosynthesis; as a consequence, plants have a variety of strategies to avoid canopy shade and compete with their neighbors for light. Collectively the response to foliar shade is called the shade avoidance syndrome (SAS). The SAS includes elongation of a variety of organs, acceleration of flowering time, and additional physiological responses, which are seen throughout the plant life cycle. However, current mechanistic knowledge is mainly limited to shade-induced elongation of seedlings. Here we use phenotypic profiling of seedling, leaf, and flowering time traits to untangle complex SAS networks. We used over-representation analysis (ORA) of shade-responsive genes, combined with previous annotation, to logically select 59 known and candidate novel mutants for phenotyping. Our analysis reveals shared and separate pathways for each shade avoidance response. In particular, auxin pathway components were required for shade avoidance responses in hypocotyl, petiole, and flowering time, whereas jasmonic acid pathway components were only required for petiole and flowering time responses. Our phenotypic profiling allowed discovery of seventeen novel shade avoidance mutants. Our results demonstrate that logical selection of mutants increased success of phenotypic profiling to dissect complex traits and discover novel components.

Author Summary

Because plants depend on light for photosynthesis, neighboring plant shade can be detrimental to survival. Many plants sense and respond to neighbor shade to compete for light. Although shade causes responses throughout the plant (collectively known as the shade avoidance syndrome or SAS), most SAS studies have been limited to single-gene analyses in seedlings. Here we move beyond these analyses by taking a multi-gene, multi-trait study of SAS across developmental stages. Recently, whole-genome studies examining large mutant collections have been exploited to determine the pathways and their interactions that combine to determine complex phenotypes. This type of analysis (phenotypic profiling) typically uses thousands of mutants and robotic phenotyping for assaying many characters

had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

in the multitude of mutant lines. In this paper, we develop a directed alternative that allows us to take a similar approach to understanding SAS. To reduce the number of mutants required for such an approach, we used a logical selection procedure to define mutants of interest by over-representation analysis of shade-responsive genes. We found at least three different subgroups of shade responses, and that each subgroup had both shared and separate pathways. Also, we found eighteen novel genes involved in SAS. Therefore, our method is useful for multi-dimensional phenotypic profiling without expensive robots.

Introduction

Plant canopy shade limits available light for photosynthesis. Because plants are sessile, this presents a particular challenge. Perhaps as a consequence plants developed a light-quality sensory system for canopy shade; perception of foliar shade and/or reflection from neighbor plants (“neighbor detection”) can induce the shade avoidance syndrome (SAS) is collection of responses to canopy shade in plants. These SAS responses can be seen in all developmental stages from seeds to adult plants [1]. Various plant organs elongate under shade, including the hypocotyl (stem) of young seedlings, and the internodes, and leaf petioles of older plants. Furthermore shade induces upward leaf movement, accelerates flowering time (the developmental transition from vegetative phase to reproductive phase), suppresses shoot branching, and alters resource allocation [1]. All of these responses can be helpful for promoting survival when there is competition for light from neighboring plants.

Foliar shade, which has reduced photosynthetically active radiation (PAR) can be detected both by cryptochrome photoreceptors due to its reduced intensity of blue light and by phytochrome photoreceptors due to its reduced ratio of red to far-red light [1]. Remarkably, plants can perceive nearby neighbors even before true shading and the concomitant reduction in PAR. This type of neighbor detections is possible because, even though PAR is not reduced, light reflected from neighbors has a reduced ratio of red to far-red light detectable by phytochromes [1,2]. Here we use a neighbor-detection protocol to focus on phytochrome-mediated responses. Detailed analysis of shade induced hypocotyl elongation has revealed that light perception activates transcription factors (TF) that, in turn, modulate plant hormone pathways to promote organ growth. For example, the PHYTOCHROME-INTERACTING FACTOR (PIF) 5 TF protein is stabilized under shade and induces transcription of genes important for synthesis of the growth-promoting hormone auxin [3,4]. Another example is that upon shade treatment PIF7 is dephosphorylated and activated to induce *YUCCA (YUC) 2*, *YUC5*, *YUC8*, and *YUC9* auxin biosynthetic genes [5].

Plant hormones regulate many aspects of development and growth. At least five plant hormone pathways are related to SAS; auxin, brassinosteroid (BR), gibberellic acid (GA), ethylene, and cytokinin (CK). Many auxin or BR responsive genes are induced by end-of-day far-red treatment (EODFR, a proxy for shade treatment) and both auxin (*big* and *shade avoidance 3 (sav3) / tryptophan aminotransferase of Arabidopsis 1 (taa1)*) and BR (*rotundifolia3 (rot3)*) mutant showed reduced shade-induced or EODFR-induced petiole elongation as well as shade-induced gene expression [6,7]. There is some evidence that GA and CK are involved in leaf SAS [8,9,10]. Shade also influences jasmonic-acid (JA) mediated plant immune system [11] and reduced volatile JA levels [12]. However, the entire network of light signaling and hormone pathways in regulation of shade avoidance are unclear. Also, the extent of shared and separate pathways for each shade avoidance response is not currently known. In part this is

because each SAS mutant has been tested under different experimental conditions making it difficult to compare the phenotypic consequences of each mutant.

Phenotypic profiling of genetic mutants with high-throughput phenotyping is a powerful method to tease out complex gene networks [13]. Systematic phenotypic profiling of multiple traits originated with bacterial studies, followed by studies on single eukaryotic cells such as yeast and cultured animal cells [14]. Multi-dimensional phenotypic profiling of gene-perturbed multi-cellular organisms had been done both in invertebrates and vertebrates [15,16,17]. In plants phenotypic profiling of recombinant inbred lines or natural population has been conducted for QTL analysis or genome-wide association studies [18], but it has not yet been applied to induced mutants in plants.

Advances of automatic and robotic technologies made it possible to conduct high-throughput phenotyping. In plant, high-throughput robotic phenotyping systems had been reported (reviewed in [19,20]), but its use in research has only recently been published [21]. Profiling of multiple phenotypes in selected gene-perturbed plants has been reported in root epidermal cell patterning study [22] and red-light signaling [23,24], which showed effectiveness of reverse-genetics approaches to recover mutants of interest from transcriptome data.

Here we extend this approach to multiple phenotypes to develop a systems-level understanding of shade avoidance. We took advantage of a newly developed semi-automated leaf shape measurement system for throughput measurement of shade avoidance in leaves [25]. Furthermore, we aimed to narrow down candidate genes involved in SAS to be screened instead of screening of entire knockout mutant collection or mutagenized population. For this purpose, we selected candidate mutants based on over-representation analysis of shade-responsive genes in leaves. Our broad phenotypic profiling of hypocotyl, leaf, and flowering time in selected mutants has allowed us to dissect the complex shade avoidance syndrome network.

Results and Discussion

Shared and separate gene sets are enriched in shade-responsive genes in *Arabidopsis* seedling and leaf/apical region

To begin to define the genes required for SAS in leaf/apical region, after the seedling stage we performed an expression profiling experiment to find gene induces or repressed by simulated shade. Previous SAS expression profiling experiments have used microarrays and focused on seedlings or specifically on the petiole or leaf blade after EODFR treatment [6,7,26]. To obtain a broader view of expression changes in older plants, we harvested leaf and shoot apex tissue and used RNAseq (statistics are presented in [S1 Table](#)) since the most common *Arabidopsis* microarray (Affymetrix ATH1) only covers about 70% of defined genes in the transcriptome. We compared gene expression in samples treated with simulated shade (white light supplemented with far-red light to achieve R/FR of 0.5 and 80–100 $\mu\text{E PAR}$) for 1 hour or 4 hours to untreated control samples (R/FR = 1.9 and 80–100 $\mu\text{E PAR}$) and found a total of 164 and 97 genes to be differentially expressed (FDR <0.001; [S2 Table](#) see [Materials and Methods](#)). Most known shade induced genes in leaves were found in our list (bold text in [S2 Table](#), 1 hour after onset of shade treatment) [3,4,7,27,28,29]. There is a high correlation of expression fold changes by shade treatment between our data and published microarray data in leaf [6] ([S1 Fig](#)), confirming that our RNA-seq based transcriptional analysis is reliable. Significant correlation of shade-responsive genes between seedlings and leaves indicates common mechanisms exist between these two organs ([S1 Fig](#)). In addition we identified 38 (1 hour treatment) and 19 (4 hour treatment) genes not present on the *Arabidopsis* ATH1 microarray, including the known shade induced gene *PHYTOCHROME-INTERACTING-FACTOR-3 LIKE 1 (PIL1)* [3,30], while 50 (1 hour) and 68

(4 hour) genes were on ATH1 but not previously found as shade-responsive genes in EODFR treated petiole or leaf (S2 Table).

GO enrichment analysis showed that plants respond to 1 hour and 4 hour shade treatment differently (Tables 1 and 2). Two GO terms common to both time points were GO:0009733 (response to auxin stimulus) and GO:0009753 (response to JA stimulus). Known shade avoidance related genes were also enriched in the 1 hour treatment, but not in the 4 hour treatment. Plant immune related pathways (GO:0009611 (response to wounding) and GO:0009617 (response to bacterium), which are related to the JA pathway, are enriched in the 4 hour treatment.

It has been described that some plant hormone pathways, such as auxin, BR, and GA, are involved in SAS [1]. To gain a better understanding of involvement of hormone pathways in SAS, over-representation analysis (ORA) was done to test if any hormone-responsive genes were enriched among the shade-regulated genes (Table 3). Consistent with ORA with previous microarray data of leaf upon EODFR treatment, auxin, BR, and JA pathways were enriched [6]. In addition, we found that ethylene and abscisic acid (ABA) pathways were enriched in our data sets. Involvement of ethylene in SAS was suggested because shade increases ambient ethylene levels in *Sorghum* [31] and tobacco [32]. However ethylene production is required only for early response to shade in petiole and stem, but not response in leaf angle [32], although ethylene induces leaf hyponasty [33], hypocotyl elongation [34], and stem elongation [35]. Involvement of ABA in SAS is known for shade-suppressed branching [36], but has not been reported to be involved in the shade-avoidance responses examined in this paper. Interestingly leaves of four-day shade treated tomato plants have increase level of ethylene precursor and ABA [37]. In summary, the significant differences in the shade-responsive transcriptome at these two time-points reflected dynamic temporal changes of early SAS signaling cascade.

Table 1. GO category analysis of 1 hour shade-responsive genes in leaf/apical region.

category	Term	over_represented_pvalue	over_represented_padjust value
GO:0009733	response to auxin stimulus	2.70e-41	6.70e-38
GO:0010583	response to cyclopentenone	7.49e-07	6.61e-04
GO:0009753	response to jasmonic acid stimulus	8.00e-07	6.61e-04
GO:0009641	shade avoidance	3.03e-06	1.88e-03
GO:0009630	gravitropism	5.48e-06	2.72e-03
GO:0009741	response to brassinosteroid stimulus	7.92e-06	3.27e-03

Plants were grown on soil under simulated sun (R/FR = 1.9) and long-day (16/8 hours) conditions for two weeks and then were transferred to simulated shade (R/FR = 0.5) or left in the sun condition. Above-ground parts excluding hypocotyl were collected at 1 hour shade treatment in leaf. Terms with adjusted p-value by Benjamini & Hochberg method [135] <0.01 were selected from GOseq analysis.

doi:10.1371/journal.pgen.1004953.t001

Table 2. GO category analysis of 4 hour shade-responsive genes in leaf/apical region.

category	Term	over_represented_pvalue	over_represented_padjust value
GO:0009753	response to jasmonic acid stimulus	5.27e-10	1.31e-06
GO:0009611	response to wounding	2.29e-09	2.84e-06
GO:0055114	oxidation-reduction process	2.75e-07	2.27e-04
GO:0080167	response to karrikin	2.08e-06	1.10e-03
GO:0009733	response to auxin stimulus	2.21e-06	1.10e-03
GO:0009617	response to bacterium	2.92e-06	1.21e-03

Same as Table 1 except 4 hour shade treatment instead of 1 hour shade treatment

doi:10.1371/journal.pgen.1004953.t002

Table 3. Over-representation analysis of shade-responsive genes and hormone responsive genes.

regulated by hormone	1 hour shade treatment		4 hour shade treatment	
	UP	DOWN	UP	DOWN
IAA UP	4.59e-75	1	1.2e-06	0.138
IAA DOWN	1	0.0642	0.152	1
BR UP	0.000269	1	1	1
BR DOWN	1.04e-10	1	0.00355	1
GA UP	1	1	1	1
GA DOWN	0.00148	1	1	1
JA UP	0.0083	0	8.34e-18	0.0567
JA DOWN	1.9e-12	1	0.000406	1
CK UP	0.00259	1	1	1
CK DOWN	1	1	1	1
ABA UP	0.0258	0.431	1.31e-07	1
ABA DOWN	0.00141	0.252	0.0064	1
ACC UP	1.58e-05	1	0.000106	1

p-values were calculated by Goseq [129] and adjusted by Benjamini & Hochberg correction [135].

doi:10.1371/journal.pgen.1004953.t003

Next we asked if the previous seedling shade transcriptome data [7] shows the same trends as our data. We found that there were both common and specific GO terms between the hypocotyl and our leaf/apical region data sets (Table 1, 2, and 4). Of the common terms, we focused on GO:0009733 (response to auxin stimulus), GO:0009741 (response to BR stimulus), and

Table 4. GO category analysis of 1 hour shade-responsive genes in hypocotyl.

category	Term	over_represented_pvalue
GO:0009719	response to endogenous stimulus	1.72e-38
GO:0010033	response to organic substance	1.46e-32
GO:0009725	response to hormone stimulus	9.18e-32
GO:0009733	response to auxin stimulus	6.02e-28
GO:0042221	response to chemical stimulus	3.04e-24
GO:0050896	response to stimulus	1.23e-17
GO:0009741	response to brassinosteroid stimulus	5.38e-16
GO:1901700	response to oxygen-containing compound	2.03e-12
GO:0033993	response to lipid	7.38e-10
GO:0097305	response to alcohol	1.25e-09
GO:0009639	response to red or far red light	2.94e-09
GO:0014070	response to organic cyclic compound	4.45e-09
GO:0065007	biological regulation	1.62e-08
GO:0009612	response to mechanical stimulus	2.55e-08
GO:0009641	shade avoidance	5.70e-08
GO:0010200	response to chitin	2.78e-07
GO:0010243	response to organic nitrogen	4.77e-07
GO:0009416	response to light stimulus	8.17e-07
GO:0009314	response to radiation	2.87e-06
GO:0009628	response to abiotic stimulus	4.51e-06

Same as Table 1 except 1 hour shade treatment in hypocotyl instead of 1 hour shade treatment in leaf. Expression data is from [7]. Terms with p-value<1e-5 were selected from amiGO analysis.

doi:10.1371/journal.pgen.1004953.t004

GO:0009641 (shade avoidance). Among the leaf/apical region-specific terms GO:0009753 (response to JA stimulus) is of particular interest because the role of JA pathways in morphological aspects of SAS is not fully understood. Based on our ORA we chose to include mutants of genes in these categories in our phenotypic profiling (see below).

Leaf phenotype profiling of fifty-nine mutant lines

We hypothesized that shade-responsive genes and/or pathways are required for proper SAS because among the 34 causal genes in known hypocotyl SAS mutants 11 (*TAA1*, *PIN3*, *YUC2*, *YUC5*, *YUC8*, *YUC9*, *BIMI*, *GAI*, *PHYB*, *PAR1*, *HAT3*) of them have shade-responsive transcripts (S3 Table). Based on our differentially expressed gene list and previous knowledge, we chose 59 mutant lines encompassing 59 mutant genes (although some lines have more than one mutant gene) from nine categories (auxin, GA, JA, BR, light signaling, shade avoidance, flowering time, leaf size, and unknown shade responsive genes) (Fig 1). We prioritized auxin and JA pathways because both pathways were enriched in both time points of our transcriptome analysis (see above).

Petiole elongation is an important component of SAS, but most leaf phenotype measurement software does not report petiole and blade length. Phenotyping of mutants/overexpressors with a pathway of interest is a direct method to test if the given gene in the pathway is involved in phenotype of your interest. At the first step towards high throughput petiole and blade phenotyping, we developed LeafJ, an ImageJ plug-in, which is more accurate and faster measurement system than manual method [25].

To normalize petiole elongation between genotypes with different leaf size, we calculated ratio of petiole length to blade length. Statistical significance between Col under sun treatment and shade treatment effects was examined by a mixed effects model (Fig 2). Mutants that showed a statistically different response to shade when compared to the corresponding wild-type ($P < 0.05$; see Methods) were considered to have a significant SAS phenotype. In Col, we could detect significant shade-induced petiole elongation and an increase of the petiole length to leaf blade length ratio (S2F Fig and S3 Table), but not in other leaf blade parameters (length, width, and area) (S2C–S2E Fig). Some other studies have reported that leaf area does increase or decrease upon shade treatment; the differences between these studies and ours may be due to differences in plant growth conditions or because the developmental stage of our leaves might be too young to show these responses [7,38,39,40]. Comparing the kinetics of leaf development under both light condition could tell us when shade-responsive organ elongation happens and should be examined in future studies.

Screening these 59 mutant lines, we found 33 mutants that showed differences in at least one trait from the background ecotype (Fig 2, and S3 Table). These 33 mutants include genes in the auxin, jasmonic acid, and light signaling pathways. Involvement of auxin in petiole elongation upon far-red light treatment has been previously reported [5], whereas there have not been any reports of JA involvement in petiole shade avoidance. Details of these mutants will be discussed below. It is important to note that we assay a leaf series from each plant where some leaves are still expanding and others are mature. As a consequence, the mutations that we have identified could be affecting leaf development itself or developmental timing (the proportion of expanding to expanded leaves).

Shade-induced acceleration of flowering time of fifty-nine mutant lines

Compared to other SAS phenotypes, only seven mutant lines for acceleration of flowering time have been described; phytochrome mutants (*phytochromeB* (*phyB*) *phyD*, *phyB/D/E*) [41,42], known flowering time mutants (*constans* (*co*) and *gigantea* (*gi*)) [43], a circadian clock

category	subcategory	mutant name	gene name (AGI num)	source
auxin	biosynthesis	<i>yuc2/5/8/9</i>	<i>YUC2</i> (AT4G13260) <i>YUC5</i> (AT5G43890) <i>YUC8</i> (AT4G28720) <i>YUC9</i> (AT1G04180)	Yunde Zhao ¹
		<i>sav3-2/taal1</i>	<i>TAA1</i> (AT1G70560)	Joanne Chory ^{2,3}
	inactivation	<i>atr4</i>	<i>ATR4</i> (AT4G31500)	Judith Bender ⁴
	signaling	<i>iaa6/19</i>	<i>IAA6</i> (AT1G52830) <i>IAA19</i> (AT3G15540)	ABRC (CS25222)
jasmonic acid (JA)	biosynthesis	<i>aos</i>	<i>AOS</i> (AT5G42650)	Katayoon Dehesh ⁵
		<i>opr3</i> (Ws)	<i>OPR3</i> (AT2G06050)	Katayoon Dehesh ⁵
		<i>jar1</i>	<i>JAR1</i> (AT2G46370)	Daniel Kliebenstein ⁵
	inactivation	<i>st2b</i>	<i>ST2B</i> (AT5G07000)	ABRC (SALK_024717C)
	receptor	<i>coil-16</i>	<i>COI1</i> (AT2G39940)	Daniel Kliebenstein ⁵
gibberellic acid (GA)	biosynthesis	<i>ga20ox1/2</i>	<i>GA20ox1</i> (AT4G25420) <i>GA20ox2</i> (AT5G51810)	Peter Hedden ⁶
		signaling	<i>dellaQ; gai rga rgl1 rgl2</i> (Ler)	<i>GAI</i> (AT1G14920) <i>RGA</i> (AT2G01570) <i>RGL1</i> (AT1G66350) <i>RGL2</i> (AT3G03450)
	receptor	<i>phyA-211 phyB-9</i> (<i>phyA/B</i>), <i>phyB-9</i>	<i>PHYA</i> (AT1G09570) <i>PHYB</i> (AT2G18790)	Chentao Lin ¹⁰
		<i>cry1/2</i>	<i>CRY1</i> (AT4G08920) <i>CRY2</i> (AT1G04400)	
		signaling	<i>arr4</i> <i>hy5-215 hyh</i> (<i>hy5 hyh</i>) (Ws) <i>pif3-1, pif3-3</i> <i>pil5-1</i> <i>pks 1/2/4</i>	
	<i>hrb1</i> <i>lzf1, LZF1-OX</i> <i>srr1</i> <i>mida9-3, mida9-4</i>	<i>HRB1</i> (AT5G49230) <i>LZF1</i> (AT1G78600) <i>SRR1</i> (AT5G59560) <i>MIDA9</i> (AT5G02760)	ABRC (SALK_087868C) Shu-Hsing Wu ¹⁴ Christian Fankhauser ¹³ ABRC (SALK_099356C, SALK_072093C)	
shade avoidance	transcription factors	<i>pif4/5</i>	<i>PIF4</i> (AT2G43010) <i>PIF5</i> (AT3G59060) <i>ATHB2</i> (AT4G16780) <i>PIL1</i> (AT2G46970) <i>HFR1</i> (AT1G02340)	Christian Fankhauser ¹³ Giltso Choi ¹¹
		<i>athb2</i> <i>pil1</i> <i>hfr1</i> <i>PAR1</i> RNAi06/08/09 <i>par2-1</i> <i>sto</i> <i>sth2-3</i> <i>bbx18-2</i>	<i>PAR1</i> (AT2G42870) <i>PAR2</i> (AT3G58850) <i>STO</i> (AT1G06040) <i>STH2</i> (AT1G75540) <i>BBX18</i> (AT2G21320)	Jaime Martinez-Garcia ¹⁵ Jaime Martinez-Garcia ¹⁵ ABRC (SALK_067473C) ABRC (SALK_107948C) ABRC (SALK_061956C)
	others	<i>SCL13</i> as1, as2 <i>pft1</i>	<i>SCL13</i> (AT4G17230) <i>PFT1</i> (AT1G25540)	Cordelia Bolle ¹⁰ Joanne Chory ^{2,3}
shade induced genes		<i>kat1-1, kat1-2</i>	<i>KAT1</i> (AT5G46240)	ABRC (SALK_127506C, SALK_093506C) ABRC (SALK_072093C)
flowering time		<i>atlea5</i>	<i>ATLEA5</i> (AT4G02380)	ABRC (SALK_072093C)
		<i>co-9</i>	<i>CO</i> (AT5G15840)	Detlef Weigel ¹⁷
		<i>gi-2</i> <i>pft1</i>	<i>GI</i> (AT1G22770) <i>PFT1</i> (AT1G25540)	ABRC (CS3397) Joanne Chory ^{2,3}
leaf size		<i>spt-11, spt-12, SPT-OX</i> (Ler) <i>rve8</i>	<i>SPT</i> (AT4G36930) <i>RVE8</i> (AT3G09600)	Hirokazu Tsukaya ¹⁸ , David Smyth ¹⁸ Stacey Harmer ⁵

- University of California San Diego
- Howard Hughes Medical Institute
- The Salk Institute for Biological Studies
- Brown University
- University of California, Davis
- Ghent University
- Tokyo University
- Rothamsted Research
- University of Oxford
- University of California, Los Angeles
- Korea Advanced Institute of Science and Technology
- University of California, Davis
- University of Lausanne
- Academia Sinica
- Centre for Research in Agricultural Genomics
- Ludwig-Maximilians-Universität München
- Max Planck Institute for Developmental Biology
- Monash University

Fig 1. Mutants used in SAS phenotypic profiling. Shade-induced genes are shown in magenta bold and shade-repressed are shown in green. Background genotype for almost all mutants is Col (other genotypes are show in parentheses).

doi:10.1371/journal.pgen.1004953.g001




Fig 2. Phenotypic profiling of 59 mutants/overexpressors. For hypocotyl phenotype, plants were grown under continuous simulated sun conditions (R/FR = 1.3) for four days and further grown for three days under either simulated sun or simulated shade (R/FR = 0.5). For leaf phenotypes, plants were grown under long day conditions (16 hour light/8 hour dark) with approximately 90 μ E PAR (R/FR = 1.9). Two week old plants were further grown for 12 days under either simulated sun (R/FR = 1.9) or simulated shade (R/FR = 0.5). For

flowering time phenotype plants were grown in the same condition with leaf phenotyping and days after stratification at bolted was used for flowering time index. For phenotype clustering heatmap, differences between shade and sun values (except flowering time) were normalized and centered on Col (i.e., Col value = 0) and visualized with color coding (magenta indicates larger response than Col while green indicates reduced response relative to Col). For flowering time, responses to shade were normalized against flowering time under sun condition to eliminate strong dependencies of response on flowering time under sun condition (see [S2H Fig](#) for normalized data and [S2G Fig](#) for non normalized data). Asterisks (*) from hypocotyl.length to flowering.time indicate a significant difference from corresponding wild type (p-value<0.05). Although SPT_ox appeared to be a SAS mutant in flowering time (residual method, [S3H Fig](#)), it was eliminated from SAS mutants in flowering time (residual method) because its background genotype (Ler) also showed a similar shift from the regression line. Colors of mutant names correspond to groups found in [Fig 1](#). Clustering of mutants according to its phenotype is shown in the dendrogram on the right. Clustering of traits is shown on the top dendrogram. Known phenotypes for hypocotyl, petiole, or flowering time are shown in colored boxes (yellow-green for less response, blue for normal response, and magenta for exaggerated response).

doi:10.1371/journal.pgen.1004953.g002

component (*early flowering 3 (elf3)*) [[44](#)], an auxin-related mutant (*big1*) [[45](#)], and a mediator complex mutant (*phytochrome and flowering time 1 (pft1)*) [[46](#)]. Our SAS phenotypic profiling provided additional mutants for shade-accelerated flowering. In our condition shade treatment accelerated flowering time about 10% in Col, which is less effective than previously reported (35–40% [[42,43,47](#)]). This small acceleration is because our shade treatment started when plants were already close to flowering in long day conditions. Perhaps because of this we observed a strong inverse correlation between flowering time shade response and flowering time in sun ([S3H Fig](#); genotypes that flowered later in sun were more responsive to shade). Given this correlation we defined two categories flowering time shade response mutants. One category (“flowering.time” in [Fig 2](#)) consists of mutants whose log₂ shade response is different from Col-0. The second category (“flowering.time.resid” in [Fig 2](#)) are those mutants whose response is significantly different from that expected based on their flowering time in the sun (see [Methods](#)). Even though our experimental conditions were suboptimal for detecting acceleration of flowering time and we could not reproduce reduced responses to shade of *pft1*, we did find six new flowering time SAS mutants; *cryptochrome (cry) 1 cry2 (cry1/2)*, *altered-tryptophan regulation 4 (atr4)*, *reveille 8/lhy-cca1-like 5 (rve8/lcl5)*, *coronatine insensitive 1 (coi1)*, *myc2/jasmonate insensitive 1 (jin1)*, and *jasmonate-zim-domain protein 5 (jaz5)* ([Fig 2](#), and [S3 Table](#)). Details of these mutants will be discussed below.

Separate and overlapping pathways for three different shade avoidance responses

To investigate how many of the genes under study are required for normal shade avoidance response in both the leaf and the hypocotyl, we assayed hypocotyl SAS in the same 59 mutant panel ([S3A Fig](#), [Fig 1](#), and [S3 Table](#)). Among the eighteen previously reported hypocotyl SAS mutants tested, we observed altered SAS phenotypes in six mutants (28%). This relatively low validation rate is probably due to differences in growth conditions such as day length, the ratio of red to far-red used for sun and shade, or whether shade was applied throughout the day or simulated by EODFR. In addition to previously described SAS mutants, we discovered seven mutant lines with previously unknown hypocotyl SAS defects (*gibberellin 20-oxidase (ga20ox) 1 ga20ox2 (ga20ox1/2)*, *phytochrome rapidly regulated (par) 2-1*, *indole-3-acetic acid inducible (iaa) 6 iaa19 (iaa6/19)*, *light-regulated zinc finger protein 1 (lzf1)*, *spatula (spt)*, *elongated hypocotyl 5 (hy5)*, *cry1/2*, and *atr4*).

Our mutant phenotypic profiling also revealed that two different indices for one phenotype (“petiole length” and ratio of “petiole blade length ratio” for SAS in petiole; “flowering.time” and “flowering.time.resid” for SAS in flowering time) were clustered together, two elongation phenotypes (hypocotyl elongation and petiole elongation) were clustered while the two

elongation phenotypes and flowering time were distinct (top dendrogram in Fig 2). Examining which genes have mutant phenotypes for each trait revealed that there are common and separate pathways for each shade avoidance response (Fig 2 and S3 Table). Auxin-related genes are required for all responses; phytochromes and three TFs (*LZF1*, *HY5*, and *PIL1*) were involved in both hypocotyl and leaf responses; cryptochromes are involved in both hypocotyl and flowering-time response; and JA related pathways were involved in petiole and flowering time responses. Twelve genes were required specifically for petiole response and two genes were required specifically for hypocotyl response (Fig 2 and S2 Table). Our results also showed that our strategy was powerful for identifying new genes required for these shade avoidance responses, including genes in light, auxin, JA, and BR pathways (S2 Table). Details of those genes will be discussed in following sections.

Differences in the sets of genes required for shade avoidance response in hypocotyls and leaves are consistent with GO analysis of shade-responsive genes in hypocotyl or leaf tissue (Table 1, 2, and 4). For example GO:0009753 (response to JA stimulus) is only found in leaf data sets and we found that mutants affecting the JA pathway only affected SAS phenotypes in adult plants. There is a previous report of JA affecting seedling SAS, but this was under extremely low R:FR (0.068) and JA was found to act by modulating phyA signaling [48]. Under the more moderate low R:FR conditions used in this study and for the seedling microarray assays [7], phyA is not involved (Fig 2) and phyB is the major receptor for shade. Thus, under moderate shade conditions, JA is likely to affect adult rather than seedling SAS.

Our findings that pathways of flowering time in response to shade were different from those of hypocotyl are consistent with previous data. For example, altered *ARABIDOPSIS THALIANA HOMEBOX PROTEIN 2* expression changed shade-avoidance responses in hypocotyl [49] but not in flowering time [43]. In addition, *suppressor of phytochrome a-105 (spa1/2/3/4)* quadruple mutant and *constitutive photomorphogenesis 1 (cop1)* show shade-induced acceleration of flowering time, but did not show shade-induced hypocotyl elongation [50].

Updated shade avoidance syndrome pathways

Among 59 tested mutant lines we could detect 33 mutants with defects in at least one shade avoidance response, including 20 new mutant lines (Fig 2). A schematic diagram of SAS signaling pathways is shown in Fig 3. SAS phenotypes with mutants used in this study and known SAS mutants are summarized in S3 Table. Below we discuss details of the pathways corresponding to each mutant category in Fig 1 and then discuss phenotypic clustering of SAS mutants (Fig 2).

Light signaling pathway. We simulated canopy shade solely by changing the R/FR ratio, which is perceived by phytochromes. Five *PHY* genes in *Arabidopsis thaliana* have partially overlapping functions [51]. Our phenotypic profiling showed *PHYB* is the major photoreceptor and *PHYA* has minor function with *PHYB* in the SAS responses of hypocotyl and leaf in low R/FR condition because *phyA* mutant phenotypes could be observed only in a *phyB* mutant background, consistent with previous data [1] (S3 Table, S3A, S3B, and S3F Fig). The situation is different for flowering time where *PHYB* does not dominate the response, instead it is redundant with *PHYD* and *PHYE* [47]. This is consistent with our data that *phyB* and even *phyA/B* showed normal shade acceleration of flowering time (Fig 2).

We found that the CRY blue-light photoreceptors were important for proper hypocotyl and flowering time SAS under simulated canopy shade. At first this is surprising because our simulated sun and shade conditions only have altered R/FR ratios but have the same blue irradiance. However, CRY1 and CRY2 blue light photoreceptors of *Arabidopsis thaliana* are known to interact genetically with phytochrome signaling in other photomorphogenic responses [52,53,54,55,56,57,58,59,60]. There are numerous molecular events known to occur upon




Fig 3. Schematic representation of proposed signal transduction for shade-avoidance syndrome. PIFs represent PIF3, PIF4, PIF5, PIF7, PIL1, and SPT. JAZs represents JAZ5 and JAZ10. Black lines represent genetic interactions, blue lines show direct interactions, and yellow lines show hormone biosynthesis/metabolism. Lines without arrowheads are for interactions where directionality is unknown. Dashed lines show hypothetical interactions. SCL13 is omitted because of its unknown function within this context.

doi:10.1371/journal.pgen.1004953.g003

phytochrome activation that may explain functional interaction between phytochromes and cryptochromes. One example is light-dependent PHYB-CRY1 interaction [61] and light-dependent CRY-COP1 E3 ligase interaction [62]. Recent studies showed COP1 and its interacting proteins SPA1/2/3/4 are also required for SAS in hypocotyl [50,63,64], so that COP1-SPAs system could be a key component of phytochrome-cryptochrome interaction observed in our data. Although the *cry1/2* double showed reduced SAS in hypocotyl and reduced acceleration of flowering time, the mode of action for these two traits are different. *cry1/2* hypocotyls showed constitutive shade avoidance, while *cry1/2* showed constitutive exaggerated “sun” phenotype for flowering time, indicating that the interaction between phytochrome and cryptochrome signaling are different between these two traits. Possible mechanisms of the differences could be related to photoperiod-dependent flowering time regulation by CRY2 [65]. CRY2 interacts with SPA in blue-light dependent manner, which prevents CO degradation by COP1-SPAs [66]. Also CRY activates a subset of bHLH TFs in a blue light dependent manner; these bHLHs in turn activate the flowering time master gene, FT [67]. Since cryptochromes are required to sense depleted blue light under canopy shade [68,69], plants may have evolved a cross-talk system between phytochrome and cryptochrome signaling systems to coordinate shade avoidance syndromes response in natural condition.

Among TFs in light signaling pathways that we tested (Fig 1), our data showed that *HY5*, *LZF1*, *PIL1*, and *SPT* were required in common between hypocotyl and petiole responses, while *PIF1*, *PIF3*, *PIF4*, *PIF5*, *PAR1*, *LONG HYPOCOTYL IN FAR-RED (HFR1)*, *SALT TOLERANCE (STO)*, and *SALT TOLERANCE HOMOLOG2/B-BOX DOMAIN PROTEIN 21 (STH2/BBX21)* were required in petiole response and *PAR2* was required in hypocotyl.

Among them *SPT* is of particular interest because of its unique spatial expression pattern at the boundary between the leaf blade and petiole, where petiole cell proliferation occurs [70]. Thus the spatial expression pattern of *SPT* could explain the reduced SAS petiole elongation phenotype of *spt-11*. Also the severe allele of *SPT* mutant (*spt-11*) showed less shade avoidance responses than mild allele (*spt-12*) (S3 Table), consistent with the reported differences of petiole length and leaf blade area phenotype between *spt-11* and *spt-12* [70]. *SPT* regulates gynoecium development by activating genes involved in shade avoidance [71]. Similar mechanisms of *SPT* regulation of shade avoidance are likely conserved in shade-induced petiole elongation.

It is well recognized that PIFs are growth-promoting proteins whose activities are modulated by a variety of environmental factors [72]. In addition to *PIF4* and *PIF5*, our data showed that both *PIF1* and *PIF3* are new positive regulators of SAS in petiole (Fig 2). *PIF1* is known to regulate hypocotyl growth [72,73] as well as non-growth related processes such as chlorophyll biosynthesis [74]. *PIF3* is also known to regulate hypocotyl growth (reviewed in [75]), but the role of either gene in regulating adult plant growth is not known. We confirmed the *pif3* result by examining a second allele, *pif3-3* [76], and found that *pif3-3* also had a significant petiole SAS defect (S4 Fig; $p < 0.01$). In hypocotyl growth, *PIF1/3/4/5* regulate overlapping and specific target genes [77], and it seems likely that many of these targets also contribute to petiole SAS.

We found that a knock-out mutant of a PP2C-type phosphatase gene, *MISREGULATED IN DARK (MIDA) 9*, showed reduced petiole shade response. *MIDA9* has been shown to be involved in hook formation and its expression is regulated by *PIF3* [23], consistent with our finding that *pif3* mutant also showed reduced petiole shade response. We previously found that *MIDA9* is induced by *PIF4* and/or *PIF5* during hypocotyl growing phase [78], suggesting that *MIDA9* is involved in growth-control networks. Another link is inactivation of H^+ -ATPase activity (required for cell wall loosening) via dephosphorylation of H^+ -ATPase by *MIDA9* phosphatase 2C-typeD [79]. It would be interesting to know if other *PIFs* and *SPT* controls *MIDA9* expression.

Auxin pathways. Previous studies have implicated auxin as being important in hypocotyl and leaf SAS [80]. Here we extend those studies to identify specific auxin signaling components important for SAS in various organs. Consistent with prior studies [4,5,6,7], both our transcriptome data and mutant phenotypic profiling showed auxin pathways were involved in hypocotyl and leaf shade avoidance responses. By assaying many auxin pathway genes we can begin to assign shade avoidance functions to specific components.

Our data showed that components of auxin pathway required for each response are different. For example, two different gene families in one auxin biosynthesis pathway contribute differently in three separate shade avoidance responses. Although *TAA1* protein catalyzes auxin biosynthesis in the same pathway as *YUC* proteins [81,82], *TAA1* was required only for hypocotyl shade avoidance response, while *YUC2/5/8/9* genes were required for all shade avoidance responses (Fig 2; see Müller-Moulé et al., submitted, for a more detailed analysis). These differences could be explained by lower gene expression level of *TAA1* in leaves, in contrast of ubiquitous expression of *YUC2/5/8/9* (S5 Fig). Among shade-induced *AUX/IAA* family genes (see S2 Table), we tested the function of *IAA6*, and *IAA19*. The *iaa6/19* double showed a defect only in hypocotyl and leaf SAS, but not in acceleration of flowering time. Therefore *IAA6/IAA19* are required for response to shade in elongation. Similar to other auxin-related phenotypes, these

shade-induced *AUX/IAA* genes likely function redundantly with other family members in the other two responses [83].

From transcriptome data it is not clear whether the auxin pathway is involved in acceleration of flowering time by shade, but phenotypic profiling indicates its involvement in flowering time control in response to shade. An auxin overproducing mutant (*atr4*) [84] and reduced auxin level mutant (*yuc2/5/8/9* and *sav3/taa1*) showed opposite flowering responses, supporting a role of auxin in this phenomenon (Fig 2).

How are light signaling and auxin pathways connected? All available data suggested that shade increases active auxin content by inducing auxin biosynthesis genes (*YUCs*) [80]. Among the *PIFs*, connection of *PIF4/5/7* and *SPT* to auxin has been reported [4,5,78,85,86,87]. At least some SAS traits are reduced in single or multiple *pif4/5/7* mutants (summarized in S3 Table). In our study, *yuc2/5/8/9* mutant is defective in all three shade-avoidance responses, which is consistent with upregulation of *YUC2/5/8/9* genes by *PIF7* [5] and *YUC5/8/9* by *PIF5* [4]. Interaction of *TERMINAL FLOWER 2 (TFL2)* protein with *IAA6* and *IAA19* could be another pathway to link SAS and auxin [88], although we were not able to test *TFL2* in this study. Nevertheless, *tfl2* showed reduced shade avoidance (EODFR response) in hypocotyl [88].

JA pathway. Evidence showing a link between light and JA pathways has been accumulating for some time (reviewed in [89]). For example JA mediated defense responses are attenuated in shade [11]. This effect is mediated by COI1-JAZ10-dependent, salicylic acid-independent mechanisms [90]. Previous work also implicated JA signaling in *phyA* mediated responses to very low R:FR [48]. In tomato, stems of plants treated with shade for four days showed reduced JA levels [37]. Our transcriptome ORA with hormone responsive genes also shows shade-attenuation of the JA pathway (Table 3).

While it is known that shade attenuates JA responses, we used JA pathway mutants (Fig 1) to probe which JA components are required for SAS. Because *PHYA* is not important for the moderately low R/FR used in our study (see “light signaling pathway” section above) this enables us to determine whether JA is important for the typical *PHYB* mediated shade responses examined here. Our phenotypic profiling of these mutants showed that the JA pathway is involved in shade avoidance responses in leaf and flowering time, but we did not find significant effects on hypocotyl response (Fig 2, and S3 Table). This is consistent with GO analysis of transcriptomes of hypocotyl and juvenile plants (Table 1, 2, and 4) where JA GO categories were over-represented in the leaf/apical region but not in the hypocotyl data set. Reduced JA mediated plant immunity by shade is found in adult plants, but it is not known if this is true for hypocotyl. Comparison of shade-responsive genes between hypocotyl and leaf/apical region and comparison of phenotypic profiling predict that JA mediated plant immunity in hypocotyl is not affected by shade. In conclusion, we found that interaction of JA pathways and SAS network were not unidirectional but bidirectional.

Interestingly, partially different subsets of JA pathway genes are required for normal leaf and flowering shade response: *COI1* and *JASMONATE RESISTANT 1 (JAR1)* for leaf response and *COI1* and *JAZ5* for flowering time response. What might drive these differences? *JAR1* encodes a protein that catalyzes conjugation of Ile to JA to produce active JA, which is required for JA mediated immunity [89]. Therefore one possibility is that JA-Ile is the active component for leaf response while other active JA-related compounds such as OPDA or cis-jasmonate [89], could be the active components for flowering time. Another possibility is that *JAZ5* acts in a flowering time specific pathway.

It is interesting that JA biosynthesis mutants (*allene oxide synthase (aos)* and *12-oxophytodienoate-reductase 3 (opr3)*) still retained weak shade-avoidance responses although a JA receptor mutant (*coi1-16*) had a defect of shade avoidance responses in petiole and flowering time (Fig 2). The fact that *opr3* mutants retaining JA biosynthesis in certain

condition [91] may explain this. Alternatively, native JA related compounds released from neighbor plants in our experiments might partially rescue our JA biosynthesis mutants.

MYC2 is a basic helix-loop-helix (bHLH) TF important for JA mediated immune responses that acts redundantly with its homologs, *MYC3* and *MYC4* [92]. Mutation in *MYC2* slightly reduced SAS in adult plants (Fig 2, and S3 Table), raising the possibility that the *MYC2/3/4* redundancy is also true for SAS. *PHYTOCHROME AND FLOWERING TIME 1 (PFT1)* encodes the conserved MED25 subunit of the mediator complex, which is involved in JA-mediated immune system as well as light-signaling pathways [46,93,94,95]. Our hypothesis that *pft1* mutant impairs SAS was not confirmed because *pft1* did not show detectable differences compared with Col in any phenotypes we tested.

Our phenotypic profiling data showed less contribution of JA related-genes to shade-induced hypocotyl elongation than expected from previous data. For example, it was previously reported that exogenous MeJA application inhibited hypocotyl growth [96] and that this effect was influenced by *PHYB* [97]. Another examples is that *coi1* mutants showed an increases response to low R/FR [48]. The inconsistency with our results could be due to differences between exogenous JA application in the previous study versus the use of JA pathway mutants in our study, or due to differences in light conditions: previous studies used continuous monochromatic red light or extremely low R/FR (0.068), while our study used low R/FR (0.5). Therefore how light quality influences JA-dependent SAS is of future interest.

How JA pathways modulate petiole elongation is also unclear. Possible mechanism is promotion of PIF activities by binding of JAZ proteins to DELLA proteins that suppress PIF protein activities) [98]. Another link between JA and growth is that exogenous MeJA delays the start of endoreduplication cycle [99]. Interaction of these factors is of future interest.

Gibberellic acid pathways. The GA pathway is involved in shade-induced hypocotyl and petiole elongation as shown by GA biosynthesis inhibitor treatments and a GA deficient mutant (*ga requiring 1 (ga1-3)*) [8]. The GA signaling *DELLA* gene family is reported to have a weak effect in hypocotyl SAS and none in petiole SAS [8]. We used another GA deficient mutant line (*ga20ox1/2*) [78,100] and a quadruple *DELLA* mutant line (*dellaQ*) [101] to test if GA is involved in other SAS traits (Fig 2). Similar to *ga1-3*, *ga20ox1/2* showed reduced response to shade in hypocotyl (Fig 2). In contrast to *ga1-3*, *ga20ox1/2* showed shade-induced petiole elongation even though both organs were about 60% of wild-type size (S3A and S3B Fig). In agreement of previous studies, *dellaQ* showed very weak influence on shade avoidance responses of hypocotyl and petioles in our condition (Fig 2, S3A, and S3B Fig). Both GA deficient and GA signaling mutants showed normal shade response for flowering time (Fig 2 and S3 Table), suggesting that GA pathway is not involved in flowering time acceleration by shade.

Circadian clock pathways. We found that over-expression of *RVE8*, a circadian clock component, caused moderately exaggerated shade responses in flowering time (Fig 2). This is consistent with the finding that natural variation of *ELF3* modulates flowering time response to shade [44]. Recent studies showed direct targets of *RVE8* TF [102]. Among them *PSEUDO RESPONSE REGULATOR5 (PRR5)*, another circadian clock component, is of interest because *PRR5* is activated by *RVE8* [102] and knocking out *PRR5* caused exaggerated shade avoidance response in petiole possibly through elevation of *PIF4* and *PIF5* expression levels [103]. Also it has been shown that circadian modulation of hypocotyl response to shade is mediated by circadian clock regulation of *PIL1* expression [30]. However, our shade avoidance assay for flowering time showed *pif4/5* and *pil1* had normal flowering time shade responses. It would be interesting to test if a *RVE8*—*PRR5* pathway contributes to shade response in flowering time.

Novel component. Our mutant analysis defined a new components of shade avoidance responses; a potassium channel gene (*POTASSIUM CHANNEL IN ARABIDOPSIS THALIANA 1, KAT1*). The *KAT1* gene encodes a potassium channel gene, which is locally expressed in

guard cells [104,105]. *KAT1* potassium channel is inactivated upon ABA treatment and mediates stomata closure [106]. *KAT1* is thought to be important for vegetative growth [107], but its molecular mechanisms are unknown. Recently ABA was shown to be involved in suppression of branching by shade [36]. It is not clear if ABA pathways were involved in shade avoidance responses described in this paper, although ABA responsive genes were enriched in shade responsive genes (see above). Interestingly *KAT1* gene expression is induced by auxin [108,109], which may be related to its role in petiole SAS.

Phenotypic-clustered SAS mutants may share common functions

In *C. elegans*, multiple phenotypic profiling showed that mutants with similar phenotypic profiles function in shared pathways [16]. In our phenotypic clusters (Fig 2), two flowering time mutant lines (*gi-2* and *co-9*) cluster together. These are known from prior studies to act in photoperiodic induction of flowering, and to have reduced shade response for flowering time but not petiole elongation [42] (cluster 7), showing proof of concept. In our data, clusters that consist of SAS mutants are of particular interest (clusters 9, 10, 11, and 12) because they could indicate shared membership in sub-networks of the shade avoidance pathway. Therefore, the following genes likely function in common sub-networks: *IAA6*, *IAA19*, *PIL1*, *LZF1*, *HY5*, and *SPT* in cluster 9 (hyp and pet), and *PAR1*, *KAT1*, *PIF4*, *PIF5*, *MIDA9*, *PIF3*, and *JAR1* in cluster 10 (pet only). Opposite effects of mutations were found in cluster 12; mutation caused reduced responses in hypocotyl and petiole, while its caused exaggerated responses in flowering time. Molecular networks within these clusters are of future interests.

Mutants affecting of flowering time regulation under simulated sun

We observed many genotypes that showed altered flowering time in sun condition (S6 Fig). In this section, we will discuss about known components and then novel components in flowering time pathways.

Known components. As reported known knock out of master regulators of flowering time (*CO* and *GI*) showed late flowering in our long-day condition [110,111,112]. Light signaling has been known to modulate flowering time [113]. Phytochromes affect flowering time by post-transcriptional regulation and cryptochromes affect flowering time by both transcriptional and post-transcriptional regulation [62]. We observed that *phyB*, *phyA/B*, and *cry1/2* mutants showed flowering time phenotype as reported. Early flowering phenotype of *hy5 hy5* homolog (*hyh*) is also reported [114]. *PIF4* has been known to induce a master regulator of flowering time, *FT* gene, upon raised temperature [115]. *pif4/5* mutant showed late flowering time, so that it is likely that *PIF4* and/or *PIF5* induce *FT* gene expression in our condition.

GA is essential for flowering time control. The GA biosynthesis double mutant (*ga20ox1/2*) showed delayed flowering in our experiment, consistent with earlier reports [100]. However, this mutant showed normal shade effects on flowering time (see above).

The circadian clock is also important for flowering time control. As reported, we found opposite phenotypes of *rve8* (early flowering) and *RVE8-OX* (late flowering) [116].

New components. It is surprising that auxin was involved in flowering time regulation, since it is not incorporated into current models of flowering time pathways (reviewed in [113]). However, exogenous auxin has been reported to delay flowering time, perhaps due to induced damage on plants [117]. This result is consistent with our auxin biosynthesis mutant flowering time data; overproduction delayed flowering time (*atr4*) while reduced production accelerated flowering time (*yuc2/5/8/9* and *taa1*). These results suggest that it is necessary to re-investigate the involvement of auxin pathways in flowering time.

It has been shown that *coi1* mutants flowered early [48,98]. We observed that other JA mutants also show altered flowering time. Not only JA biosynthesis mutants (*aos* and *opr3*) and JA receptor mutant (*coi1-16*), but also three JA signaling component mutants (*JAZ1* RNAi, *jaz5-1*, *myc2*) showed early flowering phenotypes under sun condition. Early flowering of *myc2* is inconsistent with a previous report showing its late flowering phenotype [118] possibly due to different photoperiods. It will be interesting to investigate how JA pathways regulate flowering time. One possibility is that JA pathways interact with GA pathways in control of flowering, similar to their interaction in growth control [98].

Our data showed that two SPT knock out mutants (*spt-11* and *spt-12*), *PAR1* RNAi, *sth2*, and two KAT1 knock out mutants (*kat1-1* and *kat1-2*) showed early flowering phenotype, while *SCL13* anti-sense line 1 (*scl13* as1) and *hfr1* showed late flowering phenotype. Early flowering of *spt-11* has been reported [119] and that was confirmed by another allele (*spt-12*) in our study. Early flowering phenotype of *PAR1* RNAi, *sth2*, and *kat1-1* have not been previously reported and their connections to the flowering time pathway are unknown.

Conclusion

Here we showed that RNA-seq followed by phenotypic profiling is a powerful approach for elucidating complex SAS pathways and discovery of new SAS components. A similar approach was successful in searching new components of de-etiolation of seedlings, a developmental stage with a simple architecture [24]. Our study expanded this approach to show the transcriptome-based discovery of new mutants are also effective for complex syndrome by multiple phenotypic profiling.

After our phenotypic profiling, an additional SAS mutant line has been reported which contained genes that were also in our shade-responsive genes. Specifically, we found that *BR ENHANCED EXPRESSION 3* (*BEE3*), a bHLH TF, was induced by shade (S2 Table). It was recently shown that the *bee1 bee2 bee3* triple mutant has altered hypocotyl SAS [120], perhaps due to the altered BR signaling in this triple [121]. This example is additional evidence that our strategy is effective to find novel SAS mutants. Further analysis is needed for elucidating interactions between these genes and/or pathways. Our approaches are straightforward and cost effective, so that these should be applicable to other cases in general.

We found that the effects of some mutations were context-dependent (only found for some organs or developmental stages) whereas others were ubiquitous. Those mutations that affect all organs points to shared mechanisms underlying the SAS in different organs. The mutations that have context-dependent effects could indicate unique genes functioning in the different organs or more quantitative differences in the relative importance of the components in different organs. Regardless the fact that we did find organ-specific effects suggests that we need to be cautious when generalizing conclusions from hypocotyl studies.

Materials and Methods

Light condition

For simulated sun condition, white light (cool-white fluorescent light) was supplemented with far-red light (provided by LEDs (Orbitec, inc) to obtain R/FR = 1.86. For simulated shade condition, white light was supplemented with far-red LEDs to obtain R/FR = 0.52. Both condition had 80–100 μ E of Photosynthetically Active Radiation (PAR). Plants were grown under long day condition (16 hour light/8 hour dark) at constant temperature (22°C). For hypocotyl experiments, seedlings were grown under simulated sun (R/FR = 1.3) or simulated shade condition (R/FR = 0.5) with combination of LED lights (Quantum Devices Snap-Lite) [122]. Ambient light spectrum was measured by Black-Comet (StellarNet, Florida).

Plant materials

Arabidopsis seed stocks used in this study are listed in Fig 1. To confirm genotypes of T-DNA insertion lines ordered from Arabidopsis Biological Resource Center (ABRC), genomic DNA was extracted (DNeasy Plant Mini kit, Qiagen) and subject to genomic PCR. cDNA was synthesized by direct mRNA extraction [123] and quantitative PCR (qPCR) was done with homemade SYBR green master mix with the iCycler Multicolor real-time PCR detection system (Bio-Rad). For *kat1* mutants, standard RT-PCR was done. Primers used for genomic PCR and (q)RT-PCR and their results were summarized in S4 Table. *Arabidopsis* seeds were imbibed with water on filter papers and stored them at 4°C for four days. Three days after stratification under sun condition, three germinated seeds were transferred to soil in a well of 5x10 well flat. Fourteen days after stratification, excess seedlings were removed to leave one well-grown plant per pot and the flats were transferred to either sun or shade condition. For hypocotyl growth measurements, seeds were grown on vertical square plates [124] with 1/2 MSMO, 5 mM 2-(N-morpholino) ethanesulfonic acid (MES, pH = 5.8, Sigma), and 0.8% agar (Sigma). Each plate was divided into three rows and two columns and in six spaces five or six seeds of six genotypes were sown. Genotype positions were randomized in repeated sets. 4593 seedling images were taken by a scanner and hypocotyl length was measured by ImageJ (<http://rsb.info.nih.gov/ij/>) [125].

RNA-seq library preparation and sequencing

For RNA extraction, plants were treated with shade starting at ZT 4 or left in the sun. We prepared two replicates of each sample at 1 hour and 4 hours after sun and shade treatment and five plants were pooled for each replicate. Cotyledons, hypocotyls, and roots were removed from the samples, leaving leaves and apical tissue. Total RNA from the plants was extracted using RNeasy Plant Mini kit (Qiagen) with DNase treatment (Qiagen). Five µg total RNA was used to construct mRNA library using mRNA-Seq-8 sample Prep kit (Illumina). The resulting cDNA libraries were sequenced by Illumina GAIIx with 40 bp single end mode. Basic statistics of mapping results are given in S1 Table.

Differential expression analysis and over-representation analysis (ORA)

Reads after sorting according to barcodes were subjected to removal of adaptor contamination by custom Perl scripts. Reads were mapped by TopHat [126] to *Arabidopsis* reference genome using known annotation (TAIR10). Differentially expressed genes were extracted by edgeR package [127] in R statistical environment [128] (FDR <0.001). ORA was done by Goseq package [129] in R statistical environment. GO analysis was done by using GO category database package from Bioconductor (org.At.tair.db and ANNOTATE package). For ORA of hormone responsive genes custom categories were used as defined in Supplemental Table S9 in [130] and Supplemental S1 Table in [131]. GO analysis of shade-responsive genes in hypocotyl [7] was done using the GO Web site (http://amigo.geneontology.org/cgi-bin/amigo/term_enrichment; [132]).

Phenotype measurement and analysis

For scoring leaf phenotypes, 26 day old plants were dissected and leaf images were recorded by a flatbed scanner (Epson, Perfection V700 PHOTO). Scanned images were measured using ImageJ [125] and the LeafJ plugin [25] to determine petiole length, leaf blade length, leaf blade width, and leaf blade area. Days to bolting was scored to measure flowering time. Leaf phenotypes (petiole length, leaf blade length, leaf blade width, leaf blade area) were measured from 10 sets of experiments with 1268 plants in total. For flowering time (days to bolting) measurement, 1950 plants were measured in total. Each phenotype was fitted by mixed

effects model, i.e.

$$\text{trait} = \text{plant} + \text{treatment} + \text{plant} : \text{treatment} + (\text{treatment}|\text{set}) + \epsilon$$

where plant is a mutant/overexpressor, treatment is sun or shade condition, plant:treatment is interaction of “plant” and “treatment”, (treatment|set) is the random effect associated with the treatment in set of experiments, and ϵ is the error. The model was applied to each trait to calculate coefficient (“sun” value). For leaf traits where we measured across multiple leaves (from leaf 3 to leaf 6) for a given trait we treated leaf as a random effect, using the following model

$$\text{trait} = \text{plant} + \text{treatment} + \text{plant} : \text{treatment} + (1|\text{leaf}) + (\text{treatment}|\text{set}) + \epsilon$$

Mutants were considered to have a defect in SAS when the plant:treatment term was significant ($P < 0.05$), indicating that the genotype of the plant (mutant versus wild-type) affected the response to shade.

For flowering time, days to bolting was log₂ transformed. We found that acceleration of flowering time by shade treatment was strongly correlated with days to bolting in sun condition, i.e., late flowering mutants had more shade-accelerated flowering time than Col (S2G Fig). To address this issue we regressed flowering time shade response on average sun flowering time for each genotype and calculated the residuals from the regression [122]. These residuals represent the amount of flowering time shade response that was not predicted by the sun flowering time. The residuals for shade treated plants were then used in the mixed effects model (S3H Fig).

$$\text{residuals} = \text{plant} + (1|\text{set}) + \epsilon$$

The lme4 (R package version 1.0–6) [133] and lmerTest [134] packages in R was used for these analyses. All phenotyping data is summarized in S5 Table.

Heatmaps for phenotypic clustering were drawn after scaling each trait data and centered at Col.

All R scripts for this paper and raw data are available at <https://bitbucket.org/knozue/sasphenotyping>.

Accession numbers

RNA-seq data in this study have been deposited in the NCBI SRA (Study ID PRJNA214254) and the NCBI GEO database (accession GSE66967). Mutants used in this study are listed in Fig 1.

Supporting Information

S1 Fig. Comparison of different platform and tissues with fold-changes of shade-responsive gene expression. Current RNA-seq data (juvenile plants under low R:FR labeled as “Nozue”) and microarray (leaf or petiole treated with EODFR [6] labeled as “Kozuka”), and hypocotyl treated with low R:FR (1 hour [7,26] (labeled as “Tao” and “Sessa.1h”, 4 days [26] labeled as “Sessa.4d”).

(PDF)

S2 Fig. Model checking for each trait. (A) hypocotyl, (B) petiole length, (C) leaf blade length, (D) leaf blade width, (E) leaf blade area, (F) petiole length/leaf blade length ratio, (G) flowering time, and (H) flowering time (log₂ transformed).

(PDF)

S3 Fig. Graphs of each trait. (A) hypocotyl, (B) petiole length, (C) leaf blade length, (D) leaf blade width, (E) leaf blade area, (F) petiole length/leaf blade length ratio, (G) flowering time

(log₂ transformed) and (H) flowering time (log₂ transformed residuals). Error bars in (A) to (G) represent standard errors. Genotype names in (H) indicate lines whose flowering time shade response differs significantly from prediction by regression ($p < 0.05$).

(PDF)

S4 Fig. Col and *pif3-3* petiole length. Plants were grown in simulated sun and shade using our standard conditions. Three independent experiments were performed and a total of 20 to 41 plants were examined per treatment/genotype combination. *pif3-3* has a significantly reduced response to shade ($p < 0.01$ for genotype X treatment interaction in linear regression).

(PDF)

S5 Fig. Expression pattern of *TAA1* and *YUC2/5/8/9*. Developmental expression pattern was obtained from eFP browser [136].

(TIF)

S6 Fig. Heatmap with absolute values. Values were normalized and centered on Col (i.e., Col value = 0) and visualized with color coding (magenta indicates larger value than Col while green indicates smaller value relative to Col). Colors of asterisks indicate genetic background of each mutant, i.e., Col (white), *Ws* (yellow), and *Ler* (light blue).

(TIF)

S1 Table. Statistics of RNA-seq data. In each condition two biological replicates are shown in a and b.

(XLSX)

S2 Table. A complete list of shade-responsive genes in leaf/apical region of plants. Blue letters; shade-induced genes, pink letters; shade-repressed genes, yellow box; no probes on ATH1 microarray. Bold text; known shade-responsive genes. Comparison with transcriptome data from EODFR treated leaf blade and petiole [5] are also shown.

(XLSX)

S3 Table. Summary of SAS phenotypes with known SAS mutants and mutants used in this study.

(XLSX)

S4 Table. List of primers used in genotyping and (q)RT-PCR and their results. (Nearly) homozygous lines are shown in light blue.

(XLSX)

S5 Table. Complete phenotype data after applying mixed effects models.

(CSV)

Acknowledgments

We thank Arabidopsis Biological Resource Center (ABRC), Judith Bender, Cordelia Bolle, Giltsu Choi, Joanne Chory, Katayoon Dehesh, Christian Fankhauser, Wim Grunewald, Nicholas P. Harberd, Stacey Harmer, Peter Hedden, Daniel Kliebenstein, Chentao Lin, Jaime F. Martínez-García, David Smyth, Hirokazu Tsukaya, Detlef Weigel, Shu-Hsing Wu, and Yunde Zhao for providing seeds listed in Fig 1. We also thank Patricia Mueller-Moule for bulking seeds, Cody Markelz for stimulating discussion, and Natalie Gath, Grace Pan, Ji-hoon Lee, Amanda Schrager, Leonela Carriedo, Matthew Jones, Huy Tran, Ian Andrew-Hanna Knox, Mary Zhang, and Cody Markelz for sample preparation and collection.

Author Contributions

Conceived and designed the experiments: JNM KN YI. Performed the experiments: KN MR MRM SL UKD. Analyzed the data: AVT KN JNM. Contributed reagents/materials/analysis tools: AVT UKD KN JNM. Wrote the paper: KN JNM.

References

- Casal JJ (2013) Photoreceptor signaling networks in plant responses to shade. *Annu Rev Plant Biol* 64: 403–427. doi: [10.1146/annurev-arplant-050312-120221](https://doi.org/10.1146/annurev-arplant-050312-120221) PMID: [23373700](https://pubmed.ncbi.nlm.nih.gov/23373700/)
- Ballaré CL, Sánchez RA, Scopel AL, Casal JJ, Ghersa CM (1987) Early detection of neighbour plants by phytochrome perception of spectral changes in reflected sunlight. *Plant Cell Environ* 10: 551–557.
- Lorrain S, Allen T, Duek PD, Whitelam GC, Fankhauser C (2008) Phytochrome-mediated inhibition of shade avoidance involves degradation of growth-promoting bHLH transcription factors. *Plant J* 53: 312–323. PMID: [18047474](https://pubmed.ncbi.nlm.nih.gov/18047474/)
- Hornitschek P, Kohnen MV, Lorrain S, Rougemont J, Ljung K, et al. (2012) Phytochrome interacting factors 4 and 5 control seedling growth in changing light conditions by directly controlling auxin signaling. *Plant J* 71: 699–711. doi: [10.1111/j.1365-313X.2012.05033.x](https://doi.org/10.1111/j.1365-313X.2012.05033.x) PMID: [22536829](https://pubmed.ncbi.nlm.nih.gov/22536829/)
- Li L, Ljung K, Breton G, Schmitz RJ, Pruneda-Paz J, et al. (2012) Linking photoreceptor excitation to changes in plant architecture. *Genes Dev* 26: 785–790. doi: [10.1101/gad.187849.112](https://doi.org/10.1101/gad.187849.112) PMID: [22508725](https://pubmed.ncbi.nlm.nih.gov/22508725/)
- Kozuka T, Kobayashi J, Horiguchi G, Demura T, Sakakibara H, et al. (2010) Involvement of auxin and brassinosteroid in the regulation of petiole elongation under the shade. *Plant Physiol* 153: 1608–1618. doi: [10.1104/pp.110.156802](https://doi.org/10.1104/pp.110.156802) PMID: [20538889](https://pubmed.ncbi.nlm.nih.gov/20538889/)
- Tao Y, Ferrer JL, Ljung K, Pojer F, Hong F, et al. (2008) Rapid synthesis of auxin via a new tryptophan-dependent pathway is required for shade avoidance in plants. *Cell* 133: 164–176. doi: [10.1016/j.cell.2008.01.049](https://doi.org/10.1016/j.cell.2008.01.049) PMID: [18394996](https://pubmed.ncbi.nlm.nih.gov/18394996/)
- Djakovic-Petrovic T, Wit Md, Voesenek LACJ, Pierik R (2007) DELLA protein function in growth responses to canopy signals. *Plant J* 51: 117–126. PMID: [17488236](https://pubmed.ncbi.nlm.nih.gov/17488236/)
- Kurepin LV, Emery RJN, Pharis RP, Reid DM (2007) The interaction of light quality and irradiance with gibberellins, cytokinins and auxin in regulating growth of *Helianthus annuus* hypocotyls. *Plant Cell Environ* 30: 147–155. PMID: [17238906](https://pubmed.ncbi.nlm.nih.gov/17238906/)
- Carabelli M, Possenti M, Sessa G, Ciolfi A, Sassi M, et al. (2007) Canopy shade causes a rapid and transient arrest in leaf development through auxin-induced cytokinin oxidase activity. *Genes Dev* 21: 1863–1868. PMID: [17671088](https://pubmed.ncbi.nlm.nih.gov/17671088/)
- Moreno JE, Tao Y, Chory J, Ballaré CL (2009) Ecological modulation of plant defense via phytochrome control of jasmonate sensitivity. *Proc Natl Acad Sci U S A* 106: 4935–4940. doi: [10.1073/pnas.0900701106](https://doi.org/10.1073/pnas.0900701106) PMID: [19251652](https://pubmed.ncbi.nlm.nih.gov/19251652/)
- Kegge W, Weldegergis BT, Soler R, Vergeer-Van Eijk M, Dicke M, et al. (2013) Canopy light cues affect emission of constitutive and methyl jasmonate-induced volatile organic compounds in *Arabidopsis thaliana*. *New Phytol* 200: 861–874. doi: [10.1111/nph.12407](https://doi.org/10.1111/nph.12407) PMID: [23845065](https://pubmed.ncbi.nlm.nih.gov/23845065/)
- Sozzani R, Benfey P (2011) High-throughput phenotyping of multicellular organisms: finding the link between genotype and phenotype. *Genome Biol* 12: 219. doi: [10.1186/gb-2011-12-3-219](https://doi.org/10.1186/gb-2011-12-3-219) PMID: [21457493](https://pubmed.ncbi.nlm.nih.gov/21457493/)
- Winzeler EA, Shoemaker DD, Astromoff A, Liang H, Anderson K, et al. (1999) Functional characterization of the *S. cerevisiae* genome by gene deletion and parallel analysis. *Science* 285: 901–906. PMID: [10436161](https://pubmed.ncbi.nlm.nih.gov/10436161/)
- Neumann B, Walter T, Heriche JK, Bulkescher J, Erfle H, et al. (2010) Phenotypic profiling of the human genome by time-lapse microscopy reveals cell division genes. *Nature* 464: 721–727. doi: [10.1038/nature08869](https://doi.org/10.1038/nature08869) PMID: [20360735](https://pubmed.ncbi.nlm.nih.gov/20360735/)
- Green RA, Kao H-L, Audhya A, Arur S, Mayers JR, et al. (2011) A high-resolution *C.elegans* essential gene network based on phenotypic profiling of a complex tissue. *Cell* 145: 470–482. doi: [10.1016/j.cell.2011.03.037](https://doi.org/10.1016/j.cell.2011.03.037) PMID: [21529718](https://pubmed.ncbi.nlm.nih.gov/21529718/)
- Yanik MF, Rohde CB, Pardo-Martin C (2011) Technologies for micromanipulating, imaging, and phenotyping small invertebrates and vertebrates. *Annu Rev Biomed Eng* 13: 185–217. doi: [10.1146/annurev-bioeng-071910-124703](https://doi.org/10.1146/annurev-bioeng-071910-124703) PMID: [21756142](https://pubmed.ncbi.nlm.nih.gov/21756142/)
- Atwell S, Huang YS, Vilhjalmsón BJ, Willems G, Horton M, et al. (2010) Genome-wide association study of 107 phenotypes in *Arabidopsis thaliana* inbred lines. *Nature* 465: 627–631. doi: [10.1038/nature08800](https://doi.org/10.1038/nature08800) PMID: [20336072](https://pubmed.ncbi.nlm.nih.gov/20336072/)

19. Berger B, Regt B, Tester M (2012) High-throughput phenotyping of plant shoots. In: Normanly J, editor. High-throughput phenotyping in plants: Humana Press. pp. 9–20.
20. Spalding EP, Miller ND (2013) Image analysis is driving a renaissance in growth measurement. *Curr Opin Plant Biol* 16: 100–104. doi: [10.1016/j.pbi.2013.01.001](https://doi.org/10.1016/j.pbi.2013.01.001) PMID: [23352714](https://pubmed.ncbi.nlm.nih.gov/23352714/)
21. Dornbusch T, Lorrain S, Kuznetsov D, Fortier A, Liechti R, et al. (2012) Measuring the diurnal pattern of leaf hyponasty and growth in *Arabidopsis*—a novel phenotyping approach using laser scanning. *Funct Plant Biol* 39: 860–869.
22. Bruex A, Kainkaryam RM, Wieckowski Y, Kang YH, Bernhardt C, et al. (2012) A gene regulatory network for root epidermis cell differentiation in *Arabidopsis*. *PLoS Genet* 8: e1002446. doi: [10.1371/journal.pgen.1002446](https://doi.org/10.1371/journal.pgen.1002446) PMID: [22253603](https://pubmed.ncbi.nlm.nih.gov/22253603/)
23. Sentandreu M, Martin G, Gonzalez-Schain N, Leivar P, Soy J, et al. (2011) Functional profiling identifies genes involved in organ-specific branches of the PIF3 regulatory network in *Arabidopsis*. *Plant Cell* 23: 3974–3991. doi: [10.1105/tpc.111.088161](https://doi.org/10.1105/tpc.111.088161) PMID: [22108407](https://pubmed.ncbi.nlm.nih.gov/22108407/)
24. Khanna R, Shen Y, Toledo-Ortiz G, Kikis EA, Johannesson H, et al. (2006) Functional profiling reveals that only a small number of phytochrome-regulated early-response genes in *Arabidopsis* are necessary for optimal deetiolation. *Plant Cell* 18: 2157–2171. PMID: [16891401](https://pubmed.ncbi.nlm.nih.gov/16891401/)
25. Maloof JN, Nozue K, Mumbach MR, Palmer CM (2013) LeafJ: An ImageJ plugin for semi-automated leaf shape measurement. *J Vis Exp* 71: e50028.
26. Sessa G, Carabelli M, Sassi M, Cioffi A, Possenti M, et al. (2005) A dynamic balance between gene activation and repression regulates the shade avoidance response in *Arabidopsis*. *Genes Dev* 19: 2811–2815. PMID: [16322556](https://pubmed.ncbi.nlm.nih.gov/16322556/)
27. Carabelli M, Morelli G, Whitelam G, Ruberti I (1996) Twilight-zone and canopy shade induction of the *Athb-2* homeobox gene in green plants. *Proc Natl Acad Sci U S A* 93: 3530–3535. PMID: [11607652](https://pubmed.ncbi.nlm.nih.gov/11607652/)
28. Roig-Villanova I, Bou-Torrent J, Galstyan A, Carretero-Paulet L, Portoles S, et al. (2007) Interaction of shade avoidance and auxin responses: a role for two novel atypical bHLH proteins. *EMBO J* 26: 4756–4767. PMID: [17948056](https://pubmed.ncbi.nlm.nih.gov/17948056/)
29. Sorin C, Salla-Martret M, Bou-Torrent J, Irma Roig-Villanova I, Martínez-García J (2009) *ATHB4*, a regulator of shade avoidance, modulates hormone response in *Arabidopsis* seedlings. *Plant J* 59: 266–277. doi: [10.1111/j.1365-3113X.2009.03866.x](https://doi.org/10.1111/j.1365-3113X.2009.03866.x) PMID: [19392702](https://pubmed.ncbi.nlm.nih.gov/19392702/)
30. Salter MG, Franklin KA, Whitelam GC (2003) Gating of the rapid shade-avoidance response by the circadian clock in plants. *Nature* 426: 680–683. PMID: [14668869](https://pubmed.ncbi.nlm.nih.gov/14668869/)
31. Finlayson SA, Lee IJ, Mullet JE, Morgan PW (1999) The mechanism of rhythmic ethylene production in sorghum. The role of phytochrome B and simulated shading. *Plant Physiol* 119: 1083–1089. PMID: [10069847](https://pubmed.ncbi.nlm.nih.gov/10069847/)
32. Pierik R, Cuppens MLC, Voesenek LACJ, Visser EJW (2004) Interactions between ethylene and gibberellins in phytochrome-mediated shade avoidance responses in tobacco. *Plant Physiol* 136: 2928–2936. PMID: [15448197](https://pubmed.ncbi.nlm.nih.gov/15448197/)
33. Millenaar FF, Cox MC, van Berkel YE, Welschen RA, Pierik R, et al. (2005) Ethylene-induced differential growth of petioles in *Arabidopsis*. Analyzing natural variation, response kinetics, and regulation. *Plant Physiol* 137: 998–1008. PMID: [15728343](https://pubmed.ncbi.nlm.nih.gov/15728343/)
34. Smalle J, Haegman M, Kurepa J, Van Montagu M, Straeten DVD (1997) Ethylene can stimulate *Arabidopsis* hypocotyl elongation in the light. *Proc Natl Acad Sci U S A* 94: 2756–2761. PMID: [11038610](https://pubmed.ncbi.nlm.nih.gov/11038610/)
35. Pierik R, Whitelam GC, Voesenek LACJ, de Kroon H, Visser EJW (2004) Canopy studies on ethylene-insensitive tobacco identify ethylene as a novel element in blue light and plant-plant signalling. *Plant J* 38: 310–319. PMID: [15078333](https://pubmed.ncbi.nlm.nih.gov/15078333/)
36. Reddy SK, Holalu SV, Casal JJ, Finlayson SA (2013) Abscisic acid regulates axillary bud outgrowth responses to the ratio of red to far-red light. *Plant Physiol* 163: 1047–1058. doi: [10.1104/pp.113.221895](https://doi.org/10.1104/pp.113.221895) PMID: [23929720](https://pubmed.ncbi.nlm.nih.gov/23929720/)
37. Cagnola J, Ploschuk E, Benech-Arnold T, Finlayson S, Casal JJ (2012) Stem transcriptome reveals mechanisms to reduce the energetic cost of shade-avoidance responses in tomato. *Plant Physiol* 160: 1110–1119. doi: [10.1104/pp.112.201921](https://doi.org/10.1104/pp.112.201921) PMID: [22872775](https://pubmed.ncbi.nlm.nih.gov/22872775/)
38. Nagatani A, Chory J, Furuya M (1991) Phytochrome B is not detectable in the *hy3* mutant of *Arabidopsis*, which is deficient in responding to end-of-day far-red light treatments. *Plant Cell Physiol* 32: 1119–1122.
39. Reed JW, Nagpal P, Poole DS, Furuya M, Chory J (1993) Mutations in the gene for the red/far-red light receptor phytochrome B alter cell elongation and physiological responses throughout *Arabidopsis* development. *Plant Cell* 5: 147–157. PMID: [8453299](https://pubmed.ncbi.nlm.nih.gov/8453299/)

40. Robson P, Whitelam GC, Smith H (1993) Selected components of the shade-avoidance syndrome are displayed in a normal manner in mutants of *Arabidopsis thaliana* and *Brassica rapa* deficient in phytochrome B. *Plant Physiol* 102: 1179–1184. PMID: [12231894](#)
41. Devlin PF, Robson PR, Patel SR, Goosey L, Sharrock RA, et al. (1999) Phytochrome D acts in the shade-avoidance syndrome in *Arabidopsis* by controlling elongation growth and flowering time. *Plant Physiol* 119: 909–915. PMID: [10069829](#)
42. Wollenberg AC, Strasser B, Cerdan PD, Amasino RM (2008) Acceleration of flowering during shade avoidance in *Arabidopsis* alters the balance between FLOWERING LOCUS C-mediated repression and photoperiodic induction of flowering. *Plant Physiol* 148: 1681–1694. doi: [10.1104/pp.108.125468](#) PMID: [18790998](#)
43. Kim SY, Yu X, Michaels SD (2008) Regulation of CONSTANS and FLOWERING LOCUS T expression in response to changing light quality. *Plant Physiol* 148: 269–279. doi: [10.1104/pp.108.122606](#) PMID: [18667727](#)
44. Jimenez-Gomez JM, Wallace AD, Maloof JN (2010) Network analysis identifies ELF3 as a QTL for the shade avoidance response in *Arabidopsis*. *PLoS Genet* 6: e1001100. doi: [10.1371/journal.pgen.1001100](#) PMID: [20838594](#)
45. Kanyuka K, Praekelt U, Franklin KA, Billingham OE, Hooley R, et al. (2003) Mutations in the huge *Arabidopsis* gene *BIG* affect a range of hormone and light responses. *Plant J* 35: 57–70. PMID: [12834402](#)
46. Cerdan PD, Chory J (2003) Regulation of flowering time by light quality. *Nature* 423: 881–885. PMID: [12815435](#)
47. Franklin KA, Praekelt U, Stoddart WM, Billingham OE, Halliday KJ, et al. (2003) Phytochromes B, D, and E act redundantly to control multiple physiological responses in *Arabidopsis*. *Plant Physiol* 131: 1340–1346. PMID: [12644683](#)
48. Robson F, Okamoto H, Patrick E, Harris SR, Wasternack C, et al. (2010) Jasmonate and phytochrome A signaling in *Arabidopsis* wound and shade responses are integrated through JAZ1 stability. *Plant Cell* 22: 1143–1160. doi: [10.1105/tpc.109.067728](#) PMID: [20435902](#)
49. Steindler C, Matteucci A, Sessa G, Weimar T, Ohgishi M, et al. (1999) Shade avoidance responses are mediated by the ATHB-2 HD-zip protein, a negative regulator of gene expression. *Development* 126: 4235–4245. PMID: [10477292](#)
50. Rolauffs S, Fackendahl P, Sahn J, Fiene G, Hoecker U (2012) *Arabidopsis* COP1 and SPA genes are essential for plant elongation but not for acceleration of flowering time in response to a low red light to far-red light ratio. *Plant Physiol* 160: 2015–2027. doi: [10.1104/pp.112.207233](#) PMID: [23093358](#)
51. Franklin KA, Quail PH (2010) Phytochrome functions in *Arabidopsis* development. *J Exp Bot* 61: 11–24. doi: [10.1093/jxb/erp304](#) PMID: [19815685](#)
52. Neff MM, Chory J (1998) Genetic interactions between phytochrome A, phytochrome B, and cryptochrome 1 during *Arabidopsis* development. *Plant Physiol* 118: 27–35. PMID: [9733523](#)
53. Casal JJ, Mazzella MA (1998) Conditional synergism between cryptochrome 1 and phytochrome B is shown by the analysis of *phyA*, *phyB*, and *hy4* simple, double, and triple mutants in *Arabidopsis*. *Plant Physiol* 118: 19–25. PMID: [9733522](#)
54. Casal JJ, Boccalandro H (1995) Co-action between phytochrome B and HY4 in *Arabidopsis thaliana*. *Planta* 197: 213–218. PMID: [8547813](#)
55. Ahmad M, Cashmore AR (1997) The blue-light receptor cryptochrome 1 shows functional dependence on phytochrome A or phytochrome B in *Arabidopsis thaliana*. *Plant J* 11: 421–427. PMID: [9107032](#)
56. Hennig L, Funk M, Whitelam GC, Schafer E (1999) Functional interaction of cryptochrome 1 and phytochrome D. *Plant J* 20: 289–294. PMID: [10571889](#)
57. Castillon A, Shen H, Huq E (2009) Blue light induces degradation of the negative regulator phytochrome interacting factor 1 to promote photomorphogenic development of *Arabidopsis* seedlings. *Genetics* 182: 161–171. doi: [10.1534/genetics.108.099887](#) PMID: [19255368](#)
58. Usami T, Matsushita T, Oka Y, Mochizuki N, Nagatani A (2007) Roles for the N- and C-terminal domains of phytochrome B in interactions between phytochrome B and cryptochrome signaling cascades. *Plant Cell Physiol* 48: 424–433. PMID: [17251203](#)
59. Usami T, Mochizuki N, Kondo M, Nishimura M, Nagatani A (2004) Cryptochromes and phytochromes synergistically regulate *Arabidopsis* root greening under blue light. *Plant Cell Physiol* 45: 1798–1808. PMID: [15653798](#)
60. Meijer G, Engelsma G (1965) The synergistic influence of a pre-irradiation on the photoinhibition of gherkin seedlings. *Photochem Photobiol* 4: 251–258.

61. Hughes RM, Vrana JD, Song J, Tucker CL (2012) Light-dependent, dark-promoted interaction between *Arabidopsis* cryptochrome 1 and phytochrome B proteins. *J Biol Chem* 287: 22165–22172. doi: [10.1074/jbc.M112.360545](https://doi.org/10.1074/jbc.M112.360545) PMID: [22577138](https://pubmed.ncbi.nlm.nih.gov/22577138/)
62. Liu H, Liu B, Zhao C, Pepper M, Lin C (2011) The action mechanisms of plant cryptochromes. *Trends Plant Sci* 16: 684–691. doi: [10.1016/j.tplants.2011.09.002](https://doi.org/10.1016/j.tplants.2011.09.002) PMID: [21983106](https://pubmed.ncbi.nlm.nih.gov/21983106/)
63. Crocco CD, Holm M, Yanovsky MJ, Botto JF (2010) AtBBX21 and COP1 genetically interact in the regulation of shade avoidance. *Plant J* 64: 551–562. doi: [10.1111/j.1365-313X.2010.04360.x](https://doi.org/10.1111/j.1365-313X.2010.04360.x) PMID: [21070414](https://pubmed.ncbi.nlm.nih.gov/21070414/)
64. Pacín M, Legris M, Casal JJ (2013) COP1 re-accumulates in the nucleus under shade. *Plant J* 75: 631–641. doi: [10.1111/tpj.12226](https://doi.org/10.1111/tpj.12226) PMID: [23647163](https://pubmed.ncbi.nlm.nih.gov/23647163/)
65. Guo H, Yang H, Mockler TC, Lin C (1998) Regulation of flowering time by *Arabidopsis* photoreceptors. *Science* 279: 1360–1363. PMID: [9478898](https://pubmed.ncbi.nlm.nih.gov/9478898/)
66. Zuo Z, Liu H, Liu B, Liu X, Lin C (2011) Blue light-dependent interaction of CRY2 with SPA1 regulates COP1 activity and floral initiation in *Arabidopsis*. *Curr Biol* 21: 841–847. doi: [10.1016/j.cub.2011.03.048](https://doi.org/10.1016/j.cub.2011.03.048) PMID: [21514160](https://pubmed.ncbi.nlm.nih.gov/21514160/)
67. Liu Y, Li X, Li K, Liu H, Lin C (2013) Multiple bHLH proteins form heterodimers to mediate CRY2-dependent regulation of flowering-time in *Arabidopsis*. *PLoS Genet* 9: e1003861. doi: [10.1371/journal.pgen.1003861](https://doi.org/10.1371/journal.pgen.1003861) PMID: [24130508](https://pubmed.ncbi.nlm.nih.gov/24130508/)
68. Keller MM, Jaillais Y, Pedmale UV, Moreno JE, Chory J, et al. (2011) Cryptochrome 1 and phytochrome B control shade-avoidance responses in *Arabidopsis* via partially independent hormonal cascades. *Plant J* 67: 195–207. doi: [10.1111/j.1365-313X.2011.04598.x](https://doi.org/10.1111/j.1365-313X.2011.04598.x) PMID: [21457375](https://pubmed.ncbi.nlm.nih.gov/21457375/)
69. Keuskamp DH, Sasidharan R, Vos I, Peeters AJ, Voesenek LA, et al. (2011) Blue-light-mediated shade avoidance requires combined auxin and brassinosteroid action in *Arabidopsis* seedlings. *Plant J* 67: 208–217. doi: [10.1111/j.1365-313X.2011.04597.x](https://doi.org/10.1111/j.1365-313X.2011.04597.x) PMID: [21457374](https://pubmed.ncbi.nlm.nih.gov/21457374/)
70. Ichihashi Y, Horiguchi G, Gleissberg S, Tsukaya H (2010) The bHLH transcription factor SPATULA controls final leaf size in *Arabidopsis thaliana*. *Plant Cell Physiol* 51: 252–261. doi: [10.1093/pcp/pcp184](https://doi.org/10.1093/pcp/pcp184) PMID: [20040585](https://pubmed.ncbi.nlm.nih.gov/20040585/)
71. Reymond MC, Brunoud G, Chauvet A, Martinez-Garcia JF, Martin-Magniette ML, et al. (2012) A light-regulated genetic module was recruited to carpel development in *Arabidopsis* following a structural change to SPATULA. *Plant Cell* 24: 2812–2825. doi: [10.1105/tpc.112.097915](https://doi.org/10.1105/tpc.112.097915) PMID: [22851763](https://pubmed.ncbi.nlm.nih.gov/22851763/)
72. Leivar P, Tepperman JM, Cohn MM, Monte E, Al-Sady B, et al. (2012) Dynamic antagonism between phytochromes and PIF family basic helix-loop-helix factors induces selective reciprocal responses to light and shade in a rapidly responsive transcriptional network in *Arabidopsis*. *Plant Cell* 24: 1398–1419. doi: [10.1105/tpc.112.095711](https://doi.org/10.1105/tpc.112.095711) PMID: [22517317](https://pubmed.ncbi.nlm.nih.gov/22517317/)
73. Soy J, Leivar P, Monte E (2014) PIF1 promotes phytochrome-regulated growth under photoperiodic conditions in *Arabidopsis* together with PIF3, PIF4, and PIF5. *J Exp Bot* 65: 2925–2936. doi: [10.1093/jxb/ert465](https://doi.org/10.1093/jxb/ert465) PMID: [24420574](https://pubmed.ncbi.nlm.nih.gov/24420574/)
74. Huq E, Al-Sady B, Hudson M, Kim C, Apel K, et al. (2004) Phytochrome-interacting factor 1 is a critical bHLH regulator of chlorophyll biosynthesis. *Science* 305: 1937–1941. PMID: [15448264](https://pubmed.ncbi.nlm.nih.gov/15448264/)
75. Leivar P, Monte E (2014) PIFs: Systems Integrators in Plant Development. *The Plant Cell Online* 26: 56–78. doi: [10.1105/tpc.113.120857](https://doi.org/10.1105/tpc.113.120857) PMID: [24481072](https://pubmed.ncbi.nlm.nih.gov/24481072/)
76. Monte E, Tepperman JM, Al-Sady B, Kaczorowski KA, Alonso JM, et al. (2004) The phytochrome-interacting transcription factor, PIF3, acts early, selectively, and positively in light-induced chloroplast development. *Proc Natl Acad Sci U S A* 101: 16091–16098. PMID: [15505214](https://pubmed.ncbi.nlm.nih.gov/15505214/)
77. Zhang Y, Mayba O, Pfeiffer A, Shi H, Tepperman JM, et al. (2013) A quartet of PIF bHLH factors provides a transcriptionally centered signaling hub that regulates seedling morphogenesis through differential expression-patterning of shared target genes in *Arabidopsis*. *PLoS Genet* 9: e1003244. doi: [10.1371/journal.pgen.1003244](https://doi.org/10.1371/journal.pgen.1003244) PMID: [23382695](https://pubmed.ncbi.nlm.nih.gov/23382695/)
78. Nozue K, Harmer SL, Maloof JN (2011) Genomic analysis of circadian clock-, light-, and growth-correlated genes reveals PHYTOCHROME-INTERACTING FACTOR5 as a modulator of auxin signaling in *Arabidopsis*. *Plant Physiol* 156: 357–372. doi: [10.1104/pp.111.172684](https://doi.org/10.1104/pp.111.172684) PMID: [21430186](https://pubmed.ncbi.nlm.nih.gov/21430186/)
79. Spartz AK, Ren H, Park MY, Grandt KN, Lee SH, et al. (2014) SAUR inhibition of PP2C-D phosphatases activates plasma membrane H⁺-ATPases to promote cell expansion in *Arabidopsis*. *Plant Cell* 26: 2129–2142. PMID: [24858935](https://pubmed.ncbi.nlm.nih.gov/24858935/)
80. de Wit M, Lorrain S, Fankhauser C (2014) Auxin-mediated plant architectural changes in response to shade and high temperature. *Physiol Plant* 151: 13–24. doi: [10.1111/ppl.12099](https://doi.org/10.1111/ppl.12099) PMID: [24011166](https://pubmed.ncbi.nlm.nih.gov/24011166/)
81. Mashiguchi K, Tanaka K, Sakai T, Sugawara S, Kawaide H, et al. (2011) The main auxin biosynthesis pathway in *Arabidopsis*. *Proc Natl Acad Sci U S A* 108: 18512–18517. doi: [10.1073/pnas.1108434108](https://doi.org/10.1073/pnas.1108434108) PMID: [22025724](https://pubmed.ncbi.nlm.nih.gov/22025724/)

82. Won C, Shen X, Mashiguchi K, Zheng Z, Dai X, et al. (2011) Conversion of tryptophan to indole-3-acetic acid by *TRYPTOPHAN AMINOTRANSFERASES OF ARABIDOPSIS* and *YUCCAs* in *Arabidopsis*. *Proc Natl Acad Sci U S A* 108: 18518–18523. doi: [10.1073/pnas.1108436108](https://doi.org/10.1073/pnas.1108436108) PMID: [22025721](https://pubmed.ncbi.nlm.nih.gov/22025721/)
83. Overvoorde PJ, Okushima Y, Alonso JM, Chan A, Chang C, et al. (2005) Functional genomic analysis of the *AUXIN/INDOLE-3-ACETIC ACID* gene family members in *Arabidopsis thaliana*. *Plant Cell* 17: 3282–3300. PMID: [16284307](https://pubmed.ncbi.nlm.nih.gov/16284307/)
84. Delarue M, Prinsen E, Va H, Onckelen, Caboche M, et al. (1998) *Sur2* mutations of *Arabidopsis thaliana* define a new locus involved in the control of auxin homeostasis. *Plant J* 14: 603–611. PMID: [9675903](https://pubmed.ncbi.nlm.nih.gov/9675903/)
85. Franklin KA, Lee SH, Patel D, Kumar SV, Spartz AK, et al. (2011) Phytochrome-interacting factor 4 (PIF4) regulates auxin biosynthesis at high temperature. *Proc Natl Acad Sci U S A* 108: 20231–20235. doi: [10.1073/pnas.1110682108](https://doi.org/10.1073/pnas.1110682108) PMID: [22123947](https://pubmed.ncbi.nlm.nih.gov/22123947/)
86. Nemhauser JL, Feldman LJ, Zambryski PC (2000) Auxin and ETTIN in *Arabidopsis* gynoecium morphogenesis. *Development* 127: 3877–3888. PMID: [10952886](https://pubmed.ncbi.nlm.nih.gov/10952886/)
87. Foreman J, White J, Graham I, Halliday K, Josse EM (2011) Shedding light on flower development: phytochrome B regulates gynoecium formation in association with the transcription factor *SPATULA*. *Plant Signal Behav* 6: 471–476. PMID: [21364315](https://pubmed.ncbi.nlm.nih.gov/21364315/)
88. Valdes AE, Rizzardi K, Johannesson H, Para A, Sundas-Larsson A, et al. (2011) *Arabidopsis thaliana* *TERMINAL FLOWER2* is involved in light-controlled signalling during seedling photomorphogenesis. *Plant Cell Environ* 35: 1013–1025.
89. Wasternack C, Hause B (2013) Jasmonates: biosynthesis, perception, signal transduction and action in plant stress response, growth and development. An update to the 2007 review in *Annals of Botany*. *Ann Bot* 111: 1021–1058. doi: [10.1093/aob/mct067](https://doi.org/10.1093/aob/mct067) PMID: [23558912](https://pubmed.ncbi.nlm.nih.gov/23558912/)
90. Cerrudo I, Keller MM, Cargnel MD, Demkura PV, de Wit M, et al. (2012) Low red/far-red ratios reduce *Arabidopsis* resistance to *Botrytis cinerea* and jasmonate responses via a COI1-JAZ10-dependent, salicylic acid-independent mechanism. *Plant Physiol* 158: 2042–2052. doi: [10.1104/pp.112.193359](https://doi.org/10.1104/pp.112.193359) PMID: [22371506](https://pubmed.ncbi.nlm.nih.gov/22371506/)
91. Chehab EW, Kim S, Savchenko T, Kliebenstein D, Dehesh K, et al. (2011) Intronic T-DNA insertion renders *Arabidopsis opr3* a conditional jasmonic acid-producing mutant. *Plant Physiol* 156: 770–778. doi: [10.1104/pp.111.174169](https://doi.org/10.1104/pp.111.174169) PMID: [21487047](https://pubmed.ncbi.nlm.nih.gov/21487047/)
92. Fernandez-Calvo P, Chini A, Fernandez-Barbero G, Chico JM, Gimenez-Ibanez S, et al. (2011) The *Arabidopsis* bHLH transcription factors MYC3 and MYC4 are targets of JAZ repressors and act additively with MYC2 in the activation of jasmonate responses. *Plant Cell* 23: 701–715. doi: [10.1105/tpc.110.080788](https://doi.org/10.1105/tpc.110.080788) PMID: [21335373](https://pubmed.ncbi.nlm.nih.gov/21335373/)
93. Cevik V, Kidd BN, Zhang P, Hill C, Kiddle S, et al. (2012) MEDIATOR25 acts as an integrative hub for the regulation of jasmonate-responsive gene expression in *Arabidopsis*. *Plant Physiol* 160: 541–555. doi: [10.1104/pp.112.202697](https://doi.org/10.1104/pp.112.202697) PMID: [22822211](https://pubmed.ncbi.nlm.nih.gov/22822211/)
94. Chen R, Jiang H, Li L, Zhai Q, Qi L, et al. (2012) The *Arabidopsis* mediator subunit MED25 differentially regulates jasmonate and abscisic acid signaling through interacting with the MYC2 and ABI5 transcription factors. *Plant Cell* 24: 2898–2916. doi: [10.1105/tpc.112.098277](https://doi.org/10.1105/tpc.112.098277) PMID: [22822206](https://pubmed.ncbi.nlm.nih.gov/22822206/)
95. Klose C, Buche C, Fernandez AP, Schafer E, Zwick E, et al. (2012) The mediator complex subunit PFT1 interferes with COP1 and HY5 in the regulation of *Arabidopsis* light signaling. *Plant Physiol* 160: 289–307. doi: [10.1104/pp.112.197319](https://doi.org/10.1104/pp.112.197319) PMID: [22760208](https://pubmed.ncbi.nlm.nih.gov/22760208/)
96. Hou X, Lee LY, Xia K, Yan Y, Yu H (2010) DELLAs modulate jasmonate signaling via competitive binding to JAZs. *Dev Cell* 19: 884–894. doi: [10.1016/j.devcel.2010.10.024](https://doi.org/10.1016/j.devcel.2010.10.024) PMID: [21145503](https://pubmed.ncbi.nlm.nih.gov/21145503/)
97. Chen J, Sonobe K, Ogawa N, Masuda S, Nagatani A, et al. (2013) Inhibition of *Arabidopsis* hypocotyl elongation by jasmonates is enhanced under red light in phytochrome B dependent manner. *J Plant Res* 126: 161–168. doi: [10.1007/s10265-012-0509-3](https://doi.org/10.1007/s10265-012-0509-3) PMID: [22825635](https://pubmed.ncbi.nlm.nih.gov/22825635/)
98. Yang D-L, Yao J, Mei C-S, Tong X-H, Zeng L-J, et al. (2012) Plant hormone jasmonate prioritizes defense over growth by interfering with gibberellin signaling cascade. *Proc Natl Acad Sci U S A* 109: E1192–E1200. doi: [10.1073/pnas.1201616109](https://doi.org/10.1073/pnas.1201616109) PMID: [22529386](https://pubmed.ncbi.nlm.nih.gov/22529386/)
99. Noir S, Bomer M, Takahashi N, Ishida T, Tsui TL, et al. (2013) Jasmonate controls leaf growth by repressing cell proliferation and the onset of endoreduplication while maintaining a potential stand-by mode. *Plant Physiol* 161: 1930–1951. doi: [10.1104/pp.113.214908](https://doi.org/10.1104/pp.113.214908) PMID: [23439917](https://pubmed.ncbi.nlm.nih.gov/23439917/)
100. Rieu I, Ruiz-Rivero O, Fernandez-Garcia N, Griffiths J, Powers SJ, et al. (2008) The gibberellin biosynthetic genes *AtGA20ox1* and *AtGA20ox2* act, partially redundantly, to promote growth and development throughout the *Arabidopsis* life cycle. *Plant J* 53: 488–504. PMID: [18069939](https://pubmed.ncbi.nlm.nih.gov/18069939/)
101. Achard P, Cheng H, De Grauwe L, Decat J, Schoutteten H, et al. (2006) Integration of plant responses to environmentally activated phytohormonal signals. *Science* 311: 91–94. PMID: [16400150](https://pubmed.ncbi.nlm.nih.gov/16400150/)

102. Hsu PY, Devisetty UK, Harmer SL (2013) Accurate timekeeping is controlled by a cycling activator in *Arabidopsis*. *eLife* 2: e00473. doi: [10.7554/eLife.00473](https://doi.org/10.7554/eLife.00473) PMID: [23638299](https://pubmed.ncbi.nlm.nih.gov/23638299/)
103. Takase M, Mizoguchi T, Kozuka T, Tsukaya H (2013) The unique function of the *Arabidopsis* circadian clock gene *PRR5* in the regulation of shade avoidance response. *Plant Signal Behav* 8: e23534. doi: [10.4161/psb.23534](https://doi.org/10.4161/psb.23534) PMID: [23333981](https://pubmed.ncbi.nlm.nih.gov/23333981/)
104. Schachtman D, Schroeder J, Lucas W, Anderson J, Gaber R (1992) Expression of an inward-rectifying potassium channel by the *Arabidopsis* *KAT1* cDNA. *Science* 258: 1654–1658. PMID: [8966547](https://pubmed.ncbi.nlm.nih.gov/8966547/)
105. Nakamura RL, McKendree WL Jr., Hirsch RE, Sedbrook JC, Gaber RF, et al. (1995) Expression of an *Arabidopsis* potassium channel gene in guard cells. *Plant Physiol* 109: 371–374. PMID: [7480337](https://pubmed.ncbi.nlm.nih.gov/7480337/)
106. Sutter J-U, Sieben C, Hartel A, Eisenach C, Thiel G, et al. (2007) Abscisic acid triggers the endocytosis of the *Arabidopsis* *KAT1* K⁺ channel and its recycling to the plasma membrane. *Current Biol* 17: 1396–1402. PMID: [17683934](https://pubmed.ncbi.nlm.nih.gov/17683934/)
107. Eisenach C, Chen Z-H, Grefen C, Blatt MR (2011) The trafficking protein *SYP121* of *Arabidopsis* connects programmed stomatal closure and K⁺ channel activity with vegetative growth. *Plant J* 69: 241–251. doi: [10.1111/j.1365-313X.2011.04786.x](https://doi.org/10.1111/j.1365-313X.2011.04786.x) PMID: [21914010](https://pubmed.ncbi.nlm.nih.gov/21914010/)
108. Abel S, Nguyen MD, Theologis A (1995) The PS-IAA4/5-like family of early auxin-inducible mRNAs in *Arabidopsis thaliana*. *J Mol Biol* 251: 533–549. PMID: [7658471](https://pubmed.ncbi.nlm.nih.gov/7658471/)
109. Philippark K, Ivashikina N, Ache P, Christian M, Luthen H, et al. (2004) Auxin activates *KAT1* and *KAT2*, two K⁺-channel genes expressed in seedlings of *Arabidopsis thaliana*. *Plant J* 37: 815–827. PMID: [14996216](https://pubmed.ncbi.nlm.nih.gov/14996216/)
110. Putterill J, Robson F, Lee K, Simon R, Coupland G (1995) The *CONSTANS* gene of *Arabidopsis* promotes flowering and encodes a protein showing similarities to zinc finger transcription factors. *Cell* 80: 847–857. PMID: [7697715](https://pubmed.ncbi.nlm.nih.gov/7697715/)
111. Park DH, Somers DE, Kim YS, Choy YH, Lim HK, et al. (1999) Control of circadian rhythms and photoperiodic flowering by the *Arabidopsis* *GIGANTEA* gene. *Science* 285: 1579–1582. PMID: [10477524](https://pubmed.ncbi.nlm.nih.gov/10477524/)
112. Fowler S, Lee K, Onouchi H, Samach A, Richardson K, et al. (1999) *GIGANTEA*: a circadian clock-controlled gene that regulates photoperiodic flowering in *Arabidopsis* and encodes a protein with several possible membrane-spanning domains. *EMBO J* 18: 4679–4688. PMID: [10469647](https://pubmed.ncbi.nlm.nih.gov/10469647/)
113. Song YH, Ito S, Imaizumi T (2013) Flowering time regulation: photoperiod- and temperature-sensing in leaves. *Trends Plant Sci* 18: 575–583. doi: [10.1016/j.tplants.2013.05.003](https://doi.org/10.1016/j.tplants.2013.05.003) PMID: [23790253](https://pubmed.ncbi.nlm.nih.gov/23790253/)
114. Holm M, Ma LG, Qu LJ, Deng XW (2002) Two interacting bZIP proteins are direct targets of COP1-mediated control of light-dependent gene expression in *Arabidopsis*. *Genes Dev* 16: 1247–1259. PMID: [12023303](https://pubmed.ncbi.nlm.nih.gov/12023303/)
115. Kumar SV, Lucyshyn D, Jaeger KE, Alos E, Alvey E, et al. (2012) Transcription factor PIF4 controls the thermosensory activation of flowering. *Nature* 484: 242–245. doi: [10.1038/nature10928](https://doi.org/10.1038/nature10928) PMID: [22437497](https://pubmed.ncbi.nlm.nih.gov/22437497/)
116. Rawat R, Takahashi N, Hsu PY, Jones MA, Schwartz J, et al. (2011) *REVEILLE8* and *PSEUDO-RESPONSE REGULATOR5* form a negative feedback loop within the *Arabidopsis* circadian clock. *PLoS Genet* 7: e1001350. doi: [10.1371/journal.pgen.1001350](https://doi.org/10.1371/journal.pgen.1001350) PMID: [21483796](https://pubmed.ncbi.nlm.nih.gov/21483796/)
117. Leopold AC (1958) Auxin uses in the control of flowering and fruiting. *Annu Rev Plant Physiol* 9: 281–310.
118. Gangappa SN, Chattopadhyay S (2010) *MYC2*, a bHLH transcription factor, modulates the adult phenotype of *SPA1*. *Plant Signal Behav* 5: 1650–1652. doi: [10.4161/psb.5.12.13981](https://doi.org/10.4161/psb.5.12.13981) PMID: [21512327](https://pubmed.ncbi.nlm.nih.gov/21512327/)
119. Makkena S, Lamb RS (2013) The bHLH transcription factor *SPATULA* is a key regulator of organ size in *Arabidopsis thaliana*. *Plant Signal Behav* 8: e24140. doi: [10.4161/psb.24140](https://doi.org/10.4161/psb.24140) PMID: [23470719](https://pubmed.ncbi.nlm.nih.gov/23470719/)
120. Cifuentes-Esquivel N, Bou-Torrent J, Galstyan A, Gallerni M, Sessa G, et al. (2013) The bHLH proteins *BEE* and *BIM* positively modulate the shade avoidance syndrome in *Arabidopsis* seedlings. *Plant J* 75: 989–1002. doi: [10.1111/tpj.12264](https://doi.org/10.1111/tpj.12264) PMID: [23763263](https://pubmed.ncbi.nlm.nih.gov/23763263/)
121. Friedrichsen DM, Nemhauser J, Muramitsu T, Maloof JN, Alonso J, et al. (2002) Three redundant brassinosteroid early response genes encode putative bHLH transcription factors required for normal growth. *Genetics* 162: 1445–1456. PMID: [12454087](https://pubmed.ncbi.nlm.nih.gov/12454087/)
122. Filiault DL, Maloof JN (2012) A genome-wide association study identifies variants underlying the *Arabidopsis thaliana* shade avoidance response. *PLoS Genet* 8: e1002589. doi: [10.1371/journal.pgen.1002589](https://doi.org/10.1371/journal.pgen.1002589) PMID: [22438834](https://pubmed.ncbi.nlm.nih.gov/22438834/)
123. Kumar R, Ichihashi Y, Kimura S, Chitwood DH, Headland LR, et al. (2012) A high-throughput method for Illumina RNA-Seq library preparation. *Front Plant Sci* 3: 202. doi: [10.3389/fpls.2012.00202](https://doi.org/10.3389/fpls.2012.00202) PMID: [22973283](https://pubmed.ncbi.nlm.nih.gov/22973283/)

124. Nozue K, Covington MF, Duek PD, Lorrain S, Fankhauser C, et al. (2007) Rhythmic growth explained by coincidence between internal and external cues. *Nature* 448: 358–361. PMID: [17589502](#)
125. Abramoff MD, Magalhaes PJ, Ram SJ (2004) Image processing with ImageJ. *Biophotonics International* 11: 36–42.
126. Trapnell C, Pachter L, Salzberg S (2009) TopHat: discovering splice junctions with RNA-Seq. *Bioinformatics* 25: 1105–1111. doi: [10.1093/bioinformatics/btp120](#) PMID: [19289445](#)
127. Robinson MD, McCarthy DJ, Smyth GK (2010) edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26: 139–140. doi: [10.1093/bioinformatics/btp616](#) PMID: [19910308](#)
128. R_Development_Core_Team (2005) R: A language and environment for statistical computing. R Foundation for Statistical Computing. Vienna, Austria.
129. Young MD, Wakefield MJ, Smyth GK, Oshlack A (2010) Gene ontology analysis for RNA-seq: accounting for selection bias. *Genome Biol* 11: R14. doi: [10.1186/gb-2010-11-2-r14](#) PMID: [20132535](#)
130. Nemhauser JL, Hong F, Chory J (2006) Different plant hormones regulate similar processes through largely nonoverlapping transcriptional responses. *Cell* 126: 467. PMID: [16901781](#)
131. Zentella R, Zhang Z-L, Park M, Thomas SG, Endo A, et al. (2007) Global analysis of DELLA direct targets in early gibberellin signaling in *Arabidopsis*. *Plant Cell* 19: 3037–3057. PMID: [17933900](#)
132. Carbon S, Ireland A, Mungall CJ, Shu S, Marshall B, et al. (2009) AmiGO: online access to ontology and annotation data. *Bioinformatics* 25: 288–289. doi: [10.1093/bioinformatics/btn615](#) PMID: [19033274](#)
133. Bates D, Maechler M, Bolker B, Walker S (2014) lme4: Linear mixed-effects models using Eigen and S4. <http://CRAN.R-project.org/package=lme4>.
134. Kuznetsova A, Brockhoff PB, Christensen RHB (2014) lmerTest: Tests for random and fixed effects for linear mixed effect models (lmer objects of lme4 package). <http://CRAN.R-project.org/package=lmerTest>.
135. Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate—a practical and powerful approach to multiple testing. *J Roy Statist Soc Ser B* 57: 289–300.
136. Winter D, Vinegar B, Nahal H, Ammar R, Wilson GV, et al. (2007) An "Electronic Fluorescent Pictograph Browser" for exploring and analyzing large-scale biological data sets. *PLoS ONE* 2: e718. PMID: [17684564](#)