

Seyed Mohammad Masoud
Hojjati (MD)¹
Amin Zarghami *²
Seyed Ali Hojjati (MD)³
Mobina Baes (MD)⁴

1. Department of Neurology, Babol University of Medical Sciences, Babol, Iran.
2. Student Research Committee, Babol University of Medical Sciences, Babol, Iran.
3. Shahid Beheshti University of Medical Sciences, Tehran, Iran.
4. Babol University of Medical Sciences, Babol, Iran.

* **Correspondence:**

Amin Zarghami, Student Research Committee, Babol University of Medical Sciences, Babol, Iran.

E-mail:

aminzarghami18@yahoo.com

Tel: 0098 11 32190971

Fax: 0098 11 32238284

Received: 11 April 2014

Revised: 7 June 2015

Accepted: 22 June 2015

Optic neuritis, the most common initial presenting manifestation of multiple sclerosis in northern Iran

Abstract

Background: Multiple sclerosis (MS) is an inflammatory and demyelinating disease of central nervous system (CNS). The aim of the present study was to determine the type and the frequency of initial presenting symptoms in patients with MS and their relation with demographic characteristics in Babol, northern Iran.

Methods: All patients of this study were recruited over a ten year period from 2002 to 2012 from single neurologic clinic. Diagnosis of MS was confirmed according to the McDonald criteria, demographic and clinical features. Then, all the clinical findings and demographic variables including: age, sex, marital status, age at onset, education, place of residence, disease duration, initiation pattern of disease have been collected. Expanded Disability Status Scale (EDSS) was used for the evaluation of disability at the onset of disease. Data analysis was performed by chi-square test.

Results: A total of 263 consecutive MS patients with the age range of 17 to 61 yr were examined. Optic neuritis was the most prevalent initial presenting symptom in 123 (46.8%) patients followed by sensory disturbances as the second common presenting symptom of MS. Significant difference was found between patients with or without optic neuritis and the onset age of the disease and EDSS ($p < 0.001$). The mean EDSS score at the time of initial presentation was 1.67 ± 0.77 .

Conclusion: The findings of this study indicated that optic neuritis is the most prevalent initial presentation of MS in the geographic region of northern Iran. In patients less than 30 years, development of visual disturbances justifies neurologic examination.

Keywords: Multiple sclerosis, Optic neuritis, Iran.

Citation:

Masoud Hojjati SM, Zarghami A, et al. Optic neuritis, the most common initial presenting manifestation of multiple sclerosis in northern Iran. *Caspian J Intern Med* 2015; 6(3):151-155.

Caspian J Intern Med 2015; 6(3):151-155

Multiple sclerosis (MS) is a chronic inflammatory demyelinating disease that attacks myelinated axons in the central nervous system, damaging the myelin and the axon in variable degrees and producing significant progressive physical disability in patients (1). The hallmark of MS is symptomatic episodes occurs in months or years apart and involves various anatomic locations. The etiology of MS is unknown; but it seems the combination of genetic susceptibility and environmental factors (eg. viral infection, low vitamin D levels, etc.). Has a contribution in the development of MS. Geographic variation in the incidence of MS supports the probability that environmental factors are involved in the etiology (2-3). More than 2.1 million people are affected by MS worldwide (4). MS is more common in women and the estimated incidence for the female to male ratio reached to 2.3 in 2000 (5). Although MS can occur in any age groups, it usually diagnosed in persons aged 15-45 years (6).

The disease seen in all parts of the world and in all races; but rates vary widely. Based on the Kurtzke classification, Iran is located in a low-risk zone for MS (7). However, recent studies on the prevalence of MS came up with different results and indicated a higher prevalence rate in our country (8-9). In previous studies in various countries, different frequencies reported for initial presenting symptoms of MS. Concerning the sites of involvement, there have been several reports that in eastern MS patients more commonly represent clinical evidence of more involvement of the optic nerves and spinal cord than Caucasians (10). Optic neuritis is one the common presenting symptoms of MS which developed due to the involvement of optic nerve in the process of pathogenicity of the MS. This complication causes the deficit in visual acuity mainly in one of the eye. Depending on which part of the nerve is affected and the area of demyelination, manifestations of MS can vary apparently from benign condition to incapacitating clinical symptoms (11).

Previous studies in recent decades indicated higher frequency of optic neuritis in Iranian patients with MS as compared with other countries (12, 13). Nonetheless, the data regarding the associated factors as well as the types and the frequency of the initial presenting manifestation of MS in Iran is scarce. Thus, the aim of this study was to assess the type and frequency of initial presenting symptoms of MS with regard to age at disease onset, sex and demographic features in geographic region of Babol, Northern Iran.

Methods

In this cross-sectional study we reviewed all the 263 consecutive MS patients who had been admitted to a Multiple Sclerosis Center from 2002 to 2012 in Babol, northern Iran. Although the majority of the registered patients are residents of Babol, a city in North of Iran but a number of patients were referred from other cities of Mazandaran. The McDonald criteria applied in the present study and data confirmed by a neurologist in order to diagnose MS in all the consecutive patients (14). The protocol of the study was approved by the ethical board of the research council of Babol University of medical sciences. Data were collected for demographic and clinical features of MS such as the duration of disease, age at onset of first symptom, family history, and associated factors of optic neuritis. The authors were committed to the ethical

consideration during the stages of the study and follow the Declaration of Helsinki.

After receiving the informed consent, all patients underwent a standard neurologic examination. The main manifestations assessed by the investigators based on impairment of sensory disturbance, impairment of cranial nerves, impairment of motor pathways, cerebellar symptoms (tremor, dysarthria etc.), impairment of visual pathways, brain stem function, impairment of bladder or sexual dysfunction. The diagnosis of optic neuritis was made by the recommendations for clinical practice of the International Federation of Clinical Neurophysiology (15). All patients underwent a complete ophthalmic examination. Neurologists diagnosed patients' optic neuritis according to the standard methods. We used Kurtzke Expanded Disability Status Scale (EDSS) for the evaluation of disability in patients with MS (16). Data was analyzed by SPSS Version 18 (SPSS Inc., Chicago, Ill., USA) and chi-square test for comparison of the proportions. A p-value less than 0.05 was considered significant.

Results

A total of 263 consecutive patients with MS were examined (28.5% men; 71.5% women). The mean age of participants was 34.28 ± 9.47 (age range, 17 to 61 yr). The female to male ratio was 2.5. The course was relapsing-remitting (RR) in 91% of patients and primary progressive (PP) in 9%. Optic neuritis with the prevalence of 46.8% was the most prevalent initial presenting symptoms in our patients with MS. Figure 1 shows the prevalence of different initial presenting symptoms in our patients with MS.




Figure 1: Frequency of initial symptoms in 263 MS patients

Monosymptomatic initiation pattern was seen in (161.263) 61.2% of the cases. The mean EDSS at the onset of disease was 1.67 ± 0.77 . The distribution of demographic information and features of the disease among MS patients presented with optic neuritis and without it are presented in

Table 1. Significant difference was found between patients with or without optic neuritis and onset age of the disease ($p < 0.001$); however in other variables such as gender, marital status, education, place of live and initiation pattern did not indicated any significant differences.

Table 1: Demographic and clinical profile of 263 patients with Multiple Sclerosis with or without optic neuritis

Variables	Optic neuritis		Odds Ratio	95% Confidence interval		P value
	With (n=160)	Without (n=103)		Lower	Upper	
Gender						
Male	39 (24.4%)	36 (35%)	1.66	0.96	2.86	0.06
Female	121 (75.6%)	67 (65%)				
Marital status*						
Single	47 (29.7%)	22 (21.4%)	0.64	0.35	1.14	0.13
married	111 (70.3%)	81 (78.6%)				
Place of live						
Urban	93 (58.1%)	52 (50.5%)	0.73	0.44	1.20	0.22
Rural	67 (41.9%)	51 (49.5%)				
Education						
Lower diploma	98 (61.3%)	66 (64.1%)	1.12	0.67	1.88	0.66
Higher diploma	62 (38.8%)	37 (35.9%)				
Age at onset						
≤30	130 (81.3%)	63 (61.2%)	0.36	0.20	0.63	<0.001
>30	30 (18.8%)	40 (38.8%)				
Initiation pattern						
Monosymptomatic	101 (63.1%)	60 (58.3%)	0.815	0.491	1.35	0.42
Polysymptomatic	59 (36.9%)	43 (41.7%)				

* The data of two cases was missing.

Discussion

According to our findings, optic neuritis was observed in 46.8% of the patients with MS as the most prevalent initial presenting symptom in our study population. But in contrast to similar studies in Iran, our data placed in the row of high frequencies. Saadatnia et al. reported that the most common symptom at the first episode was sensory disturbance (51.1%), followed by optic neuritis (47%) (13).

In our study 61.2% of the patients with optic neuritis - as a presenting symptom- had monosymptomatic initiation pattern of the disease. There were limited studies regarding to optic involvement in related to other initiation patterns of the disease. Several reports have demonstrated that monosymptomatic onset occurs in adults than in childhood

(17, 18). Ruggieri et al revealed that brain stem syndromes or cerebellar ataxia were the most common onset symptom in monosymptomatic children (19). While in a population-based study in Isfahan-Iran the most common onset symptom, optic neuritis (31%) was the most frequent initial presentation in early onset MS and polysymptomatic onset was more common in early-onset patients (35.6%) compared to adult-onset patients (12%) (17). In our study optic neuritis involvement in patients with the age of lower than 30 at onset of the disease was significantly more than the patients more than 30 years old. Ashtari et al. investigated the characteristics of early-onset (EOMS) Multiple Sclerosis in comparison to adult-onset (AOMS) in Isfahan, Iran. They revealed that the most common presenting symptom was

optic neuritis in the EOMS patients and paresthesia in AOMS despite of no significant difference in terms of major clinical manifestations (20).

However, it is assumed that in different populations with different social and individual characteristics, the distribution of MS might differ which leads to various reasons like genetic, environment and geographical location (21, 22). In a study of comparing the role of ethnicity on the characteristics of MS among Persians and Iranian- Arab population, the authors indicated that optic nerve(40%) involvement was the most frequent features of initial symptoms which followed by sensory problems in Persians. On the other hand, motor deficit and cerebellar symptoms are more prevalent among Iranian- Arabs than Persians(19.1% and 18.2% versus 8.5% and 9.8% respectively) (23). While reports from several Arab countries revealed that motor dysfunction was the most prevalent feature of initial presentation (24, 25). For example, Sidhom et al. in Tunisia reported motor symptoms (28%) followed by optic neuritis (20%) as the most frequent isolated onset symptoms (26). According to the close neighboring of these countries, the differences are the matter of controversy. These contrasts attributed to the socioeconomical differences rather than genetic and ethnicity(27). Reports from Latin America also revealed that motor dysfunction followed by optic neuritis and sensory dysfunction were the most prevalent initiating symptoms (28). In a study in Bengal, India based on Poser's criteria the Visual impairment (53.33%), weakness of limbs (31.11%) and sensory paresthesia (20%) were the common presenting symptoms (29). Studies from China and eastern Asia were revealed controversial frequencies of sensory disturbance, visual loss and optic neuritis (30, 31). The results of this study should be considered with limitations. Since the study population has been collected over a period of 10 years, therefore the spectrum of clinical features and the nature of MS are expected to be changed compared with new cases. Nonetheless, this study has are strength with regard to diagnostic method using a single diagnostic criteria, as well as patient characteristics which have been recruited from single neurologic clinic. The findings of this study is clinically important. Because, vitamin D deficiency has been suspected as a possible environmental risk factor of MS and vitamin D deficiency is highly frequent in the geographic locale of the study subjects particularly the young women (32). Several studies have shown an

association between vitamin D deficiency and several skeletal and nonskeletal conditions (33-35). This issue justifies a study regarding to MS and vitamin D deficiency.

In conclusion, this study indicated that optic neuritis is the most prevalent initial presenting symptoms of MS in the geographic region of Babol. Based on the results of the present study, development of ophthalmic symptoms in young patients justify neurological evaluation particularly for MS.

Acknowledgments

We would like to appreciate Dr. Ali Bijani for his contribution in this study.

Funding: This study was funded by the Chancellor for Research and Technology (Grant no: 2834).

Conflict of interest: Authors declared no conflict of interest.

References

1. Parfenov V, Schlupe M, Du Pasquier R. Assessing risks of multiple sclerosis therapies. *J Neurol Sci* 2013; 332: 59-65.
2. Kakalacheva K, Lunemann JD. Environmental triggers of multiple sclerosis. *FEBS Lett* 2011; 585: 3724-9.
3. Noseworthy JH, Lucchinetti C, Rodriguez M, Weinshenker BG. Multiple sclerosis. *N Engl J Med* 2000; 343: 938-52.
4. Noonan CW, Williamson DM, Henry JP, et al. The prevalence of multiple sclerosis in 3 US communities. *Prev Chronic Dis* 2010; 7: A12.
5. Kampman MT, Brustad M. Vitamin D: a candidate for the environmental effect in multiple sclerosis - observations from Norway. *Neuroepidemiology* 2008; 30: 140-6.
6. Kurtzke JF. Epidemiology in multiple sclerosis: a pilgrim's progress. *Brain* 2013; 136: 2904-17.
7. Kurtzke JF. Geographic distribution of multiple sclerosis: An update with special reference to Europe and the Mediterranean region. *Acta Neurol Scand* 1980; 62: 65-80.
8. Etemadifar M, Sajjadi S, Nasr Z, et al. Epidemiology of multiple sclerosis in Iran: a systematic review. *Eur Neurol* 2013; 70: 356-63.
9. Sahraian MA, Khorramnia S, Ebrahim MM, et al. Multiple sclerosis in Iran: a demographic study of 8,000

- patients and changes over time. *Eur Neurol* 2010; 64: 331-6.
10. Kalanie H, Kholghie Y, Shamsai GR, Ghorbani M. Opticospinal multiple sclerosis in Iran. *J Neurol Sci* 2009; 276: 130-2.
 11. Cruccu G, Biasiotto A, Di Rezze S, et al. Trigeminal neuralgia and pain related to multiple sclerosis. *Pain* 2009; 143: 186-91.
 12. Danesh-Sani SA, Rahimdoost A, Soltani M, et al. Clinical assessment of orofacial manifestations in 500 patients with multiple sclerosis. *J Oral Maxillofac Surg* 2013; 71: 290-4.
 13. Saadatnia M, Etemadifar M, Maghzi AH. Multiple sclerosis in Isfahan, Iran. *Int Rev Neurobiol* 2007; 79: 357-75.
 14. McDonald WI, Compston A, Edan G, et al. Recommended diagnostic criteria for multiple sclerosis: guidelines from the International Panel on the diagnosis of multiple sclerosis. *Ann Neurol* 2001; 50:121-7.
 15. Recommendations for the practice of clinical neurophysiology: guidelines of the International Federation of Clinical Neurophysiology. *Electroencephalogr Clin Neurophysiol Suppl* 1999; 52: 1-304.
 16. Hojjati SM, Hojjati SA, Baes M, Bijani A. Relation between EDSS and monosymptomatic or polysymptomatic onset in clinical manifestations of multiple sclerosis in Babol, northern Iran. *Caspian J Intern Med* 2014; 5: 5-8.
 17. Etemadifar M, Janghorbani M, Shaygannejad V, Ashtari F. Prevalence of multiple sclerosis in Isfahan, Iran. *Neuroepidemiology* 2006; 27: 39-44.
 18. Liguori M, Marrosu MG, Pugliatti M, et al. Age at onset in multiple sclerosis. *Neurol Sci* 2000; 21: S825-9.
 19. Ruggieri M, Polizzi A, Pavone L, Grimaldi LM. Multiple sclerosis in children under 6 years of age. *Neurology* 1999; 53: 478-84.
 20. Ashtari F, Shaygannejad V, Farajzadegan Z, Amin A. Does early-onset multiple sclerosis differ from adult-onset form in Iranian people. *J Res Med Sci* 2010; 15: 94-9.
 21. Ebers GC. Environmental factors and multiple sclerosis. *Lancet Neurol* 2008; 7: 268-77.
 22. Rosati G. The prevalence of multiple sclerosis in the world: an update. *Neurol Sci* 2001; 22: 117-39.
 23. Sharafaddinzadeh N, Moghtaderi A, Majdinasab N, et al. The influence of ethnicity on the characteristics of multiple sclerosis: a local population study between Persians and Arabs. *Clin Neurol Neurosurg* 2013; 115: 1271-5.
 24. Benamer HT, Ahmed ES, Al-Din AS, Grosset DG. Frequency and clinical patterns of multiple sclerosis in Arab countries: a systematic review. *J Neurol Sci* 2009; 278: 1-4.
 25. Yamout B, Barada W, Tohme RA, et al. Clinical characteristics of multiple sclerosis in Lebanon. *J Neurol Sci* 2008; 270: 88-93.
 26. Sidhom Y, Damak M, Riahi A, et al. Clinical features and disability progression in multiple sclerosis in Tunisia: do we really have a more aggressive disease course? *J Neurol Sci* 2014; 343: 110-4.
 27. Fereidan-Esfahani M, Abtahi SH, Nourian SM, Etemadifar M. Multiple sclerosis among Iranian-Arabs and Persians of Khuzestan: is there any ethnic difference? *Clin Neurol Neurosurg* 2013; 115: 2309-10.
 28. Ojeda E, Diaz-Cortes D, Rosales D, Duarte-Rey C, Anaya JM, Rojas-Villarraga A. Prevalence and clinical features of multiple sclerosis in Latin America. *Clin Neurol Neurosurg* 2013; 115: 381-7.
 29. Gangopadhyay G, Das SK, Sarda P, et al. Clinical profile of multiple sclerosis in Bengal. *Neurol India* 1999; 47: 18-21.
 30. Itoh T, Aizawa H, Hashimoto K, et al. Prevalence of multiple sclerosis in Asahikawa, a city in northern Japan. *J Neurol Sci* 2003; 214: 7-9.
 31. Cheng XJ, Xu LZ. Review of the diagnosis and clinical features of multiple sclerosis in China. *Neurosci Bull* 2009; 25: 38-42.
 32. Heidari B, Heidari P, Hajian Tilaki K. High prevalence of vitamin D deficiency in women presenting to rheumatology clinic in North of Iran: An inverse relation with age. *J Womens Health Care* 2013; 2: 123.
 33. Heidari B, Haji Mirghassemi MB. Seasonal variations in serum vitamin D according to age and sex. *Caspian J Intern Med* 2012; 3: 535-40.
 34. Heidari B, Heidari P, Tilaki KH. Relationship between unexplained arthralgia and vitamin D deficiency: a case control study. *Acta Med Iran* 2014; 52: 400-5.
 35. Monadi M, Heidari B, Asgharpour M, et al. Relationship between serum vitamin D and forced expiratory volume in patients with chronic obstructive pulmonary disease (COPD). *Caspian J Intern Med* 2012; 3: 451-5.