

Rostral anterior cingulate cortex dysfunction during error processing in schizophrenia

Kristin R. Laurens,^{1,2} Elton T. C. Ngan,¹ Alan T. Bates,² Kent A. Kiehl³ and Peter F. Liddle²

¹Department of Psychiatry, University of British Columbia, Vancouver, BC, Canada, ²Division of Psychiatry, School of Community Health Sciences, University of Nottingham, Nottingham, UK and ³Department of Psychiatry, Institute of Living, Yale University School of Medicine, New Haven, CT, USA

Correspondence to: Peter F. Liddle, Division of Psychiatry, School of Community Health Sciences, A Floor, South Block, Queen's Medical Centre, Nottingham NG7 2UH, UK
E-mail: peter.liddle@nottingham.ac.uk

Summary

Previous research has demonstrated that patients with schizophrenia have an impaired ability to monitor erroneous responses to stimuli internally. Event-related potential (ERP) studies of error-eliciting tasks indicate that, in healthy adults, the commission of an erroneous response is associated with a fronto-centrally distributed negative voltage component termed the error negativity (Ne) or error-related negativity (ERN). In patients with schizophrenia, the Ne/ERN elicited by errors of commission (EoC) is reduced in amplitude compared with that elicited in healthy participants. Functional MRI (fMRI) studies and source localization analyses of ERP data in healthy participants suggest that EoC are associated with activity in the rostral anterior cingulate cortex (ACC). Using event-related fMRI, we examined the brain activity associated with EoC in a group of 10 patients with schizophrenia and 16 matched healthy participants. Patients were stable, partially remitted, medicated out-patients recruited from the community. Participants performed a Go/NoGo task variant that

was shown previously to elicit a reduced Ne/ERN during EoC in patients with schizophrenia relative to healthy participants, as well as robust rostral ACC activation during EoC in healthy participants. Patients with schizophrenia were characterized by relative underactivity in the rostral ACC compared with healthy participants. There was also evidence for more widespread underactivity in the limbic system. In contrast to these regions of relative hypoactivity, patients with schizophrenia demonstrated hyperactivity relative to healthy participants in bilateral parietal cortex during both EoC and correctly rejected NoGo trials. Our results are consistent with previous ERP research demonstrating functional abnormalities during error processing in schizophrenia. In light of the role of the rostral ACC and other limbic structures in mediating affective and motivational behaviour, our results suggest there may be a disturbed affective or motivational response to the commission of errors in schizophrenia.

Keywords: schizophrenia; anterior cingulate cortex; error processing; limbic system; event-related functional MRI

Abbreviations: ACC = anterior cingulate cortex; CR = correct rejections; EoC = errors of commission; ERN = error-related negativity; ERP = event-related potential; fMRI = functional MRI; Ne = error negativity; Pe = error positivity; ROI = region of interest; RT = reaction time

Introduction

Schizophrenia is characterized by disordered monitoring and regulation of self-generated thoughts and behaviour (Frith and Done, 1989; Leudar *et al.*, 1994; Mlakar *et al.*, 1994; Stirling *et al.*, 1998, 2001; Johns *et al.*, 2001). Research suggests that an impaired ability to monitor internally error responses to stimuli contributes to self-monitoring problems (Malenka *et al.*, 1982, 1986; Frith and Done, 1989). Evidence for a functional abnormality associated with error monitoring in schizophrenia derives primarily from event-related poten-

tial (ERP) research investigating a fronto-central negative voltage component termed error negativity (Ne; Falkenstein, *et al.*, 1990, 1991) or error-related negativity (ERN; Gehring *et al.*, 1990, 1993). The Ne/ERN peaks ~50–150 ms after the commission of an erroneous response during tasks that necessitate speeded and accurate response choices, thus providing a physiological marker of internal error monitoring (for a review see Falkenstein *et al.*, 2000). The Ne/ERN is elicited in situations where participants know the correct

answer but fail to execute the correct response (Dehaene *et al.*, 1994), and decreases in amplitude as the participant's confidence in having committed an error decreases (e.g. in tasks in which the quality of the stimulus has been degraded by reducing the contrast between the stimulus and its background; Scheffers and Coles, 2000). Several ERP studies have demonstrated that the Ne/ERN is attenuated in patients with schizophrenia compared with healthy adults (Kopp and Rist, 1999; Mathalon *et al.*, 2002), even in paradigms in which the correct response is readily apparent and errors are easily identifiable (Bates *et al.*, 2002). These results are consistent with the hypothesis that error responses are processed abnormally in schizophrenia.

The application of error-eliciting tasks during functional MRI (fMRI) provides further neurophysiological evidence of disturbed brain function in patients with schizophrenia during error commission. Converging evidence from fMRI research (e.g. Carter *et al.*, 1998; Kiehl *et al.*, 2000; Ullsperger and von Cramon, 2001) and dipole localization analyses of dense array ERP data (e.g. Dehaene *et al.*, 1994; Miltner *et al.*, 1997; Holroyd *et al.*, 1998; Luu *et al.*, 2000b) in healthy individuals suggests that the commission of errors (and the Ne/ERN) is critically associated with activity in the anterior cingulate cortex (ACC). Carter *et al.* (2001) recently reported attenuation of the haemodynamic response during errors of commission (EoC) in patients with schizophrenia compared with healthy participants in the ACC. Taken together, the ERP and fMRI results imply that impaired error monitoring in schizophrenia relates to dysfunction in ACC. However, the nature of the error-related processes purported to underlie the Ne/ERN and ACC activity have been the subject of debate, leaving open the question as to the root of abnormal error monitoring in schizophrenia.

Early proposals that the Ne/ERN reflects the activity of a rapid, pre-conscious error detection system that compares and detects mismatch between representations of the intended response and the actual response (Gehring *et al.*, 1993; Bernstein *et al.*, 1995; Falkenstein *et al.*, 2000; Scheffers and Coles, 2000) have been challenged. fMRI evidence that tasks involving strong response competition elicit ACC activation irrespective of response accuracy led others to hypothesize that the ACC functions to detect conflict between incompatible potential responses rather than overt errors (Carter *et al.*, 1998; MacDonald *et al.*, 2000; Botvinick *et al.*, 2001). Other theorists argue that error monitoring incorporates processes related to motivation and/or affective processing of error responses (Dikman and Allen, 2000; Luu *et al.*, 2000a; Vidal *et al.*, 2000). For example, Luu *et al.* (2000a) demonstrated that the amplitude of the Ne/ERN was larger in participants who reported a propensity to experience negative affect on personality assessment scales than in participants without this propensity, and, further, that the amplitude of the Ne/ERN decreased in these participants as they affectively disengaged from the task. Also consistent with the proposal that the error response incorporates a motivational or affective processing component are a number

of fMRI studies that localized activity associated with EoC to the rostral ACC (Kiehl *et al.*, 2000; Braver *et al.*, 2001; Menon *et al.*, 2001). Activity in the rostral ACC region during errors was dissociated from activity in more superior, caudal ACC that was elicited during both response inhibition and target detection processing that included a degree of response competition. Data from a recent study reporting dipole localization of error-related ERP components are also consistent with the idea that caudal ACC activity may reflect a conflict detection component to error monitoring that is dissociated from an affective component mediated by activity in rostral ACC (Van Veen and Carter, 2002).

Structural and functional dissociation of the ACC into rostral and caudal subregions has been described on the basis of convergent evidence from cytoarchitectural, lesion, electrophysiological and neuroimaging data (Devinsky *et al.*, 1995; Bush *et al.*, 2000). The caudal ACC, termed the 'cognitive' subdivision, appears to be responsible for mediating attention and executive functions such as the detection of response conflict via strong reciprocal interconnections with dorsolateral prefrontal cortex, parietal cortex, and premotor and supplementary motor areas. The 'affective' subdivision in the rostral ACC has connections to limbic and paralimbic areas including the amygdala and hippocampus, and appears primarily involved in assessing the salience of emotional and motivational information, and in regulating emotional responses (for a review see Bush *et al.*, 2000). Functional subspecialization of the ACC is also supported by neuroimaging studies that demonstrate reciprocal suppression of rostral ACC and enhancement of caudal ACC activity during attentionally demanding cognitive tasks (e.g. divided attention, sequential learning, working memory and response competition tasks; see review by Drevets and Raichle, 1998), as well as the converse condition of suppressed caudal ACC and enhanced rostral ACC activity during tasks employing emotional stimuli (e.g. Whalen *et al.*, 1998). Carter *et al.* (2001) demonstrated that healthy participants, but not patients with schizophrenia, showed an increase in activity in caudal ACC during the commission of errors in a task that elicited strong response conflict. This result suggests that error monitoring deficits in schizophrenia may be partially associated with a more generalized dysfunction in the detection of response conflict. However, evidence that the amplitude of the Ne/ERN is modulated by the affective or motivational response to errors (Dikman and Allen, 2000; Luu *et al.*, 2000a) implies that the attenuated Ne/ERN in schizophrenia may reflect a disturbance in the affective or motivational component of error monitoring that is related to dysfunction in rostral ACC.

The idea that disturbed error-related processing in schizophrenia may be related to motivational or affective processing abnormalities is consistent with the clinical presentation of schizophrenia. An extensive range of disorders of emotion occur in schizophrenia, of which blunted affect and inappropriate affect are the most characteristic and tend to be the most persistent (Bleuler, 1908, 1950). Disruptions of

Table 1 Demographic data for patients with schizophrenia and matched healthy control participants

Variable	Healthy participants		Schizophrenic patients	
	Mean	SD	Mean	SD
Age	32.3	10.6	32.9	12.9
Parental socio-economic status (Hollingshead)	3.7	1.2	3.5	1.0
Pre-morbid intellectual functioning (NART)	116.0	6.1	116.0	7.3
Current intellectual functioning (Quick Test)*	107.0	9.7	101.0	8.2

*Data from one healthy participant were not available. NART = National Adult Reading Test.

motivation and will are reflected in weakened or disjointed volition (manifest as extended periods of underactivity and poorly organized, ill-judged, impulsive activities, respectively). PET studies have demonstrated a positive correlation between regional cerebral blood flow in the ACC and the severity of disorganization symptoms (which incorporates inappropriate affect and bizarre, erratic behaviour; Liddle *et al.*, 1992; Ebmeier *et al.*, 1993; Yuasa *et al.*, 1995). Using ERPs, Bates *et al.* (2002) reported a significant negative correlation between psychomotor poverty syndrome (which includes the symptoms of blunted affect and underactivity) and Ne/ERN amplitude. To the extent that the error detection signal derives from a motivational or emotional response to errors generated in rostral ACC, these results suggest that the motivational and/or affective response to EoC in patients with schizophrenia may be disordered.

The purpose of the present study was to employ whole-brain event-related fMRI to examine the neural response to EoC and correct non-responses in healthy participants and patients with schizophrenia. The Go/NoGo task used in the present study was employed previously in an ERP study that demonstrated attenuation of Ne/ERN in patients with schizophrenia compared with healthy participants (Bates *et al.*, 2002). In fMRI, the task was shown to elicit robust activation in rostral ACC during EoC by healthy participants (Kiehl *et al.*, 2000). In light of the ERP and fMRI evidence, we hypothesized that patients with schizophrenia would fail to show the same magnitude of rostral ACC activity during the commission of errors as is observed in healthy participants. Given the extensive connections between the rostral ACC and limbic and paralimbic structures, we further hypothesized that reduced activation in limbic and/or paralimbic areas would be elicited in patients with schizophrenia relative to healthy participants. To the extent that the Go/NoGo paradigm includes a degree of response conflict, it was also expected that patients with schizophrenia might show an attenuated response compared with healthy participants in caudal ACC during errors.

Methods

Clinical assessment of the participants

Sixteen healthy adults (12 male) and 10 patients with schizophrenia (nine male) participated in the experiment and provided written informed consent. All participants were

right-handed (as per Annett, 1970), with normal or corrected-to-normal visual acuity. All procedures complied with University and Hospital ethical requirements.

Patients were stable, partially remitted, medicated out-patients recruited from community mental health teams in Vancouver, British Columbia and out-patient programmes at the University of British Columbia Hospital. All patients met DSM-IV criteria for schizophrenia, as diagnosed by an institutional or University Hospital psychiatrist (American Psychiatric Association, 1994). Mean duration of illness was 11 years (SD 3.1 years), with a range spanning 2–30 years. All patients received stable doses of atypical antipsychotics as their primary medication over the preceding 6-month period. Two patients also received a typical antipsychotic. A trained psychiatrist evaluated the symptoms of the patients with schizophrenia on the day of scanning using the Signs and Symptoms of Psychotic Illness interview schedule (Liddle *et al.*, 2002). This schedule comprises 20 symptom items scored 0–4 according to the severity of the symptom. Mean total score was 8.3 (SD 1.6), with a range of 1–18. Syndrome scores were calculated from the items according to the three-syndrome model of schizophrenia described by Liddle (1987a, b). Mean syndrome scores for Reality Distortion (sum of two items: delusions and hallucinations), Disorganization (sum of three items: thought disorder, inappropriate affect and peculiar behaviour) and Psychomotor Poverty (sum of three items: blunted affect, poverty of speech and underactivity), respectively, were: 2.0 (SD 0.7), 0.9 (SD 0.4) and 1.0 (SD 0.5).

Healthy participants were medication-free volunteers without history of neurological or Axis I psychiatric illness. Participant groups did not differ significantly on the demographic variables of age, gender, parental socio-economic status (Hollingshead and Redlich, 1958), or on estimates of pre-morbid (National Adult Reading Test; Nelson, 1982; Sharpe and O'Carroll, 1991) and current (Quick Test; Ammons and Ammons, 1962) intellectual functioning ($P > 0.05$; see Table 1).

fMRI procedure

A single scanning run comprising 246 visual stimuli was presented to participants using a computer-controlled visual and auditory presentation package (<http://nilab.psychiatry.ubc.ca/vapp/>). Stimuli were displayed on a rear-

projection screen mounted at the entrance to the magnet bore and subtended a visual angle of $\sim 3 \times 5^\circ$. Each stimulus appeared for 240 ms in white text within a continuously displayed rectangular fixation box. Participants viewed the screen from a distance of ~ 2 m by means of a mirror system attached to the head coil. The scanning room and magnet bore were darkened to permit easy visualization of the stimuli.

Participants were instructed to respond as quickly and accurately as possible with their right index finger to each presentation of the Go stimulus (the letter 'X'; occurrence probability = 0.84). They were instructed not to respond to the NoGo stimulus (the letter 'K'; occurrence probability = 0.16). Prior to scanning, participants completed a brief practice session of ~ 10 trials to promote speeded responding to the Go stimulus and thus increase the likelihood of EoC.

The stimulus onset asynchrony (SOA) between Go stimuli varied pseudorandomly between 1000, 2000 and 3000 ms, subject to the constraint that three Go stimuli were presented within each consecutive 6-s period. In light of the protracted evolution of the haemodynamic response elicited by a single stimulus, it was anticipated that the Go stimuli would generate a sustained, relatively constant baseline haemodynamic activity. The NoGo stimuli were interspersed among the Go stimuli in a pseudorandom manner subject to three constraints: the minimum SOA between a Go and NoGo stimulus was 1000 ms; the SOA between successive NoGo stimuli was in the range 10–15 s; and NoGo stimuli had an equal likelihood of occurring at 0, 1 or 2 s after the beginning of a 3-s acquisition period. Thus, the haemodynamic response to each NoGo stimulus occurred as a perturbation set against the relatively constant haemodynamic response to Go stimuli. By jittering stimulus presentation relative to the acquisition time, the haemodynamic response to the stimuli of interest was sampled effectively at 1-s intervals.

Motor responses were recorded using a commercially available MRI-compatible fibre optic response device (Lightwave Medical, Vancouver, BC). Reaction times to Go events were computed for trials in which the participants responded within 1000 ms of stimulus onset. EoC were defined as responses that occurred within 1000 ms of the onset of a NoGo stimulus. Correctly rejected NoGo events, termed correct rejects (CR), were determined by the absence of a motor response within 1000 ms of the NoGo stimulus.

fMRI parameters

Images were acquired on a standard clinical GE 1.5 T system fitted with a Horizon Echo-speed upgrade. A custom head holder was used to prevent movement. Conventional spin-echo T_1 -weighted sagittal localizing images were acquired to view the positioning of the participant's head in the scanner and to prescribe the functional image volumes. BOLD (blood oxygen-level dependent) contrast images were collected with a gradient-echo sequence [retention time (TR)/echo time (TE) 3000/40 ms, flip angle 90° , 24×24 cm field of view, 64×64 matrix, 62.5 kHz bandwidth, $3.75 \text{ mm} \times 3.75 \text{ mm}$ in

plane resolution, 5 mm thickness, 29 slices] that effectively covered the entire brain (145 mm axial extent). A total of 142 brain volumes were acquired. Four image volumes collected prior to the presentation of stimuli were discarded from subsequent analyses in order to remove the effects of the T_1 stabilization process.

Image processing and analysis

Functional images were reconstructed offline, and realigned and motion corrected using the procedure described by Friston *et al.* (1995a) and implemented in Statistical Parametric Mapping 99 (SPM99, Wellcome Department of Cognitive Neurology, London, UK; <http://www.fil.ion.ucl.ac.uk/spm/>). Corrections for translations and rotations did not exceed 3.0 mm and 2.5° , respectively for any participant. A group (schizophrenic patients, healthy participants) \times movement (translation, rotation) \times displacement axis (x, y, z) ANOVA (analysis of variance) was conducted on maximal estimated movement parameters to ensure that the groups did not differ in extent of head motion. A mean functional image was constructed in each participant and used to derive parameters for spatial normalization into the modified Talairach stereotaxic space implemented in SPM99. Both affine and non-linear components were used in the spatial normalization (Friston *et al.*, 1995b). The spatial normalization parameters for each mean image were then applied to the corresponding functional images for each session, and the images were resampled into isotropic 4 mm voxels. The normalized images were smoothed with an 8 mm full-width at half-maximum Gaussian kernel to optimize the signal-to-noise ratio and compensate for intersubject anatomical variation. High frequency noise associated with alterations of the applied radio frequency field was removed using a 0.16 Hz low-pass fifth-order IIR butterworth filter applied to the fMRI time series at each voxel. While all coordinates in the present study are reported and displayed in the modified Talairach stereotaxic space implemented in SPM99, a transformation algorithm was applied to these coordinates in order to localize activation patterns within standard Talairach space (i.e. to identify and label functional areas: Talairach and Tournoux, 1988; see <http://www.mrc-cbu.cam.ac.uk/Imaging/mnispac.html> for the transformation algorithm).

Statistical analysis was performed within each voxel using the general linear model approach implemented in SPM99. Event-related responses to both EoC and CR on NoGo stimuli were modelled using a synthetic haemodynamic response function comprised of two gamma functions and their temporal derivatives (Josephs *et al.*, 1997; Friston *et al.*, 1998). The first gamma function modelled the haemodynamic response peak at 6 s post-stimulus, and the second gamma function modelled the small 'overshoot' of the haemodynamic response on recovery. The occurrence of the erroneous motor response determined the timing of EoC (i.e. response-locked timing), whereas the timing of CR

Table 2 Significance of post hoc Scheffé tests on RT means for the main effect of each condition.

	Reaction time condition			
	Go-Go (mean: 366 ms)	EoC-Go (mean: 365 ms)	CR-Go (mean: 329 ms)	EoC (mean: 327 ms)
Go-Go		0.9993	0.0124*	0.0082*
EoC-Go			0.0178*	0.0120*
CR-Go				0.9991

Condition Go-Go = correct hits to Go trials following another correctly hit Go trial; EoC-Go = correct hits to Go trials following an EoC to a NoGo trial; CR-Go = correct hits to Go trials following a CR NoGo trial; EoC = EoC on NoGo trials.

corresponded to the presentation of the NoGo stimulus (i.e. stimulus-locked timing). Response-locked timing for EoC was chosen for consistency with standard ERP data analysis methods. The temporal derivatives of the gamma functions were included to compensate for slight variation in the peak latency of the onset of the haemodynamic response. The response to the Go events was treated as a baseline and not modelled explicitly. A high-pass filter was applied to remove noise associated with low frequency confounds (e.g. respiratory artefact). The confounding effects of fluctuations in global signal intensity between image volumes were removed using an adjusted proportional scaling routine (Desjardins *et al.*, 2001).

Two contrast images were specified for each participant, summarizing the amplitude of the fitted response in each voxel to: (i) EoC relative to the baseline of motor responding to Go events; and (ii) CR relative to the baseline of responding to Go events. These contrast images were then entered into separate independent samples *t* tests at the second level to test the null hypothesis that there was no difference between patients with schizophrenia and healthy participants in the mean amplitude of the fitted haemodynamic response for either of these event types. The contrast images were also entered into a second level one-sample *t* test for each group to test the null hypothesis that the mean of the observations on each event type did not differ significantly from zero in either healthy participants or patients with schizophrenia.

To test the significance of the *a priori* hypothesis of reduced activation in rostral ACC in patients with schizophrenia compared with healthy participants during EoC, a correction for multiple comparisons at $P \leq 0.05$ within a predefined 12 mm diameter spherical region of interest (ROI) was applied. This ROI was centred on the rostral ACC voxel identified in Kiehl *et al.* (2000) as preferentially active in healthy participants during EoC compared with CR on NoGo trials (voxel coordinates $x, y, z = -8, 45, 15$). In addition to voxels within the rostral ACC, this ROI also incorporated voxels in the medial frontal gyrus (corresponding to Brodmann area 10). To ascertain whether patients with schizophrenia showed reduced activation compared with healthy participants in the region of caudal ACC that was preferentially active in healthy participants during EoC

compared with correctly-rejected trials in Kiehl *et al.* (2000), a second ROI in caudal ACC was specified (12 mm diameter sphere centred on voxel coordinates $x, y, z = 4, 22, 40$). To examine the specificity of the results to error trials rather than NoGo trials in general, both the rostral and caudal ROIs were also applied to the comparison of patients with schizophrenia and healthy participants on correctly-rejected trials. Non-directed searches for differences between healthy participants and patients with schizophrenia on EoC and CR across the entire brain volume were implemented at the cluster level ($P \leq 0.05$ corrected for multiple comparisons, height threshold $P \leq 0.01$ uncorrected) according to the method of Worsley (1994) implemented in SPM99.

Results

Behavioural data

Mean reaction times (RTs) for correct hits to Go trials and EoC on NoGo trials for patients with schizophrenia were 393 (SD 60) and 349 ms (SD 48), respectively; and for healthy participants were 334 (SD 41) and 306 ms (SD 45), respectively. For subsequent RT analyses, correct hits to Go trials were differentiated into three types: (i) those that followed an EoC to a NoGo trial (EoC-Go); (ii) those that followed a correctly-rejected NoGo trial (CR-Go); and (iii) those that followed another correct hit to a Go trial (Go-Go).

RT data were analysed using a group (healthy participants, schizophrenic patients) \times condition (EoC, EoC-Go, CR-Go, Go-Go) ANOVA. The analysis revealed a significant main effect of group [$F(1,24) = 9.18, P = 0.006$], indicating that healthy participants responded faster to the task stimuli than patients with schizophrenia. The slowed performance of patients with schizophrenia on this task is consistent with that typically observed on speeded RT tasks (e.g. Ngan and Liddle, 2000).

The main effect of condition was also significant [$F(3,72) = 7.40, P = 0.0002$]; however, a non-significant group \times condition interaction [$F(3,72) = 1.33, P = 0.27$] indicated that patients with schizophrenia responded more slowly than healthy participants across all trial types. The results of *post hoc* Scheffé tests conducted on the RT means for the main effect of condition are reported in Table 2. The




Fig 1 Mean magnitude of the fitted response (± 2 SEM) at the voxel of peak activation (x, y, z coordinates = $-8, 52, 16$) located within the rostral ACC/medial frontal gyrus ROI identified in the comparison of EoC on NoGo trials relative to a baseline of responding to Go trials. The magnitude of the fitted response is provided for EoC to NoGo trials (solid lines) and correctly-rejected NoGo trials (hashed lines) relative to baseline in patients with schizophrenia (right) and healthy control participants (left).

RT for Go trials following an EoC were significantly longer than those for Go trials following a correctly-rejected NoGo stimulus, demonstrating that participants modified their response behaviour after committing an error. This was particularly true for patients with schizophrenia. A planned comparison between EoC-Go trials versus CR-Go trials confirmed that the increase in RT following errors was significant within the schizophrenic patient group [$F(1,24) = 7.90, P = 0.0097$]. The RT to Go trials following correctly-rejected NoGo trials was less than that to Go trials following other correct Go trials, which may reflect the fact that NoGo trials were always followed by Go trials. As expected, our results replicate previous research demonstrating that EoC are associated with faster responses than occur on correct hits to Go trials, possibly reflecting premature or impulsive response decisions on error trials (e.g. Pailing *et al.*, 2002).

Both healthy participants and patients with schizophrenia correctly identified the majority of Go stimuli (99.6 and 98.7%, respectively), although patients responded to significantly fewer Go stimuli than healthy participants [$t(24) = 2.42, P = 0.024$]. On NoGo trials, healthy participants and patients with schizophrenia did not differ significantly on accuracy of responding [mean EoC on NoGo trials were 16.6 (SD 7.2) and 16.3 (SD 6.4) in healthy participants and patients, respectively; $t(24) = 0.94, P = 0.93$].

Imaging data

The non-significant main effect and interactions for group in the ANOVA examining head motion [main effect: $F(1,24) = 0.613, P = 0.44$] indicates that the healthy participant and schizophrenic patient groups did not differ significantly on any estimated maximum head motion parameter, suggesting that movement did not contribute differentially to the haemodynamic results across groups.

EoC responses

Within the ROI centred in the rostral ACC, direct comparison of the magnitude of the fitted response in the two groups during EoC revealed significantly greater activation in healthy participants compared with patients with schizophrenia [coordinates of voxel of peak activation: $x, y, z = -8, 52, 16$; $t(24) = 3.11, P = 0.043$ corrected for multiple comparisons within the volume of interest]. Examination of the magnitude of the fitted response at this peak voxel in the healthy participant and schizophrenic patient groups separately provided clarification of the nature of this effect (see Fig. 1, solid lines). Consistent with the findings of Kiehl *et al.* (2000), healthy participants showed significant recruitment of the rostral ACC/medial frontal gyrus during EoC [coordinates of voxel of peak activation: $x, y, z = -8, 52, 16$; $t(24) = 4.39, P = 0.009$ corrected for multiple comparisons within the volume of interest]. In




Fig 2 Illustration of limbic and paralimbic clusters in which greater haemodynamic activity was observed in healthy participants than in patients with schizophrenia during EoC to NoGo stimuli (top row) and correctly-rejected NoGo stimuli (bottom row) relative to a baseline of responding to Go stimuli. Data are presented in the modified Talairach space used in SPM99 and are rendered onto a standard reference brain. The sagittal slices illustrating clusters in the anterior and posterior cingulate cortex are located at $x = -4$; the transverse slice illustrating the cluster in the hippocampal gyrus is located at $z = -16$. The view is in neurological convention, with the left hemisphere indicated by 'L' and the right hemisphere indicated by 'R'. The image is thresholded at a height of $t(24) = 2.49$, which corresponds to a significance level of $P \leq 0.01$ uncorrected for multiple comparisons. None of the illustrated differences in cerebral activation were statistically significant after correcting for multiple comparisons across the whole brain.

contrast, in the patients with schizophrenia, EoC were associated with a failure to activate this region.

Direct comparison of the magnitude of the fitted response in the two groups during EoC in the ROI centred in caudal ACC revealed no significant difference between patients with schizophrenia and healthy participants at $P \leq 0.05$ corrected for multiple comparisons within the ROI volume. Inspection of the magnitude of the fitted response within the ROI for the healthy participant and schizophrenic patient groups separately provided some evidence that both groups recruited caudal ACC during EoC [coordinates of voxel of peak activation for healthy participants: $x, y, z = 0, 24, 36$; $t(24) = 1.88$, $P = 0.0392$ uncorrected for multiple comparisons; for patients with schizophrenia: $x, y, z = 8, 28, 40$; $t(24) = 3.20$, $P = 0.0054$ uncorrected].

A non-directed search of the entire brain to identify additional regions in which healthy participants demonstrated significantly greater activation than patients with schizophrenia during EoC failed to identify any clusters satisfying the criteria of correction for multiple comparisons. The four largest clusters of activation were located in the posterior cingulate gyrus [24 voxels; peak voxel coordinate and

statistics: $x, y, z = 0, -44, 12$; $t(24) = 4.43$, $P = 0.001$ uncorrected for multiple comparisons across the entire brain], the rostral ACC/medial frontal gyrus [24 voxels, peak voxel coordinate and statistics: $x, y, z = -12, 60, 0$; $t(24) = 3.18$, $P = 0.001$ uncorrected], the left hippocampus [14 voxels, peak voxel coordinate and statistics: $x, y, z = -32, -20, -12$; $t(24) = 3.37$, $P = 0.001$ uncorrected] and the left angular gyrus [14 voxels, peak voxel coordinate and statistics: $x, y, z = -48, -72, 32$; $t(24) = 3.21$, $P = 0.002$ uncorrected]. Although none of these clusters satisfied stringent criteria for significance after correction for multiple comparisons in the entire brain volume, it is noteworthy that the three most significant clusters were located in limbic or paralimbic cortex. These clusters are illustrated in Fig. 2.

In the converse comparison that sought to identify brain regions in which patients with schizophrenia showed significantly greater activation than healthy participants during EoC, two clusters of 165 and 157 voxels, respectively, were observed (see Table 3 and Fig. 3). These clusters were located bilaterally in the superior parietal lobule/precuneus, and the larger cluster in the left hemisphere extended into inferior parietal lobule and postcentral gyrus.

Table 3 Clusters showing significantly greater activation in patients with schizophrenia compared with healthy participants for EoC to NoGo stimuli and correctly-rejected NoGo stimuli

Cluster level statistics			Voxel level statistics				Talairach coordinates			Anatomical level (Brodmann area)*
$P_{\text{corrected}}$	K_E	$P_{\text{uncorrected}}$	$P_{\text{corrected}}$	t	Z_e	$P_{\text{uncorrected}}$	x	y	z	
Errors of commission										
0.001	165	0.000	0.888	4.42	(3.74)	0.000	-24	-72	44	Left superior and inferior parietal lobules, precuneus and postcentral gyrus (7/40/2/3)
			0.999	3.75	(3.29)	0.000	-36	-44	40	
			1.000	3.17	(2.87)	0.002	-28	-44	64	
0.001	157	0.000	0.816	4.45	(3.83)	0.000	20	-64	52	Right precuneus and superior parietal lobule (7/19)
			1.000	3.64	(3.22)	0.001	28	-80	32	
			1.000	3.64	(3.21)	0.001	16	-72	60	
Correct-rejects										
0.003	134	0.000	0.882	4.44	(3.75)	0.000	24	-12	60	Right middle frontal gyrus (6/9)
			1.000	3.65	(3.22)	0.001	28	-4	48	
			1.000	3.20	(2.89)	0.002	40	8	40	
0.007	115	0.000	0.936	4.30	(3.67)	0.000	40	-36	44	Right inferior parietal lobule, precuneus and postcentral gyrus (40/7/2)
			0.999	3.80	(3.33)	0.000	32	-40	52	
			1.000	3.57	(3.16)	0.001	20	-60	56	
0.031	86	0.001	0.999	3.82	(3.35)	0.000	-28	-56	48	Left inferior parietal lobule and precuneus (7/19)
			1.000	3.70	(3.26)	0.001	-20	-76	48	
			1.000	3.03	(2.76)	0.003	-24	-76	32	
0.039	82	0.001	0.810	4.57	(3.84)	0.000	-44	-80	0	Left middle occipital gyrus, cuneus and lingual gyrus (19/17/18)
			0.997	3.90	(3.40)	0.000	-16	-84	4	
			1.000	3.18	(2.88)	0.002	-20	-80	-4	

Random effects cluster level statistics ($P \leq 0.01$ threshold for inclusion in the cluster) are reported along with voxel level statistics from several representative maxima of activation within the cluster.

*Brodmann areas correspond to those provided in the atlas of Talairach and Tournoux (1988).




Fig 3 Illustration of the significant clusters in which greater haemodynamic activity was observed in patients with schizophrenia than in healthy participants during EoC to NoGo stimuli (top row) and correctly-rejected NoGo stimuli (bottom row) relative to a baseline of responding to Go stimuli. Data are presented in the modified Talairach space used in SPM99 and are rendered onto a standard reference brain. For EoC, the sagittal slices are located at (from left) $x = -32, -24$ and $+24$; the sagittal slices for CR are located at (from left) $x = -32, -24, +40$ and -44 . The view is in neurological convention, with the left hemisphere indicated by 'L' and the right hemisphere indicated by 'R'. The image is thresholded at a height of $t(24) = 2.49$, which corresponds to a significance level of $P \leq 0.01$ uncorrected for multiple correction criteria across the whole brain. All clusters are significant at $P \leq 0.05$ corrected for multiple comparisons.

CR trials

The ROI analysis directly comparing the magnitude of the fitted response in healthy participants and patients with

schizophrenia in rostral ACC for correctly-rejected trials did not reveal any significant voxel satisfying the correction within the specified small volume of interest. For the purposes

of comparison, the magnitude of the fitted response for correctly-rejected trials at the peak voxel identified for EoC is provided in the healthy participant and schizophrenic patient groups separately in Fig. 1 (hashed lines). The figure indicates that neither the healthy participants nor the patients with schizophrenia showed significant recruitment of the rostral ACC during correctly-rejected trials.

Similarly, the ROI analysis that directly compared the magnitude of the fitted response between groups in caudal ACC for correctly-rejected trials revealed no significant voxel satisfying correction within the volume of interest. Neither healthy participants nor patients with schizophrenia showed significant recruitment of this region of the caudal ACC during correctly-rejected trials.

The non-directed search of the entire brain for regions showing greater activation in healthy participants compared with patients with schizophrenia during correctly-rejected NoGo trials failed to identify any clusters satisfying the criteria of correction for multiple comparisons. However, as for EoC, a large cluster of 34 suprathreshold voxels was located in posterior cingulate gyrus [peak voxel $x, y, z = 0, -52, 12$; $t(24) = 3.80$, $P = 0.001$ uncorrected for multiple comparisons across the entire brain; see Fig. 2].

Four clusters were significantly more active in patients with schizophrenia than in healthy controls during correctly-rejected trials (see Table 3 and Fig. 3). As for EoC, two of these clusters were located bilaterally in the parietal cortex, with the two further clusters located in middle occipital cortex and in middle frontal cortex (premotor area).

Discussion

Our results demonstrate that schizophrenia is characterized by relative underactivity of the rostral ACC during commission of errors. While errors elicited increased haemodynamic activity in this region in healthy participants, patients with schizophrenia failed to recruit this area during EoC. The rostral ACC activation in healthy participants was not observed during correct non-response to NoGo stimuli, which is consistent with previous research indicating that this region is involved specifically in error processing (Kiehl *et al.*, 2000; Braver *et al.*, 2001; Menon *et al.*, 2001). In caudal ACC, the haemodynamic activity elicited in the patient and healthy groups did not differ significantly during either EoC or CR. While there was some evidence that both groups activated this area of the caudal ACC during EoC, no significant activity was observed in this region in either group during CR to NoGo trials. ERP research has demonstrated an attenuation in the amplitude of the fronto-centrally distributed Ne/ERN in patients with schizophrenia relative to healthy participants (Kopp and Rist, 1999; Bates *et al.*, 2002; Mathalon *et al.*, 2002). Our fMRI results suggest that a failure to activate the rostral ACC sufficiently during error commission may contribute to the attenuated Ne/ERN observed in patients with schizophrenia.

The observation of aberrant activity in rostral ACC during EoC is consistent with the hypothesis that patients with schizophrenia experience a disturbed affective and/or motivational response to having committed an error. As an interface between limbic–paralimbic areas and widespread frontal cortex, the ACC would be expected to mediate the influence of the motivational and emotional state of an individual on the processing of, and response to, sensory stimuli. Disturbances in affect and motivation are common and persistent symptoms of schizophrenia. Bates *et al.* (2002) demonstrated that the expression of psychomotor poverty symptoms (including blunted affect and underactivity) is particularly associated with the attenuation of Ne/ERN. Thus, the attenuation of rostral ACC activity during EoC in patients with schizophrenia may reflect a relative diminution of the affective or motivational response associated with their realization that an error has been committed.

Relative underactivity in patients with schizophrenia compared with healthy participants was not apparent in the caudal ACC region identified by Kiehl *et al.* (2000) as being preferentially active during EoC compared with CR in healthy participants. This finding contrasts with that of Carter *et al.* (2001), who demonstrated diminished haemodynamic activity in the caudal ACC of patients with schizophrenia during errors that were elicited in a task that used degraded stimuli to increase error rates. As well as impaired caudal ACC function, the patients with schizophrenia in that study exhibited significantly reduced slowing of RT after error commission. In healthy individuals, RTs typically increase and fewer errors are committed following an erroneous response, which is consistent with the adoption of a more conservative response strategy following detection of an error (Rabbitt, 1966). Carter *et al.* (2001) interpreted the decreased error-related activity in caudal ACC and reduced post-error performance adjustment as evidence for impaired internal monitoring function in schizophrenia. The present Go/NoGo paradigm employed stimuli that were relatively easier to discriminate than the degraded stimuli employed by Carter *et al.* (2001), and errors were easily identifiable. The behavioural data obtained in this simple task revealed that patients with schizophrenia exhibited as great an increase in RT as was exhibited by healthy participants for Go trials following EoC relative to Go trials following CR, indicating that they modified their response behaviour following the commission of an error. This observation implies that they detected their error responses appropriately. Taken together, the results of the present study and those of Carter *et al.* (2001) suggest that there may be dissociable rostral and caudal ACC contributions to error processing, the relative strength of which may be modulated by task paradigm. Impairments in an internal monitoring component in schizophrenia appear to be reflected in caudal ACC underactivity, whereas disturbance in a subjective affective error assessment process may be associated with relative decreases in rostral ACC activity in schizophrenia.

The question of the relative contributions of rostral and caudal ACC dysfunction to the attenuated Ne/ERN observed in schizophrenia is unresolved. A recent report of source localization of high-density ERP data from healthy participants modelled the Ne/ERN as having a caudal ACC generator (Van Veen and Carter, 2002). The rostral ACC was also active during error processing, but later in time, and related to a positive error-related ERP component termed the error positivity (Pe; Falkenstein *et al.*, 2000). The current temporal resolution of fMRI does not allow the identification of differential Ne/ERN- and Pe-related contributions to ACC activity, and either component might be related to the rostral ACC activity elicited in healthy participants during error responses in the present study. However, previous research has failed to observe differences in Pe amplitude between patients with schizophrenia and healthy participants, in spite of detecting a reduction in Ne/ERN amplitude in schizophrenia (Mathalon *et al.*, 2002). Unpublished ERP data from a study in our laboratory that employed the same task as in the present study also failed to identify a difference in Pe amplitude between a small sample of acutely ill patients with schizophrenia and healthy participants during EoC, in spite of observing a reduction in Ne/ERN amplitude in patients compared with healthy participants.

In addition to the error-related failure to activate rostral ACC, there was some evidence for relative underactivity in patients with schizophrenia compared with healthy participants in the hippocampus and posterior cingulate cortex during EoC. The cingulate and hippocampal gyri constitute part of the limbic lobe that regulates affective and motivational functions. Relative underactivity in patients in extended limbic cortex thus provides support for the hypothesis that the error-related abnormality in rostral ACC function is associated with a disturbed emotional or motivational reaction to errors in schizophrenia. While the hippocampal and rostral ACC underactivity in schizophrenia were associated specifically with EoC, the posterior cingulate underactivity was also observed during correct response behaviour (i.e. on correctly-rejected NoGo events). Vogt *et al.* (1992) proposed a functional dichotomy between the anterior and posterior cingulate cortices, whereby the former is involved with executive functions and the emotional regulation of behaviour, and the latter subserves evaluative events such as monitoring sensory events and the organism's own behaviour. Thus, the reported underactivity in posterior cingulate cortex in schizophrenia might reflect a generalized impairment in the ability to evaluate rare but behaviourally relevant stimuli that occur against a background of more common events signalling an alternative behavioural response. Such a breakdown in posterior evaluative functions might be expected to increase the likelihood of error responses. While the limbic lobe activations reported outside the rostral ACC did not satisfy multiple correction criteria across the whole brain and must be interpreted with caution, they are suggestive of widespread dysfunction within the limbic system in schizophrenia.

In addition to observing areas of relative underactivity in patients with schizophrenia, a number of brain areas showed greater activity in patients than in the healthy participants. On NoGo trials, regardless of the accuracy of their subsequent response, patients with schizophrenia showed a relative increase in activity bilaterally around the interparietal sulcus. Several fMRI studies employing event-related Go/NoGo paradigms in healthy adults have reported activation of the interparietal sulcus during response inhibition on NoGo trials that occurred in the context of pre-potent responding to Go stimuli (Garavan *et al.*, 1999; Braver *et al.*, 2001; Liddle *et al.*, 2001; Menon *et al.*, 2001). The interparietal sulcus region is also activated across multiple sensory modalities during the detection of salient stimuli (Downar *et al.*, 2000, 2001, 2002) and during the detection of rare, behaviourally relevant target stimuli (Kiehl *et al.*, 2001*a, b*). Together, these results suggest that the interparietal sulcus area plays an important role in assessing the relevance of incoming stimuli for the purposes of deciding whether or not a behavioural response to the stimuli is required. The relative hyperactivity of this region in patients with schizophrenia may imply that this process is more difficult for patients than healthy individuals, and hence requires relatively greater engagement of resources to perform the task.

The other clusters of increased activity in patients with schizophrenia relative to healthy participants during correctly rejected NoGo trials occurred in right premotor cortex and left unimodal visual association cortex. Premotor cortex is involved in the planning and production of movements, particularly movements guided by external stimuli. The activated region was ipsilateral to the responding hand, and the activation occurred during a trial that involved suppression, rather than commission, of a pre-potent motor act. In acts of simple motor responding, patients with schizophrenia are characterized by reduced lateralization of premotor activation compared with healthy participants (Mattay *et al.*, 1998). Our results suggest that patients with schizophrenia also show a loss of hemispheric specialization during regulation of motor activity. The explanation for the greater activation in visual association cortex is also uncertain. However, previous neuroimaging research has described an increase in the relative magnitude of signal intensity (Renshaw *et al.*, 1994) and extent of activation (Taylor *et al.*, 1997) in patients with schizophrenia relative to healthy participants in striate cortex during photic stimulation. Our results suggest that abnormalities in visual cortex function may extend beyond the initial processing of sensory stimuli in the context of minimal task requirements (i.e. maintaining fixation on visual stimuli) to disturbed function in areas concerned with evaluating the identity of visual stimuli.

The reported underactivity in rostral ACC and extended limbic-paralimbic cortex during EoC in patients with schizophrenia relative to healthy participants was observed in spite of the low levels of symptomology reported by the patients, who were out-patients living and functioning in the community. Reality distortion symptoms (i.e. delusions and

hallucinations) were the most common symptoms reported, with a variable, though relatively low occurrence of disorganized or negative symptoms reported across the patient group. Further research in a larger patient group is needed to clarify the relationship between affective and motivational disturbance (as well as other symptomology), and the relative reduction in rostral ACC activity during error commission in schizophrenia.

This study examines the cortical response to the commission of errors in a medicated patient population, which raises the possibility that some of the observed differences between groups may be attributable to the effects of antipsychotic medication. Previous neuroimaging studies that have examined frontal function in unmedicated patients with schizophrenia have demonstrated both hypofrontality (Andreasen *et al.*, 1992) and hyperfrontality (e.g. Ebmeier *et al.*, 1993) of function, with patient symptomology contributing to variability in the extent of dysfunction. ACC function may be affected differentially by type of antipsychotic medication, as Braus *et al.* (2000, 2002) report higher levels of neuronal function markers in patients receiving atypical antipsychotics than in those receiving typical antipsychotics. The present study indicates that rostral ACC activity is reduced compared with healthy participants even in a sample of patients who were receiving atypical antipsychotic medication. Further research examining error-related activity in unmedicated patients pre- and post-treatment is required in order to determine the effect of antipsychotic medication on error processing and internal monitoring in schizophrenia.

In the present study, we have demonstrated that schizophrenia is associated with relative underactivity of the rostral ACC and associated limbic–paralimbic structures during the commission of errors in a simple Go/NoGo task variant. We additionally identified several brain regions, particularly bilateral parietal cortex, in which patients were characterized by abnormal overactivity compared with healthy participants during NoGo trial processing, suggesting a disturbance in response inhibition or stimulus evaluation processes in schizophrenia. Our results indicate that during performance of the simple Go/NoGo task employed in this study, the error monitoring system of patients with schizophrenia functions sufficiently well to detect errors. However, the pattern of activation indicates that the affective evaluation or motivational response to error commission is impaired. Future research might test this hypothesis by examining whether the rostral ACC activity associated with EoC normalizes in patients with schizophrenia when the salience of response accuracy is increased (e.g. by rewarding correct responses and/or penalizing erroneous responses).

Acknowledgements

We wish to thank Tara Cairo and Cameron Anderson for assistance with participant recruitment and data collection, Athena Vouloumanos for proofreading the manuscript, MRI

technicians Jennifer McCord, Sylvia Renneberg and Trudy Shaw, and Drs Alex MacKay, Ken Whittall and Bruce Forster for fMRI scanning assistance and advice. This research was supported in part by grants from the Dr Norma Calder Foundation for Schizophrenia Research and the Medical Research Council of Canada. K.R.L. was supported by the Gertrude Langridge Graduate Scholarship in Medical Sciences and a University of British Columbia Graduate Fellowship. A.T.B. was supported by a scholarship from the Natural Science and Engineering Research Council of Canada.

References

- American Psychiatric Association. Diagnostic and statistical manual of mental disorders. DSM-IV. 4th edn. Washington (DC): American Psychiatric Association; 1994.
- Ammons RB, Ammons CH. The Quick Test (QT): provisional manual. *Psychol Rep* 1962; 11: 111–61.
- Andreasen NC, Rezaei K, Alliger R, Swayze VW, 2nd, Flaum M, Kirchner P, et al. Hypofrontality in neuroleptic-naïve patients and in patients with chronic schizophrenia. Assessment with xenon 133 single-photon emission computed tomography and the Tower of London. *Arch Gen Psychiatry* 1992; 49: 943–58.
- Annett M. A classification of hand preference by association analysis. *Br J Psychol* 1970; 61: 303–21.
- Bates AT, Kiehl KA, Laurens KR, Liddle PF. Error-related negativity and correct-response negativity in schizophrenia. *Clin Neurophysiol* 2002; 113: 1454–63.
- Bernstein PS, Scheffers MK, Coles MG. ‘Where did I go wrong?’ A psychophysiological analysis of error detection. *J Exp Psychol Hum Percept Perform* 1995; 21: 1312–22.
- Bleuler E. The prognosis of dementia praecox: the group of schizophrenias. In: Cutting J, Shepherd M, editors. *The clinical roots of the schizophrenia concept*. Cambridge: Cambridge University Press; 1908. p. 59–74.
- Bleuler E. *Dementia praecox: or, the group of schizophrenias*. New York: International Universities Press; 1950.
- Botvinick MM, Braver TS, Barch DM, Carter CS, Cohen JD. Conflict monitoring and cognitive control. *Psychol Rev* 2001; 108: 624–52.
- Braus DF, Ende G, Hubrich-Ungureanu P, Henn FA. Cortical response to motor stimulation in neuroleptic-naïve first episode schizophrenics. *Psychiatry Res* 2000; 98: 145–54.
- Braus DF, Ende G, Weber-Fahr W, Demirakca T, Tost H, Henn FA. Functioning and neuronal viability of the anterior cingulate neurons following antipsychotic treatment: MR-spectroscopic imaging in chronic schizophrenia. *Eur Neuropsychopharmacol* 2002; 12: 145–52.
- Braver TS, Barch DM, Gray JR, Molfese DL, Snyder A. Anterior cingulate cortex and response conflict: effects of frequency, inhibition and errors. *Cereb Cortex* 2001; 11: 825–36.
- Bush G, Luu P, Posner MI. Cognitive and emotional influences in anterior cingulate cortex. *Trends Cogn Sci* 2000; 4: 215–22.

- Carter CS, Braver TS, Barch DM, Botvinick MM, Noll D, Cohen JD. Anterior cingulate cortex, error detection, and the online monitoring of performance. *Science* 1998; 280: 747–9.
- Carter CS, MacDonald AW III, Ross LL, Stenger VA. Anterior cingulate cortex activity and impaired self-monitoring of performance in patients with schizophrenia: an event-related fMRI study. *Am J Psychiatry* 2001; 158: 1423–8.
- Dehaene S, Posner MI, Tucker DM. Localization of a neural system for error detection and compensation. *Psychol Sci* 1994; 5: 303–5.
- Desjardins AE, Kiehl KA, Liddle PF. Removal of confounding effects of global signal in functional MRI analyses. *Neuroimage* 2001; 13: 751–8.
- Devinsky O, Morrell MJ, Vogt BA. Contributions of anterior cingulate cortex to behaviour. [Review]. *Brain* 1995; 118: 279–306.
- Dikman ZV, Allen JJ. Error monitoring during reward and avoidance learning in high- and low-socialized individuals. *Psychophysiology* 2000; 37: 43–54.
- Downar J, Crawley AP, Mikulis DJ, Davis KD. A multimodal cortical network for the detection of changes in the sensory environment. *Nat Neurosci* 2000; 3: 277–83.
- Downar J, Crawley AP, Mikulis DJ, Davis KD. The effect of task relevance on the cortical response to changes in visual and auditory stimuli: an event-related fMRI study. *Neuroimage* 2001; 14: 1256–67.
- Downar J, Crawley AP, Mikulis DJ, Davis KD. A cortical network sensitive to stimulus salience in a neutral behavioral context across multiple sensory modalities. *J Neurophysiol* 2002; 87: 615–20.
- Drevets WC, Raichle ME. Reciprocal suppression of regional cerebral blood flow during emotional versus higher cognitive processes: implications for interactions between emotion and cognition. *Cogn Emotion* 1998; 12: 353–85.
- Ebmeier KP, Blackwood DH, Murray C, Souza V, Walker M, Dougall N, et al. Single-photon emission computed tomography with 99mTc-exametazime in unmedicated schizophrenic patients. *Biol Psychiatry* 1993; 33: 487–95.
- Falkenstein M, Hohnsbein J, Hoormann J, Blanke L. Effects of errors in choice reaction tasks on the ERP under focused and divided attention. In: Brunia CHM, Gaillard AW, Kok A, editors. *Psychophysiological brain research*. Tilburg: Tilburg University Press; 1990. p. 192–195.
- Falkenstein M, Hohnsbein J, Hoormann J, Blanke L. Effects of crossmodal divided attention on late ERP components. II. Error processing in choice reaction tasks. *Electroencephalogr Clin Neurophysiol* 1991; 78: 447–55.
- Falkenstein M, Hoormann J, Christ S, Hohnsbein J. ERP components on reaction errors and their functional significance: a tutorial. [Review]. *Biol Psychol* 2000; 51: 87–107.
- Friston KJ, Ashburner J, Frith CD, Poline JB, Heather JD, Frackowiak RS. Spatial registration and normalization of images. *Hum Brain Mapp* 1995a; 3: 165–89.
- Friston KJ, Holmes AP, Worsley KJ, Poline JB, Frith CD, Frackowiak RS. Statistical parametric maps in functional imaging: a general linear approach. *Hum Brain Mapp* 1995b; 2: 189–210.
- Friston KJ, Fletcher P, Josephs O, Holmes A, Rugg MD, Turner R. Event-related fMRI: characterizing differential responses. *Neuroimage* 1998; 7: 30–40.
- Frith CD, Done DJ. Experiences of alien control in schizophrenia reflect a disorder in the central monitoring of action. *Psychol Med* 1989; 19: 359–63.
- Garavan H, Ross TJ, Stein EA. Right hemispheric dominance of inhibitory control: an event-related functional MRI study. *Proc Natl Acad Sci USA* 1999; 96: 8301–6.
- Gehring WJ, Coles MGH, Meyer DE, Donchin E. The error-related negativity: an event-related brain potential accompanying errors [abstract]. *Psychophysiology* 1990; 27: S43.
- Gehring WJ, Goss B, Coles MG, Meyer DE, Donchin E. A neural system for error detection and compensation. *Psychol Sci* 1993; 4: 385–90.
- Hollingshead AB, Redlich FC. *Social class and mental illness: a community study*. New York: Wiley; 1958.
- Holroyd CB, Dien J, Coles MG. Error-related scalp potentials elicited by hand and foot movements: evidence for an output-independent error-processing system in humans. *Neurosci Lett* 1998; 242: 65–8.
- Johns LC, Rossell S, Frith C, Ahmad F, Hemsley D, Kuipers E, et al. Verbal self-monitoring and auditory verbal hallucinations in patients with schizophrenia. *Psychol Med* 2001; 31: 705–15.
- Josephs O, Turner R, Friston KJ. Event-related fMRI. *Hum Brain Mapp* 1997; 5: 1–7.
- Kiehl KA, Liddle PF, Hopfinger JB. Error processing and the rostral anterior cingulate: an event-related fMRI study. *Psychophysiology* 2000; 37: 216–23.
- Kiehl KA, Laurens KR, Duty TL, Forster BB, Liddle PF. An event-related fMRI study of visual and auditory oddball tasks. *J Psychophysiol* 2001a; 15: 221–240.
- Kiehl KA, Laurens KR, Duty TL, Forster BB, Liddle PF. Neural sources involved in auditory target detection and novelty processing: an event-related fMRI study. *Psychophysiology* 2001b; 38: 133–42.
- Kopp B, Rist F. An event-related brain potential substrate of disturbed response monitoring in paranoid schizophrenic patients. *J Abnorm Psychol* 1999; 108: 337–46.
- Leudar I, Thomas P, Johnston M. Self-monitoring in speech production: effects of verbal hallucinations and negative symptoms. *Psychol Med* 1994; 24: 749–61.
- Liddle PF. The symptoms of chronic schizophrenia. A re-examination of the positive–negative dichotomy. *Br J Psychiatry* 1987a; 151: 145–51.
- Liddle PF. Schizophrenic syndromes, cognitive performance and neurological dysfunction. *Psychol Med* 1987b; 17: 49–57.
- Liddle PF, Friston KJ, Frith CD, Hirsch SR, Jones T, Frackowiak RS. Patterns of cerebral blood flow in schizophrenia. *Br J Psychiatry* 1992; 160: 179–86.
- Liddle PF, Kiehl KA, Smith AM. Event-related fMRI study of response inhibition. *Hum Brain Mapp* 2001; 12: 100–9.

- Liddle PF, Ngan ET, Duffield G, Kho K, Warren AJ. Signs and Symptoms of Psychotic Illness (SSPI): a rating scale. *Br J Psychiatry* 2002; 180: 45–50.
- Luu P, Collins P, Tucker DM. Mood, personality, and self-monitoring: negative affect and emotionality in relation to frontal lobe mechanisms of error monitoring. *J Exp Psychol Gen* 2000a; 129: 43–60.
- Luu P, Flaisch T, Tucker DM. Medial frontal cortex in action monitoring. *J Neurosci* 2000b; 20: 464–9.
- MacDonald AW 3rd, Cohen JD, Stenger VA, Carter CS. Dissociating the role of the dorsolateral prefrontal and anterior cingulate cortex in cognitive control. *Science* 2000; 288: 1835–8.
- Malenka RC, Angel RW, Hampton B, Berger PA. Impaired central error-correcting behavior in schizophrenia. *Arch Gen Psychiatry* 1982; 39: 101–7.
- Malenka RC, Angel RW, Thiemann S, Weitz CJ, Berger PA. Central error-correcting behavior in schizophrenia and depression. *Biol Psychiatry* 1986; 21: 263–73.
- Mathalon DH, Fedor M, Faustman WO, Gray M, Askari N, Ford JM. Response-monitoring dysfunction in schizophrenia: an event-related brain potential study. *J Abnorm Psychol* 2002; 111: 22–41.
- Mattay VS, Callicott JH, Bertolino A, Santha AK, Van Horn JD, Tallent KA, et al. Hemispheric control of motor function: a whole brain echo planar fMRI study. *Psychiatry Res* 1998; 83: 7–22.
- Menon V, Adelman NE, White CD, Glover GH, Reiss AL. Error-related brain activation during a Go/NoGo response inhibition task. *Hum Brain Mapp* 2001; 12: 131–43.
- Miltner WHR, Braun CH, Coles MGH. Event-related brain potentials following incorrect feedback in a time-production task: evidence for a 'generic' neural system for error detection. *J Cogn Neurosci* 1997; 9: 788–98.
- Mlakar J, Jensterle J, Frith CD. Central monitoring deficiency and schizophrenic symptoms. *Psychol Med* 1994; 24: 557–64.
- Nelson HE. National Adult Reading Test (NART): test manual. Windsor (UK): NFER-Nelson; 1982.
- Ngan ET, Liddle PF. Reaction time, symptom profiles and course of illness in schizophrenia. *Schizophr Res* 2000; 46: 195–201.
- Pailing PE, Segalowitz SJ, Dywan J, Davies PL. Error negativity and response control. *Psychophysiology* 2002; 39: 198–206.
- Rabbitt PM. Errors and error correction in choice-response tasks. *J Exp Psychol* 1966; 71: 264–72.
- Renshaw PF, Yurgelun-Todd DA, Cohen BM. Greater hemodynamic response to photic stimulation in schizophrenic patients: an echo planar MRI study. *Am J Psychiatry* 1994; 151: 1493–5.
- Scheffers MK, Coles MG. Performance monitoring in a confusing world: error-related brain activity, judgments of response accuracy, and types of errors. *J Exp Psychol Hum Percept Perform* 2000; 26: 141–51.
- Sharpe K, O'Carroll R. Estimating premorbid intellectual level in dementia using the National Adult Reading Test: a Canadian study. *Br J Clin Psychol* 1991; 30: 381–4.
- Stirling JD, Hellewell JS, Quraishi N. Self-monitoring dysfunction and the schizophrenic symptoms of alien control. *Psychol Med* 1998; 28: 675–83.
- Stirling JD, Hellewell JS, Ndlovu D. Self-monitoring dysfunction and the positive symptoms of schizophrenia. *Psychopathology* 2001; 34: 198–202.
- Talairach J, Tournoux P. Co-planar stereotaxic atlas of the human brain: 3-dimensional proportional system: an approach to cerebral imaging. Stuttgart: Thieme; 1988.
- Taylor SF, Tandon R, Koeppe RA. PET study of greater visual activation in schizophrenia. *Am J Psychiatry* 1997; 154: 1296–8.
- Ullsperger M, von Cramon DY. Subprocesses of performance monitoring: a dissociation of error processing and response competition revealed by event-related fMRI and ERPs. *Neuroimage* 2001; 14: 1387–401.
- Van Veen V, Carter CS. The timing of action-monitoring processes in the anterior cingulate cortex. *J Cogn Neurosci* 2002; 14: 593–602.
- Vidal F, Hasbroucq T, Grapperon J, Bonnet M. Is the 'error negativity' specific to errors? *Biol Psychol* 2000; 51: 109–28.
- Vogt BA, Finch DM, Olson CR. Functional heterogeneity in cingulate cortex: the anterior executive and posterior evaluative regions. [Review]. *Cereb Cortex* 1992; 2: 435–43.
- Whalen PJ, Bush G, McNally RJ, Wilhelm S, McInerney SC, Jenike MA, et al. The emotional counting Stroop paradigm: a functional magnetic resonance imaging probe of the anterior cingulate affective division. *Biol Psychiatry* 1998; 44: 1219–28.
- Worsley KJ. Local maxima and the expected Euler characteristic of excursion sets of Chi², F and t fields. *Adv Appl Probability* 1994; 26: 13–42.
- Yuasa S, Kurachi M, Suzuki M, Kadono Y, Matsui M, Saitoh O, et al. Clinical symptoms and regional cerebral blood flow in schizophrenia. *Eur Arch Psychiatry Clin Neurosci* 1995; 246: 7–12.

Received July 12, 2002. Revised October 15, 2002.
Accepted October 16, 2002