

Influence of Atopic Dermatitis on Reproduction and Uterine Natural Killer Cells

Kazuhiko HAYASHI¹⁾, Ken Takeshi KUSAKABE^{1,2)*}, Satoko SUGIMOTO²⁾, Shoichi WAKITANI³⁾, Shinji SUGI¹⁾, Nobue KUNIYOSHI¹⁾, Masato HIYAMA¹⁾, Ai TAKESHITA⁴⁾, Kiyoshi KANO^{1, 2)} and Yasuo KISO^{1,2)}

¹⁾Laboratory of Basic Veterinary Science, the United Graduate School of Veterinary Science, Joint Faculty of Veterinary Medicine, Yamaguchi University, Yamaguchi 753–8515, Japan

²⁾Laboratory of Veterinary Anatomy, Joint Faculty of Veterinary Medicine, Yamaguchi University, Yamaguchi 753–8515, Japan

³⁾Laboratory of Veterinary Biochemistry and Molecular Biology, Department of Veterinary Sciences, Faculty of Agriculture, University of Miyazaki, Miyazaki, 889–2192, Japan

⁴⁾Laboratory of Laboratory Animal Science, Division of Veterinary Science, Graduate School of Life and Environmental Science, Osaka Prefecture University, Osaka 598–8531, Japan

(Received 31 October 2013/Accepted 11 February 2014/Published online in J-STAGE 27 February 2014)

ABSTRACT. The causal relationship between severe allergic conditions and successful pregnancy remains unclear. We aimed to evaluate reproductive performance in an experimental mouse model of atopic disease (AD), and the appearance of uterine natural killer (uNK) cells that have crucial roles in placental formation was examined. In the NC/Nga pregnant mice with moderate skin allergic lesions and an 8.6-fold elevation of plasma IgE, significant differences were not detected in the reproductive indices of the number of normal fetuses, abortion rate and placental size. There were few uNK cells in the placenta of AD mice, and they showed a significant decrease regarding the immature subtype as compared with controls. These findings revealed that AD disturbs uNK cell differentiation and provides disadvantageous effects on placental formation, although it does not arrest the pregnancy process. It may be possible that specific immunological conditions behind AD operate favorably to recover the reproductive performance.

KEY WORDS: atopic disease, NC/Nga mouse, placenta, uNK cell.

doi: 10.1292/jvms.13-0547; *J. Vet. Med. Sci.* 76(6): 913–916, 2014

Atopic dermatitis (AD) is a common allergic disease characterized by chronic inflammatory skin lesions and hyperproduction of IgE and Th-2-type cytokines, such as interleukin (IL)-4 and IL-13, in the blood [3, 12]. There are conflicting data from human studies regarding the relationship between AD and pregnancy. Hanzlikova *et al.* showed that blood levels of IL-4 and IgE were downregulated in patients with repeated pregnancy loss [2]. On the other hand, an epidemiological study has revealed that AD and allergic diseases were related to difficulty achieving a successful pregnancy [15].

In many mammals, specific lymphocytes appear in the uteri during pregnancy [4]. In rodents and humans, the dominant subset of lymphocytes in the pregnant uterus is natural killer (NK) cells, and they frequently appear in the decidua basalis region. The uterine NK (uNK) cells can be classified into subpopulations by surface markers. The mature population has potential functions in angiogenesis, vasodilation and local blood pressure control, suggesting that these cells contribute to vascular remodeling and placental circulation [1]. However, uNK cells have other cellular

profiles to produce cytotoxic proteins, such as perforin and Fas ligand [6, 7]. Generally, NK cells express Fcγ receptors, which can bind to the IgE-immune complex and activate antibody-dependent cell cytotoxicity (ADCC). Therefore, abnormal immunological conditions have the possibility of transforming uNK cell profiles to unmask cytotoxicity against fetal tissue.

NC/NgaTndCrIj (NC/Nga) mice are a valuable experimental model of AD. In response to parasitic ticks or hapten stimulation, these mice develop severe skin lesions, enhanced blood levels of IgE and increased production of IL-4 and IL-13 [11]. In the present study, we evaluated the influence of AD on reproductive performance and uNK cellular profiles by using pregnant NC/Nga mice with AD.

Female NC/Nga mice were purchased from Charles River Japan (Yokohama, Japan). The experimental group was housed in a conventional breeding room and received first application of 5% picryl chloride dissolved in olive oil to the head and back, followed by repeated application of 0.8% picryl chloride for 1 week, according to the recommended protocol [8]. Following confirmation of moderate skin lesions and pruritic dermatitis, NC/Nga mice were mated with ICR male mice. Control NC/Nga female mice were received applications of olive oil in the same manner as the experimental group under specific pathogen free (SPF) condition and were mated with SPF-grade ICR male mice. The morning that a vaginal plug was detected was considered day 0 of pregnancy. On days 10 and 12 of pregnancy, blood and uteri were collected. Experiments were approved by the Ethics Committee for Animal Experiments at Yamaguchi

*CORRESPONDENCE TO: KUSAKABE, K. T., Laboratory of Basic Veterinary Science, The United Graduate School of Veterinary Science, 1677–1 Yoshida, Yamaguchi 753–8515, Japan.
e-mail: kusakabe@yamaguchi-u.ac.jp

©2014 The Japanese Society of Veterinary Science

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives (by-nc-nd) License <<http://creativecommons.org/licenses/by-nc-nd/3.0/>>.

University, and all animals were treated with humane care in keeping with the Yamaguchi University Experimental Guidelines.

The plasma IgE level was measured by using a Mouse IgE ELISA kit (Bethyl Labs, Montgomery, TX, U.S.A.). Each implantation site was visually checked regarding fetal viability and fixed with Bouin's reagent. Tissue from the central region of the implantation site was cut transversally and prepared as 4 μm -thick paraffin-embedded sections. Some sections were used for periodic acid Schiff (PAS) staining, and the thickness of the placenta and each component region was measured by using a Biozero microscope system (Keyence, Osaka, Japan). Other tissue sections were used for staining with *Dolichos biflorus* agglutinin (DBA) lectin, which is a specific marker for uNK cell classification [10]. Briefly, deparaffinized sections were incubated with biotinylated-DBA lectin (Vector Labs., Burlingame, CA, U.S.A.) at 4°C overnight. Specific reactions were detected with Avidin-Biotin Complex solution (Vector Labs) and 3-3'-diaminobenzidine tetrachloride. The number of uNK cells was counted according to the criteria of differentiation stages from subtypes I to IV [10]. For statistical evaluation, data were collected from at least 3 different mice, and more than 3 measurement fields were selected from at least 3 different sections prepared from different mice. *P* values < 0.05 were considered to be significant following Fisher's exact probability test (for abortion rate) or the Student's *t*-test (for plasma IgE level, placental thickness and uNK cell number).

Mice in the experimental group showed an 8.6-fold elevation of plasma IgE levels (Fig. 1). Pregnant uteri contained spontaneously absorbed implantation sites that had shrunk in size and changed in color to dark-red. The abortion rate was higher in the experimental group at both days 10 and 12 (Table 1), although a statistically significant difference was not detected because of the large standard deviation (SD). The placentas of the experimental group showed a reduced thickness at day 12 as compared to control placentas, but the reduction was not statistically significant (Fig. 2). The ratios of each placental component area in the control and experimental groups, respectively, were as follows: (Day 10) metrial gland (MG): 27.4% and 26.3%; decidua basalis (DB): 43.1% and 45.8%; placental labyrinth (PL): 29.6% and 27.8%; (Day 12) MG: 20.7% and 28.3%; DB: 20.9% and 19.8%; PL: 58.4% and 51.9%. The DB and PL tended to be reduced in the experimental group. Apparent pathological changes were not observed in the experimental group placentas. The metrial gland and decidua basalis regions




Fig. 1. Plasma IgE levels in control and experimental groups. **: *P* < 0.01.

contained many uNK cells that were DBA-lectin positive (Fig. 3A). At most developmental stages, the number of uNK cells was smaller in the placentas of the experimental group as compared to the control group (Fig. 3B). In particular, a significantly lower density was observed on the immature type I uNK cells in the metrial gland region of experimental group placentas.

The present study showed a negative influence of AD and severe hyperimmunoglobulinemia on fetal survival, placental formation and appearance of uNK cells in placentas, but these effects seemed to be non-fatal. Insufficient uNK cell recruitment is considered to negatively affect reproductive performance in NC/Nga mice. Paffaro *et al.* suggested subtype I uNK cells as immature progenitor cells recruit from peripheral blood to pregnant uteri [10]. For vascular recruitment, plural adhesion molecules are required for uNK cell settlement in uteri [5]. Under allergic conditions, the microenvironment of adhesion molecules is modified by the IL-4 effect to elevate vascular cell adhesion molecule (VCAM)-1 and eotaxin, resulting in the activation of eosinophils and basophils [11]. Reactivity to eotaxin has been observed especially on the CD56^{dim} NK cell subset, which is a cytotoxic subtype of human NK cells, while human CD-

Table 1. The number of fetus and the abortion rate

	Day10		Day12	
	Control	Experimental	Control	Experimental
Nomal fetuses	7.00 ± 1.00	6.50 ± 3.54	7.33 ± 1.37	7.33 ± 1.53
Absorbed fetuses	0.67 ± 0.58	1.50 ± 0.71	1.00 ± 1.26	2.00 ± 2.00
Abortion rate (%)	8.47 ± 7.50	21.7 ± 16.5	11.8 ± 15.4	20.7 ± 20.0

Data represent Mean ± SD.




Fig. 2. Upper column: Thickness of placenta and each placental area (MG: metrial gland, DB: decidua basalis, PL: placental labyrinth). Lower column: representative photo of a placenta from each group. Zonation of each placental area is indicated. Scale bar=500 μm

56^{bright} NK cells, which are analogous to mouse uNK cells, hardly show responsiveness to eotaxin [14]. Modification of the adhesion molecule pattern may induce the functional decline of uNK cells and inadequate vascular reconstitution [1, 9], resulting in a less efficient blood supply and reduced fetal survival. Additionally, since some atopic dams showed a high abortion rate, hyperimmunoglobulinemia itself could possibly affect the maternal reproductive system. However, a clear causal association between pregnancy loss and atopic disease could not be found in the present study.

In normal pregnancy, placental trophoblasts stably

express IL-4, which stimulates the release of chorionic gonadotropin and subsequently promotes the production of ovarian progesterone [13]. Thus, IL-4 itself has a direct reproductive effect that contributes to normal pregnancy. It is also possible that vascular permeability, enhanced by allergic conditions through histamine and IgE-sensitized basophils, operates positively to enhance substance exchange and fetal nutrition. To protect an embryo from the maternal immune system as a semi-allograft, the development of immune privilege in the placenta is necessary. The induction of Th-2-type cytokine dominance (IL-4, 6, 10, etc.) is crucial for immune privilege in placentas. Th-2-type cytokine dominance inhibits lymphocyte cytotoxicity and promotes the release of pregnancy-essential cytokines and hormones [13]. Profiles of Th-2-type cytokines seem to be a key factor in the effect of AD on pregnancy physiology. Cytokine profiles of AD may coincidentally promote the recovery of normal reproductive ability. This idea is consistent with our finding that AD causes little damage to the normal reproductive process. The immunological specificity of AD may resemble that of the reproductive process, e.g. antigen commonality between allergens and embryonic tissue relating to the Th-2 cytokine-dominant condition. On the other hand, a study of the differences of immunological profiles between AD and pregnancy may be necessary to evaluate the pathogenesis of pregnancy disorder.

In brief, the present study raised the possibility that AD inhibits uNK cell development at an early stage and adversely influences placental formation. On the other hand, cytokine profiles evoked by AD may potentially improve the immunological conditions for adapting to reproductive physiology.

ACKNOWLEDGMENTS. This study was supported by JSPS KAKENHI, Grant No. 24380159 (to Y.K.) and No. 23580407 (to K-T.K.).




Fig. 3. Lectin staining of placenta at day 12 (A). DBA lectin reacts on the surface and cytoplasmic granules of uNK cells (arrowheads). Subtypes for uNK cell maturation were determined by cell size and degree of granular development. Scale bar=20 μm . Cell density of uNK cells at the metrial gland (B) and the decidua basalis regions (C). Open bar: control group; closed bar: experimental group. *: $P < 0.05$.

REFERENCES

1. Chen, Z., Zhang, J., Hatta, K., Lima, P. D. A., Yadi, H., Colucci, F., Yamada, A. T. and Croy, B. A. 2012. DBA-lectin reactivity defines mouse uterine natural killer cell subsets with biased gene expression. *Biol. Reprod.* **87**: 81. [Medline] [CrossRef]
2. Hanzlikova, J., Ulcova-Gallova, Z., Malkusova, I., Sefrna, F. and Panzner, P. 2009. TH1-TH2 response and the atopy risk in patients with reproduction failure. *Am. J. Reprod. Immunol.* **61**: 213–220. [Medline] [CrossRef]
3. Kay, A. B., Ying, S., Varney, V., Gaga, M., Durham, S. R., Moqbel, R., Wardlaw, A. J. and Hamid, Q. 1991. Messenger RNA expression of the cytokine gene cluster, interleukin 3 (IL-3), IL-4, IL-5, and granulocyte/macrophage colony-stimulating factor, in allergen-induced late-phase cutaneous reactions in atopic subjects. *J. Exp. Med.* **173**: 775–778. [Medline] [CrossRef]
4. Kiso, Y. and Kusakabe, K. 1998. Some aspects of granulated metrial gland cells found at the feto-maternal interface during successful pregnancy. pp. 327–336. *In: Reproductive Biology Update. Novel Tools for Assessment of Environmental Toxicity* (Miyamoto, H. and Manabe, N. eds.), Shoukado, Kyoto.
5. Kruse, A., Martens, N., Fernekorn, U., Hallmann, R. and Butcher, E. C. 2002. Alterations in the expression of homing-associated molecules at the maternal/fetal interface during the course of pregnancy. *Biol. Reprod.* **66**: 333–345. [Medline] [CrossRef]
6. Kusakabe, K., Otsuki, Y. and Kiso, Y. 2005. Involvement of Fas ligand and Fas system in apoptosis induction of mouse uterine natural killer cell. *J. Reprod. Dev.* **51**: 333–340. [Medline] [CrossRef]
7. Kusakabe, K., Li, Z. L., Kiso, Y. and Otsuki, Y. 2005. Perforin improves the morphogenesis of mouse placenta disturbed by IL-2 treatment. *Immunobiology* **209**: 719–728. [Medline] [CrossRef]
8. Matsuda, H. and Tanaka, A. 1998. Effectiveness of NC/Nga mice as a model for atopic dermatitis. *CRJ letters* **11**: 1–8 (in Japanese).
9. Monk, J. M., Leonard, S., McBey, B. A. and Croy, B. A. 2005. Induction of murine spiral artery modification by recombinant human interferon-gamma. *Placenta* **26**: 835–838. [Medline] [CrossRef]
10. Paffaro, V. A., Bizinotto, M. C., Joazeiro, P. P. and Yamada, A. T. 2003. Subset classification of mouse uterine natural killer cells by DBA lectin reactivity. *Placenta* **24**: 479–488. [Medline] [CrossRef]
11. Prussin, C. and Metcalfe, D. D. 2003. IgE, mast cells, basophils, and eosinophils. *J. Allergy Clin. Immunol.* **111**: S486–S494. [Medline] [CrossRef]
12. Reinhold, U., Wehrmann, W., Kukul, S. and Kreysel, H. W. 1990. Evidence that defective interferon-gamma production in atopic dermatitis patients is due to intrinsic abnormalities. *Clin. Exp. Immunol.* **79**: 374–379. [Medline] [CrossRef]
13. Saito, S. 2000. Cytokine network at the feto-maternal interface. *J. Reprod. Immunol.* **47**: 87–103. [Medline] [CrossRef]
14. Wilk, E., Kalippke, K., Buyny, S., Schmidt, R. E. and Jacobs, R. 2008. New aspects of NK cell subset identification and inference of NK cells' regulatory capacity by assessing functional and genomic profiles. *Immunobiology* **213**: 271–283. [Medline] [CrossRef]
15. Zac, R. I., Machado, V. M., Alberti, L. R. and Petroianu, A. 2005. Association of allergy, infertility and abortion. *Rev. Assoc. Med. Bras.* **51**: 177–180 (in Portuguese). [Medline] [CrossRef]