

A FACTORIZATION THEOREM FOR LOGHARMONIC MAPPINGS

ZAYID ABDULHADI and YUSUF ABUMUHANNA

Received 24 March 2002

We give the necessary and sufficient condition on sense-preserving logharmonic mapping in order to be factorized as the composition of analytic function followed by a univalent logharmonic mapping.

2000 Mathematics Subject Classification: 30C55, 30C62, 49Q05.

Let D be a domain of \mathbb{C} and denote by $H(D)$ the linear space of all analytic functions defined on D . A logharmonic mapping is a solution of the nonlinear elliptic partial differential equation

$$\overline{f_z} = \left(a \frac{\overline{f}}{f} \right) f_z, \quad (1)$$

where $a \in H(D)$ and $|a(z)| < 1$ for all $z \in D$. If f does not vanish on D , then f is of the form

$$f = H \cdot \overline{G}, \quad (2)$$

where H and G are locally analytic (possibly multivalued) functions on D . On the other hand, if f vanishes at z_0 , but is not identically zero, then f admits the local representation

$$f(z) = (z - z_0)^m |z - z_0|^{2\beta m} h(z) \overline{g(z)}, \quad (3)$$

where

- (a) m is a nonnegative integer,
- (b) $\beta = \overline{a(0)}(1 + a(0))/(1 - |a(0)|^2)$ and therefore $\Re \beta > -1/2$,
- (c) h and g are analytic in a neighbourhood of z_0 .

In particular, if D is a simply connected domain, then f admits a global representation of the form (3) (see, e.g., [2]). Univalent logharmonic mappings defined on the unit disk U have been studied extensively (for details, see, e.g., [1, 2, 3, 4, 5, 6]).

In the theory of quasiconformal mappings, it is proved that for any measurable function μ with $|\mu| < 1$, the solution of Beltrami equation $f_{\overline{z}} = \mu f_z$ can be factorized in the form $f = \psi \circ F$, where F is a univalent quasiconformal mapping and ψ is an analytic function (see [8]). Moreover, for sense-preserving

harmonic mappings, the answer was negative. In [7], Duren and Hengartner gave a necessary and sufficient condition on sense-preserving harmonic mapping f for the existence of such factorization. Since logharmonic mappings are preserved under precomposition with analytic functions, it is a natural question to ask whether every sense-preserving logharmonic mapping can be factorized in the form $f = F \circ \phi$ for some univalent logharmonic mapping F and some analytic function ϕ .

It is instructive to begin with two simple examples.

EXAMPLE 1. Let f be the logharmonic mapping $f(z) = z^2/|1 - z|^4$ defined on the unit disc U . Then f is sense-preserving in U with dilatation $a(z) = z$. We claim that f has no decomposition of the desired form in any neighborhood of the origin. Suppose on the contrary that $f = F \circ \phi$, where ϕ is analytic near the origin and F is univalent logharmonic mapping on the range of ϕ . Then F is sense-preserving because f is. Without loss of generality, we suppose that $\phi(0) = 0$. Then F has a representation $F = \zeta H(\zeta) \overline{G(\zeta)}$, where H and G are analytic and have power series expansion

$$H(\zeta) = \sum_{n=0}^{\infty} A_n \zeta^n, \quad G(\zeta) = \sum_{n=0}^{\infty} B_n \zeta^n, \tag{4}$$

where $|A_0| = |B_0| = 1$.

Since the analytic part of $f(z)$ is $\phi(z)H(\phi(z)) = z^2/(1 - z)^2$, the function ϕ must have an expansion of the form

$$\phi(z) = c_2 z^2 + c_3 z^3 + \dots \tag{5}$$

It follows that $G \circ \phi$ has an expansion of the form $B_0 + C_1 z^2 + C_2 z^3 + \dots$. However, the given form of f shows that $G(\phi(z)) = 1/(1 - z)^2 = 1 + 2zz + 3z^2 + \dots$, this leads to contradiction. Hence, f has no factorization of the form $f = F \circ \phi$ of the required form in any neighborhood of the origin.

EXAMPLE 2. Let $f(z) = z^2|z^2|$ be defined in the unit disc U . Now, f is sense-preserving logharmonic mapping in U since the dilatation $a(z) = 1/3$. But here f has the desired factorization $f = F \circ \phi$, with $F(\zeta) = \zeta|\zeta|$ and $\phi(z) = z^2$.

Now, we state and prove the factorization theorem.

THEOREM 3. *Let f be a nonconstant logharmonic mapping defined on a domain $D \subset \mathbb{C}$ and let a be its dilatation function. Then, f can be factorized in the form $f = F \circ \phi$, for some analytic function ϕ and some univalent logharmonic mapping F if and only if*

- (a) $|a(z)| \neq 1$ on D ;
- (b) $f(z_1) = f(z_2)$ implies $a(z_1) = a(z_2)$.

Under these conditions, the representation is unique up to a conformal mapping; any other representation $f = F_1 \circ \phi_1$ has the form $F_1 = F \circ \psi^{-1}$ and $\phi_1 = \psi \circ \phi$ for some conformal mapping defined on $\phi(D)$.

PROOF. Suppose that $f = F \circ \phi$, where F is a univalent logharmonic mapping and ϕ is an analytic function. Let $A(\zeta)$ be the dilatation function of F . Then simple calculations give that $f_z = F_w(\phi)\phi'$, $f_{\bar{z}} = F_{\bar{w}}(\phi)\overline{\phi'}$, and $a(z) = A(\phi(z))$. Since F is univalent, the Jacobian is nonzero and hence $|a(z)| = |A(\phi(z))| \neq 1$ (see [2]). Also, F is univalent and $f(z_1) = f(z_2)$ implies that $\phi(z_1) = \phi(z_2)$. Hence, $a(z_1) = a(z_2)$.

Next, suppose that the two conditions are satisfied. We want to show that f can be factorized in the form $f = F(\phi)$. This is equivalent to finding a univalent continuous function G defined on $f(D)$ so that $G \circ f$ is analytic. In view of the Cauchy-Riemann conditions, this is equivalent to

$$(G_w b + G_{\bar{w}})\overline{f_z} = 0, \tag{6}$$

where $b(z) = \overline{a(z)}(f(z)/\overline{f(z)}) = \overline{f_z}/\overline{f_{\bar{z}}}$.

Hence, $-b(f^{-1}(w)) = G_{\bar{w}}/G_w$. Let $\mu(w) = G_{\bar{w}}/G_w$. Now, we show that μ is well defined. Suppose that $f(z_1) = f(z_2) = w$. Then, as $b(z_1) = \overline{a(z_1)}(f(z_1)/\overline{f(z_1)})$, $b(z_2) = \overline{a(z_2)}(f(z_2)/\overline{f(z_2)})$, and $a(z_1) = a(z_2)$, it follows that $b(z_1) = b(z_2)$. Hence, $\mu(w)$ is well defined and $|\mu(w)| \neq 1$ for all $w \in f(D)$.

Let $\{D_n\}$ be an exhaustion of D , $\Omega_n = f(D_n)$ and let μ_n be the restriction of μ to Ω_n . Extend μ_n to $\overline{\mathbb{C}}$ by assuming that $\mu_n \equiv 0$ on $\mathbb{C} \setminus \Omega_n$. Then the Beltrami equation $G_{\bar{w}} = \mu_n G_w$ has a quasiconformal solution G_n from \mathbb{C} on \mathbb{C} , see [8]. Let $G_n(\infty) = \infty$, then G_n is a homeomorphism on $\overline{\mathbb{C}}$. Replace the solution G_n with the solution

$$H_n(w) = \frac{G_n(w) - G_n(w_0)}{G_n(w_1) - G_n(w_0)}, \tag{7}$$

where $w_0, w_1 \in \Omega_1$ and $w_1 \neq w_0$. This is possible because f is not constant on D_1 . Then, H_n is also a homeomorphic solution to the Beltrami equation, normalized to satisfy $H_n(w_0) = 0$, $H_n(w_1) = 1$, and $H_n(\infty) = \infty$. This and the fact that each H_n is K -quasiconformal mapping on Ω_j imply that H_n converges locally uniformly to a K -quasiconformal mapping H on Ω_j . It follows that H is a homeomorphism on Ω and H satisfies the equation

$$H_{\bar{w}} = \mu H_w \quad \text{on } \Omega. \tag{8}$$

Hence, $\phi = H \circ f$ is analytic in D .

Next, we show that $F = H^{-1}$ is logharmonic mapping. Note that $f = F \circ \phi$ was assumed to be logharmonic in D . Then, near any point $\zeta = \phi(z)$ where $\phi'(z) \neq 0$, we can then deduce that $F = f \circ \phi^{-1}$ is logharmonic, where ϕ^{-1} is a local inverse. But F is locally bounded, so the (isolated) images of critical points of ϕ are removable, and F is logharmonic mapping on $\phi(D)$.

Finally, we prove the uniqueness. Suppose that $f = F \circ \phi = F_0 \circ \phi_0$. If we let $G_0 = F_0^{-1}$, then $G_0 \circ f = \phi_0$ is nonconstant and analytic, and $G_{0\bar{w}} = \mu G_{0w}$. But the solution of this Beltrami equation is unique; hence, $G_0 = G$. This completes the proof of the theorem. □

REFERENCES

- [1] Z. Abdulhadi, *Close-to-starlike logharmonic mappings*, Int. J. Math. Math. Sci. **19** (1996), no. 3, 563-574.
- [2] Z. Abdulhadi and D. Bshouty, *Univalent functions in $H \cdot \bar{H}(D)$* , Trans. Amer. Math. Soc. **305** (1988), no. 2, 841-849.
- [3] Z. Abdulhadi and W. Hengartner, *Spirallike logharmonic mappings*, Complex Variables Theory Appl. **9** (1987), no. 2-3, 121-130.
- [4] ———, *Univalent harmonic mappings on the left half-plane with periodic dilatations*, Univalent Functions, Fractional Calculus, and Their Applications (Kōriyama, 1988), Ellis Horwood Ser. Math. Appl., Horwood, Chichester, 1989, pp. 13-28.
- [5] ———, *Univalent logharmonic extensions onto the unit disk or onto an annulus*, Current Topics in Analytic Function Theory, World Scientific Publishing, New Jersey, 1992, pp. 1-12.
- [6] ———, *One pointed univalent logharmonic mappings*, J. Math. Anal. Appl. **203** (1996), no. 2, 333-351.
- [7] P. Duren and W. Hengartner, *A decomposition theorem for planar harmonic mappings*, Proc. Amer. Math. Soc. **124** (1996), no. 4, 1191-1195.
- [8] O. Lehto and K. I. Virtanen, *Quasiconformal Mappings in the Plane*, 2nd ed., Springer-Verlag, New York, 1973.

Zayid Abdulhadi: Department of Computer Science, Mathematics and Statistics,
American University of Sharjah (AUS), P.O. Box 26666, Sharjah, United Arab Emirates
E-mail address: zahadi@aus.ac.ae

Yusuf Abumuhanna: Department of Computer Science, Mathematics and Statistics,
American University of Sharjah (AUS), P.O. Box 26666, Sharjah, United Arab Emirates
E-mail address: ymuhanna@aus.ac.ae

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk