

***Naegleria fowleri* Lysate Induces Strong Cytopathic Effects and Pro-inflammatory Cytokine Release in Rat Microglial Cells**

Yang-Jin Lee^{1,†}, Chang-Eun Park^{2,†}, Jong-Hyun Kim¹, Hae-Jin Sohn¹, Jinyoung Lee¹, Suk-Yul Jung²
and Ho-Joon Shin^{1,*}

¹Department of Microbiology, and Department of Molecular Science and Technology, Ajou University School of Medicine, Suwon 443-721, Korea;

²Department of Biomedical Laboratory Science, College of Health and Medicine, Namseoul University, Chonan 331-707, Korea

Abstract: *Naegleria fowleri*, a ubiquitous free-living amoeba, causes fatal primary amoebic meningoencephalitis in humans. *N. fowleri* trophozoites are known to induce cytopathic changes upon contact with microglial cells, including necrotic and apoptotic cell death and pro-inflammatory cytokine release. In this study, we treated rat microglial cells with amoebic lysate to probe contact-independent mechanisms for cytotoxicity, determining through a combination of light microscopy and scanning and transmission electron microscopy whether *N. fowleri* lysate could effect on both necrosis and apoptosis on microglia in a time- as well as dose-dependent fashion. A ⁵¹Cr release assay demonstrated pronounced lysate induction of cytotoxicity (71.5%) toward microglial cells by 24 hr after its addition to cultures. In an assay of pro-inflammatory cytokine release, microglial cells treated with *N. fowleri* lysate produced TNF- α , IL-6, and IL-1 β , though generation of the former 2 cytokines was reduced with time, and that of the last increased throughout the experimental period. In summary, *N. fowleri* lysate exerted strong cytopathic effects on microglial cells, and elicited pro-inflammatory cytokine release as a primary immune response.

Key words: *Naegleria fowleri*, microglial cell, pro-inflammatory cytokine, cytotoxicity

Free-living *N. fowleri* is a ubiquitous, soil- and water-borne amoeba that can cause fatal primary amoebic meningoencephalitis (PAM) in both humans and animal models. PAM occurs predominantly in healthy young adults and adolescents and in non-immunocompromised children [1]. PAM patients generally have a history of swimming or recreation in ponds or hot-springs, or on river banks [2-4]. The development of PAM results from the invasion of trophozoites from amoeba-contaminated water into the nasal mucosa and their passage via the olfactory nerves to the forebrain, followed by multiplication in the tissues of the central nervous system (CNS) [5,6]. However, the precise mechanisms by which they exert their pathogenicity are unknown. Previous researches have suggested that infection may evoke both contact-dependent and contact-independent CNS pathology [7-11].

The mechanics of contact-dependent *N. fowleri* pathogenicity were previously observed to proceed via formation of vigorous pseudopodia and food-cup structures by trophozoites incubated with varying target cell-types [7,8]. Murine microglial cells experienced both necrotic and apoptotic cell death when exposed to *N. fowleri* trophozoites, with amoebic cytotoxicity increasing with the length of incubation [12]. Additionally, microglia co-cultured with *N. fowleri* trophozoites secreted the pro-inflammatory cytokines, TNF- α , IL-1 β , and IL-6 [12]. Amoebic cytotoxicity and the secretion of TNF- α from microglia were both reduced in co-culture experiments with *N. fowleri* by the addition of an antibody that specifically recognized Nfa1 protein, expressed on the pathogen's food-cups [12,13].

To elucidate the contact-independent mechanism for amoebic pathogenicity, researchers have attempted to examine the cytopathic effects of incubating *N. fowleri* secreted proteins (excretory-secretory proteins; ESPs) or acanthamoebic lysate or their secreted proteins against a number of target cell types [9-11,14,15]. However, no studies have examined the specific effects of *N. fowleri* lysate on microglial cells in experimental models of PAM.

• Received 14 February 2011, revised 28 June 2011, accepted 29 June 2011.

* Corresponding author (hjshin@ajou.ac.kr)

† These authors contributed equally to this work.

© 2011, Korean Society for Parasitology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Rat microglial cells can differentiate into 3 distinct morphologies, including an ameboid form unique to embryogenesis, a ramified form found in the mature brain, and a rod-shaped form that congregates around inflammatory lesions in the CNS [16,17]. They can function as phagocytotic cells and additionally produce cytokines, including IL-1, IL-6, and TNF- α [18,19]. Thus, microglial cells are considered an important target cell-type in *Naegleria* infection of the CNS, and have consequently been extensively studied to shed light on its mechanisms of pathogenesis [9,12].

In the present study, we have utilized a combination of light microscopy and scanning (SEM) and transmission electron microscopy (TEM) to identify the morphological alterations and potential cytopathic effects exerted by *N. fowleri* lysate on primary cultured rat microglial cells. Additionally, we have endeavored to characterize the host cell immune responses by amebic lysate in culture by assaying microglial release of pro-inflammatory cytokines.

N. fowleri trophozoites (Cater NF69 strain, ATCC NO. 30215, Manassas, Virginia, USA) were axenically cultured at 37°C in Nelson's medium [20]. Amebic lysate was prepared by freeze-thaw, and the soluble protein fraction assessed for degradation by SDS-PAGE electrophoresis (15% gel) as previously described [21]. Rat microglial cells were prepared using the methods of Giulian and Baker [16] and Oh et al. [12].

To observe the morphological changes induced by *N. fowleri* lysate, microglial cells (2×10^5) harvested from rats were incubated with differing concentrations of lysate (0.1, 0.5, or 1 mg/ml) adjusted to the same volume in PBS (pH 7.4) for 3, 6, or 12 hr in 24-well culture plates. PBS (100 μ l) and lipopolysaccharide (LPS) (1 μ g/ml) incubation were used as controls. After incubation, samples were fixed with 2.5% glutaraldehyde in 0.1 M cacodylate buffer (pH 7.4), post-fixed with 1% osmium tetroxide-1.5% potassium for 1 hr, and imaged with a light microscope. For electron microscopies (SEM or TEM), harvested rat microglial cells (2×10^5) were seeded onto Lab-Tek II chamber slides (Nunc A/S, Roskilde, Belgium), and 1 mg/ml of *N. fowleri* lysate was added to the culture system. After incubation for 3, 6, or 12 hr, SEM examination was carried out using the method of Oh et al. [12]. An image analyzer program (Escan 4000, Bumi-Mi Universe Co., Ltd., Ansan, Korea) was used in SEM image capture (S800, Hitachi, Tokyo, Japan). Additionally, harvested microglial cells treated with *N. fowleri* lysate (1 mg/ml) for 3, 6, or 12 hr were prepared for TEM via published methods [12]. After the blocks were sectioned and

stained with Ultrastain 1H and 2 (Leica, Vienna, Austria), microglia were observed and imaged using a Zeiss EM 902A TEM (Leo, Oberkochen, Germany).

To quantitate the cytotoxicity of *N. fowleri* lysate against target microglial cells, a ^{51}Cr (chromium) release assay was performed according to methods published previously [12]. Labeled microglia (5×10^4) were treated with *N. fowleri* lysate (0.5 or 1 mg/ml) in 5% CO_2 for 3, 6, 12, or 24 hr. The percentage of radioisotope released by microglial cells was used as an index of lysis, as calculated by the following formula: Cytotoxicity (%) = (experimental release - spontaneous release)/(maximum release - spontaneous release) \times 100.

The amounts of TNF- α , IL-1 β , and IL-6 released by lysate-treated microglial cells were analyzed according to published methods [12]. All assays used ELISA kits (BioSource International, Camarillo, California, USA) and were performed in triplicate. Data are shown as mean \pm SD and were statistically analyzed by the Student's t-test. Values of $P < 0.05$ were considered significant.

N. fowleri lysate (1 mg/ml) exerted severe morphological changes, including disruption and breakdown of the plasma membrane on the rat microglia, and significantly promoted cell death (Fig. 1). These effects were dose-dependent (Fig. 1). Microglial lethality was almost complete within 6 hr culture with *N. fowleri* lysate, generally appearing necrotic in nature and characterized by membrane swelling and lysis of the nuclear membrane and nucleus (Fig. 2A, 2C). Some microglial cells additionally evinced hallmarks of apoptosis, such as blebbing, formation of apoptotic bodies, and nuclear chromatin condensation (Fig. 2B, 2D). In previous reports, pathogenic *Acanthamoeba culbertsoni* lysate induced cytopathic effects, including necrosis and apoptosis, in primary rat microglial cells [21,22], suggesting that *N. fowleri* contact-independent cytotoxicity proceeds similarly.

Table 1 depicts the cytotoxic effects of treatment with *N. fowleri* lysate on rat microglia. Cell death increased dependent on the length of incubation, reaching 14.6, 21.9, 38.5, and 71.5% at 3, 6, 12, and 24 hr, respectively, by a ^{51}Cr release assay. Cell death was also dose-dependent. Kim et al. [10,11] performed an LDH release assay to evaluate the cytotoxicity of *N. fowleri* ESPs on human microglial cells or Chinese hamster ovary (CHO) cells, and likewise found a time- and dose-dependent rise in cytotoxicity consistent with a pathogenic role for ESPs [10,11]. An analysis of *N. fowleri* whole-cell lysate found 2 major protein bands (13 and 21 kDa), predicted to be the Nfa1




Fig. 1. Light microscopic images of microglia treated with *N. fowleri* lysate for 12 hr. Microglial cells were treated with media only (A), PBS (B), LPS (1 µg/ml) (C), or different concentrations of *N. fowleri* lysate (0.1, 0.5 and 1 mg/ml; D-F, respectively). Microglia demonstrated significant, dose-dependent reduction in cell density. $\times 200$.

and proteinase proteins [11,21]. These may constitute efficacious cytolytic factors in amebic lysate, and an in-depth fractionation and biochemical characterization will help elucidate the pathogenicity induced by amebic lysate.

We queried release of pro-inflammatory cytokines by microglia treated with *N. fowleri* lysate. TNF- α release by microglia cultured with *N. fowleri* lysate (1 mg/ml) was 121.6, 90.4, and 81.0 pg/ml at 3, 6, and 12 hr, respectively (Fig. 3A). IL-1 β release was 88.5, 92.7, and 190.1 pg/ml, and IL-6 was 298.8, 414.9, and 251.1 pg/ml, after identical incubations (Fig. 3B,

3C). Microglial cells produce pro-inflammatory cytokines, such as TNF- α , IL-1 β , IL-3, IL-6, IL-8, IL-12, and IL-15, as a primary inflammatory immune response to host-cell damage; release of anti-inflammatory cytokines, e.g. IL-10 and TGF- β , has been reported in defense against parasitic brain infection [8,22]. We found that TNF- α and IL-6 peaked early in microglial incubation with amebic lysate, subsequently declining until the time-course's end; in contrast, IL-1 β increased throughout the duration of the experiment. Microglial cells are therefore likely involved in the early inflammatory stage of *N. fowleri* infection




Fig. 2. TEM and SEM images of microglia treated with *N. fowleri* lysate. Microglial cells were treated with *N. fowleri* lysate (1 mg/ml) for 12 hr. Necrosis could be observed concomitantly with plasma membrane lysis (arrows; A, C), and programmed cell death with accompanying changes, including apoptotic bodies (arrows) and nuclear chromatin condensation (B, D). Bar = 5 μ m.

Table 1. Cytotoxicity of *Naegleria fowleri* lysate applied to microglial cells as assessed by ^{51}Cr -release assay

Groups	Length of cultivation (hr)			
	3	6	12	24
Microglia + lysate (0.5 mg/ml)	7.9 \pm 4.0 ^a	14.4 \pm 1.5	23.8 \pm 3.3	51.5 \pm 3.7
Microglia + lysate (1.0 mg/ml)	14.6 \pm 4.5	21.9 \pm 1.3	38.5 \pm 6.5	71.5 \pm 1.4

^aValues are % cytotoxicity, which was calculated from the amount of ^{51}Cr released from radio-labelled microglial cells (mean \pm SD).

through their secretion of various inflammatory cytokines, though more detailed studies will be necessary to fully illuminate the crosstalk between *N. fowleri* and host cells.

Acanthamoeba, comprising of several species of free-living amoebae that cause chronic granulomatous amebic encephalitis (GAE) in humans, evokes cytopathic effects on hosts through amebic adhesion to and phagocytosis of target cells, facilitated by its production of proteolytic enzymes [15,23,24], including




Fig. 3. Pro-inflammatory cytokine release from microglia treated with *N. fowleri* lysate. Treatment of microglial cells with *N. fowleri* lysate for 3, 6, or 12 hr induced dose-dependent secretion of TNF- α (A), IL-1 β (B), and IL-6 (C). M: microglia, L: *N. fowleri* lysate. * $P < 0.05$, ** $P < 0.05$, *** $P < 0.05$.

serine proteases, contactdependent metalloproteases, elastases, cysteine proteases, and cytotoxic proteinases induced by mannose-mediated adhesion [14,25-27]. Similar contact-depen-

dent and independent mechanisms have been proposed to contribute to *N. fowleri* pathogenicity [7-11]. Our findings confirm that *N. fowleri* lysate induces morphological cytopathic changes in microglial cells contact-independently, evoking necrotic as well as apoptotic cytotoxicity. In addition, microglia produced pro-inflammatory cytokines as a primary immune response concomitant with exposure to *N. fowleri* lysate.

ACKNOWLEDGEMENTS

This work (2011-0015429) was supported by Mid-career Researcher Program through NRF grant funded by the MEST.

REFERENCES

- Schuster FL, Visvesvara GS. Free-living amoebae as opportunistic and non-opportunistic pathogens of humans and animals. *Int J Parasitol* 2004; 34: 1001-1027.
- Im KI, Shin HJ. Pathogenic free-living amoebae in Korea. *Korean J Parasitol* 2004; 42: 93-119.
- John DT. Primary amebic meningoencephalitis and the biology of *Naegleria fowleri*. *Annu Rev Microbiol* 1982; 36: 101-123.
- Visvesvara GS, Moura H, Schuster FL. Pathogenic and opportunistic free-living amoebae: *Acanthamoeba* spp., *Balamuthia mandrillaris*, *Naegleria fowleri*, and *Sappinia diploidea*. *FEMS Immunol Med Microbiol* 2007; 50: 1-26.
- Carter RF. Primary amoebic meningoencephalitis: An appraisal of present knowledge. *Trans R Soc Trop Med Hyg* 1972; 66: 193-213.
- Ma P, Visvesvara GS, Martinez AJ, Theodore FH, Daggett PM, Sawyer TK. *Naegleria* and *Acanthamoeba* infection: Review. *Rev Infect Dis* 1990; 12: 490-513.
- Marciano-Cabral F. Biology of *Naegleria* spp. *Microbiol Rev* 1988; 52: 114-133.
- Cho MS, Jung SY, Park S, Kim KH, Kim HI, Sohn S, Kim HJ, Im KI, Shin HJ. Immunological characterizations of a cloned 13.1-kilodalton protein from pathogenic *Naegleria fowleri*. *Clin Diagn Lab Immunol* 2003; 10: 954-959.
- Marciano-Cabral F, Ludwick C, Puffenbarger RA, Cabral GA. Differential stimulation of microglial pro-inflammatory cytokines by *Acanthamoeba culbertsoni* versus *Acanthamoeba castellanii*. *J Eukaryot Microbiol* 2004; 51: 472-479.
- Kim JH, Kim DS, Shin HJ. Contact-independent cell death of human microglial cells due to pathogenic *Naegleria fowleri* trophozoites. *Korean J Parasitol* 2008; 46: 217-221.
- Kim JH, Yang AH, Sohn HJ, Kim D, Song KJ, Shin HJ. Immunodominant antigens in *Naegleria fowleri* excretory-secretory proteins were potential pathogenic factors. *Parasitol Res* 2009; 105: 1675-1681.
- Oh YH, Jeong SR, Kim JH, Song KJ, Kim K, Park S, Sohn S, Shin

- HJ. Cytopathic changes and pro-inflammatory cytokines induced by *Naegleria fowleri* trophozoites in rat microglial cells and protective effects of an anti-Nfa1 antibody. *Parasite Immunol* 2005; 27: 453-459.
13. Jeong SR, Kang SY, Lee SC, Song KJ, Im KI, Shin HJ. Decreasing effect of an anti-Nfa1 polyclonal antibody on the in vitro cytotoxicity of pathogenic *Naegleria fowleri*. *Korean J Parasitol* 2004; 42: 35-40.
 14. Mitro K, Bhagavathiammai A, Zhou OM, Bobbett G, McKerrow JH, Chokshi R, Chokshi B, James ER. Partial characterization of the proteolytic secretions of *Acanthamoeba polyphaga*. *Exp Parasitol* 1994; 78: 377-385.
 15. Yang Z, Cao Z, Panjwani N. Pathogenesis of *Acanthamoeba keratitis*: Carbohydrate-mediated host-parasite interactions. *Infect Immun* 1997; 65: 439-445.
 16. Giulian D, Baker TJ. Characterization of ameboid microglia isolated from developing mammalian brain. *J Neurosci* 1986; 6: 2163-2178.
 17. Suzumura A, Marunouchi T, Yamamoto H. Morphological transformation of microglia in vitro. *Brain Res* 1991; 545: 301-306.
 18. Chao CC, Gekker G, Hu S, Peterson PK. Human microglial cell defense against *Toxoplasma gondii*. The role of cytokines. *J Immunol* 1994; 152: 1246-1252.
 19. Suzumura A, Sawada M, Yamamoto H, Marunouchi T. Transforming growth factor-beta suppresses activation and proliferation of microglia in vitro. *J Immunol* 1993; 151: 2150-2158.
 20. Willaert E. Isolation and in vitro culture of the amoeba of the genus *Naegleria*. *Ann Soc Belges Med Trop Parasitol Mycol* 1971; 51: 701-708.
 21. Shin HJ, Cho MS, Jung SU, Kim HI, Park S, Kim HJ, Im KI. Molecular cloning and characterization of a gene encoding a 13.1 kDa antigenic protein of *Naegleria fowleri*. *J Eukaryot Microbiol* 2001; 48: 713-717.
 22. Benedetto N, Folgore A, Romano Carratelli C, Galdiero F. Effects of cytokines and prolactin on the replication of *Toxoplasma gondii* in murine microglia. *Eur Cytokine Netw* 2001; 12: 348-358.
 23. Shin HJ, Cho MS, Jung SY, Kim HI, Park S, Seo JH, Yoo JC, Im KI. Cytopathic change in rat microglial cells induced by pathogenic *Acanthamoeba culbertsoni*: Morphology and cytokine release. *Clin Diagn Lab Immunol* 2001; 8: 837-840.
 24. Moore MB, Ubelaker JE, Marin JH, Silvano R, Dougherty JM, Meyer DR, McCulley JP. In vitro penetration of human corneal epithelium by *Acanthamoeba castellanii*: A scanning and transmission electron microscopy study. *Cornea* 1991; 10: 291-298.
 25. Hadas E, Mazur T. Proteolytic enzymes of pathogenic and non-pathogenic strains of *Acanthamoeba* spp. *Trop Med Parasitol* 1993; 44: 197-200.
 26. Khan NA, Jarroll EL, Panjwani N, Cao Z, Paget TA. Proteases as markers for differentiation of pathogenic and non-pathogenic species of *Acanthamoeba*. *J Clin Microbiol* 2000; 38: 2858-2861.
 27. Leher H, Silvano R, Alizadeh H, Huang J, Niederkorn JY. Mannose induces the release of cytopathic factors from *Acanthamoeba keratitis*. *Infect Immun* 1988; 66: 5-10.