

Cardiac autonomic regulation during hypoxic exercise

 James P. Fisher

School of Sport, Exercise and Rehabilitation Sciences, College of Life and Environmental Sciences, University of Birmingham, Edgbaston, Birmingham, United Kingdom

AUTONOMIC NERVOUS SYSTEM REGULATION of the heart and vasculature plays a key role in enabling the tight coupling between oxygen consumption of exercising skeletal muscles and oxygen delivery. The withdrawal of cardiac parasympathetic nerve activity and elevation in cardiac sympathetic nerve activity facilitate the exercise intensity-dependent increase in heart rate, ventricular contractility, stroke volume, and cardiac output (5). Robinson et al. (10) laid the foundations for the present understanding of cardiac autonomic regulation in exercising humans with the demonstration that administration of atropine (to block cholinergic muscarinic receptors) markedly diminishes elevations in heart rate only during low-to-moderate intensity dynamic exercise, whereas administration of propranolol (to block β -adrenergic receptors) principally attenuates the cardiac acceleration during strenuous dynamic exercise. In the current issue of the *American Journal of Physiology-Heart and Circulatory Physiology*, Siebenmann et al. (12) use a similar methodological approach to provide new insights into cardiac autonomic regulation during hypoxic exercise in humans.

Hypoxia elevates heart rate at rest and during submaximal exercise (2). This serves to enhance oxygen delivery by increasing cardiac output in an attempt to protect against the reduction in arterial oxygen content. The heightened activation of the sympathetic nervous system has historically been the prime candidate for this cardiac acceleration, partly because catecholamine release and muscle sympathetic nerve activity are increased during hypoxic exercise (7, 9); however, the withdrawal of cardiac parasympathetic activity has also been implicated (8). Hopkins et al. (6) observed that during exercise, the additional elevation in heart rate and cardiac output caused by hypoxia persisted with either cholinergic or β -adrenergic receptor blockade. This raised the intriguing possibility that an important alternative mechanism is involved. However, the existence of complex pre- and postsynaptic interactions between the cardiac parasympathetic and sympathetic fibers means that a compensatory action by the nonblocked portion of the autonomic nervous system cannot be ruled out.

The laudable study by Siebenmann et al. (12) addresses this issue by comparing heart rate and cardiac output responses to an exhaustive incremental exercise protocol in normoxia and hypoxia (inspired oxygen fraction = 12%) under control (no drug) conditions, separate β -adrenergic blockade (propranolol) and cholinergic blockade (glycopyrrolate), and, critically, combined β -adrenergic and cholinergic blockade. Interestingly, the hypoxia-induced increase in heart rate under control conditions (averaged across all exercise workloads) was similar to that with β -adrenergic blockade but reduced by $\approx 50\%$ with cholinergic blockade and by $\approx 60\%$ with combined β -adrenergic and cholinergic block-

ade. There are a number of implications to these findings, with two of the most significant being that the withdrawal of cardiac parasympathetic activity is an important component of the additional increase in heart rate during exercise in hypoxia and that the explanation for a significant proportion of the heart rate response to hypoxic exercise remains obscure. The former may be explained by the activation of metabolically sensitive group III and IV skeletal muscle afferents, which are better known for their sympathoexcitatory properties but may also inhibit cardiac parasympathetic activity (4). Although not an inconsiderable undertaking, future studies in which the activity of group III and IV skeletal muscle afferents is experimentally attenuated during hypoxic exercise, e.g., using intrathecal fentanyl (1), with the concomitant blockade of β -adrenergic and cholinergic receptors might provide further insight into their role.

Siebenmann et al. (12) discount several possible explanations for the hypoxia-induced increase in heart rate during exercise that occurs independently of β -adrenergic and cholinergic receptor stimulation, including incomplete cardiac autonomic blockade (although not directly tested) and the Bainbridge reflex, and instead propose the involvement of an α -adrenergic mechanism. The chronotropic responses to α -adrenergic stimulation are not widely recognized, and, ordinarily, the administration of an α -adrenoreceptor agonist evokes a pressor response that decreases heart rate via a baroreflex mechanism. However, α_1 -adrenoreceptor stimulation (phenylephrine) will evoke an increase in heart rate in young humans when administered with complete β -adrenergic and cholinergic blockade (11). It is tempting to speculate whether the addition of an α -adrenoreceptor antagonist (e.g., prazosin, phentolamine) administered during exercise along with combined β -adrenergic and cholinergic blockade would abolish the cardiac acceleration evoked by hypoxia. Aside from classical adrenergic or cholinergic transmission, the contribution of peptidic neurotransmission to the hypoxia-induced heart rate responses described cannot be ruled out (3).

The experiments undertaken by Siebenmann and colleagues (12) provide an important advance in our understanding of cardiac autonomic regulation via β -adrenergic and cholinergic receptors during hypoxic exercise. Careful studies are now needed to reveal the mechanism accounting for the unexplained portion of the increase in heart rate occurring in hypoxic exercise.

DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the author(s).

AUTHOR CONTRIBUTIONS

J.P.F. drafted, edited, revised, and approved final version of manuscript.

REFERENCES

1. Amann M, Blain GM, Proctor LT, Sebranek JJ, Pegelow DF, Dempsey JA. Group III and IV muscle afferents contribute to ventilatory and cardiovascular response to rhythmic exercise in humans. *J Appl Physiol* (1985) 109: 966–976, 2010.

Address for reprint requests and other correspondence: J. P. Fisher, School of Sport, Exercise and Rehabilitation Sciences, College of Life and Environmental Sciences, Univ. of Birmingham, Edgbaston, Birmingham, B15 2TT, England (e-mail: j.p.fisher@bham.ac.uk).

2. **Bartsch P, Gibbs JS.** Effect of altitude on the heart and the lungs. *Circulation* 116: 2191–2202, 2007.
3. **Beaulieu P, Lambert C.** Peptidic regulation of heart rate and interactions with the autonomic nervous system. *Cardiovasc Res* 37: 578–585, 1998.
4. **Fisher JP, Adlan AM, Shantsila A, Secher JF, Sorensen H, Secher NH.** Muscle metaboreflex and autonomic regulation of heart rate in humans. *J Physiol* 591: 3777–3788, 2013.
5. **Fisher JP, Young CN, Fadel PJ.** Autonomic adjustments to exercise in humans. *Compr Physiol* 5: 475–512, 2015.
6. **Hopkins SR, Bogaard HJ, Niizeki K, Yamaya Y, Ziegler MG, Wagner PD.** Beta-adrenergic or parasympathetic inhibition, heart rate and cardiac output during normoxic and acute hypoxic exercise in humans. *J Physiol* 550: 605–616, 2003.
7. **Katayama K, Ishida K, Iwamoto E, Iemitsu M, Koike T, Saito M.** Hypoxia augments muscle sympathetic neural response to leg cycling. *Am J Physiol Regul Integr Comp Physiol* 301: R456–R464, 2011.
8. **Koller EA, Drechsel S, Hess T, Macherel P, Boutellier U.** Effects of atropine and propranolol on the respiratory, circulatory, and ECG responses to high altitude in man. *Eur J Appl Physiol Occup Physiol* 57: 163–172, 1988.
9. **Mazzeo RS, Bender PR, Brooks GA, Butterfield GE, Groves BM, Sutton JR, Wolfel EE, Reeves JT.** Arterial catecholamine responses during exercise with acute and chronic high-altitude exposure. *Am J Physiol Endocrinol Metab* 261: E419–E424, 1991.
10. **Robinson BF, Epstein SE, Beiser GD, Braunwald E.** Control of heart rate by the autonomic nervous system. Studies in man on the interrelation between baroreceptor mechanisms and exercise. *Circ Res* 19: 400–411, 1966.
11. **Saitoh H, Nomura A, Osaka M, Sasabe N, Atarashi H, Hayakawa H.** Developmental changes of alpha 1-adrenergic chronotropic action on human sinus node. *Am J Cardiol* 76: 89–91, 1995.
12. **Siebenmann C, Rasmussen P, Sørensen H, Bonne TC, Zaar M, Aachmann-Andersen NJ, Nordsborg NB, Secher NH, Lundby C.** Hypoxia increases exercise heart rate despite combined inhibition of β -adrenergic and muscarinic receptors. *Am J Physiol Heart Circ Physiol* 308: 2015. doi:10.1152/ajpheart.00861–2014.

