

Cubic Spline Approximation of a Circle with Maximal Smoothness and Accuracy

Christian Apprich, Anne Dietrich,
Klaus Höllig, Esfandiar Nava-Yazdani

Stuttgarter
Mathematische
Berichte

2017/004

Fachbereich Mathematik
Fakultät Mathematik und Physik
Universität Stuttgart
Pfaffenwaldring 57
D-70 569 Stuttgart

E-Mail: preprints@mathematik.uni-stuttgart.de
WWW: <http://www.mathematik.uni-stuttgart.de/preprints>

ISSN **1613-8309**

© Alle Rechte vorbehalten. Nachdruck nur mit Genehmigung des Autors.
L^AT_EX-Style: Winfried Geis, Thomas Merkle

Abstract

We construct cubic spline approximations of a circle which are four times continuously differentiable and converge with order six.

Keywords. Bézier curve, cubic spline, geometric smoothness, high accuracy approximation of circles

Applying concepts of differential geometry to CAGD has led to a number of interesting results. An example is the surprising fact that spline curves can approximate with higher order than spline functions. In addition, smoothness constraints become less restrictive. A general statement to this effect is the following

Conjecture. For smooth curves in \mathbb{R}^d , which satisfy mild generic assumptions, there exist spline approximations of degree $\leq n$ which are $\alpha = n - 1 + \lfloor (n - 1)/(d - 1) \rfloor$ times continuously differentiable and converge with order $\alpha + 2$.

For low degree, many schemes of optimal order have been proposed (cf. [3, 7] and the references given there for an overview). However, except for the simplest case $n = d = 2$ treated in [6], the gain in accuracy has not yet been combined with a possible improvement of smoothness. This motivated this short note, which, aside from some obvious practical relevance, supports the conjecture for a canonical test case.

The construction of a, in some sense optimal, cubic spline approximation $t \mapsto p(t)$ of a circle is straightforward. We interpolate the circle at the points

$$P_k = (\cos(k\varphi), \sin(k\varphi)), \quad \varphi = 2\pi/m,$$

which separate the m cubic Bézier segments of p , and choose $k = 0, 1, \dots, m - 1$ as knots. Moreover, we match the tangents at P_k . For a completely symmetric configuration, this leaves the length of the tangent vectors as the only degree of freedom. In terms of a corresponding parameter δ , the interior control points of the curve segments are

$$P_k^\pm = P_k \pm \delta(-\sin(k\varphi), \cos(k\varphi)), \quad \delta = |p'(k)|/3.$$

In particular, as depicted in the figure,

$$(1, 0), \quad (1, \delta), \quad (\cos \varphi, \sin \varphi) - \delta(-\sin \varphi, \cos \varphi), \quad (\cos \varphi, \sin \varphi)$$

are the control points for the Bézier segment from P_0 to P_1 .

Each Bézier segment has the curvature

$$\kappa = \frac{2(1 - \cos \varphi - \delta \sin \varphi)}{3\delta^2}$$

at its endpoints. Choosing the parameter δ to match the curvature of the circle leads to the geometric Hermite interpolant [2]. While this approximation is sixth order accurate as $\varphi \rightarrow 0$, it merely has the standard smoothness $\alpha - 2 = 2$. Several other constructions have been proposed [4, 5, 1], focussing on minimizing the error while maintaining standard smoothness. The alternative, considered in this note, is to achieve the maximal order of differentiability $\alpha = 4$. To this end, we exploit the well known fact that, for geometric smoothness, the parametrization needs not to be smooth. For our purposes, the following definition is most convenient.

Geometric Smoothness. Identifying a planar curve p in a neighborhood of a point P_0 with the graph of a function f , the smoothness of p and f at P_0 are equivalent.

We apply this criterion to the spline $t \mapsto p(t) = (x(t), y(t))$ in a neighborhood of $P_0 = (1, 0)$. Since the tangent at P_0 is vertical,

$$x(t) = f(y(t)), \quad t \approx 0.$$

By symmetry, f is an even function ($f(y) = f(-y)$), implying that all its even derivatives are continuous at $y(0) = 0$. Moreover, since by construction

$$p'(0) = (x'(0), y'(0)) = (0, 3\delta)$$

is the common tangent vector of the cubic segments, joining at $t = 0$, $f'(0)$ is zero. We will now choose δ so that

$$f'''(0^+) = 0,$$

which, by symmetry, implies $f'''(0^-) = 0$, and thus yields a four times continuously differentiable spline curve; f'''' is continuous by symmetry, as remarked above. By the chain rule,

$$\begin{aligned} x' &= f'(y)y' \\ x'' &= f''(y)(y')^2 + f'(y)y'' \\ x''' &= f'''(y)(y')^3 + 3f''(y)y''y' + f'(y)y''' . \end{aligned}$$

Substituting $y(0) = 0$ and $x'(0) = 0$ yields

$$f'(0) = 0, \quad f''(0) = x''(0)/(y'(0))^2$$

and, by the third equation, $f'''(0^+) = 0$ is equivalent to

$$x'''(0) y'(0) = 3x''(0)y''(0).$$

Expressing the derivatives in terms of the components of the Bézier control points, we obtain a quadratic equation for the parameter δ :

$$3 \sin \varphi (3 + 2 \cos \varphi) \delta^2 + 4(3 \cos^2 \varphi + \cos \varphi - 4) \delta + 6 \sin \varphi (1 - \cos \varphi) = 0.$$

Both solutions

$$\delta_{\pm} = \frac{1 - \cos \varphi}{3 \sin \varphi (3 + 2 \cos \varphi)} \left(2(4 + 3 \cos \varphi) \pm \sqrt{2(5 + 3 \cos \varphi)} \right)$$

yield C^4 cubic spline curves.

We now show that the spline curve corresponding to $\delta = \delta_-$ not only has maximal smoothness $\alpha = 4$, but is also sixth order accurate. The other solution δ_+ merely yields the standard accuracy and will not be considered in the sequel. By symmetry, and in view of the Hermite interpolation conditions at the endpoints,

$$e(t) := 1 - x(t)^2 - y(t)^2 = t^3(1-t)^3 g(\varphi) + t^2(1-t)^2 h(\varphi),$$

where computer algebra provides the explicit expressions

$$\begin{aligned} g(\varphi) &= 18(1 + \cos \varphi) \delta^2 - 24\delta \sin \varphi + 8(1 - \cos \varphi), \\ h(\varphi) &= -9\delta^2 - 6\delta \sin \varphi + 6(1 - \cos \varphi). \end{aligned}$$

It is not difficult to show that e is positive on $[0, 1]$ with a unique maximum at $t = 1/2$:

$$\begin{aligned} E &:= \max_{0 \leq t \leq 1} e(t) = e(1/2) = \frac{9}{32}(\cos \varphi - 1) \delta^2 - \frac{3}{4} \delta \sin \varphi + \frac{1}{2} - \frac{1}{2} \cos \varphi \\ &= \frac{3}{4096} \varphi^6 + O(\varphi^8). \end{aligned}$$

In a final step, we can reduce the one-sided error by scaling the parametrization p to obtain $2m$ equal oscillations with radii $1 \pm E_*$. Observing that, by definition of $e(t)$, the distance of $p(t)$ from the origin varies from $\sqrt{1 - E}$ to 1, we define

$$p_{\star} = \frac{2}{1 + \sqrt{1 - E}} p$$




Fig. Bézier segment with control points (left), graphs of κ , κ' and κ'' (right)

as our optimal spline approximation with the error

$$E_{\star} = \frac{1 - \sqrt{1 - E}}{1 + \sqrt{1 - E}} = \frac{3}{16384} \varphi^6 + O(\varphi^8).$$

The elementary example presented in this note motivates the construction of general spline curves with maximal smoothness from control polygons. While, in view of the nonlinearity of the geometric smoothness constraints analytical solutions seem out of reach, with today's computing resources, numerical techniques are quite promising. We think that it is worthwhile to pursue this in the future.

References

- [1] Y. Ahn, H. Kim: *Approximation of circular arcs by Bezier curves*, J. of Comp. and Appl. Math. 81 (1997), 145–163.
- [2] C. De Boor, K. Höllig, M. Sabin: *High accuracy geometric Hermite interpolation*, Computer Aided Geometric Design, Volume 4 (1987), 269–278.
- [3] W. Degen: *Geometric Hermite Interpolation-: In memoriam Josef Hoschek*, Computer Aided Geometric Design, Volume 22 (2005), 573–592.
- [4] T. Dokken, M. Dæhlen, T. Lyche, K. Mørken: *Good approximation of circles by curvature continuous Bezier curves*, Computer Aided Geometric Design, Volume 7 (1990), 33–41.
- [5] M. Goldapp: *Approximation of circular arcs by cubic polynomials*, Computer Aided Geometric Design, Volume 8 (1991), 227–238.
- [6] K. Höllig and J. Koch: *Geometric Hermite interpolation with maximal order and smoothness*, Computer Aided Geometric Design, Volume 13 (1996), 681–695.
- [7] A. Rababah: *The Best Uniform Cubic Approximation of Circular Arcs with High Accuracy*, Communications in Mathematics and Applications, Volume 7 (2016), 37–46.

Christian Apprich
Fachbereich Mathematik, Universität Stuttgart, Pfaffenwaldring 57, 70569 Stuttgart, Germany
E-Mail: christian.apprich@mathematik.uni-stuttgart.de
WWW: <http://www.imng.uni-stuttgart.de/LstNumGeoMod/Apprich/index.html>

Anne Dietrich
E-Mail: anne.d@live.de

Klaus Höllig
Fachbereich Mathematik, Universität Stuttgart, Pfaffenwaldring 57, 70569 Stuttgart, Germany
E-Mail: klaus.hoellig@mathematik.uni-stuttgart.de
WWW: <http://www.imng.uni-stuttgart.de/institut/mitarbeiter/Hoellig.html>

Esfandiar Nava-Yazdani
Fachbereich Mathematik, Universität Stuttgart, Pfaffenwaldring 57, 70569 Stuttgart, Germany
E-Mail: navayaz@mathematik.uni-stuttgart.de

Erschienenene Preprints ab Nummer 2012-001

Komplette Liste: <http://www.mathematik.uni-stuttgart.de/preprints>

- 2017-007 *Blunck, A.; Knarr, N.; Stroppel, B.; Stroppel, M.:* Transitive groups of similitudes generated by octonions
- 2017-006 *Blunck, A.; Knarr, N.; Stroppel, B.; Stroppel, M.:* Clifford parallelisms by octonions
- 2017-005 *Knarr, N.; Stroppel, M.:* Subforms of Norm Forms of Octonion Fields
- 2017-004 *Apprich, C.; Dieterich, A.; Höllig, K.; Nava-Yazdani, E.:* Cubic Spline Approximation of a Circle with Maximal Smoothness and Accuracy
- 2017-003 *Fischer, S.; Steinwart, I.:* Sobolev Norm Learning Rates for Regularized Least-Squares Algorithm
- 2017-002 *Farooq, M.; Steinwart, I.:* Learning Rates for Kernel-Based Expectile Regression
- 2017-001 *Bauer, B.; Devroye, L.; Kohler, M.; Krzyzak, A.; Walk, H.:* Nonparametric Estimation of a Function From Noiseless Observations at Random Points
- 2016-006 *Devroye, L.; Györfi, L.; Lugosi, G.; Walk, H.:* On the measure of Voronoi cells
- 2016-005 *Kohls, C.; Kreuzer, C.; Rösch, A.; Siebert, K.G.:* Convergence of Adaptive Finite Elements for Optimal Control Problems with Control Constraints
- 2016-004 *Blaschzyk, I.; Steinwart, I.:* Improved Classification Rates under Refined Margin Conditions
- 2016-003 *Feistauer, M.; Roskovec, F.; Sändig, AM.:* Discontinuous Galerkin Method for an Elliptic Problem with Nonlinear Newton Boundary Conditions in a Polygon
- 2016-002 *Steinwart, I.:* A Short Note on the Comparison of Interpolation Widths, Entropy Numbers, and Kolmogorov Widths
- 2016-001 *Köster, I.:* Sylow Numbers in Spectral Tables
- 2015-016 *Hang, H.; Steinwart, I.:* A Bernstein-type Inequality for Some Mixing Processes and Dynamical Systems with an Application to Learning
- 2015-015 *Steinwart, I.:* Representation of Quasi-Monotone Functionals by Families of Separating Hyperplanes
- 2015-014 *Muhammad, F.; Steinwart, I.:* An SVM-like Approach for Expectile Regression
- 2015-013 *Nava-Yazdani, E.:* Splines and geometric mean for data in geodesic spaces
- 2015-012 *Kimmerle, W.; Köster, I.:* Sylow Numbers from Character Tables and Group Rings
- 2015-011 *Györfi, L.; Walk, H.:* On the asymptotic normality of an estimate of a regression functional
- 2015-010 *Gorodski, C, Kollross, A.:* Some remarks on polar actions
- 2015-009 *Apprich, C.; Höllig, K.; Hörner, J.; Reif, U.:* Collocation with WEB-Splines
- 2015-008 *Kabil, B.; Rodrigues, M.:* Spectral Validation of the Whitham Equations for Periodic Waves of Lattice Dynamical Systems
- 2015-007 *Kollross, A.:* Hyperpolar actions on reducible symmetric spaces
- 2015-006 *Schmid, J.; Griesemer, M.:* Well-posedness of Non-autonomous Linear Evolution Equations in Uniformly Convex Spaces
- 2015-005 *Hinrichs, A.; Markhasin, L.; Oettershagen, J.; Ullrich, T.:* Optimal quasi-Monte Carlo rules on higher order digital nets for the numerical integration of multivariate periodic functions
- 2015-004 *Kutter, M.; Rohde, C.; Sändig, A.-M.:* Well-Posedness of a Two Scale Model for Liquid Phase Epitaxy with Elasticity

- 2015-003 *Rossi, E.; Schleper, V.:* Convergence of a numerical scheme for a mixed hyperbolic-parabolic system in two space dimensions
- 2015-002 *Döring, M.; Györfi, L.; Walk, H.:* Exact rate of convergence of kernel-based classification rule
- 2015-001 *Kohler, M.; Müller, F.; Walk, H.:* Estimation of a regression function corresponding to latent variables
- 2014-021 *Neusser, J.; Rohde, C.; Schleper, V.:* Relaxed Navier-Stokes-Korteweg Equations for Compressible Two-Phase Flow with Phase Transition
- 2014-020 *Kabil, B.; Rohde, C.:* Persistence of undercompressive phase boundaries for isothermal Euler equations including configurational forces and surface tension
- 2014-019 *Bilyk, D.; Markhasin, L.:* BMO and exponential Orlicz space estimates of the discrepancy function in arbitrary dimension
- 2014-018 *Schmid, J.:* Well-posedness of non-autonomous linear evolution equations for generators whose commutators are scalar
- 2014-017 *Margolis, L.:* A Sylow theorem for the integral group ring of $PSL(2, q)$
- 2014-016 *Rybak, I.; Magiera, J.; Helmig, R.; Rohde, C.:* Multirate time integration for coupled saturated/unsaturated porous medium and free flow systems
- 2014-015 *Gaspoz, F.D.; Heine, C.-J.; Siebert, K.G.:* Optimal Grading of the Newest Vertex Bisection and H^1 -Stability of the L_2 -Projection
- 2014-014 *Kohler, M.; Krzyżak, A.; Walk, H.:* Nonparametric recursive quantile estimation
- 2014-013 *Kohler, M.; Krzyżak, A.; Tent, R.; Walk, H.:* Nonparametric quantile estimation using importance sampling
- 2014-012 *Györfi, L.; Ottucsák, G.; Walk, H.:* The growth optimal investment strategy is secure, too.
- 2014-011 *Györfi, L.; Walk, H.:* Strongly consistent detection for nonparametric hypotheses
- 2014-010 *Köster, I.:* Finite Groups with Sylow numbers $\{q^x, a, b\}$
- 2014-009 *Kahnert, D.:* Hausdorff Dimension of Rings
- 2014-008 *Steinwart, I.:* Measuring the Capacity of Sets of Functions in the Analysis of ERM
- 2014-007 *Steinwart, I.:* Convergence Types and Rates in Generic Karhunen-Loève Expansions with Applications to Sample Path Properties
- 2014-006 *Steinwart, I.; Pasin, C.; Williamson, R.; Zhang, S.:* Elicitation and Identification of Properties
- 2014-005 *Schmid, J.; Griesemer, M.:* Integration of Non-Autonomous Linear Evolution Equations
- 2014-004 *Markhasin, L.:* L_2 - and $S_{p,q}^r B$ -discrepancy of (order 2) digital nets
- 2014-003 *Markhasin, L.:* Discrepancy and integration in function spaces with dominating mixed smoothness
- 2014-002 *Eberts, M.; Steinwart, I.:* Optimal Learning Rates for Localized SVMs
- 2014-001 *Giesselmann, J.:* A relative entropy approach to convergence of a low order approximation to a nonlinear elasticity model with viscosity and capillarity
- 2013-016 *Steinwart, I.:* Fully Adaptive Density-Based Clustering
- 2013-015 *Steinwart, I.:* Some Remarks on the Statistical Analysis of SVMs and Related Methods
- 2013-014 *Rohde, C.; Zeiler, C.:* A Relaxation Riemann Solver for Compressible Two-Phase Flow with Phase Transition and Surface Tension

- 2013-013 *Moroianu, A.; Semmelmann, U.:* Generalized Killing spinors on Einstein manifolds
- 2013-012 *Moroianu, A.; Semmelmann, U.:* Generalized Killing Spinors on Spheres
- 2013-011 *Kohls, K.; Rösch, A.; Siebert, K.G.:* Convergence of Adaptive Finite Elements for Control Constrained Optimal Control Problems
- 2013-010 *Corli, A.; Rohde, C.; Schleper, V.:* Parabolic Approximations of Diffusive-Dispersive Equations
- 2013-009 *Nava-Yazdani, E.; Polthier, K.:* De Casteljau's Algorithm on Manifolds
- 2013-008 *Bächle, A.; Margolis, L.:* Rational conjugacy of torsion units in integral group rings of non-solvable groups
- 2013-007 *Knarr, N.; Stroppel, M.J.:* Heisenberg groups over composition algebras
- 2013-006 *Knarr, N.; Stroppel, M.J.:* Heisenberg groups, semifields, and translation planes
- 2013-005 *Eck, C.; Kutter, M.; Sändig, A.-M.; Rohde, C.:* A Two Scale Model for Liquid Phase Epitaxy with Elasticity: An Iterative Procedure
- 2013-004 *Griesemer, M.; Wellig, D.:* The Strong-Coupling Polaron in Electromagnetic Fields
- 2013-003 *Kabil, B.; Rohde, C.:* The Influence of Surface Tension and Configurational Forces on the Stability of Liquid-Vapor Interfaces
- 2013-002 *Devroye, L.; Ferrario, P.G.; Györfi, L.; Walk, H.:* Strong universal consistent estimate of the minimum mean squared error
- 2013-001 *Kohls, K.; Rösch, A.; Siebert, K.G.:* A Posteriori Error Analysis of Optimal Control Problems with Control Constraints
- 2012-013 *Diaz Ramos, J.C.; Dominguez Vázquez, M.; Kollross, A.:* Polar actions on complex hyperbolic spaces
- 2012-012 *Moroianu, A.; Semmelmann, U.:* Weakly complex homogeneous spaces
- 2012-011 *Moroianu, A.; Semmelmann, U.:* Invariant four-forms and symmetric pairs
- 2012-010 *Hamilton, M.J.D.:* The closure of the symplectic cone of elliptic surfaces
- 2012-009 *Hamilton, M.J.D.:* Iterated fibre sums of algebraic Lefschetz fibrations
- 2012-008 *Hamilton, M.J.D.:* The minimal genus problem for elliptic surfaces
- 2012-007 *Ferrario, P.:* Partitioning estimation of local variance based on nearest neighbors under censoring
- 2012-006 *Stroppel, M.:* Buttons, Holes and Loops of String: Lacing the Doily
- 2012-005 *Hantsch, F.:* Existence of Minimizers in Restricted Hartree-Fock Theory
- 2012-004 *Grundhöfer, T.; Stroppel, M.; Van Maldeghem, H.:* Unitals admitting all translations
- 2012-003 *Hamilton, M.J.D.:* Representing homology classes by symplectic surfaces
- 2012-002 *Hamilton, M.J.D.:* On certain exotic 4-manifolds of Akhmedov and Park
- 2012-001 *Jentsch, T.:* Parallel submanifolds of the real 2-Grassmannian