

**Brief Definitive Report****Cytotoxic Differentiation of Mouse Gut Thymodependent and Independent Intraepithelial T Lymphocytes Is Induced Locally. Correlation between Functional Assays, Presence of Perforin and Granzyme Transcripts, and Cytoplasmic Granules**By Delphine Guy-Grand,\* Michèle Malassis-Seris,\*  
Chantal Briottet,† and Pierre Vassalli‡

From \*INSERM U.132, Hôpital Necker-Enfants Malades, 75743 Paris, cédex 15, France; and the †Département de Pathologie, Centre Médical Universitaire, CH 1211 Genève 4, Switzerland

**Summary**

Mouse gut intraepithelial lymphocytes (IEL), whether thymodependent ( $CD4^+$  or  $CD8 \alpha/\beta^+$ ;  $TCR-\alpha/\beta^+$ ) or thymoindependent ( $CD8 \alpha/\alpha^+$ ;  $TCR-\alpha/\beta^+$  or  $-\gamma/\delta^+$ ), all display cytotoxic activity in a "redirected lysis assay" using anti-CD3 or anti-TCR  $\beta$  or  $\delta$  chains secreting hybridomas as targets; this is also observed with IEL of germ-free mice, indicating that this activity, which is absent in peripheral T lymphocytes, does not require stimulation by bacterial antigens. Perforin and granzyme transcripts are detectable in unselected gut IEL, in contrast to normal T lymphocytes of peripheral lymphoid organs. Cytological labeling (with [ $^3H$ ]DFP) of IEL smears reveals labeled granules (i.e., containing serine-esterases, presumably granzymes) in all subsets of gut IEL. This indicates that the gut micro-environment has an inductive role on the cytotoxic differentiation of lymphocytes of various origins when they reach the gut wall to become IEL.

It has recently been shown that gut intraepithelial lymphocytes (IEL) have a dual origin (1, 2). About 45% are the progeny of thymodependent lymphoblasts elicited to proliferate in Peyer's patches under antigenic stimulation and migrate through the thoracic duct (TD) lymph and the blood to seed all along the gut mucosa (3). These thymodependent IEL express  $TCR-\alpha/\beta$  and bear either CD8, or, for a small minority ( $\sim 10\%$ ), CD4 molecules. Their CD8 molecules are made of  $\alpha$  (Lyt-2) and  $\beta$  (Lyt-3) chains ( $CD8 \alpha/\beta^+$  cells), as is also the case for all peripheral CD8 T lymphocytes. About 55% are thymoindependent, arising from bone marrow progenitors migrating directly to the gut, and bear  $TCR-\alpha/\beta$  or  $-\gamma/\delta$ . These thymoindependent IEL are easily identifiable since, strikingly, they bear CD8 molecules made only of homodimeric  $\alpha$  chains, without  $\beta$  chains ( $CD8 \alpha/\alpha^+$ ). All subsets of IEL display a peculiar cytological feature: a high percentage of them (20–70%) contain intracytoplasmic granules with sulfated proteoglycans, found in all species studied: rabbits (4), mice (3), rats (5), and humans (6). The function of IEL is incompletely understood, but evidence exists that, whether obtained from normal or nude mice, they can be cytotoxic in vitro, as has been found using "redirected" lysis assays (7–9).

In this study, we compared IEL and their subpopulations with peripheral T lymphocytes from normal mice, from germ-

free mice, or from mice with an acute graft-vs.-host reaction (GVHR), a condition leading to a strong in vivo stimulation of grafted T cells. We attempted to correlate the level of cytotoxic activity of these various T lymphocytes (using as targets hybridoma cells secreting anti-CD3 or anti-TCR  $\beta$  or  $\delta$  mAbs) with that of perforin and granzyme mRNAs transcripts, serine esterase enzymatic activity, and presence of cytoplasmic granules. Perforin and granzymes are indeed thought to be involved in the cytotoxic process (for review, see reference 10).

**Materials and Methods**

**Animals and In Vivo Treatment.** C57Bl/6, C<sub>3</sub>HeB, and C<sub>3</sub>H-DBA/2 mice were raised in our animal house. Germ free (C<sub>3</sub>HeB) mice were a kind gift of C. Moreau (INRA CNRZ, Jouy an Josas, France). GVHRs were elicited in lethally irradiated C<sub>3</sub>H-DBA/2 mice with C<sub>3</sub>H lymphocytes (11).

**Lymphocytes Surface Labeling, Selection, and Cytological Procedures.** Detection of surface T cell antigens and cell separation were performed as previously described (1). Thymoindependent IEL were obtained by labeling IEL with anti-CD4 (GK1.5), anti-CD8  $\alpha/\beta$  (H35-17-2), or anti-Thy-1 mAb (TB107), incubation with goat anti-rat Ig-coated steel beads (Dynabeads-Dynal, Oslo, Norway), and sorting of the labeled cells with a magnet (MACS-Miltenyi Biotec Bergisch, Gladbach, FRG). Peripheral T lymphocytes were

obtained after incubation with goat anti-mouse Ig-coated steel beads. Detection of Di-isopropyl phosphorofluoridate (DFP) binding serine esterases was performed on paraformaldehyde-fixed smears (12) with  $10^{-5}$  M [ $^3$ H]DFP (Amersham, Burlington, Massachusetts, England) applied for 30 min. This type of labeling can be performed on cells previously stained for surface antigens by fluorescent mAbs, but is not compatible with dye-staining of the granules.


**Cytotoxic Assays and Northern Blot Analysis.** Lymphocytes were incubated with  $10^4$   $^{51}$ Cr-labeled hybridomas secreting anti-CD3 (13), anti-TCR  $\beta$  (14), or anti-TCR  $\delta$  (GL3; a kind gift of L. Lefrançois [8]) for 3 h, at varying E/T ratios. Percent lysis was calculated as  $100 \times [(cpm \text{ release with effector cells} - cpm \text{ release alone}) / (cpm \text{ release by CIH acid} - cpm \text{ release alone})]$ .

Northern blot analyses, RNA preparations, transfer to Biodyne membrane, and hybridizations were as described (15). For the detection of granzyme mRNA, an oligonucleotide, 5'TTAGTTC-TCTTGGCCTTACTCTC (complementary to a sequence starting on nucleotide 197, 309, and 346, respectively, of the three different granzymes cDNAs described as MCSP-1, -2, and -3 by Kwon et al. [16]), was end labeled with  $^{32}$ P. For the detection of perforin mRNA, a riboprobe containing the EcoRI-Hind III fragment of the mouse perforin cDNA (position 255-766 of the coding region) was used (17) (kind gift of Drs. E. Podack and C. Mueller).

## Results and Discussion

In "redirected" cytotoxic assays involving the TCR-CD3 complex, using anti-CD3, anti-TCR  $\beta$ , or anti-TCR  $\delta$  mAb-secreting hybridomas as targets (Fig. 1), total IEL and their thymoindependent subset (obtained by depletion of CD4 $^+$  and CD8  $\alpha/\beta^+$  cells or of all Thy-1 $^+$  cells) are cytotoxic against the three hybridomas. The cytotoxicity of the thymoindependent population is not altered when it is depleted of its Thy-1 $^+$  component, which represents a minority of this population (10-40%; reference 1); this is in contrast to reports that only Thy-1 $^+$  IEL are cytotoxic in comparable assays (7, 8). In the thymoindependent population, TCR- $\alpha/\beta^+$  and TCR- $\gamma/\delta^+$  IEL have comparable cytotoxic abilities. Since TCR- $\gamma/\delta^+$  IEL are found only in the thymoindependent CD8  $\alpha/\alpha^+$  population (1), thymus-derived IEL should not be cytotoxic for the anti- $\delta$  hybridomas cells; this population, however, cannot be adequately separated. When IEL were obtained from mice with acute GVHR induced by the injection of foreign peripheral T lymphocytes, a condition in which all IEL are of donor origin and thus of the thymus-derived variety (1, 11), they indeed displayed cytotoxicity against the anti-CD3 and the anti-TCR  $\beta$ , but not against the anti-TCR  $\delta$  hybridomas (Fig. 1). In contrast, IEL obtained from germ-free mice, which are in small amounts but predominantly thymoindependent (CD8  $\alpha/\alpha^+$ ) and TCR- $\gamma/\delta^+$  (1) are strongly cytotoxic against anti- $\delta$  and slightly against anti- $\beta$  hybridomas (Fig. 1). Thus, contrarily to prior suggestion (7, 8), IEL do not need an in vivo exogenous stimulation to develop cytotoxic capabilities.


Contrarily to IEL, normal circulating T lymphocytes (thoracic duct lymphocytes or TDL, which contain the precursors of thymus-derived CD8  $\alpha/\beta^+$  and CD4 $^+$  IEL; references 1 and 3) do not show any cytotoxic activity (Fig. 1). However, when TDL are obtained during a GVHR and thus are, as



**Figure 1.** Cytotoxic activity of various populations of gut IEL compared with that of TDL, in redirected lysis assays using secreting hybridomas as targets (1). Results are shown at a 33:1 E/T ratio (see Materials and Methods). Standard deviations were calculated from the results of three to five experiments. The other percentages are the mean of two experiments with close results. For germ-free mice, a single experiment was carried out. (2) TCR- $\gamma/\delta^+$  cells among IEL:  $22 \pm 7.5\%$ . (3) Cytotoxicity against the anti-TCR  $\delta$  hybridoma does not increase much although there are more TCR- $\gamma/\delta^+$  in this population (1), but it appears that there is some nonspecific cell loss during the isolation procedure. (4) Residual host cells detected with an anti-H2 antibody: <4%. Cells containing TCR- $\gamma/\delta^+$ : 0%. Dye-stained cytoplasmic granules in these experiments: 34% and 49%. (5) IEL pooled from eight germ-free mice, with 13% TCR- $\alpha/\beta^+$  and 80% TCR- $\gamma/\delta^+$  cells. (6) TDL are deprived of B cells. TCR- $\gamma/\delta^+$  lymphocytes in the population used: 1.2%. (7) Residual host cells: <1%. TCR- $\gamma/\delta^+$ : 0%. Cells containing dye-stained cytoplasmic granules in these experiments: 4.8% and 1%.

the GVHR IEL described above, of donor origin and cytotoxic against the host alloantigens (11), they are cytotoxic against anti-CD3 and anti- $\beta$ , but not anti- $\delta$  hybridomas (Fig. 1).

Since perforin and granzymes are detected in cytotoxic lymphocytes in culture or in immunopathological reactions and are localized in cytoplasmic granules (10), the presence of perforin and of granzyme transcripts in IEL was explored by Northern blot analysis. Perforin and granzyme mRNAs were detectable in gut IEL but not in peripheral lymph node T cells nor in TDL, except in conditions of GVHR; i.e., when stimulated cytotoxic grafted cells represent almost all of the T lymphocytes present in the circulation, the lymphoid organs,



**Figure 2.** Northern blots of RNA extracts from IEL and TDL using perforin c-mRNA (P) or granzyme DNA  $^{32}\text{P}$  labeled (G) probes and showing bands of  $\sim 3$  kb (perforin) and  $\sim 900$  bp (granzyme).

and gut IEL (Fig. 2). Since granzymes are serine-esterases; i.e., enzymes characterized by their ability to react with DFP (18), smears of IEL and other T cells were exposed to  $^3\text{H}$ DFP. IEL fixed  $^3\text{H}$ DFP but TD lymphocytes did not, even the TD blasts that are the precursors of thymus derived IEL (3). The localization and frequency of  $^3\text{H}$ DFP silver grains coincided with that of the granules: (a) silver grains were distributed in clusters, frequently localized at one pole of the cell, near a nuclear invagination containing the dye-stained granules (Fig. 3); (b) the percentages of granule containing IEL and that of  $^3\text{H}$ DFP-labeled cells were very close (Table 1). Combination of  $^3\text{H}$ DFP binding with determination of the IEL subsets by immunofluorescence showed the presence of silver grains in all subsets:  $\text{CD4}^+$  and  $\text{CD8 } \alpha/\beta^+$  thymus-derived IEL as well as thymoindependent  $\text{CD8 } \alpha/\alpha^+$  IEL, which were the most frequently granulated (81%) and bore especially numerous silver grains. In contrast to normal TDL,  $^3\text{H}$ DFP grains were also detected in  $\sim 25\%$  of GVHR TD blasts, but they were not concentrated in clusters, and granules are rare or absent in GVHR TD blasts (Fig. 3; Table 1). This observation probably indicates that perforin and granzyme storing granules occur late in differentiation rather than in still rapidly dividing cells.

In conclusion, it appears that the gut microenvironment exerts a differentiation influence on lymphocytes becoming located within the epithelium, whatever their origin, their


**Table 1.** Granular Differentiation in IEL and Circulating Lymphocytes Labeling with  $^3\text{H}$ DFP

	Normal mice		Mice with GVHR	
	IEL	TDL	IEL	TDL
Percent giemsa-stained granules	69	<0.5	22	<1
Percent $^3\text{H}$ -DFP-labeled cells*				
Clustered labeling	67	0	21	<1
Dispersed labeling	<2	0	4	24

\* See Materials and Methods.

TCR type, or the presence of Thy-1 molecules, leading to synthesis and storage of perforin and granzymes, accompanied by the development of a cytotoxic activity that is not found in peripheral T lymphocytes, except when they are activated.

What could be the role of IEL with this newly acquired cytotoxic ability? They may insure the defense of the epithelia integrity by rapidly eliminating altered enterocytes; the rapidity of epithelium renewal, which increases in pathological conditions such as GVHR (11), may compensate for cell destruction. Since in normal conditions IEL are not cytotoxic when binding to targets is obtained by PHA (data not shown), recognition of the targets must involve TCR activation. This may occur in a classical MHC-restricted way during viral infections (19); or this may occur in "unclassical" ways, involving other types of specificity, since we have found that IEL (but not unactivated peripheral T cells) are cytotoxic against xenogenic targets bearing enterotoxins with "superantigens" properties (our unpublished observation).



**Figure 3.** Intracytoplasmic granules and  $^3\text{H}$ DFP labeling patterns in IEL and TDL in normal mice and in mice with GVHR.

This work was supported by grants from INSERM and the Swiss National Foundation (31-28866.90).

Address correspondence to Delphine Guy-Grand, INSERM U.132, Hôpital Necker-Enfants Malades, 75743 Paris, cédex 15, France.

Received for publication 25 February 1991 and in revised form 18 March 1991.

## References

- Guy-Grand, D., N. Cerf-Bensussan, B. Malissen, M. Malassis-Seris, C. Briottet, and P. Vassalli. 1991. Two gut intraepithelial CD8<sup>+</sup> lymphocyte populations with different T cell receptors: a role for the gut epithelium in T cell differentiation. *J. Exp. Med.* 173:471.
- Rocha, B., P. Vassalli, and D. Guy-Grand. 1991. The V $\beta$  repertoire of mouse gut homodimeric  $\alpha$  CD8<sup>+</sup> intraepithelial T cell receptor  $\alpha/\beta$ <sup>+</sup> lymphocytes reveals a major extrathymic pathway of T cell differentiation. *J. Exp. Med.* 173:483.
- Guy-Grand, D., C. Griscelli, and P. Vassalli. 1978. The mouse gut T lymphocyte, a novel type of T cell. Nature, origin, and traffic in mice in normal and graft-versus-host conditions. *J. Exp. Med.* 148:1661.
- Rudzik, O., and J. Bienenstock. 1974. Isolation and characteristics of gut mucosal lymphocytes. *Lab. Invest.* 30:260.
- Mayrhofer, G. 1980. Fixation and staining of granules in mucosal mast cells and intraepithelial lymphocytes in the rat jejunum, with special reference to the relationship between the acid glycosaminoglycans in the two cell types. *Histochem. J.* 12:513.
- Cerf-Bensussan, N., D. Guy-Grand, and C. Griscelli. 1985. Intraepithelial lymphocytes of human gut: isolation, characterization and study of natural killer activity. *Gut.* 26:81.
- Lefrançois, L., and T. Goodman. 1989. In vivo modulation of cytolytic activity and Thy-1 expression in TCR- $\gamma\delta$ <sup>+</sup> intraepithelial lymphocytes. *Science (Wash. DC)*. 243:1716.
- Goodman, T., and L. Lefrançois. 1989. Intraepithelial lymphocytes. Anatomical site, not T cell receptor form, dictates phenotype and function. *J. Exp. Med.* 170:1569.
- Viney, J.L., P.J. Kilshaw, and T.T. MacDonald. 1990. Cytotoxic  $\alpha/\beta$ <sup>+</sup> and  $\gamma/\delta$ <sup>+</sup> T cells in murine intestinal epithelium. *Eur. J. Immunol.* 20:1623.
- Tschopp, J., and M. Nabholz. 1990. Perforin-mediated target cell lysis by cytolytic T lymphocytes. *Annu. Rev. Immunol.* 8:279.
- Guy-Grand, D., and P. Vassalli. 1986. Gut injury in mouse graft-versus-host reaction. Study of its occurrence and mechanisms. *J. Clin. Invest.* 77:1584.
- Young, J.D.E., L.G. Leong, C.C. Liu, A. Daminao, D.A. Wall, and Z.A. Cohn. 1986. Isolation and characterization of a serine esterase from cytolytic T cell granules. *Cell.* 47:183.
- Leo, O., M. Foo, D.H. Sach, L.E. Samelson, and J.A. Bluestone. 1987. Identification of a monoclonal antibody specific for a murine T3 polypeptide. *Proc. Natl. Acad. Sci. USA.* 84:1374.
- Kubo, R.T., W. Born, J.W. Kappler, P. Marrack, and M. Pigeon. 1989. Characterization of a monoclonal antibody which detects all murine  $\alpha/\beta$  T cell receptors. *J. Immunol.* 142:2736.
- Collart, M.A., D. Belin, J.D. Vassalli, S. de Kossodo, and P. Vassalli. 1986.  $\gamma$  Interferon enhances macrophage transcription of the tumor necrosis factor/cachectin, interleukin 1, and urokinase genes, which are controlled by short-lived repressors. *J. Exp. Med.* 164:2113.
- Kwon, B.S., D. Kestler, E. Lee, M. Wakulchik, and J.D.E. Young. 1988. Isolation and sequence analysis of serine protease cDNAs from mouse cytolytic T lymphocytes. *J. Exp. Med.* 168:1839.
- Lowrey, D.M., T. Aebischer, K. Olsen, M. Lichtenheld, F. Rupp, H. Hengartner, and E.R. Podack. 1989. Cloning, analysis, and expression of murine perforin 1 cDNA, a component of cytolytic T-cell granules with homology to complement component C9. *Proc. Natl. Acad. Sci. USA.* 86:247.
- Masson, D., and J. Tschopp. 1987. A family of serine esterases in lytic granules of cytolytic T lymphocytes. *Cell.* 49:679.
- London, S.D., J.J. Cebra, and D.H. Rubin. 1989. Intraepithelial lymphocytes contain virus-specific, MHC-restricted cytotoxic cell precursors after gut mucosal immunization with reovirus serotype 1/Lang. *Reg. Immunol.* 2:98.