

Short Communication

A Novel *orf108* Co-Transcribed with the *atpA* Gene is Associated with Cytoplasmic Male Sterility in *Brassica juncea* Carrying *Moricandia arvensis* Cytoplasm

Ashutosh^{1,4}, Pankaj Kumar^{1,4}, V. Dinesh Kumar², Prakash C. Sharma³, Shyam Prakash¹ and Shripad R. Bhat^{1,*}

¹ National Research Centre on Plant Biotechnology, Indian Agricultural Research Institute, New Delhi-110012, India

² Directorate of Oilseeds Research, Rajendranagar, Hyderabad-500030, India

³ Guru Gobind Singh Indraprastha University, Kashmere Gate, Delhi-110006, India

Mitochondrial *atpA* transcripts were examined in cytoplasmic male sterile (CMS) and fertility restorer lines of CMS (*Moricandia arvensis*) *Brassica juncea*. Male sterile flowers had longer *atpA* transcripts than male fertiles. The mitochondrial *atpA* region of the CMS line was cloned and sequenced. The 5' and 3' ends of the *atpA* transcripts of the CMS and the fertility restorer lines were mapped and full-length transcripts were cloned and sequenced. A novel *orf108* (open reading frame 108) co-transcribed with the *atpA* gene was found in the male sterile flowers. In the fertility restorer line, the transcript was cleaved within *orf108* to yield mono-cistronic *atpA* transcripts.

Keywords: *atpA* — *Brassica juncea* — Cytoplasmic male sterility — Fertility restoration — *orf108* — Transcript mapping.

Abbreviations: cRT-PCR: circular reverse transcription-PCR; CMS: cytoplasmic male sterility; ORF: open reading frame; TAP: tobacco acid pyrophosphatase.

Mitochondria carry their own genetic information but are heavily dependent on nuclear-encoded proteins for their replication and correct expression of the information. Similarly, cellular mitochondrial activity can influence expression of nuclear genes and is termed retrograde regulation (Liu and Butow 2006). Thus a well-orchestrated expression of mitochondrial and nuclear genes is regarded as essential for normal growth and development of eukaryotic organisms. As mitochondria carry out vital cellular functions, any impairment of their function is likely to be lethal. Hence there are very few phenotypes linked to mitochondrial gene mutations in plants. Maternally inherited male sterility in angiosperms, referred to as cytoplasmic male sterility (CMS), is known to result from mitochondrial dysfunction. This defect, however, can be overcome by

the action of appropriate nuclear gene(s) called restorer of fertility (*Rf*). Thus CMS is one of the rare traits that is governed by nuclear-mitochondrial gene interactions and serves as a model to study such interactions (Hanson and Bentolila 2004).

Detailed molecular analyses of a number of CMS systems in various plant species have revealed that male sterility is associated with either the appearance of novel, often chimeric, transcripts or altered expression, usually in a tissue-specific manner, of some essential mitochondrial gene (Schnable and Wise 1998, Hanson and Bentolila 2004). Such changes in mitochondrial gene expression usually result from rare mitochondrial genome mutations, alien mitochondrial substitutions following wide hybridization or mitochondrial recombination in somatic hybrids. However, each CMS system appears to be unique with respect to the mitochondrial transcript associated with male sterility. Further, the action of the restorer gene is also specific and restoration of male fertility is associated with suppression of cognate chimeric transcripts/proteins and the appearance of normal mitochondrial transcripts. For example, in Texas CMS maize, the chimeric *urf13* has been shown to be responsible for male sterility. Similarly, in *Brassica*, a number of novel open reading frames (ORFs) such as *orf224* (polima CMS), *orf222* (nap CMS) and *orf138* (ogura CMS) have been found to be associated with male sterility. Cloning of restorer genes of CMS systems of petunia, radish, rice, etc. in recent years has revealed that they encode proteins belonging to the PPR family (see Horn 2006), which are implicated in organellar transcript processing (Small and Peeters 2000) and thus match their expected function.

We have earlier reported synthesis of a CMS system in *Brassica juncea* carrying the mitochondrial genome of *Moricandia arvensis* (Prakash et al. 1998). Floral-specific alteration in mitochondrial *atpA* expression was found to be associated with male sterility in this system

⁴These authors contributed equally to this work.

*Corresponding author: E-mail, srbhat22@rediffmail.com; Fax, +91-11-25843984.




Fig. 1 Expression of the mitochondrial *atpA* gene in sepals (S), petals (P) and anthers (A) of the male sterile (rr) (Sterile) and the fertility-restored (Rr) (Fertile) plants. Northern blot was prepared using about 10 µg of total RNA and probed with the *atpA* coding sequence.

(Gaikwad et al. 2006). Also, we have tagged the *Rf* gene, which acts gametophytically to restore male fertility (Ashutosh et al. 2006). The present study was undertaken to characterize this CMS system further with respect to the *atpA* gene. We report here the identification of a novel *orf108*, which is co-transcribed with the *atpA*. In fertility-restored plants, the bicistronic *orf108-atpA* transcripts are specifically cleaved within the *orf108* to yield monocistronic *atpA* transcripts.

A previous study showed that *atpA* transcripts in flower bud tissues of the CMS plants are longer as compared with normal *B. juncea*, *M. arvensis* and the fertility-restored plants (Gaikwad et al. 2006). However, genetic studies revealed that the *Rf* gene acts gametophytically to confer male fertility to the CMS plants. To verify whether the *Rf* gene was active in all floral tissues or was restricted to the anthers, we examined the transcript pattern of *atpA* in sepals, petals and anthers of the CMS and the fertility-restored (i.e. F₁) plants. Northern analysis revealed a 2,800 nt long *atpA* transcript in all the three whorls of the CMS flower, whereas in the fertility-restored plants, a shorter (1,800 nt) *atpA* transcript was found in sepals and petals (Fig. 1). Anthers of the fertility-restored plants expressed both 2,800 and 1,800 nt transcripts, but the intensity of the longer transcript was weak. Thus the restorer gene action was detected in both sporophytic and gametophytic tissues of the flower, and was inferred to be dominant. When restoration is gametophytic, it is difficult to determine the dominant/recessive nature of gene action through genetic studies. In maize, Kamps et al. (1996) generated tetraploid restorer lines and showed that the diploid gametes of R/r constitution are viable, thereby confirming the dominance of the restorer. The present study demonstrated that Northern analysis of individual whorls of flowers could be helpful in determining the dominance relationship of the gametophytic restorers.

We had found distinct restriction fragment length polymorphism (RFLP) patterns between the CMS and normal *B. juncea* lines with the *atpA* probe (Gaikwad et al. 2006). To know the exact sequence differences in the *atpA* region, we cloned and sequenced the *atpA*-containing 6.0 and 5.8 kb *Hind*III fragments from the CMS and euplasmic lines, respectively (male sterile line genomic clone, GenBank accession No. EF483940). The *atpA* coding sequences (1,524 bp) in both clones were identical. Sequences downstream of *atpA* also did not show a significant difference between the two clones. On the other hand, no significant homology was found between the CMS and *B. juncea* beyond 52 bp upstream of the *atpA* initiation site. Northern hybridization with the sequences from the upstream region of *atpA* (1,728–2,173 of the 6.0 kb clone) had revealed a single transcript of 2,800 nt only in the CMS line, indicating that the *atpA* transcript of the CMS line differs from normal *B. juncea* in the 5' region (unpublished results). Hence, we first tried to determine the 5' end of the transcripts. Based on genomic DNA sequence information, we designed PCR primers that could amplify various upstream regions of the *atpA* gene (Table 1, Fig. 2A). To identify the location of the 5' ends of the *atpA* transcripts, we performed reverse transcription-PCR (RT-PCR) using different sets of primers. RNA from the male steriles gave amplicons of the expected size when primers located up to 1,019 nt upstream of the *atpA* start codon were used in RT-PCR (i.e. up to the F8 primer), but no amplicon was obtained with the primers R6 + F9 (Fig. 2B). This indicated that the 5' end of the transcript is located between the F9 and F8 region (i.e. –1,411 and –1,019 of the *atpA* start codon). In the male fertiles, primers R3 + F3 gave a 170 bp RT-PCR product, while no amplicon was obtained with R4 + F5 primers. Hence, the 5' end of the *atpA* transcript of the male fertile line was inferred to lie between –503 and –260 nt upstream of the *atpA* start codon (Fig. 2C).

The circular RT-PCR (cRT-PCR) method (Khun and Binder 2002) was employed to identify the exact 5' and 3' ends of the *atpA* transcripts. To distinguish between the primary and processed transcripts, two sets of cRT-PCRs were set up. In one, RNA was treated with tobacco acid pyrophosphatase (TAP) prior to self-ligation whereas the other was subjected to self-ligation without TAP treatment. A linear RT-PCR was also included to serve as negative control. The absence of amplicons in linear RT-PCR ruled out DNA contamination and also confirmed that only circularized RNA molecules served as templates for DNA amplification. In the male sterile line, a 1,050 bp, intense cRT-PCR amplicon was obtained only in the treatment where RNA was treated with TAP (Fig. 3). Identical cRT-PCR products were observed in both TAP(+) and TAP(–) treatments in the male fertile line (Fig. 3). Since transcripts with 5' triphosphate cannot undergo self-ligation,

Table 1 Details of primers used in various experiments

Primer	Sequence (5'-3')	Position ^a
R1	ATATGGTATCGATAGCAATAG	3,259–3,238
R2	CTAATTCGACTTTCGAATAGATTCG	2,759–2,735
R3	CCGCCCCGTTAAATTGATAATATTGAA	2,615–2,590
R4	CAAGGTCATATCATCTCCAAG	2,520–2,500
R5	CGGGTTTCACGATTAGATTA	2,281–2,262
R6	GCTAACTCCTGTAAGCAGCT	1,919–1,900
F1	GGAATTATGGAATTATCTCCCAGAG	2,698–2,722
F2	ACCTTGGGAAATGTTGCTTC	2,515–2,534
F3	ACCATAATGTTCTTTTGCCCCG	2,444–2,464
F4	CCCGAAAATCAACTTCTACTTATGAATAC	2,278–2,306
F5	CTTCACTACTCCTAGAGGCT	2,201–2,220
F6	GATCTCTATGCCCCCTGTTCTTGG	2,126–2,149
F7	GCTCCCTCCAAGTGTGG	1,801–1,818
F8	CTTGCAGACCTACTCGGAAC	1,685–1,704
F9	CTCAGCCAGATGGAAGGTCA	1,293–1,312
F10	CCACTAGACAGAATCTCTCAAT	4,081–4,102

^aWith reference to GenBank sequence No. EF483940.




Fig. 2 Identification of the *atpA* transcript initiation region in the male sterile and the male fertile lines. (A) Schematic representation of the mitochondrial *atpA* region of the *Brassica juncea* CMS line carrying *Moricandia arvensis* mitochondria. The position and direction of various primers used in the study are indicated with arrows (figure not to scale). (B) Gel photograph of RT-PCR with various primer combinations using RNA from the male-sterile flowers. M, 100 bp DNA ladder, Bangalore Genei, India; 1, negative control (PCR with R2 + F2 primers without a reverse transcription step); 2, R2 + F2 (58); 3, R2 + F3 (60); 4, R4 + F4 (58); 5, R4 + F6 (58); 6, R6 + F7 (56); 7, R6 + F8 (58); 8, R6 + F9 (58). (C) RT-PCR results obtained using RNA from the male fertile (RR) flowers with primers (1) R4 + F5 (56) and (2) R3 + F3 (60). M, 100 bp DNA ladder (Bangalore Genei, India). The primer annealing temperature (°C) is given in parentheses.

the appearance of identical cRT-PCR products in both TAP(+) and TAP(-) treatments indicated that *atpA* transcripts are processed in the male fertile line. These results show that the male steriles contain mostly the primary *atpA* transcript while male fertiles predominantly carry the processed transcript. Additional weak amplicons observed in both TAP(+) and TAP(-) treatments may have arisen from processed or partially degraded RNA. Cloning and sequencing of the longest and intense amplicons from the male sterile and the male fertile lines helped to identify

the 5' and 3' ends of *atpA* transcripts. In the male steriles, the 5' end of the *atpA* transcript was determined to be the T at 1,200 nt upstream of the *atpA* start codon while in the male fertiles it was the T nucleotide situated 390 nt upstream of the *atpA* ATG codon (Fig. 4A). In both the CMS and the fertility restorer lines, the *atpA* transcripts terminated at the same position, which was located 160 nt downstream of the *atpA* stop codon. The complete *atpA* transcripts of both the lines were amplified by RT-PCR and sequenced (GenBank accession Nos. EF483941 and EF483942).




Fig. 3 cRT-PCR amplification of *atpA* from the male sterile and the fertility restorer lines for the identification of transcript extremities. M, 1 kb DNA ladder (Fermentas); LRT, linear RT-PCR; CRT(+), circular RT-PCR with TAP treatment; CRT(-), circular RT-PCR without TAP treatment.

The *atpA* transcript in the CMS was 2,884 nt long, whereas in the fertility restorer line it was 2,074 nt long (Fig. 4A). These results confirmed our earlier size estimates of *atpA* in the two lines. Further, the 3' ends of the *atpA* transcripts of the CMS and the fertility restorer lines were not polyadenylated and thus conformed to the norm for mitochondrial transcripts.

Comparative analysis of the *atpA* transcripts of the CMS and the fertility restorer lines revealed a novel *orf108* upstream of the *atpA* region in the male sterile line (Fig. 4A). In the fertility restorer line, the *orf108* was abolished as the transcript was cleaved at 12 nt downstream of the *orf108* initiation codon. Thus the *atpA* transcript of the male sterile line was bicistronic, whereas the transcript of the male fertile line was monocistronic, capable of coding for only *atpA* protein. When primers F4 and R3, capable of yielding a product only from the male sterile transcript, were used in RT-PCR, a 300 bp amplicon was obtained

A ttgttctggacgagagagtaggaataggtacgtcttcaaccatagaaagtcaacagaggggtatcctgcttaggttgaaaac
 ctaactcgggaatgacgaggttaagaagaggagatagaaggaaggggtactgattaaccaactgtaatacaaaagtacg
 tgatcctcctaggtgacagactactcggaaactgtggaagaagcaagatattaaacggggcttgcgccctgcttag
 ctgcactgacagagtagctacagtcgagcaggtccttctcctcctccaagtgttggttagctacgatagaacacggata
 ataaacagataaattgcagtggaaggaagaaatgacagagaataatgataatgataagaagctcttacagagtagccagctca
 ctccctctcgaatgggagatagataaaagaacgaaactctataaaatcattaagagaagaattctcattccagcttaaa
 taagtaagactgactcttctcacttctttctcgtccttctggcgaagaagcggcatcaaaaataaagaagaagcaaaa
 tctcattcaattgaaatataagagactctatgccctgtcttctggttctccatgctttgtgtcaacaaccaaccacaacttc
 tatagttctcactactctagagccttgacggagtgaagctgtctggagggaattgtgaaatcaattaactaactgtaaacccg
 aaaaatcaactctacttATGAATACTATCAAA▼TCACTTCCCAAGACATCGAAGAGTTAAAAA
 AAATGGAAATCTCTTCGATGTCTCCGATCAGTTGTATAACAACGGCTTGTAGCGTAAC
 TGAAGCACTGACCGGGACGGTCCAACGCCCTTTTACAGCACCATAATGTTCTTTTGGCCG
 AAAATTGGACCCACGCAAAATAGTACAAGATATACTTGGAGATGATATGACCTTGGGA
 AATGTTGCTTCTGTATTGAGCAACATCACTCAGTTGGGACTCGCAAGTGACGAGTTTCA
 CAAATCTCAATATTATCAATTTAACGGGGCGGGGGTTTAA Gttaaggtaactatccgaaggtatac
 ttttttctaaagcggatattcctctcctctgaattcaaaaatggaattatcctctagagctcggaaactaacgaatc
 tattcgaagtcgaattaggaacttttacgcaatttcaagtgatgagatcggtcagtgctcagttggagatgggattg
 cacaagttatgattgaacgagatcaagctgggaaatggtcttttggccaacgggtgaaaggaatggccttgaatttg
 agaatgagaatgctgggattgtcttgggtgatccgctataaaagaagagatcttgaagcgcactggatctattg
 tggatgtlcccgggaaaggctatctaggggcatgtggcagcgcgatgggagctactattgatgaaaggggctctaa
 gcatcacgagcaaaagacgtgtcgaagtaaaagcccctgggattctgaactgaaatcagtgacagcctatgcaaaaca
 gggtaaaagcggtagatagcctgtctctatagcccgtggcaacgagaacttcaatcgggacgcaaaactggaaa
 aacgactattgctatgataccatataaaacaaagcaataaactcaagggccactctgagagtgagacaatgattgt
 gtctatgtagcattgacagaacgctcactgtgggacaataaacttgaagaagcgaatgcttggaaatattc
 cattctgtagcagccaccctcggatcctgctcctctgcaattttggcccataattccgggtgctccatggggaaatattc
 gcgataatggaatgcagcattataaataatctatgatcttagtaaacaggcgggtggcatatcgacaatgtcattattgta
 gccgaccaccagccgtgaggcttccaggtgatgtttctatttaccctctcttagaagaagcggctaaacgatcgg
 accagacaggtgcaggtagctgaccgcttaccctgattgaaacacaagctggagacgtatcggectatattccacca
 atgtgatctcattactgatgacaatctgttggaaacagagctctttatcgggaattagactgctattaacgtcggctta
 tctgctagtcgctgggtctgcccctcagttgaaagctatgaaacaagatgctggtagttcaaaactgggaattggcacaata
 tcgcaagtgggccttctgcaatttggctcagactgactgagcactaggcattactcaatagaggtgcaaggctga
 cagaagtaccgaaacacacaaatgaccacttcaattgaaaacaaataactagctcattatgctcagctgcaatggctta
 ctgtgatcgaatgccactagacagaatctcaatagagaagccattcaaatagtgcaaacctgaaactactcaagccc
 ttaaaggtgattactcaacgaaagaaatggaactagatgcttcttaaaagaagagctttagctttaaattgaccttcatc
 acattaggaatcctcaatcagtaagaagaaaggttcttaagcaaaagactggagagcccccgtcagatgtagtaagtagct
 ctccaactcggagactgaggagaatgggattgtgggtgaggggtgcatctccccaggccct




Fig. 4 (A) Complete cDNA sequence of the *orf108/atpA* transcript of the male sterile line. Coding sequences of *orf108* (upper case) and *atpA* (bold italics) are highlighted. The triangle indicates the 5' site of the *atpA* transcript in the male-fertile line. (B) RT-PCR amplification of *orf108* using RNA from flower buds of the male-sterile (rr), fertility-restored (Rr) and fertility restorer (RR) lines. M, 100 bp ladder (Fermentas); C(-), negative control; C(+), positive control for intact RNA in the fertility restorer (RR) with primers F1 + R1 (56°C).

References

- Ashutosh, Sharma, P.C., Prakash, S. and Bhat, S.R. (2006) Identification of AFLP markers linked to the male fertility restorer gene of CMS (*Moricandia-arvensis*) *Brassica juncea* and conversion to SCAR marker. *Theor. Appl. Genet.* 114: 385–392.
- Brown, G.G. (1999) Unique aspects of cytoplasmic male sterility and fertility restoration in *Brassica napus*. *J. Hered.* 90: 351–356.
- Dombrowski, S., Hoffmann, M., Guha, C. and Binder, S. (1999) Continuous primary sequence requirements in the 18-nucleotide promoter of dicot plant mitochondria. *J. Biol. Chem.* 274: 10094–10099.
- Gaikwad, K., Baldev, A., Kirti, P.B., Mohapatra, T., Bhat, S.R., Chopra, V.L. and Prakash, S. (2006) Organization and expression of the mitochondrial genome in CMS (*Moricandia*) *Brassica juncea*: nuclear–mitochondrial incompatibility results in differential expression of the mitochondrial *atpx* gene. *Plant Breed.* 125: 623–628.
- Hanson, M.R. and Bentolila, S. (2004) Interactions of mitochondrial and nuclear genes that affect male gametophyte development. *Plant Cell* 16: S154–S169.
- Horn, R. (2006) Cytoplasmic male sterility and fertility restoration in higher plants. *Prog. Bot.* 67: 31–52.
- Kamps, T.L., McCarty, D.R. and Chase, C.D. (1996) Gametophyte genetics in *Zea mays* L.: dominance of a restoration-of-fertility allele (*Rf3*) in diploid pollen. *Genetics* 42: 1001–1007.
- Kuhn, J. and Binder, S. (2002) RT-PCR analysis of 5' to 3'-end-ligated mRNAs identifies the extremities of *cox2* transcripts in pea mitochondria. *Nucleic Acids Res.* 30: 439–346.
- Liu, Z. and Butow, R.A. (2006) Mitochondrial retrograde signaling. *Annu. Rev. Genet.* 40: 159–185.
- L'Homme, Y., Stahl, R.J., Li, X-Q, Hameed, A. and Brown, G.G. (1997) *Brassica nap* cytoplasmic male sterility is associated with expression of a mt DNA region containing a chimeric gene similar to the *pol* CMS associated *orf 224* gene. *Curr. Genet.* 31: 325–335.
- Pathania, A., Bhat, S.R., Dinesh Kumar, V., Ashutosh, Kirti, P.B., Prakash, S. and Chopra, V.L. (2003) Cytoplasmic male sterility in alloplasmic *Brassica juncea* carrying *Diptotaxis catholica* cytoplasm: molecular characterization and genetics of fertility restoration. *Theor. Appl. Genet.* 107: 455–461.
- Prakash, S., Kirti, P.B., Bhat, S.R., Gaikwad, K., Kumar, V.D. and Chopra, V.L. (1998) A *Moricandia arvensis*-based cytoplasmic male sterility and fertility restoration system in *Brassica juncea*. *Theor. Appl. Genet.* 97: 488–492.
- Schnable, P.S. and Wise, R.P. (1998) The molecular basis of cytoplasmic male sterility and fertility restoration. *Trends Plant Sci.* 3: 175–180.
- Small, I.D. and Peeters, N. (2000) The PPR motif—a TPR-related motif prevalent in plant organellar proteins. *Trends Biochem. Sci.* 25: 46–47.
- Wang, Z., Zou, Y., Li, X., Zhang, Q., Chen, L., et al. (2006) Cytoplasmic male sterility of rice with Boro II cytoplasm is caused by a cytotoxic peptide and is restored by two related PPR motif genes via distinct modes of mRNA silencing. *Plant Cell* 18: 676–687.
- Xiao, H., Zhang, F. and Zheng, Y. (2006) The 5' stem-loop and its role in mRNA stability in maize S cytoplasmic male sterility. *Plant J.* 47: 864–872.

(Received October 8, 2007; Accepted December 22, 2007)