

Cryptic diversity in the *Antherospora vaillantii* complex on *Muscari* species

Marcin Piątek¹, Matthias Lutz², and Arthur O. Chater³

¹Department of Mycology, W. Szafer Institute of Botany, Polish Academy of Sciences, Lubicz 46, PL-31-512 Kraków, Poland; corresponding author e-mail: m.piatek@botany.pl

²Evolutionäre Ökologie der Pflanzen, Institut für Evolution und Ökologie, University of Tübingen, Auf der Morgenstelle 1, D-72076 Tübingen, Germany

³Windover, Penyrangor, Aberystwyth, SY23 1BJ, UK

Abstract: The anther smut fungi in the ustilaginomycetous genus *Antherospora* (*Floromycetaceae*, *Urocystidales*) that infect monocots, are currently placed in nine species. Against the background of the generally observed high host specificity in smut fungi, the broad host range reported for some of the species suggests much higher diversity. *Antherospora vaillantii* s. lato includes anther smuts on different *Muscari* species. In this study, specimens of anther smuts on *Muscari armeniacum*, *M. botryoides*, *M. comosum*, and *M. tenuiflorum* were analysed by rDNA sequences and morphology to determine whether they represented one polyphagous or several host specific species. The molecular phylogeny revealed three distinct lineages that were correlated with host plants, yet had only slight morphological differences. These lineages are assigned to three cryptic species: *Antherospora hortensis* sp. nov. on *Muscari armeniacum*, *A. muscari-botryoidis* comb. nov. (syn. *Ustilago muscari-botryoidis*) on *M. botryoides*, and *A. vaillantii* s. str. on *M. comosum* and *M. tenuiflorum*. All species on *Muscari* form a monophyletic group within *Antherospora*, and the phylogenetic relations within this group coincide well with the subgeneric classification of the respective host species. This indicates a common ancestry of *Muscari* anther smuts and co-evolution as a driver of their diversification.

Key words:

Basidiomycota
Cryptic species
Molecular analysis
Muscari
Phylogeny
Plant pathogens
Smut fungi

Article info: Submitted: 27 January 2013; Accepted: 28 February 2013; Published: 4 April 2013.

INTRODUCTION

The anther smut fungus *Ustilago vaillantii* was described by the Tulasne brothers at the beginning of systematic mycology (Tulasne & Tulasne 1847). The specific epithet honoured the pre-Linnean botanist Sébastien Vaillant (1669–1722), the first collector of anther-infected *Muscari comosum*. In the protologue, the Tulasne brothers included three host species of *Ustilago vaillantii* in the following order: *Muscari comosum*, *Scilla anthericoides*, and *Scilla maritima*. The two latter names are synonyms of the species currently named *Charybdis maritima* (Speta 1988, Euro+Med 2006-). The type host was attributed to *Muscari comosum* by Liro (1924) and Ciferri (1928), who reasonably explained this selection.

Subsequent to its description, *Ustilago vaillantii* became a catch-all for different smut specimens sporulating in the anthers of monocotyledonous plants that currently are placed in *Asparagaceae* subfam. *Scilloideae* (syn. *Hyacinthaceae*, APG III 2009, Chase *et al.* 2009). Some attempts to describe separate taxa within this complex, such as *Ustilago vaillantii* var. *tourneuxii* on *Bellevalia trifoliata* (Fischer von Waldheim 1880), *U. albucae* on *Albuca* sp. (Sydow & Sydow 1914a), *U. peglerae* on *Ornithogalum lacteum* (Sydow & Sydow 1914b),

U. muscari-botryoidis on *Muscari botryoides* (Ciferri 1928), *U. scillae* on *Scilla bifolia* (Ciferri 1931) or *U. urgineae* on *Charybdis maritima* (Maire 1931), were based mostly on small morphological differences and/or supposed host specificity, and received moderate acceptance by smut researchers (Zundel 1953). In general, most authors recognised anther smuts on monocots as one species *Ustilago vaillantii* (Liro 1924, Vánky 1985) and such an approach was adopted until very recently (Vánky 1994).

The first classification of smut fungi based on ultrastructure and molecular phylogeny (Bauer *et al.* 1997, Begerow *et al.* 1997, Bauer *et al.* 2001) restricted the genus *Ustilago* to species on *Poaceae*. Consequently, *Ustilago vaillantii* was moved to the genus *Vankya*, typified by the foliicolous *V. ornithogali* (Ershad 2000). Molecular analyses revealed that the anther smuts on monocots were only distantly related to the foliicolous representatives of *Vankya*, and a new genus *Antherospora*, typified by *A. vaillantii*, was established for the former group (Bauer *et al.* 2008). The genus *Vankya* is currently restricted to host plants in *Liliaceae* subfam. *Lilioideae* (Vánky 2009). In addition to describing a new genus, Bauer *et al.* (2008) demonstrated that the evolution of the anther smuts on monocots is related to host taxonomy,

© 2013 International Mycological Association

You are free to share - to copy, distribute and transmit the work, under the following conditions:

Attribution: You must attribute the work in the manner specified by the author or licensor (but not in any way that suggests that they endorse you or your use of the work).

Non-commercial: You may not use this work for commercial purposes.

No derivative works: You may not alter, transform, or build upon this work.

For any reuse or distribution, you must make clear to others the license terms of this work, which can be found at <http://creativecommons.org/licenses/by-nc-nd/3.0/legalcode>. Any of the above conditions can be waived if you get permission from the copyright holder. Nothing in this license impairs or restricts the author's moral rights.

and consequently a narrow species concept within this group was warranted, despite limited access to fresh specimens for molecular analyses. The anther smuts on monocots are uncommon and difficult to detect because the sori are often hidden by the perianths, at least in some host genera (e.g. *Bellevalia*, *Muscari*). In comparison, the anther smuts on dicots (Lutz *et al.* 2005, 2008, Roets *et al.* 2008, Curran *et al.* 2009, Hood *et al.* 2010, Kemler *et al.* 2009, 2013, Piątek *et al.* 2012, 2013) are more often collected.

The species concept adopted by Bauer *et al.* (2008) narrowed *Antherospora vaillantii* to smuts infecting *Muscari* species (Vánky 2012). However, molecular analyses revealed genetic divergences between smut collections from different *Muscari* species (Bauer *et al.* 2008), which suggest that host specialization may be higher than to the genus level and that more than one species may have radiated on *Muscari* species. Indeed, in addition to *Antherospora vaillantii* on *Muscari comosum*, a second species was described for anther smuts on *Muscari*, namely *Ustilago muscari-botryoidis* on *Muscari botryoides*, although in recent monographs it was considered a synonym of the former species (Vánky 1985, 1994, 2012, Scholz & Scholz 1988, Karatygin & Azbukina 1989, Denchev 2001). The history of discovery and description of *Ustilago muscari-botryoidis* well illustrates the approach leading to the descriptions of different biological species by taxonomists working at the beginning of 20th century, prior to establishment of molecular methods. In 1921, Ciferri (1928) discovered an anther-infected population of *Muscari botryoides* in northern Italy. The co-occurring *Muscari comosum*, flowering later, was healthy. Subsequently, Ciferri (1928) made a three-year observation of a co-cultivation of infected *Muscari botryoides* with non-infected *M. comosum* and *M. neglectum* (as *M. racemosum*). After the three years, *Muscari botryoides* was still infected while *M. comosum* and *M. neglectum* remained healthy. Consequently, the anther smut on *Muscari botryoides* was described as a distinct biological species. In the study of Bauer *et al.* (2008), anther smut collections on *Muscari botryoides* were not available and the *Antherospora vaillantii* species complex on *Muscari*, including the identity of *Ustilago muscari-botryoidis*, remained unresolved. Misidentification of some hosts may have added further confusion.

The present study is aimed at resolving the *Antherospora vaillantii* species complex on *Muscari*, and testing whether molecular phylogenetic lineages could be defined by morphological and/or ecological characteristics. To achieve this goal molecular phylogenetic analyses using rDNA sequences were applied as well as light and scanning electron microscope examination of different anther smut specimens, including collections from type hosts of all species known to infect *Muscari*.

MATERIALS AND METHODS

Specimen sampling and documentation

This study is based on specimens of the *Antherospora vaillantii* complex from four different *Muscari* species, originating from Germany, Israel, Slovenia, and the United Kingdom. The voucher specimens are deposited in GLM, KR, KRAM F, TUB, HAI and H.U.V. (Table 1). The latter abbreviation refers

to the personal collection of Kálmán Vánky, "Herbarium *Ustilaginales Vánky*" currently held at his home (Gabriel-Biel-Strasse 5, D-72076 Tübingen, Germany). The following host plants previously misidentified (Bauer *et al.* 2008) were re-identified during the present study using *Flora Europaea* (Tutin *et al.* 1980): (1) the specimen from Slovenia (H.U.V. 21337) identified as *M. neglectum* is *M. botryoides*; (2) the specimen from Germany (TUB 15838) identified as *M. comosum* is *M. armeniacum*; (3) the specimen from Germany (H.U.V. 21046) identified as *M. neglectum* is *M. armeniacum*. Nomenclatural novelties were registered in MycoBank (www.MycoBank.org, Crous *et al.* 2004). The genotype concept follows the proposal of Chakrabarty (2010).

Morphological examination

Dried fungal teliospores of the investigated specimens were mounted in lactic acid, heated to boiling point, and then examined under a Nikon Eclipse 80i light microscope at a magnification of $\times 1000$, using Nomarski optics (DIC). Spores were measured using NIS-Elements BR 3.0 imaging software. The extreme measurements were adjusted to the nearest 0.5 μm . The spore size range, mean and standard deviation of 50 spore measurements from each specimen are shown in Table 1. The spore size values are presented in a scatter diagram to show the distribution of the values. The species descriptions include combined values from all measured specimens. The specimens of the *Antherospora vaillantii* complex measured in previous work in lactophenol (Bauer *et al.* 2008) were measured again in lactic acid to minimize the error caused by different mounting media. LM micrographs were taken with a Nikon DS-Fi1 camera. The spores of *Antherospora* on *Muscari armeniacum* (KRAM F-49437), on *M. botryoides* (KR 27962), on *M. comosum* (KRAM F-49438 = HAI 2857), and on *M. tenuiflorum* (GLM 48095) were studied using scanning electron microscopy (SEM). For this purpose, dry spores were mounted on carbon tabs and fixed to an aluminium stub with double-sided transparent tape. The tabs were sputter-coated with carbon using a Cressington sputter-coater and viewed with a Hitachi S-4700 scanning electron microscope, with a working distance of ca. 12 mm. SEM micrographs were taken in the Laboratory of Field Emission Scanning Electron Microscopy and Microanalysis at the Institute of Geological Sciences, Jagiellonian University, Kraków (Poland).

DNA extraction, PCR, and sequencing

For methods of isolation and crushing of fungal material, DNA extraction, amplification of ITS 1 and ITS 2 regions of the rDNA including the 5.8S rDNA (ITS) and the 5'-end of the nuclear large subunit ribosomal DNA (LSU), purification of PCR products, sequencing, and processing of the raw data see Lutz *et al.* (2004), and Piątek *et al.* (2011). The DNA sequences determined for this study were deposited in GenBank (GenBank accession number are given in Table 1 and Fig. 1).

Phylogenetic analyses

In addition to the ITS and LSU sequences of *Antherospora* sp. on *Muscari* spp. obtained in this study, sequences from GenBank of the following species were used for molecular

phylogenetic analyses (Bauer *et al.* 2008, Piątek *et al.* 2011): *Antherospora scillae*, *A. tractemae*, *A. vaillantii*, and *A. vindobonensis* (GenBank accession numbers are included in Fig. 1).

To elucidate the phylogenetic position of the *Antherospora* specimens from *Muscari* spp. their concatenated ITS+LSU sequences were analysed within a dataset covering all *Antherospora* species available in GenBank. If present in

Table 1. List of *Antherospora* specimens examined, with host plants, GenBank accession numbers, spore size range, mean spore sizes with standard deviation, and reference specimens.

Species	Host	GenBank acc. no.	Spore size range (µm)	Mean spore size with standard deviation (µm)	Reference specimens ¹
<i>A. hortensis</i>	<i>M. armeniacum</i>	ITS: EF653982 LSU: EF653964	(6.0–)7.0–10.0(–13.0) × (5.5–)6.0–8.0	8.6 ± 1.7 × 6.8 ± 0.7	Germany, Baden-Württemberg, Tübingen, Gabriel-Biel-Strasse 5, 17 Apr. 2005, C. & K. Vánky, H.U.V. 21046
<i>A. hortensis</i>	<i>M. armeniacum</i>	ITS: KC175333 LSU: KC175326	(6.0–)7.5–10.5(–12.5) × (5.5–)6.0–8.5(–9.0)	8.9 ± 1.2 × 7.5 ± 0.7	Germany, Baden-Württemberg, Tübingen, garden, Stöcklestrasse 39, 22 Apr. 2011, M. Lutz, KR 27970
<i>A. hortensis</i>	<i>M. armeniacum</i>	ITS: EF653997 LSU: EF653979	(5.5–)6.0–10.5(–12.5) × 5.5–7.5(–8.0)	7.8 ± 1.5 × 6.5 ± 0.7	Germany, Baden-Württemberg, Kirchheim/Teck, garden Römersteinstrasse 12, 2 May 2006, N. Böhling, TUB 15838
<i>A. hortensis</i>	<i>M. armeniacum</i>	ITS: KC175334 LSU: KC175327	(6.5–)7.0–10.5(–12.5) × 6.0–8.5(–9.5)	9.0 ± 1.4 × 7.6 ± 0.8	UK, Wales, Ceredigion, Aberystwyth, Penglais Road, SN-590-818 [grid reference on UK national grid], 7 Apr. 2009, A.O. Chater, KRAM F-49434
<i>A. hortensis</i>	<i>M. armeniacum</i>	ITS: KC175335 LSU: KC175328	(5.5–)6.0–10.5(–11.0) × 5.0–7.5(–8.5)	7.9 ± 1.4 × 6.7 ± 0.7	UK, Wales, Ceredigion, Aberystwyth, Plas Crug cemetery, SN-591-812 [grid reference on UK national grid], 16 Apr. 2010, A.O. Chater, KRAM F-49435
<i>A. hortensis</i>	<i>M. armeniacum</i>	ITS: KC175336 LSU: KC175329	6.0–10.0(–11.0) × (5.5–)6.0–8.5	8.1 ± 1.2 × 7.1 ± 0.7	UK, Wales, Ceredigion, Aberystwyth, Cliff Terrace, garden of Northfield, SN-585-826 [grid reference on UK national grid], 15 Apr. 2010, A.O. Chater, KRAM F-49436
<i>A. hortensis</i>	<i>M. armeniacum</i>	ITS: KC175337 LSU: KC175330	6.0–10.5(–12.5) × 5.5–8.5(–9.0)	8.1 ± 1.7 × 6.9 ± 0.9	UK, Wales, Ceredigion, Llanbadarn Fawr, University campus, SN-604-811 [grid reference on UK national grid], 20 Apr. 2010, A.O. Chater, KRAM F-49437 – holotype
<i>A. muscari-botryoidis</i>	<i>M. botryoides</i>	ITS: KC175332 LSU: KC175325	(5.0–)6.0–9.5(–11.0) × (4.5–)5.0–7.5(–8.0)	7.7 ± 1.4 × 6.3 ± 0.8	Germany, Baden-Württemberg, Hechingen, B32 in Richtung Burladingen, 22 Apr. 2011, M. Lutz, KR 27962 – neotype
<i>A. muscari-botryoidis</i>	<i>M. botryoides</i>	ITS: EF653998 LSU: EF653980	(6.5–)7.0–10.5(–11.0) × (5.5–)6.0–8.5(–9.0)	8.5 ± 1.1 × 7.2 ± 0.8	Slovenia, Kras, 7 km ESE from Sezana, 30 Apr. 2006, C. & K. Vánky, H.U.V. 21337
<i>A. vaillantii</i>	<i>M. comosum</i>	ITS: KC175338 LSU: KC175331	(5.5–)6.0–10.5 × 5.0–7.5(–8.0)	7.9 ± 1.4 × 6.3 ± 0.8	Israel, Haifa district, Carmel National Park, 2 Feb. 2011, K.G. Savchenko, HAI 2857, KRAM F-49438
<i>A. vaillantii</i>	<i>M. tenuiflorum</i>	ITS: EF653986 LSU: EF653968	(5.5–)6.0–10.5(–13.0) × 5.0–7.5(–9.0)	8.3 ± 1.6 × 6.9 ± 0.8	Germany, Sachsen-Anhalt, Saalkreis, Löbejün, Schiedsberg, 25 May 2000, H. Jage, GLM 47396
<i>A. vaillantii</i>	<i>M. tenuiflorum</i>	ITS: EF653987 LSU: EF653969	6.0–10.0(–11.0) × 5.0–7.5(–8.0)	7.9 ± 1.2 × 6.5 ± 0.6	Germany, Sachsen-Anhalt, Saalkreis, Löbejün, Schiedsberg, 29 May 1999, U. Richter, GLM 48095
<i>A. vaillantii</i>	<i>M. tenuiflorum</i>	ITS: EF653988 LSU: EF653970	6.0–10.5(–12.0) × 6.0–8.5	8.3 ± 1.4 × 7.0 ± 0.6	Germany, Sachsen-Anhalt, Saalkreis, Löbejün, Schiedsberg, 24 May 1999, W. Durka, GLM 50411

¹ GLM – Herbarium of the Senckenberg Museum für Naturkunde Görlitz, Germany; HAI – Herbarium of the Institute of Evolution, University of Haifa, Israel; H.U.V. – Herbarium *Ustilaginales* Vánky, Gabriel-Biel-Str. 5, D-72076 Tübingen, Germany; KR – Herbarium of the Staatliches Museum für Naturkunde Karlsruhe, Germany; KRAM F – Mycological Herbarium of the W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków, Poland; TUB – Herbarium of the Eberhard-Karls-Universität Tübingen, Germany.




Fig. 1. Hypothesis on phylogenetic relationships within the sampled *Antherospora* specimens based on Maximum Likelihood analysis of an alignment of concatenated ITS+LSU base sequences using raxmlGUI invoking the GTRCAT and the rapid bootstrap option with 1000 replicates. The topology was rooted with *Antherospora scillae*, *A. tractemae*, and *A. vindobonensis*. ML bootstrap support values are indicated on branches before slashes, estimates for *a posteriori* probabilities are indicated after slashes. A. = *Antherospora*, M. = *Muscari*.

GenBank, the sequences of the respective type specimens were used.

Sequence alignment was obtained using MAFFT 6.853 (Kato *et al.* 2002, 2005, Kato & Toh 2008) using the L-INS-i option. The resulting alignment was used for both the detection of species specific autapomorphic nucleotide differences and the estimation of genetic distances within species (intraspecific) and between species (interspecific) using the software MEGA (Tamura *et al.* 2011). We calculated p-distances and report distances as percentage genetic distances. Gaps or different length of sequences were not used for calculations as we chose the pair-wise deletion option in MEGA.

To ensure reproducible phylogenetic analyses, manipulation of the alignment by hand as well as manual exclusion of ambiguous sites were avoided as suggested by Giribet & Wheeler (1999) and Gatesy *et al.* (1993), respectively. Highly divergent portions of the alignment were omitted using GBlocks 0.91b (Castresana 2000) with

the following options: 'Minimum Number of Sequences for a Conserved Position' to 10, 'Minimum Number of Sequences for a Flank Position' to 10, 'Maximum Number of Contiguous Non-conserved Positions' to 8, 'Minimum Length of a Block' to 5 and 'Allowed Gap Positions' to 'With half'.

The resulting alignment [new number of positions: 1336 (73% of the original 1810 positions) number of variable sites: 35] was used for phylogenetic analyses using Maximum Likelihood (ML) and a Bayesian Approach (BA). ML analysis (Felsenstein 1981) was conducted with the RAxML 7.2.6 software (Stamatakis 2006), using raxmlGUI (Silvestro & Michalak 2012), invoking the GTRCAT and the rapid bootstrap option (Stamatakis *et al.* 2008) with 1000 replicates.

For BA a Bayesian approach to phylogenetic inference using a Markov chain Monte Carlo technique was used as implemented in the computer program MrBayes 3.2.1 (Ronquist *et al.* 2012). Two runs over 5 000 000 generations, each consisting of four chains, were implemented using the general time reversible model of DNA substitution with

gamma distributed substitution rates and estimation of invariant sites, random starting trees and default starting parameters of the DNA substitution model as recommended by Huelsenbeck & Rannala (2004). Trees were sampled every 100th generation, resulting in a sampling of 50 001 trees for each run. From these, the first 25 % of trees from each run were discarded (burnin). The remaining trees were used to compute a 50% majority rule consensus tree to obtain estimates for the *a posteriori* probabilities of groups of species. This Bayesian approach to phylogenetic analysis was repeated five times to test the independence of the results from topological priors (Huelsenbeck *et al.* 2002). Trees were rooted with *Antherospora scillae*, *A. tractemae*, and *A. vindobonensis*.

RESULTS

Phylogenetic analyses

Species specific autapomorphic nucleotide differences are included in the species descriptions, intraspecific and interspecific genetic distances of the ITS+LSU rDNA are given in Table 2.

The different runs of BA that were performed and the ML analyses yielded consistent topologies. To illustrate the results, the phylogenetic hypothesis resulting from the ML analysis is presented in Fig. 1. ML bootstrap support values are indicated on branches before slashes, estimates for *a posteriori* probabilities are indicated after slashes.

In all analyses, the *Antherospora* species included in previous studies (Bauer *et al.* 2008, Piątek *et al.* 2011) were inferred with high support values, and phylogenetic relationships were as in Bauer *et al.* (2008), Piątek *et al.* (2011), and Vánky *et al.* (2008). All *Antherospora* specimens from *Muscari* spp. clustered together forming three well supported subgroups: one contained all sampled specimens from *M. botryoides*, another all specimens from *M. armeniacum*, and the third specimens from both *M. comosum* and *M. tenuiflorum*. The former two subgroups were revealed as sister taxa.

Morphology

The morphology of the *Antherospora* specimens is presented in accordance with the lineages obtained in the phylogenetic analyses, namely *Antherospora* on *Muscari armeniacum*, on *M. botryoides* and on *M. comosum*+*M. tenuiflorum*. The specimens of *Antherospora* on *Muscari* spp. shared a similar appearance of macroscopic symptoms of the infection as well as a similar morphology of spores. In all specimens, the olivaceous sori formed in all anthers of fertile flowers, and the sori were enclosed by floral perianths. The spore ornamentation and shape were regular in all collections: the spores were finely verrucose or verruculose and usually globose, subglobose, broadly ellipsoidal, or occasionally allantoid, ovoid, pyriform, or tear-shaped. The spore sizes of *Antherospora* specimens on *Muscari armeniacum*, *M. comosum* and *M. tenuiflorum* were similar and grouped in one “cloud” on the scatter diagram, while the spore sizes of specimens on *Muscari botryoides* were shorter compared to the specimens from the remaining hosts (Table 1, Fig. 2).

The mean spore length was variable between collections of *Antherospora* on *Muscari armeniacum*, ranging from 7.8–9.0 μm , less variable in specimens on *M. botryoides* (7.7–8.5 μm), and quite uniform in specimens on *M. comosum* and *M. tenuiflorum* (7.9–8.3 μm). Likewise, the mean spore width was variable between collections of *Antherospora* on *Muscari armeniacum*, ranging from 6.5–7.6 μm , and less variable in specimens on *M. botryoides* (6.3–7.2 μm), and in specimens on *M. comosum* and *M. tenuiflorum* (6.3–7.0 μm) (Table 1). The combined mean spore length and width were similar in *Antherospora* on *Muscari botryoides* and *Antherospora* on *M. comosum*+*M. tenuiflorum*, while both values were somewhat larger in *Antherospora* on *M. armeniacum*. The detailed morphological characteristics of the anther smuts on *Muscari* spp. are included in the species descriptions and depicted in Figs 3–5.

TAXONOMY

***Antherospora hortensis* M. Piątek & M. Lutz, sp. nov.**
Mycobank MB803427
(Fig. 3)

Etymology: In reference to the occurrence of this fungus on *Muscari armeniacum* in cultivated gardens, where all specimens examined in this study were collected.

Type: **UK**: Wales: Ceredigion, Llanbadarn Fawr, University campus, on *Muscari armeniacum*, 20 Apr. 2010, A.O. Chater (KRAM F-49437 – **holotype**)

Description: Parasitic on *Muscari armeniacum*. Sori in the anthers, producing a dark olive-brown, powdery mass of spores inside the pollen sacs, enclosed by the perianths. Infection systemic, all anthers in all fertile flowers in an inflorescence are infected. Spores globose, subglobose, broadly ellipsoidal, sometimes allantoid, tear-shaped or asymmetrical, (5.5–)6.0–10.5(–13.0) \times 5.0–8.5(–9.5) μm [av. \pm SD, 8.3 \pm 1.5 \times 7.0 \pm 0.9 μm , n = 350/7], olive-brown, sometimes lighter coloured on one side; wall even, ca. 0.5–0.7 μm thick, finely, densely verruculose in LM, spore profile almost smooth or finely wavy, wall finely verrucose in SEM. Autapomorphic nucleotide differences in the ITS at the positions 192, 193, and 437 and in the LSU at the position 1311 in the alignment.

The ITS/LSU hologenotype sequences are deposited in GenBank as KC175337/KC175330, respectively.

Additional specimens examined: **Germany**: Baden-Württemberg: Tübingen, Gabriel-Biel-Strasse 5, on *Muscari armeniacum* (as *M. neglectum*), 17 Apr. 2005, C. & K. Vánky (H.U.V. 21046); Tübingen, Stöcklestrasse 39, on *M. armeniacum*, 22 Apr. 2011, M. Lutz 2338 (KR 27970); Kirchheim/Teck, garden of Römersteinstrasse 12, on *M. armeniacum* (as *M. comosum*), 2 May 2006, N. Böhling (TUB 15838). – **UK**: Wales: Ceredigion, Aberystwyth, Penglais Road, on *M. armeniacum*, 7 Apr. 2009, A.O. Chater (KRAM F-49434); Ceredigion, Aberystwyth, Plas Crug cemetery, on *M. armeniacum*, 16 Apr. 2010, A.O. Chater (KRAM F-49435); Ceredigion, Aberystwyth,

Table 2. Intraspecific and interspecific genetic distances of the ITS+LSU rDNA in %.

	<i>A. hortensis</i>	<i>A. muscari-botryoidis</i>	<i>A. vaillantii</i>	<i>A. vindobonensis</i>	<i>A. tractemae</i>	<i>A. scillae</i>
<i>A. hortensis</i>	0	0.4-0.5	0.9	1.9	1	1.9
<i>A. muscari-botryoidis</i>		0.1	0.7	1.8-1.9	1.1-1.2	1.8-1.9
<i>A. vaillantii</i>			0	1.8	1.1	1.8
<i>A. vindobonensis</i>				0	1.7	0.3
<i>A. tractemae</i>					0	1.6
<i>A. scillae</i>						0




Fig. 2. A scatter diagram showing the spore size distribution in *Antherospora hortensis* (blue squares), *A. muscari-botryoidis* (black spots), and *A. vaillantii* (yellow triangles). Note that one point may include measurements of the respective length/width value from more than one spore.

Cliff Terrace, garden of Northfield, on *M. armeniacum*, 15 Apr. 2010, A.O. Chater (KRAM F-49436).

Host and distribution: On *Muscari armeniacum* (*Muscari* subgen. *Botryanthus*, *Asparagaceae*), Germany, UK. – *Antherospora vaillantii* s. lato on *M. armeniacum* was included in Vánky (2012), but no specific details are given for the source of this record. The anther smut on *Muscari schliemannii* (syn. *M. armeniacum*) from the Berlin Botanical Garden (Magnus 1896, Scholz & Scholz 1988) and on *M.*

cyaneoviolaceum (syn. *M. armeniacum*) from the Royal Botanic Gardens Kew (Kirk & Cooper 2013) most probably belongs to this species.

Comments: This species is morphologically similar to *Antherospora vaillantii* s. str., differing by a somewhat larger (longer and wider) average size of spores and host plant in *Muscari* subgen. *Botryanthus*. *Antherospora hortensis* and *A. vaillantii* ITS+LSU sequences differ in 32–35 positions in 16 loci. Among the sequence differences there




Fig. 3. *Antherospora hortensis* sp. nov. on *Muscari armeniacum*. **A.** General habit of infected plant. **B–C.** Healthy (to the left) and infected (to the right) inflorescences, with fungus sporulating in the anthers. **D.** Close-up of the anthers with fungal spores. Note that ovary is not destroyed. **E–I.** Spores seen by LM, median and superficial views (KRAM F-49437). **J–L.** Ornamentation of spores seen by SEM (KRAM F-49437). Bars: A–C = 5 mm, D = 1 mm, E–I = 10 μ m, J–L = 5 μ m.

are autapomorphic nucleotide differences in the ITS at the positions 192, 193, and 437 and in the LSU at the position 1311 in the alignment for *Antherospora hortensis*.

***Antherospora muscari-botryoidis* (Cif.) M. Piątek & M. Lutz, comb. nov.**
Mycobank MB803428
(Fig. 4)




Fig. 4. *Antherospora muscari-botryoidis* comb. nov. on *Muscari botryoides*. **A.** General habit of infected plant. **B–C.** Infected inflorescences, with fungus sporulating in the anthers. **D–G.** Spores seen by LM, median and superficial views (KR 27962). **H–J.** Ornamentation of spores seen by SEM (KR 27962). Bars: B–C = 5 mm, D–H = 10 μ m, I–J = 5 μ m.

Basionym: *Ustilago muscari-botryoidis* Cif., *Ann. Mycol.* **26**: 14 (1928).

Type: **Italy:** Province of Cuneo: Alba, Moretta, towards Santa Rosalia, on *Muscari botryoides*, [year and collector unclear but probably 1921 and R. Ciferri, respectively] (type not located, probably does not exist). – **Germany:** Baden-Württemberg: Hechingen, B32 in Richtung Burladingen, 48°20'55"N, 09°00'12"E, on *Muscari botryoides*, 22 Apr. 2011, M. Lutz (KR 27962 – **neotype designated here**).

Synonym: *Antherospora neglecta* Vánky, in Lutz & Vánky, *Lidia* **7**(2–3): 35 (2009), *nom. nud.*

Original material: **Slovenia:** Kras, 7 km ESE from Sezana, on *Muscari botryoides* (as *M. neglectum*), 30 Apr. 2006, C. & K. Vánky (H.U.V. 21337).

Parasitic on *Muscari botryoides*. Sori in the anthers, producing a dark olive-brown, powdery mass of spores inside the pollen sacs, enclosed by the perianths. Infection systemic, all anthers in all fertile flowers in an inflorescence are infected. Spores globose, subglobose, broadly ellipsoidal, sometimes tear-shaped or asymmetrical, (5.0–)6.0–10.5(–11.0) × (4.5–)5.0–8.5(–9.0) µm [av. ± SD, 8.1 ± 1.3 × 6.7 ± 0.9 µm, n = 100/2], olive-brown; wall even, ca. 0.5–0.8 µm thick, finely, densely verruculose in LM, spore profile almost smooth or finely wavy, wall finely verrucose in SEM. Autapomorphic nucleotide difference in the LSU at the position 1078 in the alignment.

The ITS/LSU neogenotype sequences are deposited in GenBank as KC175332/KC175325, respectively.

Host and distribution: On *Muscari botryoides* (*Muscari* subgen. *Botryanthus*, *Asparagaceae*), Germany, Italy, and Slovenia. This species was described from Italy (Ciferri 1928), and the sequenced specimens are from Germany and Slovenia. The different anther smut records and collections on *Muscari botryoides* from Europe (Bulgaria, Germany, UK) (Scholz & Scholz 1988, Denchev 2001, Kirk & Cooper 2013) and Australasia (New Zealand) (Vánky & McKenzie 2002) probably belong to this species. It was introduced to areas outside the natural range of *Muscari botryoides*, for example to UK and New Zealand.

Comments: This species shares with *Antherospora hortensis* the systematic position of its host plant in *Muscari* subgen. *Botryanthus*, but differs by somewhat shorter spores and a different host species (*M. botryoides* vs. *M. armeniacum*). *Antherospora muscari-botryoidis* and *A. hortensis* ITS+LSU sequences differ in 11–12 positions in 8 loci. Among the sequence differences there is an autapomorphic nucleotide difference in the LSU at the position 1078 in the alignment for *Antherospora muscari-botryoidis*.

This species differs from *Antherospora vaillantii* s. str. by somewhat shorter spores and its host plant in *Muscari* subgen. *Botryanthus*. The host plants of *Antherospora vaillantii* s. str. belong to *Muscari* subgen. *Leopoldia*. *Antherospora muscari-botryoidis* and *A. vaillantii* s. str.

ITS+LSU sequences differ in 22–26 positions in 11–12 loci. Among the sequence differences there is an autapomorphic nucleotide difference in the LSU at the position 1078 in the alignment for *Antherospora muscari-botryoidis*.

The name “*Antherospora neglecta* Vánky, in prep.” was assigned by Vánky to the Slovenian specimen in the checklist of smut fungi from Slovenia (Lutz & Vánky 2009). This name is a *nomen nudum* as it was never validated. The host plant was originally identified as *Muscari neglectum*. However, its re-identification revealed that it is *Muscari botryoides* (flowers ±globose, not elongated). Consequently, *Antherospora neglecta* is listed as a synonym of *A. muscari-botryoidis*.

Antherospora vaillantii (Tul. & C. Tul.) R. Bauer et al., *Mycol. Res.* **112**: 1304 (2008).

(Fig. 5)

Basionym: *Ustilago vaillantii* Tul. & C. Tul., *Ann. Sci. Nat., Bot., sér. 3*, **7**: 90 (1847).

Synonyms: *Yenia vaillantii* (Tul. & C. Tul.) Liou, *Contrib. Inst. Bot. Natl. Acad. Peiping* **6**: 45 (1949).

Vankya vaillantii (Tul. & C. Tul.) Ershad, *Rostaniha* **1**: 69 (2000).

Type: **France:** on *Muscari comosum*, S. Vaillant (lectotype designated by Lindeberg 1959: 141, but apparently without seeing the herbarium specimen; despite the statement of Vánky 2012: 8, who indicated that he studied the lectotype from PC, such a specimen was not located in PC in the course of the present study).

Parasitic on *Muscari comosum* and *M. tenuiflorum*. Sori in the anthers, producing a dark olive-brown, semi-powdery to powdery mass of spores inside the pollen sacs, enclosed by the perianths. Infection systemic, all anthers in all fertile flowers in an inflorescence are infected, flowers sometimes deformed. Spores globose, subglobose, broadly ellipsoid, sometimes ovoid, pyriform or asymmetrical, (5.5–)6.0–10.5(–13.0) × 5.0–8.5(–9.0) µm [av. ± SD, 8.1 ± 1.4 × 6.7 ± 0.8 µm, n = 200/4], olive-brown, sometimes lighter coloured on one side; wall even, ca. 0.5–0.7 µm thick, often darker than the rest of the spore, finely, densely verruculose in LM, spore profile almost smooth or finely wavy, wall finely verrucose in SEM. Autapomorphic nucleotide differences in the ITS at the positions 95, 443–452, and 575 and in the LSU at the positions 1292, and 1309 in the alignment.

Specimens examined: **Germany:** Sachsen-Anhalt: Saalkreis, Löbejün, Schiedsberg, on *Muscari tenuiflorum*, 25 May 2000, H. Jage (GLM 47396); Saalkreis, Löbejün, Schiedsberg, on *M. tenuiflorum*, 29 May 1999, U. Richter (GLM 48095); Saalkreis, Löbejün, Schiedsberg, on *M. tenuiflorum*, 24 May 1999, W. Durka (GLM 50411). – **Israel:** Haifa district: Carmel National Park, on *M. comosum*, 2 Feb. 2011, K.G. Savchenko (HAI 2857, KRAM F-49438 – representative specimens).

Hosts and distribution: On *Muscari comosum* and *M. tenuiflorum* (*Muscari* subgen. *Leopoldia*, *Asparagaceae*), France, Germany, and Israel. – The type material was collected in France (Tulasne & Tulasne 1847), and the sequenced specimens are from Israel (Savchenko et al.




Fig. 5. *Antherospora vaillantii* s. str. on *Muscari comosum* and *M. tenuiflorum*. **A–B.** Spores seen by LM, median and superficial views (from *M. comosum*, KRAM F-49438). **C–F.** Spores seen by LM, median and superficial views (from *M. tenuiflorum*, GLM 48095). **G–J.** Ornamentation of spores seen by SEM (G – from *M. tenuiflorum*, GLM 48095, H–J – from *M. comosum*, KRAM F-49438). Bars: A–F = 10 μ m, G–J = 5 μ m.

2011) and Germany (Jage & Richter 2011). Most likely the numerous *Antherospora* reports on *Muscari comosum* and *M. tenuiflorum* belong to *Antherospora vaillantii* s. str. The anther smut reports on *Muscari comosum* are from Europe (Austria, Belgium, Bosnia and Herzegovina, Bulgaria, Croatia, “Czechoslovakia”, France, Germany, Hungary,

Italy, Poland, Romania, Russia – European part, Serbia, Slovenia, Switzerland, Ukraine) (e.g. Lindtner 1950, Zundel 1953, Kochman & Majewski 1973, Vánky 1985, Săvulescu 1955, Scholz & Scholz 1988, Karatygin & Azbukina 1989, Denchev 2001, Zwetko & Blanz 2004, Savchenko & Heluta 2012), from Africa (Algeria, Egypt, and Morocco)

(Zundel 1953, Zambettakis 1970), and from North America (USA – Massachusetts) (Zundel 1953). The anther smut reports on *Muscari tenuiflorum* are from Europe (Austria, Bulgaria, “Czechoslovakia”, Germany, Poland, Romania, Serbia, Montenegro, and Ukraine) (Lindtner 1950, Zundel 1953, Săvulescu 1955, Kochman & Majewski 1973, Vánky 1985, Scholz & Scholz 1988, Karatygin & Azbukina 1989, Denchev 2001, Zwetko & Blanz 2004). The records from outside the natural range of the two *Muscari* species, for example in the USA, Belgium (*M. comosum*) or Poland (*M. tenuiflorum*), are the result of the introduction of host plant and fungus.

Comments: The type material of *Ustilago vaillantii* could not be located, and the typification of this smut species needs some clarification. In the protologue, Tulasne & Tulasne (1847) included three host species in the following order: *Muscari comosum*, *Scilla anthericoides*, and *Scilla maritima*, but from the text it is apparent that they directly studied only three smut specimens on *Muscari comosum* collected by Vaillant, Delastre and Lèveillé, respectively. The type host was assigned to *Muscari comosum* by Liro (1924), and accepted as such by Ciferri (1928). Furthermore, Liro (1924) studied the Delastre specimen included in the protologue, which was then deposited in the Persoon Herbarium in Leiden (L), and confirmed that the host plant was indeed *Muscari comosum*. That specimen should be treated as a syntype, and cannot be considered as lectotype since it was not indicated with the term “type” or an equivalent by Liro (1924), thus not meeting the criteria of Article 7.10 of the ICN. Liro (1924) only selected the type host of *Ustilago vaillantii* from the three hosts mentioned by the Tulasne brothers. Lindeberg (1959) designated Vaillant’s collection as a lectotype of *Ustilago vaillantii*, but apparently she did not see this material – the problem is that the specimen probably does not now exist.

Vánky (2012) indicated that he studied the lectotype specimen from PC (he included the symbol “!” at the typification entry of *Antherospora vaillantii*) selected by Lindeberg (1959). However, the examination of all materials from PC labelled as *Ustilago vaillantii* made in the course of the present study revealed no authentic specimen of this smut fungus matching the data from the protologue (Tulasne & Tulasne 1847). Likewise, the Delastre syntype specimen mentioned in the protologue and later studied by Liro (1924) was not found in the Persoon herbarium in L (G. Thijsse, pers. comm.). It is reasonable to assume that no authentic collection of *Ustilago vaillantii* is currently preserved and that a neotype should be selected for this species. However, the neotype of *Antherospora vaillantii* based on the Israeli specimen analysed in the present study is not designated here, as in our opinion it is better to select the neotype from a fresh European collection, which is not available at present (most herbarium collections are old and not useful for molecular analyses). Thus, in line with Hawksworth (2012), the Israeli collection of *Antherospora vaillantii* s. str. on the type host *Muscari comosum* is referred to as a “representative specimen”.

DISCUSSION

In this study, morphological and molecular phylogenetic analyses were performed in order to resolve the *Antherospora vaillantii* species complex on different *Muscari* species. The specimen sampling covers four of the six *Muscari* species reported as hosts for *Antherospora vaillantii* s. lato in the recent world monograph of smut fungi (Vánky 2012¹), including the type hosts for both species that were described for *Muscari* anther smuts, namely *Ustilago vaillantii* (*M. comosum*) and *U. muscari-botryoidis* (*M. botryoides*).

In accordance with previous molecular phylogenetic analyses based on a smaller sampling of *Muscari* anther smuts (Bauer *et al.* 2008, Piątek *et al.* 2011), the present analyses revealed that all anther smut specimens on *Muscari* spp. form a monophyletic group within the *Antherospora* clade. The phylogenetic relations within this lineage agree well with the classification of the respective host species: the basal group contains anther smuts on two hosts belonging to *Muscari* subgen. *Leopoldia* (*M. comosum*, *M. tenuiflorum*), while the derived group includes anther smuts on hosts belonging to *Muscari* subgen. *Botryanthus* that is further subdivided into two lineages correlated with the host plant species *M. armeniacum* and *M. botryoides*, respectively. That indicates a common ancestry of the *Muscari* anther smuts and co-evolution as a driver of their diversification. Results showing similar patterns in anther smuts on *Scilla* (Bauer *et al.* 2008) and *Tractema* (Piątek *et al.* 2011) support this hypothesis. However, confirmation of the monophyletic origin of the *Muscari* anther smuts with a larger sampling of anther smut specimens on diverse host plants, especially on the related genus *Bellevalia*, is desirable.

The genetic divergence between three lineages of *Muscari* anther smuts obtained does not correspond to differences in phenotypic characteristics. The morphology of all anther smut specimens was quite similar and, without support of non-morphological data, the small differences in spore size or average spore sizes observed between them could easily be treated as phenotypic variability of one single species. In contrast to the study of Bauer *et al.* (2008), where the spore size of one collection on *Muscari tenuiflorum* (GLM 47396) had much larger spores and deviated from the remaining specimens on this host, the repeated measurement of the same specimen in this study did not confirm such deviation. They were similar to values obtained from the remaining specimens on *Muscari tenuiflorum*. The different mounting media (lactophenol used by Bauer *et al.* 2008 vs. lactic acid used in this study) cannot explain this discrepancy because the repeated measurements of the remaining specimens used in the previous study did not reveal such differences. This discordance may have resulted from measuring spores from different sori. The examples from other studies indicate that in some smut samples, the spores measured from different sori within a collection may have a different size or morphology, for example in *Anthracoidea*

¹*M. alpinum* and *M. schliemannii* listed in this monograph are excluded here as both are synonyms of other species (Euro+Med 2006-)

hostianae (Piątek & Mutenko 2010) or *Farysia ugandana* (Vánky 2004a). Such anomalies between collections are also known in other groups of basidiomycetes, for example in polypores (Niemelä *et al.* 2001), and in ascomycetes, for example in *Calonectria* (Crous *et al.* 2006). The large spore size values of anther smut on *Muscari tenuiflorum* are not included in this study. Using this example, Vánky (2008) introduced the term *pseudomorphospecies* for smuts with morphological differences showing no genetic differences. This concept, however, does not take into account variation of morphological characters within specimens of the same species, as well as possible anomalies in spore morphology such as discussed above.

Molecular phylogenetic analyses revealed all anther smut specimens on *Muscari botryoides*, on *M. armeniacum*, and on *M. comosum* and *M. tenuiflorum*, respectively, to be in strongly supported monophyletic groups forming independent lineages. Furthermore, each lineage had autapomorphic nucleotide differences in the ITS and/or the LSU in the alignment. In addition, pairwise genetic distances between the lineages were found sufficient to support their separation as distinct species. They ranged from 0.4–0.9 % (within the lineages pairwise genetic distances ranged from 0–0.1 %). Thus, in spite of weak morphological differences, the genetic divergence and strict correlation with host species taxonomy indicate that the three lineages of the *Muscari* anther smuts in fact represent three different cryptic species. The host plant species appear to be good taxonomic markers for this group of parasites, as the collections from the same host (*Muscari armeniacum*, *M. botryoides*) or closely related hosts (*M. comosum*+*M. tenuiflorum*) grouped together, irrespective of their geographic origin. This is in agreement with similar results obtained for other smut fungi (Hendrichs *et al.* 2005, Lutz *et al.* 2005, 2008, Piątek *et al.* 2012). Because of some uncertainties concerning the effective description of species on the basis of molecular characters (Tripp & Lendemer 2012), the descriptions of the three recognized *Antherospora* species on *Muscari* include, in addition to the host plant name and morphology, the species specific autapomorphic nucleotides of the ITS and LSU.

The three lineages of anther smuts on *Muscari* correspond to *Antherospora vaillantii* s. str., *Ustilago muscari-botryoidis* and an undescribed species. The basal lineage containing the collection on *Muscari comosum* from Israel is assigned to *Antherospora vaillantii* s. str. The molecular results indicate that this species is additionally able to infect the closely related *Muscari tenuiflorum*. The clade represented by two specimens on *Muscari botryoides* is referred to a second species, for which the name *Ustilago muscari-botryoidis* is available (Ciferri 1928). Because this is a species of *Antherospora*, a new combination in this genus is introduced. In contrast to other monographers who synonymized it with *Antherospora vaillantii* (e.g. Vánky 1985, 1994, 2012, Scholz & Scholz 1988, Karatygin & Azbukina 1989, Denchev 2001), the present study confirms that Ciferri (1928) was correct in describing *Ustilago muscari-botryoidis*. Compared to the remaining anther smuts on *Muscari*, *Antherospora muscari-botryoidis* has somewhat smaller spores. However, more specimens should be analysed both morphologically and genetically to confirm this feature as stable and taxonomically informative for this smut.

The third lineage, containing specimens on *Muscari armeniacum*, represents a previously undescribed species. The discovery of this species is unexpected as all anther-infected *Muscari armeniacum* plants were found in gardens outside the natural range of the host species. This special habitat occurrence is reflected in the proposed epithet for this species, *Antherospora hortensis*. Except for the brief information in Vánky (2012), *Muscari armeniacum* was not reported in any publication as a host for anther smuts, at least under this species name. However, the old records of *Ustilago vaillantii* on *Muscari schliemannii* from Berlin Botanical Garden (Magnus 1896), where it was observed during the years 1892–1897 (Scholz & Scholz 1988), most probably belong to the same pathogen. *Muscari schliemannii* is a synonym of *M. armeniacum* (Euro+Med 2006-). Likewise, the 1931 collection of *Ustilago vaillantii* on *Muscari cyaneoviolaceum* from Kew Gardens (Kirk & Cooper 2013) probably also belongs to *Antherospora hortensis*. *Muscari cyaneoviolaceum* is a synonym of *M. armeniacum* (Euro+Med 2006-). This indicates that *Antherospora hortensis* persisted on cultivated plants in Germany and UK for a long time and the recent findings in Baden-Württemberg and Wales are probably only the result of special collecting activity directed to smut fungi. The concentration of localities around Tübingen in Germany as well as around Aberystwyth in UK may also reflect local transmission of infected plants between gardens by gardeners. The anther smuts on monocots may propagate by bulbs or even by infected seeds (Massee 1914, Kochman 1936). The current data suggest that the distribution of *Antherospora hortensis* is as yet localized. This could be also supported by the observation that *Antherospora hortensis* was not found in Kraków in Poland despite that *Muscari armeniacum* is commonly cultivated in gardens and has been examined by one of us (MP) for anther smuts every year since 2007. At present *Antherospora hortensis* does not pose a significant disease threat, although this may change since ongoing climate change allows many pathogens to spread to new geographical areas.

The origin of *Antherospora hortensis* is unclear. The native occurrence of *Muscari armeniacum* is in the Balkan Peninsula, but no smutted specimens have been reported from this area. In this region there are reports of anther smuts on *Muscari neglectum* (Lindtner 1950, Zundel 1953, Denchev 2001). These two *Muscari* species were often misidentified, and according to Tutin *et al.* (1980), many records of *M. neglectum* from the Balkan Peninsula in fact represent *M. armeniacum*. The occurrence of anther smuts on *Muscari neglectum* should be re-examined, preferably with fresh collections as drying often affects flower colour, which is a diagnostic character of *M. neglectum*.

Besides the host plants included in the current study or discussed above, two further species have been reported for *Antherospora vaillantii* s. lato in the recent world monograph (Vánky 2012), namely *Muscari alpinum* and *M. moschatum*. The identity of *Muscari alpinum* is uncertain since there are two names with the same epithet: *M. alpinum* J. Gay ex Baker and *M. alpinum* Szafer ex Racib., which are synonyms of *M. tenuiflorum* and *M. botryoides*, respectively (Euro+Med 2006-). Although Vánky (2012) attributed this name to “J. Gay”, in the original report the species name is given without

any authorities (Bubák 1916). The anther smut on *Muscari moschatum* that was reported from Georgia (Karatygin & Azbukina 1989), is a putative fourth distinct species within the *Antherospora vaillantii* complex, a hypothesis based on the isolated position of the host in *Muscari* subgen. *Muscari*. Fresh materials for molecular and morphological analyses are desirable to confirm this hypothesis.

In addition to resolving the species within the *Antherospora vaillantii* complex on *Muscari*, this study gives further evidence that host specificity may reveal cryptic diversity in some smut fungi, to a greater degree than morphology. Past assumptions were that host specificity was restricted to the host family (Fischer & Shaw 1953), subtribe (Vánky 2004b) or, mostly, genus level (Vánky 1994, 2012). Recent molecular studies indicate that host specialization is, in most cases, at the species level (Kemler *et al.* 2009, Lutz *et al.* 2008, 2012, Piątek *et al.* 2012, 2013, Savchenko *et al.* 2013). This is not an absolute rule since some smut species may infect two or more host species from the same genus (Castlebury & Carris 1999, Lutz *et al.* 2005, Curran *et al.* 2009, Kemler *et al.* 2013), or two or more species from different but usually phylogenetically closely related genera (Carris *et al.* 2007, Vánky & Lutz 2007). Thus, the level of host specificity in different smut species reported from multiple host species and genera, including the other members of the genus *Antherospora*, remains to be tested and is a challenge for future studies.

ACKNOWLEDGEMENTS

We thank Michael Weiß, Sigisfredo Garnica, and Robert Bauer (Tübingen, Germany) for providing facilities for molecular analyses, Kyrylo G. Savchenko (Haifa, Israel) for sending a material, the Curators of herbaria: Herbert Boyle (GLM), Bart Buyck (PC), Markus Scholler (KR), and Kálmán Vánky (H.U.V.) for loan of specimens, and Anna Łatkiewicz (Kraków, Poland) for her help with the SEM pictures. Gerard Thijssse (Leiden, Netherlands) tried to locate the Delastre collection of *Ustilago vaillantii* in the Persoon herbarium (L) and we greatly appreciate his endeavours.

REFERENCES

- APG III (2009) An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III. *Botanical Journal of the Linnean Society* **161**: 105–121.
- Bauer R, Begerow D, Oberwinkler F, Piepenbring M, Berbee ML (2001) *Ustilaginomycetes*. In: *The Mycota*. Vol. VII, Part B. *Systematics and Evolution* (DJ McLaughlin, EG McLaughlin, PA Lemke, eds): 57–83. Berlin: Springer-Verlag.
- Bauer R, Lutz M, Begerow D, Piątek M, Vánky K, Bacigálová K, Oberwinkler F (2008) Anther smut fungi on monocots. *Mycological Research* **112**: 1297–1306.
- Bauer R, Oberwinkler F, Vánky K (1997) Ultrastructural markers and systematics in smut fungi and allied taxa. *Canadian Journal of Botany* **75**: 1237–1314.
- Begerow D, Bauer R, Oberwinkler F (1997) Phylogenetic studies on nuclear LSU rDNA sequences of smut fungi and related taxa. *Canadian Journal of Botany* **75**: 2045–2056.
- Bubák F (1916) Die Pilze Böhmens. II. Teil. Brandpilze (Hemibasidii). *Archiv für die Naturwissenschaftliche Landesdurchforschung von Böhmen* **15**(3): 1–81.
- Carris LM, Castlebury LA, Huang GM, Alderman SC, Luo JF, Bao XD (2007) *Tilletia vankyi*, a new species of reticulate-spored bunt fungus with non-conjugating basidiospores infecting species of *Festuca* and *Lolium*. *Mycological Research* **111**: 1386–1398.
- Castlebury LA, Carris LM (1999) *Tilletia walkeri*, a new species on *Lolium multiflorum* and *L. perenne*. *Mycologia* **91**: 121–131.
- Castresana J (2000) Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Molecular Biology and Evolution* **17**: 540–552.
- Chakrabarty P (2010) Genetypes: a concept to help integrate molecular phylogenetics and taxonomy. *Zootaxa* **2632**: 67–68.
- Chase MW, Reveal JL, Fay MF (2009) A subfamilial classification for the expanded asparagalean families *Amaryllidaceae*, *Asparagaceae* and *Xanthorrhoeaceae*. *Botanical Journal of the Linnean Society* **161**: 132–136.
- Ciferri R (1928) Quarta contribuzione allo studio degli *Ustilaginales*. *Annales Mycologici* **26**: 1–68.
- Ciferri R (1931) Quinta contribuzione allo studio degli *Ustilaginales*. *Annales Mycologici* **29**: 1–74.
- Crous PW, Gams W, Stalpers JA, Robert V, Stegehuis G (2004) MycoBank: an online initiative to launch mycology into the 21st century. *Studies in Mycology* **50**: 19–22.
- Crous PW, Groenewald JZ, Risède JM, Simoneau P, Hyde KD (2006) *Calonectria* species and their *Cylindrocladium* anamorphs: species with clavate vesicles. *Studies in Mycology* **55**: 213–226.
- Curran HR, Roets F, Dreyer LL (2009) Anther-smut fungal infection of South African *Oxalis* species: spatial distribution patterns and impacts on host fecundity. *South African Journal of Botany* **75**: 807–815.
- Denchev CM (2001) *Classis Ustomycetes (Ordines Tilletiales, Ustilaginales et Graphiolales)*. Sofia: Pensoft.
- Ershad D (2000) *Vankya*, a new genus of smut fungi. *Rostaniha* **1**: 65–72.
- Euro+Med (2006-) *Euro+Med PlantBase - the information resource for Euro-Mediterranean plant diversity*. Published on the Internet <http://ww2.bgbm.org/EuroPlusMed/>.
- Felsenstein J (1981) Evolutionary trees from DNA sequences: a maximum likelihood approach. *Journal of Molecular Evolution* **17**: 368–376.
- Fischer GW, Shaw CG (1953) A proposed species concept in the smut fungi, with application to North American species. *Phytopathology* **43**: 181–188.
- Fischer von Waldheim A (1880) Über zwei neue aussereuropäische Brandpilze. *Verhandlungen des Botanischen Vereins der Provinz Brandenburg* **22**: 65.
- Gatesy J, DeSalle R, Wheeler W (1993) Alignment-ambiguous nucleotide sites and the exclusion of systematic data. *Molecular Phylogenetics and Evolution* **2**: 152–157.
- Giribet G, Wheeler WC (1999) On gaps. *Molecular Phylogenetics and Evolution* **13**: 132–143.
- Hawksworth DL (2012) Addressing the conundrum of unavailable name-bearing types. *IMA Fungus* **3**: 155–158.
- Hendrichs M, Begerow D, Bauer R, Oberwinkler F (2005) The genus *Anthracoidea* (*Basidiomycota*, *Ustilaginales*): a molecular phylogenetic approach using LSU rDNA sequences. *Mycological Research* **109**: 31–40.

- Hood ME, Mena-Ali JI, Gibson AK, Oxelman B, Giraud T, Yockteng R, Arroyo MTK, Conti F, Pedersen AB, Gladieux P, Antonovics J (2010) Distribution of the anther-smut pathogen *Microbotryum* on species of the *Caryophyllaceae*. *New Phytologist* **187**: 217–229.
- Huelsenbeck JP, Larget B, Miller RE, Ronquist F (2002) Potential applications and pitfalls of Bayesian inference of phylogeny. *Systematic Biology* **51**: 673–688.
- Huelsenbeck JP, Rannala B (2004) Frequentist properties of Bayesian posterior probabilities of phylogenetic trees under simple and complex substitution models. *Systematic Biology* **53**: 904–913.
- Jage H, Richter U (2011) 10 Jahre Exkursionen zum Erkunden phytoparasitischer Kleinpilze (2000–2009) in Deutschland. *Zeitschrift für Mykologie* **77**: 243–258.
- Karatygin IV, Azbukina ZM (1989) *Opredelitel' gribov SSSR. Porjadok Golovnevye*. Vyp. 1. *Semejstvo Ustilagovye (Definitorium fungorum URSS. Ordo Ustilaginales. Fasc. 1. Familia Ustilaginaceae)*. Leningrad: Nauka.
- Katoh K, Kuma K, Toh H, Miyata T (2005) MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acids Research* **33**: 511–518.
- Katoh K, Misawa K, Kuma K, Miyata T (2002) MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research* **30**: 3059–3066.
- Katoh K, Toh H (2008) Recent developments in the MAFFT multiple sequence alignment program (outlines version 6). *Briefings in Bioinformatics* **9**: 286–298.
- Kemler M, Lutz M, Göker M, Oberwinkler F, Begerow D (2009) Hidden diversity in the non-caryophyllaceous plant-parasitic members of *Microbotryum* (*Pucciniomycotina: Microbotryales*). *Systematics and Biodiversity* **7**: 297–306.
- Kemler M, Martín MP, Tellería MT, Schäfer AM, Yurkov A, Begerow D (2013) Contrasting phylogenetic patterns of anther smuts (*Pucciniomycotina: Microbotryum*) reflect phylogenetic patterns of their caryophyllaceous hosts. *Organisms Diversity and Evolution* (in press).
- Kirk P, Cooper J (2013) *The fungal records database of Britain and Ireland*. <http://www.fieldmycology.net/FRDBI/FRDBI.asp>.
- Kochman J (1936) *Ustilaginales Poloniae. Planta Polonica* **4**: 1–161.
- Kochman J, Majewski T (1973) *Basidiomycetes, Ustilaginales*. In: *Flora Polska. Rośliny zarodnikowe Polski i ziem ościennych. Grzyby* (J Kochman, A Skirgiełło, eds) **5**: 1–271. Warszawa–Kraków: Państwowe Wydawnictwo Naukowe.
- Lindeberg B (1959) *Ustilaginales of Sweden (exclusive of the Cintractias on Caricoideae)*. *Symbolae Botanicae Upsalienses* **16**(2): 1–175.
- Lindtner V (1950) *Ustilaginales Jugoslaviae. Bulletin du Muséum d'Historie Naturelle du Pays Serbe, Série B* **3–4**: 1–110.
- Liro JI (1924) Die Ustilagineen Finnlands I. *Annales Academiae Scientiarum Fennicae, Series A* **17**(1): 1–636.
- Lutz M, Bauer R, Begerow D, Oberwinkler F, Triebel D (2004) *Tuberculina*, rust relatives attack rusts. *Mycologia* **96**: 614–626.
- Lutz M, Göker M, Piątek M, Kemler M, Begerow D, Oberwinkler F (2005) Anther smuts of *Caryophyllaceae*: molecular characters indicate host-dependent species delimitation. *Mycological Progress* **4**: 225–238.
- Lutz M, Piątek M, Kemler M, Chlebicki A, Oberwinkler F (2008) Anther smuts of *Caryophyllaceae*: molecular analyses reveal further new species. *Mycological Research* **112**: 1280–1296.
- Lutz M, Vánky K (2009) An annotated checklist of smut fungi (*Basidiomycota: Ustilaginomycotina and Microbotryales*) in Slovenia. *Lidia* **7**(2–3): 33–78.
- Lutz M, Vánky K, Bauer R (2012) *Melanoxa*, a new genus in the *Urocystidales (Ustilaginomycotina)*. *Mycological Progress* **11**: 149–158.
- Magnus P (1896(1895) Die Ustilagineen (Brandpilze) der Provinz Brandenburg. *Verhandlungen des Botanischen Vereins der Provinz Brandenburg* **37**: 66–97.
- Maire R (1931) Champignons parasites africains nouveaux ou peu connus. In: *Recueil de Travaux Cryptogamiques dédiés à Louis Mangin*: 355–360. Paris.
- Massee I (1914) Observations on the life-history of *Ustilago Vaillantii* Tul. *The Journal of Economic Biology* **9**: 9–14.
- Niemelä T, Wagner T, Fischer M, Dai YC (2001) *Phellopilus* gen. nov. and its affinities within *Phellinus* s. lato and *Inonotus* s. lato (*Basidiomycetes*). *Annales Botanici Fennici* **38**: 51–62.
- Piątek M, Lutz M, Kemler M (2013) *Microbotryum silenes-saxifragae* sp. nov. sporulating in the anthers of *Silene saxifraga* in southern European mountains. *IMA Fungus* **4**: 29–40.
- Piątek M, Lutz M, Ronikier A, Kemler M, Świdorska-Burek U (2012) *Microbotryum heliospermae*, a new anther smut fungus parasitic on *Heliosperma pusillum* in the mountains of the European Alpine System. *Fungal Biology* **116**: 185–195.
- Piątek M, Lutz M, Smith PA, Chater AO (2011) A new species of *Antherospora* supports the systematic placement of its host plant. *IMA Fungus* **2**: 135–142.
- Piątek M, Muleńko W (2010) The calcareous mires in South-East Poland are home to two rare *Anthracoidea* species. *Acta Mycologica* **45**: 175–184.
- Roets F, Dreyer LL, Wingfield MJ, Begerow D (2008) *Thecaphora capensis* sp. nov., an unusual new anther smut on *Oxalis* in South Africa. *Persoonia* **21**: 147–152.
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* **61**: 539–542.
- Savchenko KG, Heluta VP (2012) Smut fungi of Ukraine, a checklist. *Sydowia* **64**: 281–300.
- Savchenko KG, Heluta VP, Wasser SP, Nevo E (2011) A contribution to the study of smut fungi of Israel. *Mycotaxon* **118**: 303–309.
- Savchenko KG, Lutz M, Piątek M, Heluta VP, Nevo E (2013) *Anthracoidea caricis-meadii* is a new North American smut fungus on *Carex* sect. *Panicaceae*. *Mycologia* **105**: 181–193.
- Săvulescu T (1955) *Die Ustilaginales der Rumänischen Volksrepublik*. Bucureşti: Akademie Verlag.
- Scholz H, Scholz I (1988) Die Brandpilze Deutschlands (*Ustilaginales*). *Englera* **8**: 1–691.
- Silvestro D, Michalak I (2012) raxmlGUI: a graphical front-end for RAxML. *Organisms Diversity and Evolution* **12**: 335–337.
- Speta F (1988) Systematische Analyse der Gattung *Scilla* L. s.l. (*Hyacinthaceae*). *Phyton* **38**: 1–141.
- Stamatakis A (2006) RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* **22**: 2688–2690.
- Stamatakis A, Hoover P, Rougemont J (2008) A rapid bootstrap algorithm for the RAxML web servers. *Systematic Biology* **57**: 758–771.
- Sydow H, Sydow P (1914a) Fungi. In: *Wissenschaftliche Ergebnisse der Deutschen Zentral-Afrika-Expedition 1907–1908 unter*

- Führung Adolf Friedrichs, Herzogs zu Mecklenburg. Band II. *Botanik* (J Mildbraed, ed): 95–101. Leipzig: Verlag von Klinkhardt und Biermann.
- Sydow H, Sydow P (1914b) Beschreibungen neuer Südafrikanischer Pilze. III. *Annales Mycologici* **12**: 263–267.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* **28**: 2731–2739.
- Tripp EA, Lendemer JC (2012) Request for binding decisions on the descriptive statements associated with *Mortierella sigyensis* (fungi: *Mortierellaceae*) and *Piromyces cryptodigmaticus* (fungi: *Neocallimastigaceae*). *Taxon* **61**: 886–888.
- Tulasne LR, Tulasne C (1847) Mémoire sur les Ustilaginées comparées aux Urédinées. *Annales des Sciences Naturelles, Botanique, Série 3*, **7**: 12–127+Pls. 2–7.
- Tutin TG, Heywood VH, Burges NA, Valentine DH, Walters SM, Webb DA (1980) *Flora Europaea*. Vol. 5. *Alismataceae* to *Orchidaceae* (Monocotyledones). Cambridge: Cambridge University Press.
- Vánky K (1985) Carpathian *Ustilaginales*. *Symbolae Botanicae Upsalienses* **24**(2): 1–309.
- Vánky K (1994) *European Smut Fungi*. Stuttgart-Jena-New York: G. Fischer Verlag.
- Vánky K (2004a) Taxonomic studies on *Ustilaginomycetes*. 24. *Mycotaxon* **89**: 55–118.
- Vánky K (2004b) The smut fungi (*Ustilaginomycetes*) of *Boutelouinae* (*Poaceae*). *Fungal Diversity* **16**: 167–198.
- Vánky K (2008) Smut fungi (*Basidiomycota* p.p., *Ascomycota* p.p.) of the world. Novelties, selected examples, trends. *Acta Microbiologica et Immunologica Hungarica* **55**: 91–109.
- Vánky K (2009) The genus *Vankya* (*Urocystidaceae*) revisited. *Mycologia Balcanica* **6**: 73–78.
- Vánky K (2012) *Smut Fungi of the World*. St Paul, MN: American Phytopathological Society Press.
- Vánky K, Lutz M (2007) Revision of some *Thecaphora* species (*Ustilaginomycotina*) on *Caryophyllaceae*. *Mycological Research* **111**: 1207–1219.
- Vánky K, Lutz M, Bauer R (2008) *Floromyces*, a new genus of *Ustilaginomycotina*. *Mycotaxon* **104**: 171–184.
- Vánky K, McKenzie EHC (2002) Smut fungi of New Zealand. [Fungi of New Zealand, vol. 2.] Hong Kong: Fungal Diversity Press.
- Zambettakis C (1970) Recherches sur les *Ustilaginales* d'Afrique. *Bulletin de la Société Mycologique de France* **86**: 305–692.
- Zundel GL (1953) The *Ustilaginales* of the world. *Pennsylvania State College School of Agriculture Department of Botany Contribution* **176**: 1–410.
- Zwetko P, Blanz P (2004) *Die Brandpilze Österreichs* [Catalogus Florae Austriae vol. III (3).] Wien: Österreichische Akademie der Wissenschaften.