

## *Erratum*

# **Erratum to “Overexpression of PGC-1 $\alpha$ Increases Fatty Acid Oxidative Capacity of Human Skeletal Muscle Cells”**

**Nataša Nikolić,<sup>1</sup> Magdalena Rhedin,<sup>2</sup> Arild C. Rustan,<sup>1</sup> Len Storlien,<sup>3</sup>  
G. Hege Thoresen,<sup>1</sup> and Maria Strömstedt<sup>2</sup>**

<sup>1</sup> Department of Pharmaceutical Biosciences, School of Pharmacy, University of Oslo, P.O. Box 1068 Blindern, N-0316 Oslo, Norway

<sup>2</sup> AstraZeneca Research and Development, SE-43185 Mölndal, Sweden

<sup>3</sup> Boden Institute of Obesity, Nutrition and Exercise, University of Sydney, Sydney, NSW 2006, Australia

Correspondence should be addressed to G. Hege Thoresen; [hege.thoresen@farmasi.uio.no](mailto:hege.thoresen@farmasi.uio.no)

Received 3 March 2013; Accepted 7 March 2013

Copyright © 2013 Nataša Nikolić et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Unfortunately there was a mistake in Figure 7. The primer used for mRNA expression was actually MYH1, which regulates expression of MHC type IIx muscle fibers and not MHC type I as stated in the figure. However, we have repeated the experiment with the correct primer MYH7 (acc\_no. NM000257.2, F: CTCTGCACAGGGAAAATCTGAA, R: CCCCTGGAGACTTTGTCTCATT), and the new Figure 7 is shown here. This makes no difference to the conclusions of the paper. There was no significant change in mRNA of MHCI (MYH7), neither was the MHCI/MHCIIa mRNA ratio significantly increased. However, the last sentence in Section 3 (page 7) should be slightly changed: “Thus, the MHCI/MHCIIa mRNA ratio was approximately doubled in cells overexpressing PGC-1 $\alpha$  compared to control cells infected with empty vector (from 4.5 to 9.2, resp.)”




FIGURE 7