

RESEARCH ARTICLE

Mechanisms for type-II vitellogenesis-inhibiting hormone suppression of vitellogenin transcription in shrimp hepatopancreas: Crosstalk of GC/cGMP pathway with different MAPK-dependent cascades



Ting Chen^{1,2}, Chunhua Ren^{1,2}, Xiao Jiang¹, Lvping Zhang¹, Hongmei Li^{1,3}, Wen Huang^{1,2*}, Chaoqun Hu^{1,2*}

1 CAS Key Laboratory of Tropical Marine Bio-resources and Ecology (LMB); Guangdong Provincial Key Laboratory of Applied Marine Biology (LAMB), South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou, China, **2** South China Sea Bio-Resource Exploitation and Utilization Collaborative Innovation Center, Guangzhou, China, **3** University of the Chinese Academy of Sciences, Beijing, China

* hucq@scsio.ac.cn (CH); huangwen@scsio.ac.cn (WH)

OPEN ACCESS

Citation: Chen T, Ren C, Jiang X, Zhang L, Li H, Huang W, et al. (2018) Mechanisms for type-II vitellogenesis-inhibiting hormone suppression of vitellogenin transcription in shrimp hepatopancreas: Crosstalk of GC/cGMP pathway with different MAPK-dependent cascades. PLoS ONE 13(3): e0194459. <https://doi.org/10.1371/journal.pone.0194459>

Editor: Gao-Feng Qiu, Shanghai Ocean University, CHINA

Received: November 17, 2017

Accepted: March 2, 2018

Published: March 28, 2018

Copyright: © 2018 Chen et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by grants from the National Natural Science Foundation of China (31402287 to Dr. Ting Chen and 31602135 to Dr. Wen Huang), the Guangdong Province Program (2016A020210062 to Dr. Ting Chen, 2017B030314052 to Dr. Wen Huang,

Abstract

Vitellogenesis is the process of yolk formation via accumulating vitellin (Vn) with nutrients in the oocytes. Expression of vitellogenin (Vg), the precursor of Vn, is one of the indicators for the start of vitellogenesis. In Pacific white shrimp (*Litopenaeus vannamei*), the type-II vitellogenesis-inhibiting hormone (VIH-2) effectively suppresses hepatopancreatic *Vg* mRNA expression. In this study, we demonstrate the increasing transcript levels of hepatopancreatic *Vg* during *L. vannamei* ovarian development, suggesting that the hepatopancreas-derived *Vg*/*Vn* may also contribute to vitellogenesis in this species. Using a combination of *in vivo* injections and *in vitro* primary cell cultures, we provide evidences that the inhibition of VIH-2 on hepatopancreatic *Vg* gene expression is mediated through a functional coupling of the GC/cGMP pathway with different MAPK-dependent cascades in female shrimp. In VIH-2 signaling, the NO-independent GC/cGMP/PKG cascades were upstream of the MAPKs. Activations of the MAPK signal by VIH-2 include the phosphorylation of JNK and the mRNA/protein expression of P³⁸MAPK. Additionally, the cAMP/PKA pathway is another positive intracellular signal for hepatopancreatic *Vg* mRNA expression but is independent of its VIH-2 regulation. Our findings establish a model for the signal transduction mechanism of *Vg* regulation by VIH and shed light on the biological functions and signaling of the CHH family in crustaceans.

Introduction

Ovarian maturation in crustaceans is an energetically expensive reproductive process that can be divided into several phases, in which the last phase is referred to as vitellogenesis [1]. During

2017A030310429 to Dr. Xiao Jiang, 2013B091300020 to Prof. Chaoqun Hu, 2014B030301064 to Prof. Chaoqun Hu, 2015B020231007 to Prof. Chaoqun Hu and 2016A030310112 to Prof. Chaoqun Hu), the Science & Technology Promoting Projects for Oceanic & Fishery in Guangdong Province (A201601A03 to Prof. Chaoqun Hu), the National High Technology Research & Development Program of China (863 Program) (2012AA10A404-4 to Ms. Chunhua Ren), the Science and Technology Service Network Initiative of the Chinese Academy of Sciences (KJF-SW-STS-146 to Prof. Chaoqun Hu) and the National Share & Service Platform Construction Program for Aquaculture Germplasm and Resources (2017DKA30470 to Dr. Lvping Zhang). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

vitellogenesis, the period characterized by significant increase in oocyte diameter, the oocytes deposit and accumulate the major yolk protein vitellin (Vn), a glycolipoprotein having properties of sugar, fat and protein, to build up a nutrient storage [2–4]. The mature Vn is derived from its protein precursor vitellogenin (Vg) by a post-translated proteolysis [5]. Therefore, the Vg gene expression is an indicators for the start of the gonadal maturity in adult female crustaceans [6–8].

Generally, the expression of Vg in crustaceans is delicately regulated by an endogenous inhibitor, the vitellogenesis-inhibiting hormone (VIH), also known as the gonad-inhibiting hormone (GIH), which was first isolated from the eyestalk extracts of lobster *Homarus americanus* and was characterized by its prominent role as a negative regulator in reproduction [9, 10]. According to phylogenetic analysis, VIH is considered a member of the crustacean hyperglycemic hormone (CHH) family [11, 12]. In crustaceans, VIH is predominantly synthesized in the X-organ/sinus gland complex in eyestalks [9, 10, 13], with the exceptions that in some species, the brain is another major source of VIH [14–16]. Therefore, ablation of the eyestalk has been extensively practiced in commercial shrimp culture as a technique to accelerate ovarian maturation, based on the assumption of diminishing VIH production in the eyestalk [17–19]. Further studies have shown that adding the VIH by injection of purified [20, 21] or recombinant protein [15, 21–23] and removing the VIH by RNA interference (RNAi) [14, 24–26] or antisera [22] may induce and reduce Vg production, respectively, followed by the respective acceleration or deceleration of ovarian development.

Pacific white shrimp (*Litopenaeus vannamei*) is the most widely cultured and productive crustacean species in shrimp aquaculture globally [27]. In *L. vannamei*, several peptides belonging to the CHH family that identified from eyestalk sinus gland extracts have been found to have inhibitory effects on Vg transcript expression in cultured ovarian fragments and are designated sinus gland peptides (SGP)-A, -B, -C, -E, -F, and -G [20]. Subsequently, the cDNA sequence of SGP-G, the most abundant peptide of the SGPs previously extracted, was cloned and is considered as a candidate for the *L. vannamei* VIH [23]. *L. vannamei* SGP-G is phylogenetically clustered into the type-I CHH subfamily and is mostly produced in the eyestalk sinus gland with an expressed dynamics related to both molting and reproductive cycles [28]. After that, our group identified another candidate for the *L. vannamei* VIH from the shrimp eyestalk, which belonging to the type-II CHH subfamily and possessing vitellogenesis-inhibiting activity [15, 16]. Given that the first and second VIH candidates in *L. vannamei* belong to the type-I and type-II CHH subfamilies, respectively [15, 23, 29], they are logically designated the type-I VIH and the type-II VIH (VIH-2). As opposed to the type-I VIH (SGP-G), the newly cloned VIH-2 in *L. vannamei* is predominately generated in both the brain and the eyestalk [15, 16]. VIH-2 transcript levels have been found to increase significantly as fertilized eggs develop into juveniles, peak at sexual differentiation [16], under a negative estrogen regulation [30], and decrease in the brain after unilateral eyestalk ablation [15]. The recombinant VIH-2 protein is effective in inhibiting the hepatopancreatic Vg mRNA expression in female shrimp by both *in vitro* primary cell culture and *in vivo* injection approaches, and in reversing ovarian growth induced by eyestalk ablation [15]. On the other hand, VIH-2 RNAi can increase the ovarian Vg mRNA levels and the oocytes diameters [25].

The VIH receptor has not yet been identified, and the signal transduction mechanism for the VIH regulation on Vg is still not completely documented [11]. In silkworm *Bombyx mori*, several G protein-coupled receptors (GPCRs) have been described as putative receptors for ion transport peptide (ITP), another peptide member of the CHH family [31]. Subsequently, the first CHH receptor was found in crab *Eriocheir sinensis*, in which two potential phosphorylation sites are observed for cyclic adenosine monophosphate (cAMP)- and cyclic guanosine monophosphate (cGMP)-dependent protein kinase [32]. cGMP, but not cAMP, has been

shown to serve as a second messenger for CHHs in different crustacean species [33, 34]. Regarding molt-inhibiting hormone (MIH), another member of the type-II CHH, although there is consensus that cyclic nucleotides mediate their actions on ecdysteroidogenesis in the Y-organs, the relative importance of cAMP and cGMP differs among different crustacean species [35–39]. In the incubated ovarian fragments of kuruma prawn *Marsupenaeus japonicus*, activation of cGMP, cAMP or Ca^{2+} signaling may decrease the Vg mRNA levels [40]. Although both the type-I [23] and type-II [15] VIHs inhibit Vg gene expression and ovarian development in *L. vannamei*, only the second messengers for the type-I VIH (SGP-G) but not the type-II VIH (VIH-2) signaling have been investigated [23]. In this case, increasing intracellular cGMP and Ca^{2+} may inhibit Vg mRNA expression, while SGP-G treatment increases cGMP but not cAMP in *L. vannamei* ovaries [41]. In contrast, no research has been developed to date regarding the *L. vannamei* VIH-2 signaling pathway.

In crustaceans, the process of vitellogenesis includes endogenous vitellogenesis and exogenous vitellogenesis [42–44]. During exogenous vitellogenesis, the hepatopancreas is the major site for Vg synthesis in some crustacean species [5, 45, 46]. Compared to ovarian Vg, Vg expression in the hepatopancreas is poorly characterized in crustaceans, despite its being a major Vg synthesis site in *L. vannamei* [47]. Our group previously established an isolation and *in vitro* culture system for shrimp primary hepatopancreatic cells [15]. The endocrinological factors that successfully regulate Vg mRNA levels in the hepatopancreatic primary cells e.g. VIH-2 [15] and MIH-2 [29] are effective for controlling ovarian maturation in *in vivo* experiments. In this study, we first characterized the change of Vg mRNA levels in *L. vannamei* hepatopancreas during ovarian development. To investigate the signal transduction mechanisms for VIH-2 inhibition on Vg transcript expression in the shrimp hepatopancreas, we detected the involvement of cGMP and cAMP pathways, and we examined the participation of mitogen-activated protein kinase (MAPK)-dependent cascades. We further described the crosstalk of the cGMP-/dependent and MAPK-dependent pathways to develop a regulatory model for VIH-2 signaling in Vg gene expression in the shrimp hepatopancreas.

Materials and methods

Ethics statement

Shrimp experiments were carried out according to the Care and Use of Agricultural Animals in Agricultural Research and Teaching and approved by the Science and Technology Bureau of China. All animal experiments were conducted in accordance with the guidelines and approval of the Ethics Committees of the South China Sea Institute of Oceanology, Chinese Academy of Sciences.

Animals and test substances

Pacific white shrimp (*L. vannamei*), with a body weight of 48.4 ± 1.6 g and a body length of 18.3 ± 0.3 cm, were collected from the Jinyang shrimp culture center, Maoming, China, and maintained in seawater at 28°C under a 12D:12L photoperiod. Previtellogenic female shrimp, with the gonadosomatic index (GSI) of $0.62 \pm 0.10\%$ were selected for *in vivo* injection and *in vitro* cell cultural experiments. Additionally, the Vg mRNA levels were detected in the hepatopancreas of female shrimp in different ovarian developmental stages, including stage O (immature, GSI of $<0.3\%$), stage I (previtellogenesis, GSI of 0.4–0.8%), stage II (primary vitellogenesis, GSI of 0.9–2.4%), stage III (secondary vitellogenesis, GSI of 2.7–4.5%) and stage IV (maturation, GSI of $>5.2\%$), were determined by using the paraffin sections stained with hematoxylin/eosin (H/E) staining as previously described before [48, 49]. For comparison, the

Vg mRNA level was also examined in hepatopancreas of sexually immature male shrimp (GSI of <0.3%).

The *L. vannamei* VIH-2 recombinant proteins were expressed in *Escherichia coli* and purified by using immobilized metal ion affinity chromatography (IMAC) as previously described [15]. For pharmacological study, NOC-18, A-350619, 8-Br-cGMP, L-NAME, Zn(II)PPIX, Rp-8-Br-PET-cGMPS, SB02190, SP600125 and BI-87G3 were purchased from Sigma, while IBMX, forskolin, 8-Br-cAMP, PD169316, PD98059 and U0126 were acquired from Calbiochem. Test substances were prepared as 1 mM or 10 mM frozen stocks in distilled water or DMSO in small aliquots and diluted with a pre-warmed culture medium to appropriate concentrations 15 min prior to drug treatments.

VIH injection in shrimp

VIH-2 recombinant protein was diluted in PBS (33‰ salinity). Only the previtellogenic female shrimp were used for VIH injection experiments. Each shrimp was injected with 100 µL of VIH-2 in PBS, and PBS injection alone was used as a negative control. Hepatopancreatic samples ($n = 6$) from each group were randomly sampled at 0–12 h and 0–24 h post injection to measure second messenger molecules, and mRNA and protein samples, respectively.

Isolation, primary culture and static incubation of shrimp hepatopancreatic cells

The *L. vannamei* hepatopancreatic cells were prepared as previously described [15]. Briefly, hepatopancreas from previtellogenic female shrimp (GSI = 0.62±0.10%, $n = 10$) were excised and washed three times in pre-chilled $\text{Ca}^{2+}/\text{Mg}^{2+}$ -free Hanks balanced salt solution (HBSS, Sigma; containing 4 mM NaHCO₃, 9 mM HEPES, 154 mM NaCl, 100 U/mL penicillin and 100 µg/mL streptomycin, pH 7.63). Hepatopancreatic fragments were diced into 0.5-mm thickness, incubated in $\text{Ca}^{2+}/\text{Mg}^{2+}$ -free HBSS with EDTA (1 mM) at room temperature for 5 min, and digested with collagenase type IV (1 mg/mL, Invitrogen) and DNase II (0.01 mg/mL, Sigma) in $\text{Ca}^{2+}/\text{Mg}^{2+}$ -free HBSS at 28 °C for 30 min. Next, shrimp hepatopancreatic fragments were mechanically dispersed into single cells by gentle pipetting. Dispersed cells were then separated from the remaining fragments by filtration through a sterile mesh (30 µm pore size) and harvested by centrifugation at 100×g for 5 min at 4 °C. The cells obtained were resuspended in a Dulbecco-modified Eagle medium (DMEM)/F-12 (1:1, Gibco BRL; containing 14 mM NaHCO₃, 2 g/L bovine serum albumin, 154 mM NaCl, 100 U/mL penicillin and 100 µg/mL streptomycin, pH 7.63). Only the cell preparations with greater than 95% viability that assessed by using a trypan blue exclusion assay were used in subsequent experiments. Dispersed hepatopancreatic cells were diluted in DMEM/F-12 to a final density of 0.4×10^6 cells/mL and cultured in polyethylenimine (PEI, 5 µg/mL, Sigma) pre-coated 24-well plates with 5% CO₂ and saturated humidity at 28 °C overnight for recovery.

On the second day after cell preparation, test substances prepared in DMEM/F12 medium were gently overlaid onto hepatopancreatic cells after the removal of the old culture medium. In pharmacological studies, the cells were incubated with test substances for another 12 h as previously validated [15].

Measurement of target gene mRNA levels by real-time PCR

For measurement of *L. vannamei* Vg (AY321153), *p38 mitogen-activated protein kinases* (*P³⁸MAPK*, JX990130), *extracellular regulated protein kinases* (ERK, JN035901) and *c-Jun N-terminal kinase* (JNK, JN035903) gene expressions in the hepatopancreas and in the hepatopancreatic primary cells, total RNA was isolated by using the TRIzol reagent (Invitrogen),

digested with DNase I (Invitrogen), and reversely transcribed by using a PrimeScript RT kit (TaKaRa). Transcriptional expression of target genes were detected using SYBR Premix Ex Taq II (TaKaRa) in the RotorGene RG-3000 real-time PCR System (Qiagen) with primers and PCR conditions as shown in [S1 Table](#). Serially diluted plasmid DNAs containing the ORF sequences for the target genes were used as the standards for the real-time PCRs. After reactions, the identities of the PCR products were routinely confirmed by analysis of the melting curve. In this study, real-time PCR of *L. vannamei* β -actin (JF288784) was used as an internal control because no significant changes were noted for β -actin mRNA in our studies. The raw data of the target gene transcripts were expressed in terms of fmol detected per tube, and the data were routinely normalized as the ratio of β -actin mRNA detected in the same sample.

Measurement of NO, cGMP and cAMP production

VIH-2 effects on NO, cGMP and cAMP production were tested on hepatopancreatic tissues and primary cells from previtellogenic female *L. vannamei*. The hepatopancreatic tissues were sampled and frozen in liquid nitrogen immediately for further experiments. The hepatopancreatic cells were cultured at DMEM/F12 medium with a density of 1×10^6 cells/mL in PEI-coated 6-well plates. To measure NO, after an overnight recovery, the old medium was replaced with a fresh medium containing the VIH-2 protein. NO production was estimated by spectrophotometrically measuring the levels of nitrite as a proxy of NO by using a Griess reagent system (Promega) as we previously described [50]. For measurement of GMP and cAMP, after an overnight recovery, the old medium was replaced with 0.9 mL fresh medium supplemented with IBMX (0.1 mM) and incubated at 28 °C for 15 min, and then 0.1 mL 10×stock solutions of VIH-2 was added. The cGMP and cAMP content were quantified using the Cyclic-GMP XP® Assay Kit (Cell Signaling) and the Cyclic-AMP XP® Assay Kit (Cell Signaling), respectively, as previously described [51].

Western blot of tissue and primary cell lysates

Western blots were performed on the samples of previtellogenic female *L. vannamei* hepatopancreatic tissues and primary cells. For tissue samples, the shrimp hepatopancreas were collected and frozen in liquid nitrogen immediately, then dissolved in RIPA Lysis Buffer I (Sangon Biotech) with complete protease and phosphatase inhibitor cocktails (Roche) by extraction with a 23-gauge needle attached to a 1-mL syringe on ice. For cell samples, the shrimp hepatopancreatic cells were seeded in PEI-coated 6-well plates at a density of 1×10^6 cells/mL/well. After incubating overnight, the culture medium was replaced with new medium containing the appropriate concentrations of test substances and cultured at 28 °C. The VIH incubation lasted from 0 to 24 h, and for other pharmacological studies, the treatment time was fixed at 1 h. Then, the cells were rinsed with PBS (pH 7.4) and cell lysate were prepared in RIPA Lysis Buffer IV (Sangon Biotech) with complete protease and phosphatase inhibitor cocktails by freezing and thawing for three times. The tissue and cell lysates were resolved on a 15% gel by using SDS-PAGE and transblotted onto a PVDF membrane (Roche) at 20 V for 1 h with a Trans-Blot SD Electrophoretic Cell (Bio-Rad). The membrane was then blocked using 3% BSA in TBST (Sangon Biotech) and incubated overnight at 4 °C with antibodies for Phospho-P³⁸MAPK (1:500, Cat#9211, Cell Signaling), total P³⁸MAPK (1:500, Cat#8690, Cell Signaling), Phospho-ERK (1:500, Cat#9102, Cell Signaling), total ERK (1:1000, Cat#9101, Cell Signaling), Phospho-JNK (1:1000, Cat#9251, Cell Signaling), total JNK (1:2000, Cat#9252, Cell Signaling) and β -actin (1:2000, Cat#D190606, Sangon Biotech). On the following day, HRP-conjugated goat anti-rabbit (1:2000, Cat#DC03L, Calbiochem) or rabbit anti-mouse IgG (1:2000, Cat#AP160P, Calbiochem) was added and Immobilon Western (Pierce) was used as an HRP substrate for signal development.

Data transformation and statistical analysis

The data expressed as the mean \pm S.E. and were analyzed by using Student's *t*-test or one-way ANOVA followed by Fisher's least significant difference (LSD) test with SPSS (IBM Software).

Results

Expression of hepatopancreatic Vg transcript during shrimp ovary development

Vg transcript expressions were detected in the hepatopancreas from female shrimp with different ovarian maturation stages (stages I-IV), and the expressions in the hepatopancreas from sexually immature female (stage O) and male shrimp were used as the controls. The results showed that the hepatopancreatic Vg mRNA in sexually immature female shrimp were higher than in sexual immature male shrimp (Fig 1A). The hepatopancreatic Vg mRNA expression of female shrimp increased acutely and continuously during ovarian maturation (stages I-III) after eyestalk ablation (Fig 1B). The peak of Vg mRNA expression occurred at the stage III, followed by a rapid decrease at stage IV (Fig 1B).

Involvement of cGMP and cAMP pathways in VIH-2-regulated Vg gene expression

The effects of VIH-2 on nitric oxide (NO), cGMP and cAMP production were measured in both the *in vivo* and *in vitro* experiments. By using the *in vivo* injection approaches, VIH-2




Fig 1. Vg mRNA expression in the hepatopancreas of *L. vannamei* during ovarian development. A: Hepatopancreatic Vg transcript levels of sexually immature male shrimp and female shrimp. The data presented are expressed as the mean \pm S.E. (n = 6), and significant differences between male and female shrimp were assessed using Student's *t*-test (* P<0.05, ** P<0.01 and *** P<0.001). B: Hepatopancreatic Vg transcript levels of female shrimp in different ovarian developmental stages. The data presented are expressed as the mean \pm S.E. (n = 6), and experimental groups denoted by the same letter represent a similar level (P>0.05, ANOVA followed by Fisher's LSD test). C: Histological observation of ovarian developmental stages based on H/E staining. The selected ovarian maturation stages of female shrimp include: stage I (previtellogenesis), stage II (primary vitellogenesis), stage III (secondary vitellogenesis), and stage IV (maturation).

<https://doi.org/10.1371/journal.pone.0194459.g001>

showed the ability to elevate the concentrations of cGMP (Fig 2B) but not NO (Fig 2A) or cAMP (Fig 2C) in the shrimp hepatopancreas. In the *in vitro* experiments that performed on the hepatopancreatic primary cells, VIH-2 administration was capable of inducing cGMP production within 1.5–6 h, with a peak of 3 h (Fig 2E). Meanwhile, this peptide was not able to change the production of NO (Fig 2D) or cAMP (Fig 2F). In a parallel experiment incubated with an increased concentration of substrates, VIH-2 showed a dose-dependent effect on cGMP production (Fig 2H), but no effects on NO (Fig 2G) or cAMP (Fig 2I). As positive controls, an NO donor (NOC-18), a soluble guanylyl cyclase (GC) activator (A-350619) and an adenylate cyclase (AC) activator (forskolin) effectively stimulated NO (Fig 2G), cGMP (Fig 2H) and cAMP (Fig 2I) production, respectively, in the hepatopancreatic primary cells.

The roles of the cGMP signaling pathway in VIH-2-regulated Vg gene expression in shrimp hepatopancreas were further investigated using pharmacological approaches. In this case, incubation of the NO donor (NOC-18), the GC activator (A-350619) or the membrane-permeable cGMP analog (cpt-cGMP) consistently reduced Vg mRNA expression in dose-dependent ways ([Fig 3A–3C](#)). On the other hand, when co-incubated with the inhibitors of GC [Protoporphyrin IX zinc(II)] or protein kinase G (PKG, Rp-8-Br-PET-cGMPS) but not with the non-specific inhibitor of NOS (L-NAME), the inhibitory effect of VIH-2 on Vg mRNA expression was abolished ([Fig 3D–3F](#)). Combined with the results above, VIH-2 may downregulate Vg gene expression in shrimp hepatopancreas via the cGMP/PKG signal pathway without NO participation.

The functions of cAMP signaling in hepatopancreatic *Vg* gene expression were also developed. Interestingly, incubation of both the AC activator (forskolin) and the membrane-permeable cAMP analog (8-Br-cAMP) increased the mRNA levels of *Vg* in dose-dependent ways (Fig 4A & 4B). Additionally, co-incubation of the AC activator (forskolin) or the cAMP analog




Fig 2. Effects of VIH-2 administration on NO, cGMP and cAMP production in the hepatopancreas of *L. vannamei*. A-C: The NO (A), cGMP (B) and cAMP (C) production in the shrimp hepatopancreas after VIH-2 (300 ng/g bwt) injection. D-F: The time-dependent NO (D), cGMP (E) and cAMP (F) productions by VIH-2 treatment in shrimp hepatopancreatic cell cultures. G-I: Dose-dependent NO (G), cGMP (H) and cAMP (I) productions by VIH-2 treatment in shrimp hepatopancreatic cell cultures. For *in vitro* experiments, hepatopancreatic cells were pretreated with IBMX (0.1 mM) at ~20 min before the substrates were added. In this study, the data presented are expressed as the mean \pm S.E. ($n = 6$ and 4 for *in vivo* and *in vitro* experiments, respectively). Experimental groups denoted by the same letter represent a similar level ($P>0.05$, ANOVA followed by Fisher's LSD test).

<https://doi.org/10.1371/journal.pone.0194459.g002>




Fig 3. Involvement of cGMP-dependent signaling in VIH-2 suppressed Vg mRNA expression in the hepatopancreas of *L. vannamei*. A-C: The effects of the NO donor (A), the GC activator (B) and the cGMP analog (C) on Vg mRNA expression in shrimp hepatopancreatic cell cultures. The hepatopancreatic cells were incubated for 12 h with increasing doses of NOC-18 (0.01–100 μM), A-350619 (0.01–100 μM) or 8-Br-cGMP (0.1–1000 nM). D-F: Effects of NOS inhibitor (D), GC inhibitor (E) or PKG inhibitor (F) on VIH-2-inhibited Vg mRNA expression in shrimp hepatopancreatic cell cultures. The hepatopancreatic cells were incubated for 12 h with VIH-2 (1 μM) in the presence or absence of L-NAME (300 μM), Zn(II)PPIX (5 μM) or Rp-8-Br-PET-cGMPS (5 μM). In this study, the data presented are expressed as the mean±S.E. (n = 4). Experimental groups denoted by the same letter represent a similar level ($P>0.05$, ANOVA followed by Fisher's LSD test).

<https://doi.org/10.1371/journal.pone.0194459.g003>

(8-Br-cAMP) reversed the inhibitory effects of VIH-2 on Vg gene expression (Fig 4C). In sum, the AC/cAMP cascade is a positive signaling for Vg gene expression in the shrimp hepatopancreas. However, this signal pathway is independent of the VIH-2 regulation on Vg expression.

Involvement of MAPK (P38/ERK/JNK)-dependent cascades in VIH-2 regulated-Vg expression

The involvements of MAPK signal pathways including P³⁸MAPK, ERK and JNK in Vg regulation by VIH-2 in shrimp hepatopancreas were evaluated in this study. In both *in vivo* injection




Fig 4. Involvement of cAMP-dependent signaling in VIH-2 suppressed Vg mRNA expression in the hepatopancreas of *L. vannamei*. A & B: The effects of the AC activator (A) and the cAMP analog (B) on Vg mRNA expressions in shrimp hepatopancreatic cell cultures. The hepatopancreatic cells were incubated for 12 h with increasing doses of forskolin (0.01–1000 nM) or 8-Br-cAMP (0.1–1000 nM). C: Interaction of cAMP-dependent signal pathways and VIH-2 on Vg mRNA expression in shrimp hepatopancreatic cell cultures. The hepatopancreatic cells were incubated for 12 h with VIH-2 (1 μM) in the presence or absence of forskolin (1 μM) or 8-Br-cAMP (1 μM). In this study, the data presented are expressed as the mean±S.E. (n = 4). Experimental groups denoted by the same letter represent a similar level ($P>0.05$, ANOVA followed by Fisher's LSD test).

<https://doi.org/10.1371/journal.pone.0194459.g004>

and *in vitro* primary cell incubation experiments, VIH-2 was effective at triggering JNK phosphorylation in a time-dependent manner without affecting the total JNK content (Fig 5A & 5B). In contrast, the enhanced P³⁸MAPK phosphorylation by VIH-2 was based on the increase of P³⁸MAPK content (Fig 5A & 5B). Furthermore, VIH-2 treatment changed neither ERK's phosphorylation nor content in both *in vivo* and *in vitro* experiments (Fig 5A & 5B).




Fig 5. Effects of VIH-2 administration on the phosphorylation and mRNA expression of three MAPKs in the hepatopancreas of *L. vannamei*. A: Effects of VIH-2 injection on the phosphorylation of P³⁸MAPK, ERK and JNK in shrimp hepatopancreas. B: Effects of VIH-2 treatment on the phosphorylation of P³⁸MAPK, ERK and JNK in shrimp hepatopancreatic cell cultures. C-E: Effects of VIH-2 injection on the P³⁸MAPK (C), ERK (D) and JNK (E) mRNA expressions in the shrimp hepatopancreas. F-H: Effects of VIH-2 treatment on the P³⁸MAPK (F), ERK (G) and JNK (H) mRNA expression in shrimp hepatopancreatic cell cultures. For *in vivo* experiments, the hepatopancreas were sampled at 0, 1.5, 3, 6, 12 and 24 h after injection of VIH-2 (300 ng/g bwt). For *in vitro* experiments, the hepatopancreatic cells were treated with VIH-2 (1000 nM) for 0, 0.5, 1, 1.5, 3, 6, 12 and 24 h. In this study, the data presented are expressed as the mean±S.E. (n = 6 and 4 for *in vivo* and *in vitro* experiments, respectively), and significant differences between treated and untreated groups were assessed using Student's t-test (* $P<0.05$, ** $P<0.01$ and *** $P<0.001$).

<https://doi.org/10.1371/journal.pone.0194459.g005>




Fig 6. Involvement of MAPK-dependent signaling in VIH-2 suppressed Vg mRNA expression in the hepatopancreas of *L. vannamei*. A-C: Effects of the P³⁸MAPK inhibitor (A), the ERK inhibitor (B) or the JNK inhibitor (C) on VIH-2-inhibited Vg mRNA expression in shrimp hepatopancreatic cell cultures. The hepatopancreatic cells were incubated for 12 h or 36 h with VIH-2 (1 μM) in the presence or absence of PD169316 (100 nM), or SB02190 (100 nM), PD98059 (10 μM) or U0126 (1 μM), and SP600125 (10 μM) or BI-87G3 (10 μM). D: Interaction of the P³⁸MAPK inhibitor and the JNK inhibitor on VIH-2 inhibited Vg mRNA expression in shrimp hepatopancreatic cell cultures. The hepatopancreatic cells were incubated for 12 h or 36 h with VIH-2 (1 μM) in the presence or absence of PD169316 (100 nM) or SP600125 (10 μM) alone, or PD169316 (100 nM) and SP600125 (10 μM) together. In this study, the data presented are expressed as the mean±S.E. (n = 4). Experimental groups denoted by the same letter represent a similar level ($P>0.05$, ANOVA followed by Fisher's LSD test).

<https://doi.org/10.1371/journal.pone.0194459.g006>

In parallel experiments, the mRNAs of shrimp *P³⁸MAPK*, *ERK* and *JNK* were measured after VIH-2 treatment. In this case, VIH-2 increased the expression only of *P³⁸MAPK* mRNA in both *in vivo* (Fig 5C) and *in vitro* (Fig 5F) experiments; the expressions of *ERK* (Fig 5D & 5G) and *JNK* (Fig 5E & 5H) transcripts were not affected. Therefore, it is logically to speculate that VIH-2 may enhance *P³⁸MAPK* phosphorylation by increasing the mRNA expression followed by protein production. However, *JNK* phosphorylation triggered by VIH-2 is independent of the regulation at the transcript level.

Further pharmacological experiments were performed on the hepatopancreatic primary cells at short and long time points, namely, 12 h and 36 h, respectively, to test the effect of three MAPK cascades blockage on the VIH-2-inhibited Vg gene expression. In this case, co-incubation of the *P³⁸MAPK* inhibitors (PD169316 and SB02190) partially reversed the inhibitory action of VIH-2 on Vg gene expression, with a more potent effects at 36 h than at 12 h (Fig 6A); while the JNK inhibitors (SP600125 and BI-87G3) partially reversed the VIH-2 action but were more potent at 12 h than at 36 h (Fig 6C). In contrast, the ERK inhibitors (PD98059 and U0126) did not change the Vg mRNA levels when VIH-2 was added (Fig 6B). In addition, when the *P³⁸MAPK* inhibitor (PD169316) and the JNK inhibitor (SP600125) were added together, the inhibitory role of VIH-2 on Vg gene expression was completely blocked at both 12 h and 36 h time (Fig 6D), suggesting that the inhibitory effect of VIH-2 on Vg gene expression was mediated by a combination of the *P³⁸MAPK* and *JNK* cascades.




Fig 7. Crosstalk of the GC/cGMP pathway with MAPK-dependent cascades in VIH-2 suppressed Vg mRNA expression in the hepatopancreas of *L. vannamei*. A: Effects of the GC activator and the cGMP analog on the JNK phosphorylation in shrimp hepatopancreatic cell cultures. The hepatopancreatic cells were incubated for 1 h with increasing doses of A-350619 (0.01–100 μM) or 8-Br-cGMP (0.1–1000 nM). B: Effects of the GC inhibitor and the PKG inhibitor on the VIH-2-induced JNK phosphorylation in shrimp hepatopancreatic cell cultures. The hepatopancreatic cells were incubated for 1 h with VIH-2 (1 μM) in the presence or absence of Zn(II)PPIX (5 μM) or Rp-8-Br-PET-cGMPS (5 μM). C & D: Effects of GC activator (C) and cGMP analog (D) on the *P³⁸MAPK* mRNA expression in shrimp hepatopancreatic cell cultures. The hepatopancreatic cells were incubated for 3 h with increasing doses of A-350619 (0.01–100 μM) or 8-Br-cGMP (0.1–1000 nM). E & F: Effects of the GC inhibitor and the PKG inhibitor on the VIH-2 induced *P³⁸MAPK* mRNA expression in shrimp hepatopancreatic cell cultures. The hepatopancreatic cells were incubated for 3 h with VIH-2 (1 μM) in the presence or absence of Zn(II)PPIX (5 μM) or Rp-8-Br-PET-cGMPS (5 μM). G-I: Effects of the *P³⁸MAPK* inhibitor (G), the ERK inhibitor (H) and the JNK inhibitor (I) on the VIH-2-induced cGMP productions in shrimp hepatopancreatic cell cultures. The hepatopancreatic cells were incubated for 3 h with VIH-2 (1 μM) in the presence or absence of PD169316 (100 nM), or SB2190 (100 nM), PD98059 (10 μM) or U0126 (1 μM), and SP600125 (10 μM) or BI-87G3 (10 μM). In this study, the data presented are expressed as the mean±S.E. (n = 4). Experimental groups denoted by the same letter represent a similar level ($P>0.05$, ANOVA followed by Fisher's LSD test).

<https://doi.org/10.1371/journal.pone.0194459.g007>

Crosstalk of the GC/cGMP/PKG and MAPK (P38/JNK) pathways in VIH-2 signaling

The relationship between the GC/cGMP/PKG and MAPK cascades in the signaling of VIH-2 regulation on *Vg* gene expression was determined further using pharmacological approaches. In the hepatopancreatic primary cells, incubation of the GC activator (A-350619) and the cGMP analog (cpt-cGMP) mimicked VIH-2's effect on JNK phosphorylation (Fig 7A). On the other hand, when the GC and PKG activities were blocked by GC inhibitor [Protoporphyrin IX zinc(II)] and PKG inhibitor (Rp-8-Br-PET-cGMPS), respectively, the phosphorylation triggered by VIH-2 treatment might be eliminated (Fig 7B).

The mediated role of GC/cGMP/PKG on *P³⁸MAPK* mRNA expression was similar to that on JNK phosphorylation. Both the GC activator (A-350619) and the cGMP analog (cpt-cGMP) could dose-dependently stimulate *P³⁸MAPK* transcript expression (Fig 7C & 7D). In parallel experiment, the VIH-2-induced *P³⁸MAPK* gene expression could be blocked by co-treatment of the GC inhibitor [Protoporphyrin IX zinc(II), Fig 7E] and PKG inhibitor (Rp-8-Br-PET-cGMPS, Fig 7F).

In contrast, co-incubation of all inhibitors of MAPK cascades including PD169316 and SB202190 for P³⁸MAPK ([Fig 7G](#)), PD98059 and U0126 for ERK ([Fig 7H](#)), and SP600125 and BI-87G3 for JNK ([Fig 7I](#)) were not able to block the stimulation of VIH-2 on cGMP production in the hepatopancreatic primary cells. The sum of these results implies that in the VIH-2 signaling, the GC/cGMP/PKG cascades are upstream and the MAPK cascades are downstream.

Discussion

During the ovarian development of *L. vannamei*, a significant increased trend of hepatopancreatic Vg mRNA levels was observed ([Fig 1B](#)), indicating that the Vg/Vn from the hepatopancreas may also participate to the process of vitellogenesis in this crustacean species. Similarly, high hepatopancreatic Vg mRNA levels were observed at the early and late exogenous vitellogenic stages in *M. japonicas* [44]. A recent study pointed out the importance of endogenous Vg/Vn in *L. vannamei* ovarian development [52], and our current study showed that exogenous Vg/Vn play the same role in this process. Our study also provided new evidence that hepatopancreatic primary cell culture is an applicable method for testing the regulators of Vg/Vn synthesis in *L. vannamei* [15, 29], in addition to the previously reported usage of cultured ovarian fragments [23, 40, 41].

Our study found that administration of VIH-2 in both *in vivo* and *in vitro* experiments might increase the content of cGMP but not NO in the hepatopancreas of *L. vannamei*. ([Fig 2A, 2B, 2D, 2E, 2G & 2H](#)). However, the Vg mRNA were decreased with incubations of the NO donor, the GC activator and the cGMP analog in hepatopancreatic cell cultures ([Fig 3A–3C](#)). Meanwhile, only the inhibitors for downstream GC or PKG, but not the inhibitor for upstream NO, reversed the suppression of Vg mRNA expression by VIH-2 treatment ([Fig 3D–3F](#)). cGMP is an intracellular second messenger that converted from guanosine triphosphate (GTP) by GC, which includes the membrane-bound and soluble forms [53]. In land crab *Gecarcinus lateralis*, three GC genes namely, an NO-sensitive GC, a membrane receptor GC and an NO-insensitive soluble GC have been distinguished [54], and the counterparts for each GC gene in *L. vannamei* were found in a transcriptome we previously reported [55]. Combined with the results we presented, it could be logically speculated that VIH-2 induces cGMP production in the shrimp hepatopancreas via a membrane NO-insensitive but not an NO-sensitive GC. Subsequently, the downstream PKG is activated to transduce the signaling on control of Vg gene expression.

Our study also showed that VIH-2 treatment could not increase cAMP content in the hepatopancreas of *L. vannamei* by either *in vivo* or *in vitro* approaches ([Fig 2C, 2F & 2I](#)). In the incubated *L. vannamei* ovarian fragments, type-I VIH (SGP-G) administration might stimulate the production of both cAMP and cGMP when the phosphodiesterase inhibitor was added [41]. In the hindgut of green shore crab *Carcinus maenas*, studies showed that the CHH actions were mediated by second messengers, including cAMP and cGMP [33]. In *M. japonicus*, cGMP rather than cAMP was found to be the second messenger for CHH family hormones [34]. For MIH, studies of several crustacean species concluded that cGMP is the predominant signaling molecule responsible for the roles of this peptide in the regulation of ecdysteroidogenesis, while cAMP also regulates acute ecdysteroidogenesis to some extent [38]. Given that in *L. vannamei*, type-I VIH (SGP-G) and CHH belong to the type-I CHH subfamily, while type-II VIH and MIH belong to the type-II CHH subfamily, we speculate that the intracellular signaling of type-II VIH (VIH-2) is more like that of MIH. In addition, incubation of the AC activator or the cAMP analog might increase Vg transcript levels in the hepatopancreatic cell culture ([Fig 4A & 4B](#)). However, in the *M. japonicus* ovarian fragment culture, the




Fig 8. Working model for the signal transduction mechanism of VIH-2 suppressed Vg mRNA expression in the shrimp hepatopancreas. In this case, a membrane receptor GC activation by VIH-2 can generate cGMP and subsequently activate PKG. The downstream MAPK-dependent cascades are then activated by cooperation of JNK phosphorylation and P³⁸MAPK content increase, to inhibit the gene expression of Vg in the hepatopancreas of *L. vannamei*.

<https://doi.org/10.1371/journal.pone.0194459.g008>

effects of the high dosage-cAMP analog were found to inhibit Vg gene expression [40], indicating that the regulatory mechanisms for Vg may be different between the shrimp hepatopancreas and ovary. Moreover, activation of the cAMP/protein kinase A (PKA) signal pathway could partially reverse the inhibition of Vg gene expression by VIH-2 treatment (Fig 4C), implying that these two signaling are independent but they interacted in regulation of Vg expression in the shrimp hepatopancreas.

We further tested the roles of three major MAPK-dependent cascades (P³⁸MAPK, ERK and JNK) in VIH-2-regulated Vg gene expression in the shrimp hepatopancreas. Unexpectedly, the *L. vannamei* VIH-2 inhibits Vg transcript expression in two ways, which include triggering JNK phosphorylation and increasing P³⁸MAPK content, while the expected ERK-dependent pathway is not involved (Figs 5 & 6). Previous studies have shown that the P³⁸MAPK- [56], ERK- [57], JNK-dependent cascades [58, 59] may be activated (including phosphorylation and/or gene expression) during infection of pathogens. It would be logical to speculate that in shrimp, a complicated signaling complex may be activated in response to pathogenic infection

[60, 61], which would include the MAPK-dependent cascades, with the side effect of inhibiting vitellogenesis, to save energy and nutrition for against the invasive pathogens.

Meanwhile, we analyzed the crosstalk of GC/cGMP and MAPK-dependent cascades in VIH-2 inhibition on Vg gene expression. In this case, the VIH-2-induced activation of JNK and P³⁸MAPK could be eliminated by inhibition of GC or PKG (Fig 7B, 7E & 7F), but not vice versa (Fig 7G–7I). Taken together, these results show that in VIH-2 signaling, the GC/cGMP/PKG signal sits upstream of different MAPK-dependent cascades, while the JNK and P³⁸MAPK pathways have parallel participations in the regulation and play a major role in short-term and long-term regulation, respectively (Fig 8). In different Decapoda crustacean species, other signaling cascades, such as Ca²⁺/PKC [40] and PI3K/Akt/mTOR [62], have been reported to be involved in the regulation of Vg gene expression. Therefore, the involvement of more signaling pathways, as well as their roles in signaling interactions, in VIH-2-regulated Vg gene expression should be examined in the further studies.

In summary, by using a combination of *in vivo* and *in vitro* experiments, we provide evidences for a comprehensive model of *L. vannamei* VIH-2-reduced Vg gene expression in the shrimp hepatopancreas. This inhibitory effect is mediated through a functional coupling of the GC/cGMP pathway with different MAPK-dependent cascades. The process of vitellogenesis in crustaceans is delicately regulated by multiple peptide or non-peptide endocrinial factors [11, 20, 29, 50, 62], and among them, VIH is the most effective one [15, 23–25]. Combined with previous studies [40, 41, 62], our study reveals that production of Vg, the main protein component of vitellogenesis, is mediated by VIH via different signaling in different target tissues and/or species. Our findings also shed light on the biological functions and signal transduction mechanisms of the CHH family hormones in crustaceans.

Supporting information

S1 Table. Primer sequences and real-time PCR conditions used in this study.
(DOCX)

Acknowledgments

This work was supported by grants the National Natural Science Foundation of China (31402287 and 31602135), the Guangdong Province Program (2016A020210062, 2017B030314052, 2017A030310429, 2013B091300020, 2014B030301064, 2015B020231007 and 2016A030310112), the Science & Technology Promoting Projects for Oceanic & Fishery in Guangdong Province (A201601A03), the National High Technology Research & Development Program of China (863 Program) (2012AA10A404-4), the Science and Technology Service Network Initiative of the Chinese Academy of Sciences (KJF-SW-STS-146) and the National Share & Service Platform Construction Program for Aquaculture Germplasm and Resources (2017DKA30470). We also thank Prof. M. N. Wilder (the University of Tokyo) for her constructive suggestion about the naming of two types of VIH in *L. vannamei*.

Author Contributions

Conceptualization: Ting Chen, Chaoqun Hu.

Data curation: Ting Chen, Chunhua Ren, Xiao Jiang.

Formal analysis: Lvping Zhang, Hongmei Li.

Funding acquisition: Ting Chen, Wen Huang, Chaoqun Hu.

Investigation: Ting Chen, Chunhua Ren, Xiao Jiang, Hongmei Li.

Methodology: Ting Chen, Chunhua Ren, Xiao Jiang, Lvping Zhang, Wen Huang.

Project administration: Wen Huang.

Resources: Ting Chen, Lvping Zhang, Wen Huang, Chaoqun Hu.

Software: Ting Chen.

Supervision: Chaoqun Hu.

Validation: Ting Chen, Chunhua Ren, Hongmei Li, Wen Huang.

Visualization: Ting Chen, Chunhua Ren, Wen Huang.

Writing – original draft: Ting Chen, Wen Huang.

Writing – review & editing: Ting Chen, Wen Huang, Chaoqun Hu.

References

1. Tsukimura B. Crustacean vitellogenesis: Its role in oocyte development. *Integr Comp Biol.* 2001; 41(3):465–76. <https://doi.org/10.1093/icb/41.3.465> WOS:000173211400011.
2. Quackenbush LS. Yolk synthesis in the marine shrimp, *Penaeus vannamei*. *Integr Comp Biol.* 2001; 41(3):458–64. <https://doi.org/10.1093/icb/41.3.458> WOS:000173211400010.
3. Avarre JC, Lubzens E, Babin PJ. Apolipoprotein, formerly vitellogenin, is the major egg yolk precursor protein in decapod crustaceans and is homologous to insect apolipoporphin II/I and vertebrate apolipoprotein B. *Bmc Evol Biol.* 2007; 7(3). Epub 2007/01/24. <https://doi.org/10.1186/1471-2148-7-3> PMID: 17241455; PubMed Central PMCID: PMC21783640.
4. Wu LT, Hui JH, Chu KH. Origin and evolution of yolk proteins: expansion and functional diversification of large lipid transfer protein superfamily. *Biol Reprod.* 2013; 88(4):102. Epub 2013/02/22. <https://doi.org/10.1095/biolreprod.112.104752> PMID: 23426435.
5. Kung SY, Chan SM, Hui JH, Tsang WS, Mak A, He JG. Vitellogenesis in the sand shrimp, *metapenaeus ensis*: the contribution from the hepatopancreas-specific vitellogenin gene (MeVg2). *Biol Reprod.* 2004; 71(3):863–70. Epub 2004/04/30. <https://doi.org/10.1095/biolreprod.103.022905> PMID: 15115717.
6. Okuno A, Yang WJ, Jayasankar V, Saido-Sakanaka H, Huong DT, Jasmani S, et al. Deduced primary structure of vitellogenin in the giant freshwater prawn, *Macrobrachium rosenbergii*, and yolk processing during ovarian maturation. *J Exp Zool.* 2002; 292(5):417–29. Epub 2002/02/22. <https://doi.org/10.1002/jez.10083> PMID: 11857476.
7. Arcos FG, Ibarra AM, Racotta LS. Vitellogenin in hemolymph predicts gonad maturity in adult female *Litopenaeus (Penaeus) vannamei* shrimp. *Aquaculture.* 2011; 316(1–4):93–8. <https://doi.org/10.1016/j.aquaculture.2011.02.045> WOS:000291337400014.
8. Tsutsui N, Saido-Sakanaka H, Yang WJ, Jayasankar V, Jasmani S, Okuno A, et al. Molecular characterization of a cDNA encoding vitellogenin in the coonstriped shrimp, *Pandalus hypsinotus* and site of vitellogenin mRNA expression. *Journal of experimental zoology Part A, Comparative experimental biology.* 2004; 301(10):802–14. Epub 2004/09/28. <https://doi.org/10.1002/jez.a.53> PMID: 15449343.
9. Soyez D, Vandenberghe JE, Martin M. Isolation and Characterization of a Vitellogenesis-Inhibiting Factor from Sinus Glands of the Lobster, *Homarus-Americanus*. *J Exp Zool.* 1987; 244(3):479–84. <https://doi.org/10.1002/jez.1402440314> WOS:A1987L363700013.
10. de Kleijn DP, Sleutels FJ, Martens GJ, Van Herp F. Cloning and expression of mRNA encoding prepro-gonad-inhibiting hormone (GIH) in the lobster *Homarus americanus*. *FEBS letters.* 1994; 353(3):255–8. Epub 1994/10/24. PMID: 7957869.
11. Webster SG, Keller R, Dirksen H. The CHH-superfamily of multifunctional peptide hormones controlling crustacean metabolism, osmoregulation, moulting, and reproduction. *Gen Comp Endocrinol.* 2012; 175(2):217–33. Epub 2011/12/08. <https://doi.org/10.1016/j.ygcen.2011.11.035> PMID: 22146796.
12. Chan SM, Gu PL, Chu KH, Tobe SS. Crustacean neuropeptide genes of the CHH/MIH/GIH family: implications from molecular studies. *Gen Comp Endocrinol.* 2003; 134(3):214–9. Epub 2003/11/26. [https://doi.org/10.1016/S0016-6480\(03\)00263-6](https://doi.org/10.1016/S0016-6480(03)00263-6) PMID: 14636627.
13. Azzouna A, Philippe M, Jarry T, Greve P, Martin G. Localization of crustacean hyperglycemic and vitellogenesis-inhibiting hormones in separate cell types in the protocerebrum of the woodlouse *Armadillidium vulgare* (Crustacea, Isopoda). *Gen Comp Endocrinol.* 2003; 131(2):134–42. Epub 2003/04/08. [https://doi.org/10.1016/S0016-6480\(02\)00645-7](https://doi.org/10.1016/S0016-6480(02)00645-7) PMID: 12679090.

14. Treerattrakool S, Panyim S, Chan SM, Withyachumnarnkul B, Udomkit A. Molecular characterization of gonad-inhibiting hormone of *Penaeus monodon* and elucidation of its inhibitory role in vitellogenin expression by RNA interference. *The FEBS journal*. 2008; 275(5):970–80. Epub 2008/01/29. <https://doi.org/10.1111/j.1742-4658.2008.06266.x> PMID: 18221491.
15. Chen T, Zhang LP, Wong NK, Zhong M, Ren CH, Hu CQ. Pacific white shrimp (*Litopenaeus vannamei*) vitellogenesis-inhibiting hormone (VIH) is predominantly expressed in the brain and negatively regulates hepatopancreatic vitellogenin (VTG) gene expression. *Biol Reprod*. 2014; 90(3):47. Epub 2014/01/24. <https://doi.org/10.1095/biolreprod.113.115030> PMID: 24451988.
16. Li GL, Deng SP, Jiang SN, Ye M, Chen HP, Chan SMF, et al. cDNA cloning, expression, and immunolocalization of gonad-inhibiting hormone (GIH) in *Litopenaeus vannamei*. *Genes Genom*. 2015; 37(10):883–92. <https://doi.org/10.1007/s13258-015-0315-0> WOS:000362020200009.
17. Hopkins PM. The eyes have it: A brief history of crustacean neuroendocrinology. *Gen Comp Endocrinol*. 2012; 175(3):357–66. Epub 2011/12/27. <https://doi.org/10.1016/j.ygcen.2011.12.002> PMID: 22197211.
18. Okumura T. Effects of bilateral and unilateral eyestalk ablation on vitellogenin synthesis in immature female kuruma prawns, *Marsupenaeus japonicus*. *Zoological science*. 2007; 24(3):233–40. Epub 2007/06/07. <https://doi.org/10.2108/zsj.24.233> PMID: 17551243.
19. Bae SH, Okutsu T, Kang BJ, Wilder MN. Alterations of pattern in immune response and vitellogenesis during induced ovarian development by unilateral and bilateral ablation in *Litopenaeus vannamei*. *Fisheries Sci*. 2013; 79(6):895–903. <https://doi.org/10.1007/s12562-013-0652-3> WOS:000327080300003.
20. Tsutsui N, Ohira T, Kawazoe I, Takahashi A, Wilder MN. Purification of sinus gland peptides having vitellogenesis-inhibiting activity from the whiteleg shrimp *Litopenaeus vannamei*. *Mar Biotechnol (NY)*. 2007; 9(3):360–9. Epub 2007/03/16. <https://doi.org/10.1007/s10126-006-6151-0> PMID: 17357858.
21. Ohira T, Okumura T, Suzuki M, Yajima Y, Tsutsui N, Wilder MN, et al. Production and characterization of recombinant vitellogenesis-inhibiting hormone from the American lobster *Homarus americanus*. *Pep-tides*. 2006; 27(6):1251–8. Epub 2005/12/03. <https://doi.org/10.1016/j.peptides.2005.10.019> PMID: 16321455.
22. Vrinda S, Jasmin C, Sivakumar KC, Jose S, Jose B, Philip R, et al. Regulating gonad inhibition and vitellogenin/vitellicin induction in *Penaeus monodon* using mature GIH fusion protein and polyclonal antisera. *Comparative Biochemistry and Physiology a-Molecular & Integrative Physiology*. 2017; 203:167–78. <https://doi.org/10.1016/j.cbpa.2016.09.007> WOS:000390718800021. PMID: 27642094
23. Tsutsui N, Ohira T, Okutsu T, Shinji J, Bae SH, Kang BJ, et al. Molecular cloning of a cDNA encoding vitellogenesis-inhibiting hormone in the whiteleg shrimp *Litopenaeus vannamei* and preparation of its recombinant peptide using an *E. coli* expression system. *Fisheries Sci*. 2013; 79(3):357–65. <https://doi.org/10.1007/s12562-013-0603-z> WOS:000318840400001.
24. Treerattrakool S, Panyim S, Udomkit A. Induction of ovarian maturation and spawning in *Penaeus monodon* broodstock by double-stranded RNA. *Mar Biotechnol (NY)*. 2011; 13(2):163–9. Epub 2010/03/25. <https://doi.org/10.1007/s10126-010-9276-0> PMID: 20333425.
25. Feijo RG, Braga AL, Lanes CF, Figueiredo MA, Romano LA, Klosterhoff MC, et al. Silencing of Gonad-Inhibiting Hormone Transcripts in *Litopenaeus vannamei* Females by use of the RNA Interference Technology. *Mar Biotechnol (NY)*. 2016; 18(1):117–23. Epub 2015/11/18. <https://doi.org/10.1007/s10126-015-9676-2> PMID: 26573611.
26. Qiao H, Xiong Y, Zhang W, Fu H, Jiang S, Sun S, et al. Characterization, expression, and function analysis of gonad-inhibiting hormone in Oriental River prawn, *Macrobrachium nipponense* and its induced expression by temperature. *Comp Biochem Physiol A Mol Integr Physiol*. 2015; 185:1–8. Epub 2015/03/17. <https://doi.org/10.1016/j.cbpa.2015.03.005> PMID: 25770669.
27. Chen T, Ren CH, Wang YH, Gao Y, Wong NK, Zhang LP, et al. Crustacean cardioactive peptide (CCAP) of the Pacific white shrimp (*Litopenaeus vannamei*): Molecular characterization and its potential roles in osmoregulation and freshwater tolerance. *Aquaculture*. 2016; 451:405–12. <https://doi.org/10.1016/j.aquaculture.2015.10.005> WOS:000364854800051.
28. Kang BJ, Okutsu T, Tsutsui N, Shinji J, Bae SH, Wilder MN. Dynamics of vitellogenin and vitellogenesis-inhibiting hormone levels in adult and subadult whiteleg shrimp, *Litopenaeus vannamei*: relation to molting and eyestalk ablation. *Biol Reprod*. 2014; 90(1):12. Epub 2013/12/18. <https://doi.org/10.1095/biolreprod.113.112243> PMID: 24337313.
29. Luo X, Chen T, Zhong M, Jiang X, Zhang L, Ren C, et al. Differential regulation of hepatopancreatic vitellogenin (VTG) gene expression by two putative molt-inhibiting hormones (MIH1/2) in Pacific white shrimp (*Litopenaeus vannamei*). *Peptides*. 2015; 68:58–63. Epub 2014/12/03. <https://doi.org/10.1016/j.peptides.2014.11.002> PMID: 25447412.
30. Li GL, Chen HP, Deng SP, Ye M, Jiang S, Chan SF, et al. In vivo and in vitro inhibitory action of 17beta-estradiol and environmental estrogen 4-nonylphenol on gonad-inhibiting hormone (GIH) expression in

- the eyestalks of *Litopenaeus vannamei*. *Genet Mol Res.* 2015; 14(4):14056–65. Epub 2015/11/05. <https://doi.org/10.4238/2015.October.29.25> PMID: 26535720.
31. Nagai C, Mabashi-Asazuma H, Nagasawa H, Nagata S. Identification and characterization of receptors for ion transport peptide (ITP) and ITP-like (ITPL) in the silkworm *Bombyx mori*. *J Biol Chem.* 2014; 289(46):32166–77. Epub 2014/10/04. <https://doi.org/10.1074/jbc.M114.590646> PMID: 25278025; PubMed Central PMCID: PMC4231692.
 32. Li R, Tian JZ, Zhuang CH, Zhang YC, Geng XY, Zhu LN, et al. CHHBP: a newly identified receptor of crustacean hyperglycemic hormone. *The Journal of experimental biology.* 2016; 219(Pt 8):1259–68. Epub 2016/02/21. <https://doi.org/10.1242/jeb.133181> PMID: 26896539.
 33. Chung JS, Webster SG. Binding sites of crustacean hyperglycemic hormone and its second messengers on gills and hindgut of the green shore crab, *Carcinus maenas*: a possible osmoregulatory role. *Gen Comp Endocrinol.* 2006; 147(2):206–13. Epub 2006/03/01. <https://doi.org/10.1016/j.ygcen.2006.01.002> PMID: 16504184.
 34. Nagai C, Asazuma H, Nagata S, Nagasawa H. Identification of a second messenger of crustacean hyperglycemic hormone signaling pathway in the kuruma prawn *Marsupenaeus japonicus*. *Ann N Y Acad Sci.* 2009; 1163:478–80. Epub 2009/05/22. <https://doi.org/10.1111/j.1749-6632.2009.04452.x> PMID: 19456392.
 35. Zheng J, Nakatsuji T, Roer RD, Watson RD. Studies of a receptor guanylyl cyclase cloned from Y-organs of the blue crab (*Callinectes sapidus*), and its possible functional link to ecdysteroidogenesis. *Gen Comp Endocrinol.* 2008; 155(3):780–8. Epub 2007/12/21. <https://doi.org/10.1016/j.ygcen.2007.10.009> PMID: 18093588.
 36. Pitts NL, Mykles DL. Localization and expression of molt-inhibiting hormone and nitric oxide synthase in the central nervous system of the green shore crab, *Carcinus maenas*, and the blackback land crab, *Gecarcinus lateralis*. *Comp Biochem Physiol A Mol Integr Physiol.* 2017; 203:328–40. Epub 2016/12/19. <https://doi.org/10.1016/j.cbpa.2016.10.012> PMID: 27989866.
 37. Zmora N, Sagi A, Zohar Y, Chung JS. Molt-inhibiting hormone stimulates vitellogenesis at advanced ovarian developmental stages in the female blue crab, *Callinectes sapidus* 2: novel specific binding sites in hepatopancreas and cAMP as a second messenger. *Saline Systems.* 2009; 5:6. Epub 2009/07/09. <https://doi.org/10.1186/1746-1448-5-6> PMID: 19583849; PubMed Central PMCID: PMC2714851.
 38. Covi JA, Chang ES, Mykles DL. Conserved role of cyclic nucleotides in the regulation of ecdysteroidogenesis by the crustacean molting gland. *Comp Biochem Physiol A Mol Integr Physiol.* 2009; 152(4):470–7. Epub 2009/01/13. <https://doi.org/10.1016/j.cbpa.2008.12.005> PMID: 19135164.
 39. Mykles DL, Adams ME, Gade G, Lange AB, Marco HG, Orchard I. Neuropeptide action in insects and crustaceans. *Physiol Biochem Zool.* 2010; 83(5):836–46. Epub 2010/06/17. <https://doi.org/10.1086/648470> PMID: 20550437; PubMed Central PMCID: PMC3844688.
 40. Okumura T. Effects of cyclic nucleotides, calcium ionophore, and phorbol ester on vitellogenin mRNA levels in incubated ovarian fragments of the kuruma prawn *Marsupenaeus japonicus*. *Gen Comp Endocrinol.* 2006; 148(2):245–51. Epub 2006/04/21. <https://doi.org/10.1016/j.ygcen.2006.03.005> PMID: 16624310.
 41. Bae SH, Okutsu T, Tsutsui N, Kang BJ, Chen HY, Wilder MN. Involvement of second messengers in the signaling pathway of vitellogenesis-inhibiting hormone and their effects on vitellogenin mRNA expression in the whiteleg shrimp, *Litopenaeus vannamei*. *Gen Comp Endocrinol.* 2017; 246:301–8. Epub 2017/01/08. <https://doi.org/10.1016/j.ygcen.2017.01.006> PMID: 28062305.
 42. Jasmani S, Kawazoe I, Shih TW, Suzuki Y, Aida K. Hemolymph vitellogenin levels during ovarian development in the kuruma prawn *Penaeus japonicus*. *Fisheries Sci.* 2000; 66(3):535–9. <https://doi.org/10.1046/j.1442-2906.2000.00084.x> WOS:000088134000018.
 43. Okumura T, Yamano K, Sakiyama K. Vitellogenin gene expression and hemolymph vitellogenin during vitellogenesis, final maturation, and oviposition in female kuruma prawn, *Marsupenaeus japonicus*. *Comp Biochem Physiol A Mol Integr Physiol.* 2007; 147(4):1028–37. Epub 2007/04/24. <https://doi.org/10.1016/j.cbpa.2007.03.011> PMID: 17449309.
 44. Tsutsui N, Kawazoe I, Ohira T, Jasmani S, Yang WJ, Wilder MN, et al. Molecular Characterization of a cDNA Encoding Vitellogenin and Its Expression in the Hepatopancreas and Ovary during Vitellogenesis in the Kuruma Prawn, *Penaeus japonicus*. *Zoological science.* 2000; 17(5):651–60. Epub 2008/06/04. <https://doi.org/10.2108/zsj.15.651> PMID: 18517301.
 45. Wang W, Wu X, Liu Z, Zheng H, Cheng Y. Insights into hepatopancreatic functions for nutrition metabolism and ovarian development in the crab *Portunus trituberculatus*: gene discovery in the comparative transcriptome of different hepatopancreas stages. *Plos One.* 2014; 9(1):e84921. Epub 2014/01/24. <https://doi.org/10.1371/journal.pone.0084921> PMID: 24454766; PubMed Central PMCID: PMC3890295.

46. Li K, Chen L, Zhou Z, Li E, Zhao X, Guo H. The site of vitellogenin synthesis in Chinese mitten-handed crab *Eriocheir sinensis*. Comparative biochemistry and physiology Part B, Biochemistry & molecular biology. 2006; 143(4):453–8. Epub 2006/02/17. <https://doi.org/10.1016/j.cbpb.2005.12.019> PMID: 16480909.
47. Raviv S, Parnes S, Segall C, Davis C, Sagi A. Complete sequence of *Litopenaeus vannamei* (Crustacea: Decapoda) vitellogenin cDNA and its expression in endocrinologically induced sub-adult females. Gen Comp Endocrinol. 2006; 145(1):39–50. Epub 2005/08/27. <https://doi.org/10.1016/j.ygcen.2005.06.009> PMID: 16122741.
48. Wongprasert K, Asuvapongpatana S, Poltana P, Tiensuwan M, Withyachumnarnkul B. Serotonin stimulates ovarian maturation and spawning in the black tiger shrimp *Penaeus monodon*. Aquaculture. 2006; 261(4):1447–54. <https://doi.org/10.1016/j.aquaculture.2006.08.044> WOS:000243289400039.
49. Yamano K, Qiu GF, Unuma T. Molecular cloning and ovarian expression profiles of thrombospondin, a major component of cortical rods in mature oocytes of penaeid shrimp, *Marsupenaeus japonicus*. Biol Reprod. 2004; 70(6):1670–8. Epub 2004/02/10. <https://doi.org/10.1095/biolreprod.103.025379> PMID: 14766728.
50. Chen T, Wong NK, Jiang X, Luo X, Zhang L, Yang D, et al. Nitric oxide as an antimicrobial molecule against *Vibrio harveyi* infection in the hepatopancreas of Pacific white shrimp, *Litopenaeus vannamei*. Fish & shellfish immunology. 2015; 42(1):114–20. Epub 2014/12/03. <https://doi.org/10.1016/j.fsi.2014.10.042> PMID: 25449376.
51. Yan AF, Chen T, Chen S, Tang DS, Liu F, Jiang X, et al. Signal transduction mechanism for glucagon-induced leptin gene expression in goldfish liver. Int J Biol Sci. 2016; 12(12):1544–54. Epub 2016/12/21. <https://doi.org/10.7150/ijbs.16612> PMID: 27994518; PubMed Central PMCID: PMC5166495.
52. Ventura-Lopez C, Galindo-Torres PE, Arcos FG, Galindo-Sanchez C, Racotta IS, Escobedo-Fregoso C, et al. Transcriptomic information from Pacific white shrimp (*Litopenaeus vannamei*) ovary and eye-stalk, and expression patterns for genes putatively involved in the reproductive process. Gen Comp Endocrinol. 2017; 246:164–82. Epub 2016/12/15. <https://doi.org/10.1016/j.ygcen.2016.12.005> PMID: 27964922.
53. Francis SH, Corbin JD. Cyclic nucleotide-dependent protein kinases: intracellular receptors for cAMP and cGMP action. Crit Rev Clin Lab Sci. 1999; 36(4):275–328. Epub 1999/09/16. <https://doi.org/10.1080/10408369991239213> PMID: 10486703.
54. Lee SG, Bader BD, Chang ES, Mykles DL. Effects of elevated ecdysteroid on tissue expression of three guanyl cyclases in the tropical land crab *Gecarcinus lateralis*: possible roles of neuropeptide signaling in the molting gland. Journal of Experimental Biology. 2007; 210(18):3245–54. <https://doi.org/10.1242/jeb.007740> WOS:000249517400018. PMID: 17766302.
55. Huang W, Ren CH, Li HM, Huo D, Wang YH, Jiang X, et al. Transcriptomic analyses on muscle tissues of *Litopenaeus vannamei* provide the first profile insight into the response to low temperature stress. Plos One. 2017; 12(6). <https://doi.org/10.1371/journal.pone.0178604> WOS:000402624300071. PMID: 28575089.
56. Shi H, Ruan L, Yan X, Yao D, Xu X. The role of *Litopenaeus vannamei* p38 in white spot syndrome virus infection. Dev Comp Immunol. 2014; 44(1):180–5. Epub 2013/12/24. <https://doi.org/10.1016/j.dci.2013.12.005> PMID: 24361691.
57. Shi H, Yan X, Xu X, Ruan L. Molecular cloning and characterization of a cDNA encoding extracellular signal-regulated kinase from *Litopenaeus vannamei*. Fish & shellfish immunology. 2012; 33(4):813–20. Epub 2012/08/14. <https://doi.org/10.1016/j.fsi.2012.07.008> PMID: 22884486.
58. Shi H, Yan X, Ruan L, Xu X. A novel JNK from *Litopenaeus vannamei* involved in white spot syndrome virus infection. Dev Comp Immunol. 2012; 37(3–4):421–8. Epub 2012/03/21. <https://doi.org/10.1016/j.dci.2012.03.002> PMID: 22430647.
59. Yao DF, Ruan LW, Xu X, Shi H. Identification of a c-Jun homolog from *Litopenaeus vannamei* as a downstream substrate of JNK in response to WSSV infection. Developmental and Comparative Immunology. 2015; 49(2):282–9. <https://doi.org/10.1016/j.dci.2014.12.012> WOS:000349880500010. PMID: 25530093
60. Li F, Xiang J. Signaling pathways regulating innate immune responses in shrimp. Fish & shellfish immunology. 2013; 34(4):973–80. Epub 2012/09/13. <https://doi.org/10.1016/j.fsi.2012.08.023> PMID: 22967763.
61. Xu D, Liu W, Alvarez A, Huang T. Cellular immune responses against viral pathogens in shrimp. Dev Comp Immunol. 2014; 47(2):287–97. Epub 2014/08/12. <https://doi.org/10.1016/j.dci.2014.08.004> PMID: 25111591.
62. Huang X, Feng B, Huang H, Ye H. In vitro stimulation of vitellogenin expression by insulin in the mud crab, *Scylla paramamosain*, mediated through PI3K/Akt/TOR pathway. Gen Comp Endocrinol. 2017; 250:175–80. Epub 2017/06/26. <https://doi.org/10.1016/j.ygcen.2017.06.013> PMID: 28647319.