

RESEARCH ARTICLE

Effect of Inorganic and Organic Carbon Enrichments (DIC and DOC) on the Photosynthesis and Calcification Rates of Two Calcifying Green Algae from a Caribbean Reef Lagoon

Friedrich W. Meyer¹✉, Nadine Schubert²✉, Karen Diele^{3,5}, Mirta Teichberg¹, Christian Wild^{1,4}, Susana Enríquez²✉*

1 Leibniz Center for Tropical Marine Ecology (ZMT), Bremen, Germany, **2** Laboratory of Photobiology, Unidad Académica de Sistemas Arrecifales Puerto Morelos (ICMyL), Universidad Nacional Autónoma de México (UNAM), Cancún, Mexico, **3** Edinburgh Napier University, School of Life, Sport and Social Sciences, EH11 4BN Edinburgh, United Kingdom, **4** Faculty of Biology & Chemistry, University of Bremen, Bibliothekstraße 1, Bremen, Germany, **5** St Abbs Marine Station, St Abbs, Berwickshire, TD14 5QF, United Kingdom

✉ These authors contributed equally to this work.

✉ Current address: Departamento de Geociências, Universidade Federal de Santa Catarina (UFSC), Florianópolis, Brasil

* enriquez@cmarl.unam.mx; susana.enriquezdominguez@gmail.com



CrossMark
click for updates

OPEN ACCESS

Citation: Meyer FW, Schubert N, Diele K, Teichberg M, Wild C, Enríquez S (2016) Effect of Inorganic and Organic Carbon Enrichments (DIC and DOC) on the Photosynthesis and Calcification Rates of Two Calcifying Green Algae from a Caribbean Reef Lagoon. PLoS ONE 11(8): e0160268. doi:10.1371/journal.pone.0160268

Editor: Chaolun Allen Chen, Academia Sinica, TAIWAN

Received: November 24, 2015

Accepted: July 15, 2016

Published: August 3, 2016

Copyright: © 2016 Meyer et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This study was funded by two research projects DGAPA (IN206710) and a SEP-CONACYT (129880) to SE and also by the Leibniz Association to CW. The MASTS pooling initiative (The Marine Alliance for Science and Technology for Scotland) with the grant reference HR09011 to KD is also gratefully acknowledged. Support to NS was provided through two post-doctoral fellowships: UNAM-DGAPA and SEP-CONACYT (129880). The funders had no

Abstract

Coral reefs worldwide are affected by increasing dissolved inorganic carbon (DIC) and organic carbon (DOC) concentrations due to ocean acidification (OA) and coastal eutrophication. These two stressors can occur simultaneously, particularly in near-shore reef environments with increasing anthropogenic pressure. However, experimental studies on how elevated DIC and DOC interact are scarce and fundamental to understanding potential synergistic effects and foreseeing future changes in coral reef function. Using an open mesocosm experiment, the present study investigated the impact of elevated DIC (pH_{NBS}: 8.2 and 7.8; pCO₂: 377 and 1076 µatm) and DOC (added as 833 µmol L⁻¹ of glucose) on calcification and photosynthesis rates of two common calcifying green algae, *Halimeda incrassata* and *Udotea flabellum*, in a shallow reef environment. Our results revealed that under elevated DIC, algal photosynthesis decreased similarly for both species, but calcification was more affected in *H. incrassata*, which also showed carbonate dissolution rates. Elevated DOC reduced photosynthesis and calcification rates in *H. incrassata*, while in *U. flabellum* photosynthesis was unaffected and thalys calcification was severely impaired. The combined treatment showed an antagonistic effect of elevated DIC and DOC on the photosynthesis and calcification rates of *H. incrassata*, and an additive effect in *U. flabellum*. We conclude that the dominant sand dweller *H. incrassata* is more negatively affected by both DIC and DOC enrichments, but that their impact could be mitigated when they occur simultaneously. In contrast, *U. flabellum* can be less affected in coastal eutrophic waters by

role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

elevated DIC, but its contribution to reef carbonate sediment production could be further reduced. Accordingly, while the capacity of environmental eutrophication to exacerbate the impact of OA on algal-derived carbonate sand production seems to be species-specific, significant reductions can be expected under future OA scenarios, with important consequences for beach erosion and coastal sediment dynamics.

Introduction

The rise of oceanic $p\text{CO}_2$ caused by increasing CO_2 concentrations in the atmosphere is leading to significant changes in the ocean carbonate system, which are primarily reflected in an increase in bicarbonate concentration and a decrease in seawater pH (ocean acidification- OA) [1, 2]. These changes also induce a significant decline in the saturation state of the different crystallization forms of calcium carbonate in the marine environment, which will facilitate the dissolution of existing calcium carbonate deposits and cause severe impacts on marine calcifiers. Many coral reef habitats and their lagoons are particularly threatened by ocean acidification. Studies conducted at natural low pH sites have shown that under OA the reef framework is less stable [3], and reef accretion is compromised [4], as are the ecosystem services provided by the reef [5].

Local impacts associated with nutrient enrichment, pollution and overfishing have also increased in the last decades, leading to so called “phase shifts” in many parts of the Caribbean and coral reefs worldwide [6, 7]. One of the main drivers of “phase shifts” is related to inorganic and organic nutrient inputs derived from untreated or poorly treated sewage. The impact of elevated DOC concentrations on coral reef health is currently of major concern in coral reef research [8–10], as elevated DOC has been associated with enhanced bacterial growth and other processes that lead to oxygen depletion and the accumulation of toxic substances, and ultimately to an increase in coral mortality [9, 11, 12]. High concentrations of DOC, predominantly in the form of dissolved carbohydrates, can also enter the coral reef system in the form of exudates released by the benthic community [13, 14]. Previous results have shown minimal or no significant differences in the DOC concentrations released by benthic calcifying algae (*Halimeda opuntia*) compared to coral exudates (*Porites lobata*) [10, 15], although it has been postulated that bacterial growth is primarily triggered by algal-derived DOC rather than DOC released by corals [10, 15].

The sandy bottom of Caribbean reef lagoons are commonly colonized by rhizophytic calcareous green algae (Siphonales) of the genera *Halimeda*, *Udotea*, *Penicillus* and *Rhipocephalus*, which are associated with the seagrass habitat builder *Thalassia testudinum* [16–18]. Calcareous green algae produce an important fraction of coral reef carbonate production in the form of calcareous sand, essential to support reef accretion [19–22]. Most of the studies that have investigated the responses of marine macrophytes to OA and other local threats have focused on species of the genus *Halimeda*, due to this genus is considered one of the most productive. Limited attention has been given to other important reef calcifiers, such as species from the genus *Udotea*, *Penicillus*, *Rhipocephalus*. For experimental studies focused on *Halimeda* spp., it has been concluded that this genus displays a large species-specific variation to increasing levels of dissolved inorganic carbon (DIC). Some species reduce their photosynthetic rates [23, 24], while others have shown positive [25] or no effect on algal photosynthesis [26–28]. Similarly, large inter-specific variation has been documented for the response of calcification of *Halimeda* spp. to DIC increases [24, 26–33] indicating that some species may be more tolerant to OA than others. Altered skeletal structure of different *Halimeda* spp. in response to OA conditions has also been reported [34, 35], being indicative of potential needle dissolution [29] and/or the formation of more slender crystals during exposure to reduced pH [25]. Yet, alteration of skeletal structure may also affect the contribution of species from the genus *Halimeda* to sediment

carbonate production under different OA scenarios, irrespective of the severity of the impact detected on algal physiology.

In contrast to OA, nutrient enrichment enhances *Halimeda* spp. production and growth [23, 36, 37], with the exception of phosphate enrichment, for which a large species-specific variation has been also reported [36]. An analysis on the combined effect of inorganic nutrient enrichment and reduced pH on *Halimeda opuntia* has shown decreased enhancement in algal production under nutrient enrichment and reduced pH, relative to the estimated values for ambient pH [38]. Meyer et al. [28] have recently shown the negative effects of increased DOC concentration on the photosynthesis of two *Halimeda* species from the Great Barrier Reef, *H. opuntia* and *H. macroloba*, but no effect was found on algal calcification rates under illumination. These authors further investigated the combined effect of elevated DOC and DIC concentrations, and documented an adverse impact on thallus photosynthesis for both species, while only *H. opuntia* showed a negative effect on dark calcification rates. These findings support the large species-specific component of the response of marine algae to the combined effects of OA and increased DOC, and the importance of increasing the number of this type of experimental studies for other sites and species.

Halimeda incrassata (J. Ellis) J.V. Lamouroux and *Udotea flabellum* (J. Ellis & Solander) M. Howe are two abundant rhizophytic species of the macrophyte community of shallow seagrass habitats dominated by the species *Thalassia testudinum*. In the Puerto Morelos reef lagoon, Mexican Caribbean, both species are among the most abundant calcareous algae. Estimates of *H. incrassata* primary production for this lagoon ($0.2\text{--}0.5\text{ g dwt m}^{-2}\text{ day}^{-1}$) [22] are lower than the reported values for seagrass leaf production ($0.9\text{--}1.2\text{ g dwt m}^{-2}\text{ day}^{-1}$) [39]. Interestingly, annual carbonate production for *H. incrassata* from this area is in the same range or even lower (between 0.5 and $1.0\text{ kg CaCO}_3\text{ m}^{-2}\text{ y}^{-1}$) [22] than estimates of annual carbonate production recently documented for the dominant seagrass *T. testudinum* (between 0.5 and $5.63\text{ kg CaCO}_3\text{ m}^{-2}\text{ y}^{-1}$) by Enríquez and Schubert [40]. To our knowledge, no estimates for *Udotea* spp. annual carbonate production are yet available for this area or other locations. The only study on the daily carbonate production of *U. flabellum* [41] has documented that this species produces about 45% less carbonate per day than *H. incrassata*.

The Mexican Caribbean (Cancún and the Riviera Maya) has experienced a 4.3-fold population increase during the period from 2000–2009 [18], which is associated with the rapid coastal development of large tourist complexes. These changes have severely affected the reef habitat, particularly the benthic macrophyte community associated with seagrass beds, which is shifting towards an increased presence of fleshy macroalgae [18], and increased biomass of green calcifiers [42]. To understand the combined effect of these local impacts with the predicted negative effect of OA on marine calcifiers, this study investigated the direct and combined effects of experimental increases in DIC and DOC on the physiological performance of *Halimeda incrassata* and *Udotea flabellum*. This multi-factorial study aims to analyze a more realistic scenario that may be also very useful for other areas affected by similar coastal eutrophication derived from different anthropogenic impacts. As calcareous macroalgae are considered important contributors to reef carbonate budgets, this study can also contribute to improve our understanding of future impacts caused by the combined effects of global and local threats on reef accretion and the stability of the reef system.

Materials and Methods

Algal Collection and Maintenance

Several individuals of *H. incrassata* and *U. flabellum* were collected by SCUBA diving from the Puerto Morelos reef lagoon, Mexican Caribbean ($20^{\circ} 52' \text{ N}$, $86^{\circ} 52' \text{ W}$), in March 2012 at 3–3.5

m depth, and transported in mesh-covered ziplog bags to the mesocosm facilities of the Universidad Nacional Autónoma de México (UNAM). To minimize physiological variability among replicates associated with age and photoacclimatory condition of the thallus, thalli of similar size and position were selected for the experimental analysis: 5–7 apical segments for *H. incrassata*, and *U. flabellum* individuals of 3–5 cm height.

The selected individuals were acclimated for 5 days to the experimental conditions by placing them in 50 L experimental tanks receiving filtered (~50 µm) ambient seawater (~28°C, pH 8.2) from the lagoon, with continuous flow of 1 L min⁻¹. Irradiance levels at mesocosms were adjusted using neutral density shade mesh to simulate light conditions at collection depth (51% of surface irradiance, E_s). E_s was calculated using surface irradiance data and the down-welling light attenuation coefficient of the reef lagoon estimated for the sampling period (February–March 2012) of K_d = 0.2 m⁻¹, which was similar to previous values reported [43]. Variation in diurnal irradiance was continuously recorded throughout the experiment using a cosine-corrected light sensor (LI-190SA; LI-COR, Lincoln, NE, USA) connected to a data logger (LI-1400; LI-COR, Lincoln, NE, USA), located at the mesocosm system.

After the pre-acclimation period (5 days), initial measurements of photosynthesis, respiration and calcification rates were performed (n = 6) as described below, and 12 individuals of each species (n = 12) were randomly positioned into each tank.

Experimental Treatments

The experiment was conducted over 10 days in an open flow-through system, which consisted of 12 tanks of 50 L each. To enhance water movement in the tanks and prevent carbon limitation of algal photosynthesis, aquaria pumps (500 L h⁻¹) were located in each tank and connected to a rectangular PVC frame surrounding the tank with holes facing inward to create homogenous flow conditions according to Cayabyab and Enríquez [44]. We used three replicate tanks per treatment and placed tanks with the different treatment levels (ambient and increased, see below) in alternating order.

The treatments comprised ambient and elevated pCO₂ concentrations in order to simulate OA changes in CO₂ availability from 380 to 1000 µatm, respectively, as well as ambient and increased DOC conditions (see Table 1). The increase in pCO₂ concentration was achieved by pH manipulation via CO₂ gas injection by a potentiometric pH sensor controlled pH stat system (IKS Aquaristic Products, Karlsbad, Germany). pH-reading by the pH-stat system was continuous (every other second) to adjust pH levels in the system, and pH sensors were calibrated every other day according to values measured by a WTW Multi 3430 probe (WTW, Weilheim, Germany). pH in the tanks was maintained at 8.2 for the control treatment, and reduced to 7.8 for the high DIC treatments (see Table 1). The elevated pCO₂ levels used here were selected considering the Representative Concentration Pathways 8.5 (RCP8.5), which predicts a decrease of seawater pH between 7.7–7.8 [45].

Table 1. Variation in the experimental conditions. Values for the carbonate system parameters were calculated using CO2SYS with temperature, salinity, total alkalinity (TA) and pH_{NBS} as input parameters (n = 3). Additionally, the Biological Oxygen Demand (BOD) of the seawater in each treatment are given, determined through oxygen consumption using dark incubations over 24 h and referring the estimated changes to the water volume of the incubation (n = 3). Data represent mean ± SD (n = 6) and the results of a one-way ANOVA (p<0.05) performed to determine significant differences in BOD between treatments, are indicated by different letters.

Treatment	pH [NBS]	DOC (µmol L ⁻¹)	T°C	Salinity (ppt)	TA (µmol kgSW ⁻¹)	pCO ₂ (µatm)	HCO ₃ (µmolkg SW ⁻¹)	Ω _{Ar}	BOD (mg O ₂ L ⁻¹ h ⁻¹)
Control	8.22 ±0.02	171	28.1±0.1	35.9 ±0.1	2414 ±8	377 ±27	1776 ±26	4.2 ±0.2	0.63 ±0.6 ^a
High DIC	7.84 ±0.04	171	28.0±0.2	36 ±0.1	2409 ±6	1076 ±101	2097 ±24	2.0 ±0.1	1.03 ±0.3 ^a
High DOC	8.19 ±0.03	550	28.1±0.2	35.9 ±0.1	2414 ±8	415 ±31	1811 ±31	4.0 ±0.2	1.35 ±0.3 ^a
High DOC & DIC	7.82 ±0.02	550	28.0±0.1	36 ±0.1	2413 ±7	1135 ±55	2114 ±14	2.0 ±0.1	1.62 ±0.4 ^a

doi:10.1371/journal.pone.0160268.t001

DOC treatment levels were adjusted to concentrations described in previous studies on coral communities, using glucose and lactose in high concentrations as DOC [8, 9, 46]. In this study, the DOC treatment in the form of highly bioavailable DOC was achieved by additions of $833 \mu\text{mol L}^{-1}$ DOC (Glucose, D-Glucose, Sigma Aldrich) twice daily at 08:00 and 20:00 to each of the six high-DOC treatment tanks, simulating sudden DOC enrichments events, common in nature associated with strong rain. To quantify the resulting DOC treatment conditions, DOC concentration were measured over a 12-hour cycle (Fig 1), showing an average DOC concentration of $550 \mu\text{mol L}^{-1}$ (Fig 1). Samples for TOC were filtered through $0.45 \mu\text{m}$ GFF filters (Whatman), acidified with $150 \mu\text{L}$ fuming HCl and frozen at -20°C until analysis using a Shimadzu TOC-5000A (Shimadzu, USA).

Salinity and temperature were also monitored for each tank twice daily throughout the experiment (WTW Multiprobe 3430), and total alkalinity (TA) every second day. Water samples from the tanks were filtered through $0.45 \mu\text{m}$ GFF filters and stored with a drop of chloroform at 4°C until TA analysis. These parameters, together with the pH values on the NBS scale taken every two days with a multiprobe (WTW3430, Weilheim, Germany), were used to calculate the carbonate system using the CO2SYS excel spreadsheet software, with the constants from Mehrbach et al. [47] (Table 1).

Biological Oxygen Demand (BOD)

To evaluate potential enhancements in microbial respiration rates in seawater, BOD was measured at the end of the experiment. We incubated 150 mL of unfiltered seawater in Winkler bottles for 24 h in the dark ($n = 3$), under constant temperature conditions (28°C). Oxygen concentration (mg L^{-1} and % saturation), as well as salinity and temperature were also measured before and after incubations. O_2 consumption rates in $\text{mg O}_2 \text{ L}^{-1} \text{ h}^{-1}$ were calculated and corrected for water volume and length of incubation.

Assessment of Maximum Photosynthetic Quantum Efficiency

Maximum photochemical efficiency of photosystem II—PSII (F_v/F_m) of experimental organisms was measured every evening at 20:00 on the apical segments of the organisms, using a




Fig 1. DOC concentrations in the high DOC-treatments measured during a 12-hour cycle, from 8:00 a. m. to 20:00 pm, after addition of $833 \mu\text{mol L}^{-1}$ DOC as glucose (indicated by an arrow). Filled-solid circles represent the average of two sampling points ($n = 2$) of the high DOC-treatments, and the open-white circle indicates the ambient DOC concentration in the non-DOC-enriched treatments.

doi:10.1371/journal.pone.0160268.g001

Pulse Amplitude Modulated fluorometer (Diving-PAM, Walz, Germany). At this time, one hour after sunset, algal thalli had already achieved the maximum F_v/F_m of the day, as all the non-photochemical quenching processes were relaxed, and the maximum PSII recovery of the day had been already reached (see [44, 48]).

Quantification of Photosynthesis, Respiration and Light Calcification Rates

For physiological measurements, young 4–5 apical segments of *H. incrassata* thalli and the uppermost 2 cm of *U. flabellum* thalli were selected (two organisms per tank and species) to reduce the variation among replicates in the thallus physiological condition, due to age, photo-acclimation, abundance of epiphytes and/or accumulation of damage. The segments were separated from the parent plant at least 2 h before physiological determinations were started in order to allow complete wound healing [49].

Before and at the end of the experiment, photosynthesis and calcification rates were simultaneously determined by incubating algal thalli for 30 min under a saturating light intensity of $500 \mu\text{mol quanta m}^{-2} \text{s}^{-1}$ (three times the E_k of the species, data not shown), in freshly filtered seawater obtained from the respective treatment tanks. The incubation water (17 mL) was collected at the beginning and at the end of the light incubation to determine the alkalinity changes induced by algal activity (see below). The samples were incubated in darkness for another 10–15 min to determine the post-illuminatory respiration rate (R_L). Oxygen evolution rates were measured polarographically in water-jacked chambers (DW3, Hansatech Instruments Ltd., Norfolk, UK), using Clark-type O_2 electrodes (Hansatech). A circulating bath with a controlled temperature system (RTE-100/RTE 101LP; Neslab Instruments Inc., Portsmouth, NH, USA) allowed maintenance of a constant temperature of 28°C (treatment temperature) during the incubation. The electrodes were calibrated with air- and N_2 -saturated filtered seawater. Freshly filtered seawater ($0.45 \mu\text{m}$) from the respective treatment tank was used for the incubations, with DIC and DOC concentrations corresponding to the treatment conditions (see Table 1). Data were captured with a computer equipped with an analog/digital converter using DATACAN V software (Sable Systems, Inc., Las Vegas, NV, USA). Gross photosynthesis was calculated adding to the net photosynthesis determined in the incubations, the oxygen consumption through post-illuminatory respiration.

Calcification rates were determined using the alkalinity anomaly principle based on the ratio of two equivalents of total alkalinity for each mole of precipitated CaCO_3 [50]. For alkalinity measurements, a modified spectrophotometer procedure as described by [51] and [40] was used. For quality control, a certified reference material of known total alkalinity (CRM, Scripps Institution of Oceanography, USA) was used to calibrate the method.

Quantification of Algal Surface Area

For normalization of the measured metabolic rates, the surface area of each algal segment was determined by scanning the thalli and analyzing the digital images using ImageJ software.

Statistical Analyses

Data were tested for normality using the Shapiro-Wilk test, and for equal variance using the Levene median test. Analyses of variance (ANOVA) allowed for the determination of significant differences ($p < 0.05$) between the different descriptors used to characterize the physiological response of the species. A one-way ANOVA was used to compare initial photosynthetic, respiratory and calcifications rates; the calcification / photosynthesis ratio; and for the comparison of BOD values between each treatment and the control. A t-student test was used to

evaluate significant differences between initial and final F_v/F_m values with respect to the control organisms. To analyze whether F_v/F_m , photosynthesis, respiration and calcification rates differed significantly between treatments, two-way-ANOVA tests were used, considering the DIC and DOC treatments as fixed factors to test for direct effects, as well as the interaction (DIC x DOC). For the comparison of differences between individuals and treatment combinations, a Newman-Keuls Post-hoc test was used. The statistical analyses were conducted using Statistica 12.0.

Results

Halimeda incrassata showed significantly higher photosynthetic and calcification rates than *Udotea flabellum*, with no difference in post-illuminatory respiratory rates (Table 2). These differences were also reflected in their calcification to photosynthesis ratios. While *H. incrassata* was able to precipitate $0.57 (\pm 0.08)$ mol CaCO_3 per mol O_2 evolved in photosynthesis, *U. flabellum* (ANOVA, $p = 0.038$) only precipitates $0.25 (\pm 0.07)$ mol CaCO_3 per mol O_2 produced (Table 2).

In addition, the two species showed contrasting responses to experimental DIC and DOC treatments. The response of maximum photosynthetic rates was similar in both species, but more pronounced than indicated by the F_v/F_m response. The variation in F_v/F_m was closely related to the diurnal variation in solar radiation. Control organisms showed a similar pattern of variation in both species, with a slight, but non-significant decline over time (t-test, $p = 0.528$ for *H. incrassata*, $p = 0.560$ for *U. flabellum*) compared to initial values (Fig 2A and 2B). When comparing final F_v/F_m values, *H. incrassata* showed a significant decline in the DOC treatments, under both ambient and elevated DIC concentrations (Fig 2C, Table 3), while *U. flabellum* only showed a negative response of F_v/F_m under elevated DIC (Fig 2D, Table 2). Significant reductions in P_{\max} were estimated for *H. incrassata* in all treatments, when compared to control organisms. P_{\max} reductions ranged from -30% (elevated DIC) to -43% (elevated DOC; Fig 2E), and showed a significant effect in the combined treatment (Table 3). In contrast, *U. flabellum* experienced a significant reduction in P_{\max} under elevated DIC compared to the control (high DIC: -33%; high DIC+high DOC: -21%), while elevated DOC did not cause any effect on thallus photosynthesis (Fig 2F, Table 3). Thallus respiratory rates were not affected by any experimental treatment in any species (Fig 2E and 2F).

The response of thallus calcification to the experimental treatments also showed large differences between species. While *H. incrassata* showed full suppression of thallus calcification and even dissolution of CaCO_3 after exposure to elevated DIC, thallus calcification was still positive albeit significantly reduced in *U. flabellum* after exposure to the same treatment (-36% compared to control; Fig 3). The opposite response was observed for elevated DOC, as we found a significant decline in calcification rates of *H. incrassata* (-68%) with respect to control organisms (yet positive values), while no calcification but dissolution of CaCO_3 (negative values) was measured for *U. flabellum* (Fig 3). The inhibition of thallus calcification by elevated DOC

Table 2. Comparison of the initial values (day 0) of gross maximum photosynthetic rates (P_{\max}), post-illumination respiration (R_L), maximum calcification rates (G_{\max}), and the ratio of calcification:photosynthesis ($G_{\max}:P_{\max}$) of *Halimeda incrassata* and *Udotea flabellum*. Data represent mean \pm SE (n = 6) and significant differences between species (one-way ANOVA, $p < 0.05$) are indicated by different letters.

Metabolic rates	<i>Halimeda incrassata</i>	<i>Udotea flabellum</i>
P_{\max}	2.33 ± 0.08^a	1.56 ± 0.06^b
R_L	0.39 ± 0.02^a	0.47 ± 0.05^a
G_{\max}	1.32 ± 0.2^a	0.40 ± 0.16^b
$G_{\max}:P_{\max}$	0.57 ± 0.08^a	0.25 ± 0.07^b

doi:10.1371/journal.pone.0160268.t002




Fig 2. Photosynthetic responses of *H. incrassata* (a, c, e) and *U. flabellum* (b, d, f) to the experimental DIC and DOC treatments. (a, b) Daily variation in the maximum quantum efficiency of photosystem II (F_v/F_m) along the experiment and the corresponding daily integrated values in light exposure in mol quanta $m^{-2} day^{-1}$ (grey bars); (c, d) F_v/F_m for each treatment at the end of the experiment; and (e, f) gross photosynthesis, P_{max} , (light grey bars) and respiration, R_L , (black bars) rates at the end of the experiment. Data represent mean \pm SE ($n = 3$) and significant differences between treatments (ANOVA, Newman-Keuls, $p < 0.05$) are indicated by different superscript letters.

doi:10.1371/journal.pone.0160268.g002

concentration was further exacerbated in *U. flabellum* in the combined treatment, due to the addition of the negative effect of DIC (Fig 3B), as no significant interactive effect was found for the response of thallus calcification in this species (Table 3). In contrast, the combined treatment did not show any significant impact on *H. incrassata* calcification, notwithstanding the significant negative direct effects of elevated DOC and DIC (Fig 3A). These findings support

Table 3. Two-way ANOVA analyses performed to determine significant differences in the physiological responses of the apical segments of *Hali-medea incrassata* and *Udotea flabellum* exposed to four experimental treatments: control, high DIC concentration, high DOC concentration, and the combined treatment ($n = 3$ for each treatment). DIC and DOC were considered fixed factors and DIC x DOC show the interaction between both factors.

Response variable	Species	Source of variation	DF	SS	MS	F-value	p-value
Maximum quantum yield (F_v/F_m)	<i>H. incrassata</i>	DIC	1	0.0005	0.0005	0.69	0.4275
		DOC	1	0.0121	0.0121	16.36	0.0037*
		DIC x DOC	1	0.0011	0.0011	1.45	0.2632
		Residual	8	0.0059	0.0007		
	<i>U. flabellum</i>	DIC	1	0.0104	0.0104	5.85	0.0419*
		DOC	1	0.0006	0.0006	0.33	0.5790
		DIC x DOC	1	0.0093	0.0093	5.21	0.0519
		Residual	8	0.0142	0.0018		
Gross	<i>H. incrassata</i>	DIC	1	0.162	0.162	6.267	0.0368*
Photosynthesis (P_{max})		DOC	1	0.863	0.863	33.37	0.00042*
		DIC x DOC	1	0.590	0.590	22.82	0.00140*
		Residual	8	0.207	0.026		
		<i>U. flabellum</i>	DIC	1	0.232	0.232	25.96
		DOC	1	0.006	0.006	0.68	0.4349
		DIC x DOC	1	0.027	0.027	2.99	0.1222
		Residual	8	0.071	0.009		
		Respiration (R_L)	<i>H. incrassata</i>	DIC	1	0.0043	0.0043
		DOC	1	0.0039	0.0039	0.98	0.3506
		DIC x DOC	1	0.0048	0.0048	1.21	0.3034
		Residual	8	0.0316	0.0039		
		<i>U. flabellum</i>	DIC	1	0.0061	0.0061	1.14
		DOC	1	0.0016	0.0016	0.30	0.5987
		DIC x DOC	1	0.0010	0.0010	0.19	0.6689
		Residual	8	0.0425	0.0053		
		Light calcification (G_{max})	<i>H. incrassata</i>	DIC	1	1.16	1.16
		DOC	1	0.495	0.495	9.98	0.0134*
		DIC x DOC	1	4.38	4.38	88.28	0.000013*
		Residual	8	0.397	0.0496		
		<i>U. flabellum</i>	DIC	1	0.018	0.018	6.184
		DOC	1	0.399	0.399	140.94	0.000002*
		DIC x DOC	1	0.002	0.002	0.778	0.4034
		Residual	8	0.023	0.0028		

Significant results ($p < 0.05$) are marked in bold and with an asterisk (*).

doi:10.1371/journal.pone.0160268.t003

the antagonistic effect between elevated DIC and DOC and their combined effect on the calcification process of *H. incrassata* (Table 3).

Additionally to the physiological responses of the organisms, measurements of BOD for the different treatment waters were performed to determine potential changes in bacterial respiration. Although increased BOD was detected in the treatments with elevated DOC, these changes were not significant (Table 1).

Discussion

Large differences between the two species investigated were found in photosynthesis and calcification rates, in agreement with previous findings [41]. Our study further revealed significant differences between *Halimeda incrassata* and *Udotea flabellum* in the calcification:photosynthesis ratio (G_{\max}/P_{\max}), as *H. incrassata* was able to precipitate twice as much CaCO_3 per mol O_2 evolved in photosynthesis than *U. flabellum*.

Elevated DIC and DOC treatments caused adverse impacts on the physiology of both species, but significant differences were observed in the severity of this impact. For example, elevated DIC resulted in a decline in F_v/F_m and photosynthesis rates in both species. However, elevated DOC only caused a similar response in *H. incrassata*, as non-significant changes were observed for the organisms of *U. flabellum* exposed to the similar treatment. Control organisms did not show the progressive reduction in F_v/F_m observed for organisms exposed to DIC and DOC enrichments throughout the experiment. This lack of change in F_v/F_m after an initial reduction during the first four days, in spite of the maintenance of high light conditions for the last five experimental days, indicates that experimental conditions were optimal for both species and did not induce significant accumulation of photodamage (i.e., F_v/F_m decline), or positive F_v/F_m recovery due to light limitation. Thus, the observed reductions in F_v/F_m and thallus photosynthesis of the organisms exposed to elevated DOC and/or DIC can be attributed to a direct negative impact of these treatments on the photosynthetic process. Thallus photosynthesis in *U. flabellum* showed a more robust response to elevated DOC, while *H. incrassata* was equally sensitive to both organic and inorganic carbon enrichments. A similar negative impact of elevated DIC on algal photosynthesis has been previously reported for other species from the genus *Halimeda* [26, 29, 30], but the causes of this decline have not yet been elucidated. Price et al. [26] suggested that the increase in dissolved CO_2 under reduced seawater pH may affect the expression of different carbon-concentrating mechanisms (CCMs), causing algal photosynthesis to rely on passive CO_2 diffusion, and thus becoming more susceptible to photosynthesis carbon limitation. The maintenance of high proton- H^+ permeability of the plasma membrane, for example, which is key for photosynthetic bicarbonate assimilation [52, 53], declines at reduced external pH [54].

In addition to the impact of DIC on algal photosynthesis, we also found a negative effect of elevated DIC on calcification rates of both species. *Udotea flabellum* showed a similar -30% reduction in photosynthesis and calcification (-36%; Figs 2F and 3B), but *H. incrassata* experienced larger declines in calcification (-155%) compared to a -30% reduction in photosynthesis (Figs 3A and 2E). With respect to the response to elevated DOC concentration, a greater impact was observed on *H. incrassata* photosynthesis and calcification rates when acting in isolation. Negative effects of elevated DOC concentrations have been recently documented for the photosynthesis rates of two *Halimeda* species from the Great Barrier Reef [28]. In contrast to our findings, calcification under illumination was not significantly affected by elevated DOC in these species. Large differences for the response of thallus calcification to elevated DIC have been already documented among *Halimeda* spp. [26, 29, 30], and this is the first time that similar inter-specific differences were also observed for the response to elevated DOC. Some




Fig 3. Calcification (G_{max}) response of *H. incrassata* (a) and *U. flabellum* (b) to the experimental DIC and DOC treatments. Data represent mean \pm SE ($n = 3$) and significant differences between treatments (ANOVA, Newman-Keuls, $p < 0.05$) are indicated by different letters.

doi:10.1371/journal.pone.0160268.g003

authors have suggested that the large inter-specific component shown by the calcification process in the genus *Halimeda* may rely on thallus morphology [26]. This genus displays large variation in the internal anatomy of algal thallus, and these anatomical characteristics are good proxies for species membership when compared to molecular data [55], what may support our interpretation. However, more work is still needed to elucidate the potential implications of the variation in thallus anatomy within the *Halimeda* genus on the species-specific sensitivity of thallus calcification to environmental changes.

Photosynthesis and calcification rates are tightly coupled in calcareous siphonal algae. Photosynthesis promotes algal calcification by removing CO_2 or bicarbonate from the calcification site, which increases the local pH and, thus, facilitates CaCO_3 precipitation [56]. Photosynthesis can also support a high fraction of the energetic costs of the biomineralization process. Therefore, any negative effect on the photosynthetic process would be reflected in a decline in algal calcification, as recently shown for coralline algae [48]. Inter-specific differences in the calcification process may explain the diversity of responses observed. For example, while *H. incrassata* only calcifies in the intercellular spaces, calcification in *U. flabellum* represents a transition between intercellular and sheath mineralization (e.g. *Penicillus*, *Rhipocephalus*) [57]. The CaCO_3 precipitation in *H. incrassata* occurs in a semi-isolated space, where CO_2 diffusion from the external environment can cause a decrease in local pH, and thus a reduction in calcification rates. Therefore, a more efficient isolation from surrounding seawater of the biomineralization site of *U. flabellum*, allows carbonate precipitation to be less dependent on the external variation of DIC and, thus, better suitable to be controlled by the physiology of the organism. The occurrence of a stronger control over the CaO_3 precipitation process by *U. flabellum* is supported by the findings of Ries [41].

Calcareous green algae are able to release DOC, but cannot incorporate organic carbon [58–59]. Thus, although DOC decline in the enriched experimental treatments was primarily due to

water turnover rates in the tanks, part of this DOC enrichment could have likely been assimilated by bacteria (Fig 1). Furthermore, the experimental addition of DOC in the form of glucose stimulates microbial respiration and growth [60]. Such enhancement in bacterial activity explains the lower O₂ concentrations observed in DOC-treatments compared to control- and DIC-treatments, as reported previously [46, 61]. Little information is available about the interaction between these epibacterial communities and algal physiology, and the potential effects of environmental changes on these communities and their interactions (i.e., [62–64]). It has been documented for *Halimeda copiosa* that the abundance of thallus surface-associated bacteria increases under organic nutrient enrichments [65]. In addition to increases in bacterial abundance, shifts in the bacterial community towards non-beneficial or even harmful bacteria have been suggested to occur for corals under increasing DOC concentrations [8]. As no increases in algal respiratory rates were observed in the DOC-treatments (Fig 2E and 2F), the negative responses on photosynthetic and calcification rates were most likely related to alterations in the bacterial community than in their abundance. Benthic reef algae have been shown to differ in the microbial communities associated with their tissue [66], therefore, part of the observed differences in the DOC response of *H. incrassata* and *U. flabellum* might be related to differences in the response of their respective epibacterial communities to the experimental DOC enrichment, as well as species-specific effects on bacterial-algal interactions. The antagonistic effect found for the combined elevated DIC and DOC treatment in *H. incrassata*, could also be due to a differential effect of each factor on the bacterial-algal interactions. More studies focusing on the seaweed holobiont are necessary to fully understand the relevance of these indirect effects on algal performance.

Ecological Perspective

According to our results, the DIC concentrations expected by the year 2100 [45] may significantly reduce photosynthesis and carbonate production in *H. incrassata*, while *U. flabellum* production may experience relative lower declines. However, when accompanied by increased concentrations of high labile DOC, the impact of elevated DIC will be alleviated for *H. incrassata* but exacerbated for *U. flabellum*. Furthermore, the effect of DIC and DOC could be even more severe when considering their impact at night on algal calcification, as has been documented for dark calcification rates that are also negatively affected in *Halimeda* spp. [28, 67]. Thus, considering the impacts both during the light and night on thallus calcification rates, the net algal carbonate production can be reduced even further.

Potential sources of high labile DOC for this particular reef lagoon are the seagrass and macroalgal beds themselves [59], human waste water discharge via groundwater [68, 69], and storm events [70], which are all predicted to increase in the future providing more labile DOC to this coastal ecosystem. For the carbon budget of the Puerto Morelos reef lagoon [71], this DOC enrichment may lead to a significant reduction of the contribution of calcifying green algae to the overall primary production and/or carbonate reef accretion. This impact will produce severe consequences on the macrophyte community, habitat structure, and ultimately, on the organic carbon fluxes of the ecosystem due to altered contributions to labile DOC and POC pools [59]. Significant reductions in carbonate sand production from algal derived sediments can alter the volume of sand deposits in coastal tropical areas, with important consequences for beach erosion and coastal sediment dynamics of reef environments. On the other hand, considering that seagrasses and fleshy algae may prosper under higher DIC conditions [72–74], and that seagrasses can modulate the OA response of calcareous algae [75–77], a deeper understanding of the changes in the macrophyte community and on species interactions will be fundamental to enhance our capacity to foresee the severity of the impact of predicted environmental changes on carbonate sand production by calcareous green algae.

Acknowledgments

The authors would like to thank the staff of UNAM, Puerto Morelos, especially Román Vázquez Elizondo and Tim Scheufen, for field and laboratory support. In addition, we would like to thank Matthias Birkicht for the support in TA and TOC analysis. This study was funded by two research projects DGAPA (IN206710) and a SEP-CONACYT (129880) to SE and also by the Leibniz Association to CW. The MASTS pooling initiative (The Marine Alliance for Science and Technology for Scotland) with the grant reference HR09011 to KD is also gratefully acknowledged. Support to NS was provided through two post-doctoral fellowships: UNAM-DGAPA and SEP-CONACYT (129880). We thank Roberto Iglesias-Prieto for his valuable support and J. E. Escalante Mancera and F. G. Ruiz Rentería for providing SST data from the Oceanographically and Meteorological Monitoring Academic Service (UASA, ICMyL, UNAM). KD received funding from the MASTS pooling initiative (The Marine Alliance for Science and Technology for Scotland) and its support is gratefully acknowledged. MASTS is funded by the Scottish Funding Council (grant reference HR09011) and contributing institutions.

The experiments performed in this work comply with the current laws of Mexico. The study was supported by the Mexican permit: “*Permiso de Pesca de Fomento*” No DGOPA.07342.010810.4121, to SE, issued by the Secretaría de Agricultura, Ganadería, Desarrollo rural, Pesca y Alimentación of the United States of Mexico, to support the development of the project: “*Evaluación del efecto de la limitación de carbono sobre diferentes productores primarios de la laguna arrecifal de Puerto Morelos: Importancia de la calcificación*”.

Author Contributions

Conceived and designed the experiments: CW SE.

Performed the experiments: FWM NS.

Analyzed the data: FWM NS SE.

Contributed reagents/materials/analysis tools: CW SE.

Wrote the paper: FWM NS CW SE.

Co-supervisors of the PhD of FWM: KD MT.

Provided interesting suggestions to the final manuscript to improve the text: KD MT.

References

1. Zeebe RE, Wolf-Gladrow DA (2001) CO₂ in seawater: equilibrium, kinetics, isotopes. Elsevier Science. Available: http://books.google.com/books?hl=en&lr=&id=VrumU3XvQ-gC&oi=fnd&pg=PP2&dq=Seawater:+Equilibrium,+Kinetics,+Isotopes,+&ots=0JJD0a2k3_&sig=6G0E8INEEUyMZQByGa-CV_RNao.
2. Doney SC, Fabry VJ, Feely RA, Kleypas JA. (2009) Ocean Acidification: The Other CO₂ Problem. *Ann Rev Mar Sci* 1: 169–192. Available: <http://www.annualreviews.org/eprint/QwPqRGcRzQM5ffhPjAdT/full/10.1146/annurev.marine.010908.163834>. PMID: 21141034
3. Manzello D, Kleypas J (2008) Poorly cemented coral reefs of the eastern tropical Pacific: Possible insights into reef development in a high-CO₂ world. *Proc Natl Acad Sci USA* 105: 10450–10455. Available: <http://www.pnas.org/content/105/30/10450.abstract>. doi: 10.1073/pnas.0712167105 PMID: 18663220
4. Perry CT, Murphy GN, Kench PS, Smithers SG, Edinger EN, Steneck RS, et al. (2013) Caribbean-wide decline in carbonate production threatens coral reef growth. *Nature Comm* 4: 1402–1408. Available: <http://www.nature.com/ncomms/journal/v4/n1/abs/ncomms2409.html>.

5. Buddemeier RW, Kleypas JA, Aronson RB (2004) Potential contributions of climate change to stresses on coral reef ecosystems. Coral reefs and global climate change. Pew Center on Global Climate Change, Virginia, USA, 56 pp. Available: <http://www.c2es.org/publications/coral-reefs-amp-global-climate-change>.
6. Hughes T (1994) Catastrophes, phase shifts, and large-scale degradation of a Caribbean coral reef. *Sci Pap Ed* 265: 1547–1551. Available: http://bio.classes.ucsc.edu/bio160/Bio160readings/Catastrophes_PhaseShifts.pdf.
7. Mumby PJ (2009) Phase shifts and the stability of macroalgal communities on Caribbean coral reefs. *Coral Reefs* 28: 761–773. Available: <http://link.springer.com/10.1007/s00338-009-0506-8>.
8. Kuntz NM, Kline DI, Sandin SA, Rohwer F (2005) Pathologies and mortality rates caused by organic carbon and nutrient stressors in three Caribbean coral species. *Mar Ecol Prog Ser* 294: 173–180. Available: <http://www.int-res.com/abstracts/meps/v294/p173-180/>.
9. Kline DI, Kuntz NM, Breitbart M, Knowlton N, Rohwer F (2006) Role of elevated organic carbon levels and microbial activity in coral mortality. *Mar Ecol Prog Ser* 314: 119–125. Available: <http://www.int-res.com/abstracts/meps/v314/p119-125/>.
10. Haas AF, Nelson C, Kelly L, Carlson C (2011) Effects of coral reef benthic primary producers on dissolved organic carbon and microbial activity. *PLoS One* 6(11): e27973. Available: <http://dx.plos.org/10.1371/journal.pone.0027973>. doi: [10.1371/journal.pone.0027973](https://doi.org/10.1371/journal.pone.0027973) PMID: [22125645](https://pubmed.ncbi.nlm.nih.gov/22125645/)
11. Smith JE, Shaw M, Edwards RA, Obura DO, Pantos O, et al. (2006) Indirect effects of algae on coral: algae-mediated, microbe-induced coral mortality. *Ecol Lett* 9: 835–845. Available: <http://www.ncbi.nlm.nih.gov/pubmed/16796574>. PMID: [16796574](https://pubmed.ncbi.nlm.nih.gov/16796574/)
12. Gregg A, Hatay M, Haas AF, Robinett N, Barott KL, et al. (2013) Biological oxygen demand optode analysis of coral reef-associated microbial communities exposed to algal exudates. *PeerJ* 1: e107. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3719127&tool=pmcentrez&rendertype=abstract>. doi: [10.7717/peerj.107](https://doi.org/10.7717/peerj.107) PMID: [23882444](https://pubmed.ncbi.nlm.nih.gov/23882444/)
13. Haas AF, Wild C (2010) Composition analysis of organic matter released by cosmopolitan coral reef-associated green algae. *Aquat Biol* 10. Available: <http://www.int-res.com/abstracts/ab/v10/n2/p131-138/>.
14. Nelson CE, Goldberg SJ, Wegley-Kelly L, Haas AF, Smith JE, Rohwer F, et al. (2013) Coral and macroalgal exudates vary in neutral sugar composition and differentially enrich reef bacterioplankton lineages. *ISME J* 7: 962–979. Available: <http://www.ncbi.nlm.nih.gov/pubmed/23303369>. doi: [10.1038/ismej.2012.161](https://doi.org/10.1038/ismej.2012.161) PMID: [23303369](https://pubmed.ncbi.nlm.nih.gov/23303369/)
15. Haas AF, Nelson CE, Rohwer F, Wegley-Kelly L, Quistad SD, et al. (2013) Influence of coral and algal exudates on microbially mediated reef metabolism. *PeerJ* 1: e108. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3719129&tool=pmcentrez&rendertype=abstract>. doi: [10.7717/peerj.108](https://doi.org/10.7717/peerj.108) PMID: [23882445](https://pubmed.ncbi.nlm.nih.gov/23882445/)
16. Collado-Vides L, Rutten LM, Fourqurean JW (2005) Spatiotemporal variation of the abundance of calcareous green macroalgae in the Florida Keys: A study of synchrony within a macroalgal functional-form group. *J Phycol* 41: 742–752. Available: <http://onlinelibrary.wiley.com/doi/10.1111/j.1529-8817.2005.00099.x/epdf>.
17. Cruz-Palacios V, van Tussenbroek BI (2005) Simulation of hurricane-like disturbances on a Caribbean seagrass bed. *J Exp Mar Biol Ecol* 324:44–60. Available: <http://www.sciencedirect.com/science/article/pii/S002209810500170X>.
18. Rodríguez-Martínez RE, Ruíz-Rentería F, van Tussenbroek B, Barba-Santos G, Escalante-Mancera E, et al. (2010) Environmental state and tendencies of the Puerto Morelos CARICOMP site, Mexico. *Rev Biol Trop* 58 Suppl 3: 23–43. Available: <http://www.ncbi.nlm.nih.gov/pubmed/21299094>. PMID: [21299094](https://pubmed.ncbi.nlm.nih.gov/21299094/)
19. Bach SD (1979) Standing crop, growth and production of calcareous Siphonales (Chlorophyta) in a South Florida lagoon. *Bull Mar Sci* 29: 191–201. Available: <http://www.ingentaconnect.com/contentone/umrsmas/bullmar/1979/00000029/00000002/art00005?crawler=true>.
20. Hillis L (1997) Coralgall reefs from a calcareous green alga perspective, and a first carbonate budget. 1, 761–766. *Proc. 8th Int. Coral Reef Symp.* Available: http://www.reefbase.org/resource_center/publication/pub_9604.aspx.
21. Neumann AC, Land LS (1975) Lime mud deposition and calcareous algae in the Bight of Abaco, Bahamas: A budget. *J Sediment Petrol* 45: 763–786. Available: <http://archives.datapages.com/data/sepm/journals/v42-46/data/045/045004/0763.htm>.
22. van Tussenbroek BI, van Dijk JK (2007) Spatial and Temporal Variability in Biomass and Production of Psammophytic *Halimeda Incrassata* (Bryopsidales, Chlorophyta) in a Caribbean Reef Lagoon. *J Phycol* 43: 69–77. Available: <http://doi.wiley.com/10.1111/j.1529-8817.2006.00307.x>.
23. Hofmann LC, Heiden J, Bischof K, Teichberg M (2014) Nutrient availability affects the response of the calcifying chlorophyte *Halimeda opuntia* (L.) J.V. Lamouroux to low pH. *Planta* 239: 231–242.

Available: <http://www.ncbi.nlm.nih.gov/pubmed/24158465>. doi: [10.1007/s00425-013-1982-1](https://doi.org/10.1007/s00425-013-1982-1) PMID: [24158465](https://pubmed.ncbi.nlm.nih.gov/24158465/)

24. Johnson MD, Price NN, Smith JE (2014) Contrasting effects of ocean acidification on tropical fleshy and calcareous algae. *PeerJ* 2: e411. Available: http://www.researchgate.net/publication/263014583_Contrasting_effects_of_ocean_acidification_on_tropical_fleshy_and_calcareous_algae. doi: [10.7717/peerj.411](https://doi.org/10.7717/peerj.411) PMID: [24918033](https://pubmed.ncbi.nlm.nih.gov/24918033/)
25. Peach KE, Koch M, Blackwelder P (2016) Effects of elevated $p\text{CO}_2$ and light on growth, photosynthesis and calcification in *Halimeda discoidea*. *Mar Ecol Prog Ser* 544: 143–158. Available: <http://www.int-res.com/articles/meps2016/544/m544p143.pdf>.
26. Price N, Hamilton S, Tootell J, Smith J (2011) Species-specific consequences of ocean acidification for the calcareous tropical green algae *Halimeda*. *Mar Ecol Prog Ser* 440: 67–78. Available: <http://www.int-res.com/abstracts/meps/v440/p67-78/>.
27. Campbell JE, Fisch J, Langdon C, Paul VJ (2015) Increased temperature mitigates the effects of ocean acidification in calcified green algae (*Halimeda* spp.). *Coral Reef* 35(1): 357–368. Available: <http://link.springer.com/article/10.1007%2Fs00338-015-1377-9>.
28. Meyer FW, Vogel N, Teichberg M, Uthicke S, Wild C (2015) The physiological response of two green calcifying algae from the Great Barrier Reef towards high dissolved inorganic and organic carbon (DIC and DOC) availability. *PlosOne* 10(8): e0133596. Available: <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0133596>.
29. Sinutok S, Hill R, Doblin MA, Wuhrer R, Ralph PJ (2011) Warmer more acidic conditions cause decreased productivity and calcification in subtropical coral reef sediment-dwelling calcifiers. *Limnol Oceanogr* 56: 1200–1212. Available: <http://cat.inist.fr/?aModele=afficheN&cpsid=24362064>.
30. Sinutok S, Hill R, Doblin MA, Kühl M, Ralph PJ (2012) Microenvironmental changes support evidence of photosynthesis and calcification inhibition in *Halimeda* under ocean acidification and warming. *Coral Reefs* 31: 1201–1213. Available: <http://link.springer.com/article/10.1007%2Fs00338-012-0952-6>.
31. Comeau S, Edmunds PJ, Spindel NB, Carpenter RC (2013) The responses of eight coral reef calcifiers to increasing partial pressure of CO_2 do not exhibit a tipping point. *Limnol Oceanogr* 58: 388–398. Available: <http://onlinelibrary.wiley.com/doi/10.4319/lo.2013.58.1.0388.pdf>.
32. Comeau S, Edmunds PJ, Spindel NB, Carpenter RC (2014) Fast coral reef calcifiers are more sensitive to ocean acidification in short-term laboratory incubations. *Limnol Oceanogr* 59: 1081–1091. Available: <http://onlinelibrary.wiley.com/doi/10.4319/lo.2014.59.3.1081/>.
33. Campbell J, Craft J, Muehlehner N, Langdon C, Paul V (2014) Responses of calcifying algae (*Halimeda* spp.) to ocean acidification: implications for herbivores. *Mar Ecol Prog Ser* 514: 43–56. Available: <http://www.int-res.com/abstracts/meps/v514/p43-56/>.
34. Robbins LL, Knorr PO, Hallock P (2009) Response of *Halimeda* to ocean acidification: field and laboratory evidence. *Biogeosciences Discuss* 6: 4895–4918. Available: <http://www.biogeosciences-discuss.net/6/4895/2009/>.
35. Wizemann A, Meyer FW, Hofmann LC, Wild C, Westphal H (2015) Ocean acidification alters the calcareous microstructure of the green macro-alga *Halimeda opuntia*. *Coral Reefs* 34: 941–954. Available: <http://link.springer.com/10.1007/s00338-015-1288-9>.
36. Littler MM, Littler DS, Lapointe BE (1988) A comparison of nutrient- and light-limited photosynthesis in psammophytic versus epilithic forms of *Halimeda* (Caulerpales, Halimediaceae) from the Bahamas. *Coral Reefs* 6: 219–225. Available: <http://link.springer.com/article/10.1007/BF00302018>.
37. Delgado O, Lapointe BE (1994) Nutrient-limited productivity of calcareous versus fleshy macroalgae in a eutrophic, carbonate-rich tropical marine environment. *Coral Reefs* 13:151–159. Available: <http://link.springer.com/article/10.1007%2FBF00301191>.
38. Hofmann LC, Bischof K, Baggini C, Johnson A, Koop-Jakobsen K, Teichberg M (2015) CO_2 and inorganic nutrient enrichment affect the performance of a calcifying green alga and its noncalcifying epiphyte. *Oecologia* 177: 1157–1169. Available: <http://link.springer.com/article/10.1007%2Fs00442-015-3242-5>. doi: [10.1007/s00442-015-3242-5](https://doi.org/10.1007/s00442-015-3242-5) PMID: [25648647](https://pubmed.ncbi.nlm.nih.gov/25648647/)
39. van Tussenbroek BI (1995) *Thalassia testudinum* leaf dynamics in a Mexican Caribbean coral reef lagoon. *Mar Biol* 122: 33–40. Available: <http://link.springer.com/article/10.1007%2FBF00349275>.
40. Enríquez S, Schubert N (2014) Direct contribution of the seagrass *Thalassia testudinum* to lime mud production. *Nature Comm* 5: 3835. Available: <http://www.nature.com/ncomms/2014/140522/ncomms4835/full/ncomms4835.html>.
41. Ries JB (2009) Effects of secular variation in seawater Mg/Ca ratio (calcite-aragonite seas) on CaCO_3 sediment production by the calcareous algae *Halimeda*, *Penicillus* and *Udotea*—evidence from recent experiments and the geological record. *Terra Nova* 21: 323–339. Available: <http://onlinelibrary.wiley.com/doi/10.1111/j.1365-3121.2009.00899.x/epdf>.

42. Mumby PJ, Flower J, Chollett I, Box SJ, Bozec Y-M, Fitzsimmons C, et al. (2014) Towards Reef Resilience and Sustainable Livelihoods: A handbook for Caribbean coral reef managers. University of Exeter, Exeter. 172 pages.
43. Enríquez S, Pantoja-Reyes NI (2005) Form-function analysis of the effect of canopy morphology on leaf self-shading in the seagrass *Thalassia testudinum*. *Oecologia* 145: 235–243. Available: <http://link.springer.com/article/10.1007%2Fs00442-005-0111-7>. PMID: 15942763
44. Cayabyab NM, Enríquez S (2007) Leaf acclimatory responses of the tropical seagrass *Thalassia testudinum* under mesocosm conditions: a mechanistic scaling-up study. *New Phytol* 176: 108–123. Available: <http://onlinelibrary.wiley.com/doi/10.1111/j.1469-8137.2007.02147.x/full>. PMID: 17696981
45. Stocker TF, Qin D, Plattner G-K, Tignor MMB, Allen SK, Bischung J, et al. (2013) Climate Change 2013. The physical science basis. Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Available: <http://www.ipcc.ch/report/ar5/wg1/>.
46. Haas AF, Al-Zibdah M, Wild C (2009) Effect of inorganic and organic nutrient addition on coral–algae assemblages from the Northern Red Sea. *J Exp Mar Bio Ecol* 380: 99–105. Available: <http://linkinghub.elsevier.com/retrieve/pii/S0022098109003712>.
47. Orr JC, Epitalon J-M, Gattuso J-P (2014) Comparison of seven packages that compute ocean carbonate chemistry. *Biogeosciences Discuss* 11: 5327–5397. Available: <http://www.biogeosciences-discuss.net/11/5327/2014/>.
48. Vásquez-Elizondo RM, Enríquez S (2016) Coralline algal physiology is more adversely affected by elevated temperature than reduced pH. *Sci Rep* 6:19030. Available: <http://www.nature.com/articles/srep19030>. doi: 10.1038/srep19030 PMID: 26740396
49. Drew EA, Abel KM (1990) Studies on Halimeda. III. A Daily Cycle of Chloroplast Migration within Segments. *Bot Mar* 33: 31–46. Available: <http://www.reference-global.com/doi/abs/10.1515/botm.1990.33.1.31>.
50. Smith S, Kinsey D (1978) Calcification and organic carbon metabolism as indicated by carbon dioxide. *Coral reefs Res methods*, pp. 469–484 Available: [http://scholar.google.com/scholar_lookup?title=Calcification and organic carbon metabolism as indicated by carbon dioxide&author=Smith&publication_year=1978#0](http://scholar.google.com/scholar_lookup?title=Calcification+and+organic+carbon+metabolism+as+indicated+by+carbon+dioxide&author=Smith&publication_year=1978#0).
51. Yao W, Byrne RH (1998) Simplified seawater alkalinity analysis. *Deep Sea Res Part I Oceanogr Res Pap* 45: 1383–1392. Available: <http://www.sciencedirect.com/science/article/pii/S0967063798000181>.
52. Prins HBA, Snel JFH, Zanstra PE, Helder RJ (1982) The mechanism of bicarbonate assimilation by the polar leaves of *Potamogeton* and *Elodea*. CO₂ concentrations at the leaf surface. *Plant Cell Environ.* 5: 207–214.
53. Elzenga JTM, Prins HBA (1989) Light-induced polar pH changes in leaves of *Elodea canadensis*. I. Effects of carbon concentration and light intensity. *Plant Physiol* 91: 62–67. PMID: 16667044
54. Miedema H, Prins HBA (1991) pH-dependent proton permeability of the plasma membrane is a regulating mechanism of polar transport through the submerged leaves of *Potamogeton lucens*. *Can J Bot* 69: 1116–1122.
55. Verbruggen H, De Clerck O, Cocquyt E, Kooistra WHCF, Coppejans E (2005) Morphometric taxonomy of siphonous green algae: A methodological study within the genus Halimeda (Bryopsidales). *J Phycol* 41: 126–139. Available: [https://www.researchgate.net/profile/Wiebe_Kooistra/publication/32251297_Morphometric_Taxonomy_of_Siphonous_Green_Algae_A_Methodological_Study_within_the_Genus_Halimeda_\(Bryopsidales\)/links/02bfe50ea8a1287b0e000000.pdf](https://www.researchgate.net/profile/Wiebe_Kooistra/publication/32251297_Morphometric_Taxonomy_of_Siphonous_Green_Algae_A_Methodological_Study_within_the_Genus_Halimeda_(Bryopsidales)/links/02bfe50ea8a1287b0e000000.pdf).
56. Borowitzka MA, Larkum AWD (1976) Calcification in the Green Alga *Halimeda*. *J Exp Bot* 27: 894–907. Available: <http://jxb.oxfordjournals.org/content/27/5/894.short>.
57. Böhm L, Fütterer D, Kaminski E (1978) Algal calcification in some Codiaceae (Chlorophyta): Ultrastructure and location of skeletal deposits. *J Phycol* 14: 486–493. Available: <http://onlinelibrary.wiley.com/doi/10.1111/j.1529-8817.1978.tb02474.x>.
58. Wild C, Haas AF, Naumann MS, Mayr C, El-Zibdah M (2008) Comparative investigation of organic matter release by corals and benthic reef algae—implications for pelagic and benthic microbial metabolism. 11th International Coral Reef Symposium. pp. 7–11. Available: <http://www.nova.edu/ncri/11icrs/proceedings/files/m25-10.pdf>.
59. Haas AF, Jantzen C, Naumann MS, Iglesias-Prieto R, Wild C (2010) Organic matter release by the dominant primary producers in a Caribbean reef lagoon: implication for in situ O₂ availability. *Mar Ecol Prog Ser* 409: 27–39. doi: 10.3354/meps08631
60. Cole JJ (1982) Interactions Between Bacteria and Algae in Aquatic Ecosystems. *Ann Rev Ecol Sys* 13: 291–314. Available: <http://www.annualreviews.org/doi/abs/10.1146/annurev.es.13.110182.001451>.
61. Cárdenas A, Meyer FW, Schwieder H, Wild C, Gärdes A (2015) The formation of aggregates in coral reef waters under elevated concentrations of dissolved inorganic and organic carbon: A mesocosm

- approach. *Mar Chem* 175: 47–55. Available: <http://www.sciencedirect.com/science/article/pii/S0304420315000754>.
62. Goecke F, Labes A, Wiese J, Imhoff JF (2010) Chemical interactions between marine macroalgae and bacteria. *Mar Ecol Prog Ser* 409: 267–300. Available: <http://www.int-res.com/articles/meps2010/409/m409p267.pdf>.
 63. Egan S, Harder T, Burke C, Steinberg P, Kjellberg S, Thomas T (2013) The seaweed holobiont: understanding seaweed-bacteria interactions. *FEMS Microbiol Rev* 37: 462–476. Available: <http://hdl.handle.net/10453/23947>. doi: [10.1111/1574-6976.12011](https://doi.org/10.1111/1574-6976.12011) PMID: [23157386](https://pubmed.ncbi.nlm.nih.gov/23157386/)
 64. Singh RP, Reddy CRK (2013) Seaweed-microbial interactions: key functions of seaweed-associated bacteria. *FEMS Microbiol Ecol* 88: 213–230. Available: <http://femsec.oxfordjournals.org/content/femsec/88/2/213.full.pdf>.
 65. Jensen PR, Kauffman CA, Fenical W (1996) High recovery of culturable bacteria from the surfaces of marine algae. *Mar Biol* 126: 1–7. Available: <http://link.springer.com/article/10.1007%2FBF00571371>.
 66. Barott KL, Rodríguez-Brito B, Janouskovec J, Marhaver KL, Smith JE, Keeling P, et al. (2011) Microbial diversity associated with four functional groups of benthic reef algae and the reef-building coral *Montastraea annularis*. *Environm Microbol* 13:1192–1204. Available: <http://onlinelibrary.wiley.com/doi/10.1111/j.1462-2920.2010.02419.x/epdf>.
 67. Vogel N, Meyer FW, Wild C, Uthicke S. (2015) Decreased light availability can amplify negative impacts of ocean acidification on calcifying coral reef organisms. *Mar Ecol Prog Ser* 521:4 9–61. Available: <http://www.int-res.com/abstracts/meps/v521/p49-61/>.
 68. Carruthers TJB, van Tussenbroek BI, Dennison WC (2005) Influence of submarine springs and wastewater on nutrient dynamics of Caribbean seagrass meadows. *Estuar Coast Shelf Sci* 64: 191–199. Available: <http://www.sciencedirect.com/science/article/pii/S0272771405000491>.
 69. Null KA, Knee KL, Crook ED, de Sieyes NR, Rebolledo-Vieyra M, et al. (2014) Composition and fluxes of submarine groundwater along the Caribbean coast of the Yucatan Peninsula. *Cont Shelf Res* 77: 38–50. Available: <http://www.sciencedirect.com/science/article/pii/S0278434314000247>.
 70. Naumann MS, Haas AF, Jantzen C, Iglesias-Prieto R, Wild C (2012) Benthic-pelagic coupling in a Caribbean reef lagoon affected by hurricane "Dolly". *Proc 12th Int Coral Reef Symp* 12. Available: http://www.icrs2012.com/proceedings/manuscripts/ICRS2012_4C_4.pdf.
 71. Naumann MS, Jantzen C, Haas AF, Iglesias-Prieto R, Wild C (2013) Benthic primary production budget of a Caribbean reef lagoon (Puerto Morelos, Mexico). *PLoS One* 8: e82923. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3867400&tool=pmcentrez&rendertype=abstract>. doi: [10.1371/journal.pone.0082923](https://doi.org/10.1371/journal.pone.0082923) PMID: [24367570](https://pubmed.ncbi.nlm.nih.gov/24367570/)
 72. Hall-Spencer JM, Rodolfo-Metalpa R, Martin S, Ransome E, Fine M, et al. (2008) Volcanic carbon dioxide vents show ecosystem effects of ocean acidification. *Nature* 454: 96–99. Available: <http://www.ncbi.nlm.nih.gov/pubmed/18536730>. doi: [10.1038/nature07051](https://doi.org/10.1038/nature07051) PMID: [18536730](https://pubmed.ncbi.nlm.nih.gov/18536730/)
 73. Kroeker KJ, Kordas RL, Crim RN, Singh GG (2010) Meta-analysis reveals negative yet variable effects of ocean acidification on marine organisms. *Ecol Lett* 13: 1419–1434. Available: doi: <http://onlinelibrary.wiley.com/doi/10.1111/j.1461-0248.2010.01518.x/pdf> PMID: [20958904](https://pubmed.ncbi.nlm.nih.gov/20958904/)
 74. Kroeker KJ, Kordas RL, Crim R, Hendriks IE, Ramajo L, Singh GS, et al. (2013) Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. *Global Change Biol* 19: 1884–1896. Available: <http://onlinelibrary.wiley.com/doi/10.1111/gcb.12179/epdf>.
 75. Semesi IS, Beer S, Björk M (2009) Seagrass photosynthesis controls rates of calcification and photosynthesis of calcareous macroalgae in a tropical seagrass meadow. *Mar Ecol Prog Ser* 382: 41–47. Available: <http://www.int-res.com/abstracts/meps/v382/p41-47/>.
 76. Barry SC, Frazer TK, Jacoby CA (2013) Production and carbonate dynamics of *Halimeda incrassata* (Ellis) Lamouroux altered by *Thalassia testudinum* Banks and Soland ex König. *J Exp Mar Biol Ecol* 444:73–80. Available: <http://www.sciencedirect.com/science/article/pii/S0022098113001202>.
 77. Hendriks IE, Olsen YS, Ramajo L, Basso L, Steckbauer A, Moore TS, et al (2014) Photosynthetic activity buffers ocean acidification in seagrass meadows. *Biogeosciences* 11:333–346. Available: <http://www.biogeosciences.net/11/333/2014/bg-11-333-2014.pdf>.