

## An Epidemiological Study of *Hypoderma* Infection and Control Using Ivermectin in Yaks in Qinghai Province, China

Wei LI<sup>1</sup>, Yong FU<sup>1</sup>, Hong DUO<sup>1</sup>, Zhihong GUO<sup>1,2</sup>, Xiuying SHEN<sup>1</sup>, Fuqiang HUANG<sup>1</sup>, Kai FENG<sup>1</sup>, Zhisheng DANG<sup>1</sup>, Peng MAO<sup>1</sup>, Fang WANG<sup>3</sup>, Tetsuo NASU<sup>3</sup> and Nariaki NONAKA<sup>2</sup>\*

<sup>1</sup>Academy of Animal and Veterinary Medicine, University of Qinghai, No. 1, Weier road, Sci-biological Industry Areas, Xining, Qinghai, The People of Republic of China

<sup>2</sup>Laboratory of Veterinary Parasitic Diseases, Interdisciplinary Graduate School of Medicine and Veterinary Medicine, University of Miyazaki, 1-1 Gakuen-Kibanadai-Nishi, Miyazaki 889-2192, Japan

<sup>3</sup>Laboratory of Veterinary Anatomy, Department of Veterinary Sciences, Faculty of Agriculture, University of Miyazaki, 1-1 Gakuen-Kibanadai-Nishi, Miyazaki 889-2192, Japan

(Received 10 June 2013/Accepted 26 September 2013/Published online in J-STAGE 10 October 2013)

**ABSTRACT.** The prevalence of *Hypoderma* spp. in yaks grazed in the east of Qinghai province was investigated in 2008. In this area, the prevalence in young yaks (1- to 3-year-old) was very high at 82.2–98.7%, whilst in adult yaks (4-year-old and older), the prevalence was 42.4–50.6%. The seasonal development and migration pattern of *Hypoderma* larvae in yak bodies was found to be similar for different locations in this area. The numbers of first, second and third instar larvae detected in yak bodies peaked in October, December and March, respectively. Different doses of ivermectin (125 to 500 µg/kg body weight) almost completely dewormed the larvae from yaks, suggesting that using a quarter of the prescribed dose (500 µg/kg body weight) was effective. In October of each year between 2009 and 2012, ivermectin (125 µg/kg body weight) was administered to a total of 562,995 yaks grazed in four counties in Qinghai province, and the prevalence of *Hypoderma* larval infection in yaks was reduced to 0.5–1.0%.

**KEY WORDS:** China, *Hypoderma*, ivermectin, Qinghai, yak.

doi: 10.1292/jvms.13-0299; *J. Vet. Med. Sci.* 76(2): 183–188, 2014

Qinghai province is located in the northeast Qinghai-Tibetan plateau in west China. Since the average elevation is higher than 3,000 meters, nomadic grazing of high-altitude adaptive animals, such as yak and Tibetan sheep, is very popular. In this province, more than 4.8 million yaks are grazed at an elevation of 3,000 to 4,500 m.

Warble flies (*Hypoderma* spp.) are commonly found in yaks grazed in the province. Hypodermosis is a serious disease affecting livestock throughout the world, and two species of *Hypoderma*, *H. bovis* and *H. lineatum*, mainly affect cattle [10]. However, in yaks, a third species, *H. sinense*, is also commonly found [12, 15]. The overall prevalence of *Hypoderma* larval infection in yak is around 70–95%, reaching 100% in some areas [24]. Hypodermosis has also been found in cattle, wild animals, such as pikas [23] and chirus (Tibetan antelopes) [6, 8, 17], and occasionally in sheep, deer and horses in Qinghai province [24]. Moreover, infection in humans, especially in nomads who have regular contact with yaks, has been frequently recognized [24]. We previously

conducted a preliminary control study in Qinghai province to compare the effect of ivermectin administration (500 µg/kg body weight) to yaks in different months and found that treatment of yaks in October was most effective [13].

Hypodermosis is a subcutaneous myiasis which causes significant economic losses in the animal production industry [2, 4, 10] due to reduced weight gain, milk production and leather quality [24]. Even though the economic and zoonotic problems of warble flies are quite significant, systematic control measures against the parasite infection have not been used in Qinghai province. To establish a potential control scheme, we investigated the seasonal migration pattern of *Hypoderma* larvae in yaks in different locations in the province and conducted a large-scale anthelmintic control program targeting the first stage larvae in four counties (Maqin, Yushu, Guinan and Haiyan counties) in Qinghai province between 2008 and 2012.

### MATERIALS AND METHODS

**Study sites:** The study sites were Yushu county at Yushu state and Maqin county at Guoluo state located in southeast Qinghai; Guinan county at Hainan state in middle-east Qinghai; and Haiyan county at Haibei state in Qinghai Lake area (Fig. 1).

**Investigation of the prevalence of hypodermosis in yaks:** The investigation was carried out in Yushu, Maqin and Haiyan counties between March and May in 2008. The skin surfaces of the neck, shoulder, chest and back of 25,769 live yaks were examined for *Hypoderma* lumps by hand

\*CORRESPONDENCE TO: NONAKA, N., Laboratory of Veterinary Parasitic Diseases, Interdisciplinary Graduate School of Medicine and Veterinary Medicine, University of Miyazaki, 1-1 Gakuen-Kibanadai-Nishi, Miyazaki 889-2192, Japan.  
e-mail: nnonaka@cc.miyazaki-u.ac.jp

©2014 The Japanese Society of Veterinary Science

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives (by-nc-nd) License <<http://creativecommons.org/licenses/by-nc-nd/3.0/>>.



Fig. 1. The study sites in Qinghai, China.

palpation. The lumps were counted, and the prevalence of *Hypoderma* larval infection and the mean number of lumps in different age groups (1-, 2- and 3-year-old and adult ( $\geq 4$ -year-old)) of yaks were evaluated for each county.

*Investigation of seasonal pattern of Hypoderma larval development and migration in yak bodies:* The investigation was performed in two counties. One was Yushu county with an average elevation of 4,000 m, and the other was Haiyan county with an average elevation of 3,000 m. Every month from April 2008 to March 2009, five one-year-old yaks were selected from the herd grazed in the counties and autopsied to check for *Hypoderma* larvae in visceral organs, such as esophagus, rumen, jejunum, colon, diaphragm, spinal canal and in the skin of back. The larvae collected were fixed in 10% formalin and counted, and their instar larval stage was determined as previously described [10, 15].

*Determination of an effective dose of ivermectin for Hypoderma larvae:* In Yushu, Maqin and Haiyan counties, different doses of ivermectin (Beijing Zhongnong Huawei Pharmaceutical Co., Ltd., Beijing, China) were administered to yaks younger than 2-year-old in October 2008. A liquid ivermectin formula was put into a syringe and splashed onto the yaks back. The yaks were divided into four groups (1,000–1,500 head/group in each county); the full dose group was given a single treatment with 500  $\mu\text{g}/\text{kg}$  body weight (bw) which is the dose recommended for the treatment of *Hypoderma* larvae by the commercial suppliers; the half dose group was given 250  $\mu\text{g}/\text{kg}$  bw, the quarter dose group was given 125  $\mu\text{g}/\text{kg}$  bw and the control group remained untreated. Between February and April 2009, the yak's skin surface was hand-palpated to check and count *Hypoderma* lumps. The neck, shoulder, chest and back were examined.

*Conducting a large-scale control trial:* Between 2009 and 2011, a large-scale control trial against *Hypoderma* larval

infection in yaks was carried out in four counties (Maqin, Yushu, Guinan and Haiyan). The number of yaks treated was 562,995. In each county, four age groups (1-, 2- and 3-year-old and adult ( $\geq 4$ -year-old)) of yaks were treated with 125  $\mu\text{g}/\text{kg}$  bw of ivermectin by splashing onto the back of yaks in October. Then, in March and May of the next year, approximately 10% of the treated yaks were selected randomly, and their skin surfaces were hand-palpated to check and count *Hypoderma* lumps. The neck, shoulder, chest and back were examined. The proportion of yaks used for this study was about 2 to 5% of the total population of the grazed yak herd in each study site.

*Statistics:* Ninety-five percent confidence intervals (95% CI) of prevalence were calculated on the basis of binomial distributions using the software program R [19].

## RESULTS

*Prevalence of hypodermosis in yaks:* The prevalence in different age groups of yaks in Yushu, Maqin and Haiyan counties is shown in Table 1. All counties showed a similar pattern of age difference in prevalence. In all counties, the prevalence in 1- and 2-year-old yaks was very high (more than 94%). In comparison, the prevalence in adult yaks was significantly lower (42 to 51%). The number of *Hypoderma* lumps detected per yak was also higher in young (1- to 3-year-old) yaks than that in adult yaks. It is recognized that the prevalence and the mean number of lumps were relatively higher in Haiyan county than other counties.

*Seasonal pattern of Hypoderma larval development and migration in the yak body:* The number of larvae and their instar larval stages in the yaks at Yushu and Haiyan counties in each month from April 2008 to March 2009 are shown in Table 2. A quite similar pattern was observed in both

Table 1. Prevalence of *Hypoderma* larval infection in yaks of different ages in three counties of Qinghai

County	Age group	No. examined	No. infected	Prevalence (%) (95% CI <sup>a</sup> )	Mean No. lumps <sup>b</sup>
Yushu	1-year-old	1,978	1,902	96.2 (95.2–96.9)	13.5
	2-year-old	2,246	2,124	94.6 (93.6–95.4)	11.2
	3-year-old	2,580	2,156	83.6 (82.1–84.9)	10.9
	Adult	3,264	1,430	43.8 (42.1–45.5)	9.8
	Total	10,068	7,612	75.6 (74.8–76.4)	11.4
Maqin	1-year-old	2,007	1,911	95.2 (94.2–96.1)	12.4
	2-year-old	2,151	2,023	94.0 (93.3–95.0)	10.2
	3-year-old	2,080	1,710	82.2 (80.5–83.8)	11.4
	Adult	2,667	1,130	42.4 (40.5–44.3)	8.2
	Total	8,905	6,774	76.1 (75.2–76.9)	10.8
Haiyan	1-year-old	2,340	2,250	96.2 (95.3–96.9)	14.4
	2-year-old	1,880	1,855	98.7 (98.0–99.1)	15.5
	3-year-old	1,553	1,386	89.2 (87.6–90.7)	15.2
	Adult	1,023	518	50.6 (47.6–53.7)	9.8
	Total	6,796	6,009	88.4 (87.6–89.2)	14.5

a) 95% confidence interval. b) Mean number of *Hypoderma* lumps found per yak.

Table 2. Monthly change in the mean number of 1st, 2nd and 3rd instar *Hypoderma* larvae found in yaks in two counties of Qinghai between April 2008 and March 2009

Year	Month	Yushu county			Haiyan county		
		1st instar	2nd instar	3rd instar	1st instar	2nd instar	3rd instar
2008	Apr	0	0	45.8	0	0	46.6
	May	0	0	30.2	0	0	32.8
	June	0	0	2.2	0	0	3.4
	July	0	0	0	0	0	0
	Aug	26.2	0	0	25.0	0	0
	Sep	40.2	0	0	38.8	0	0
	Oct	116.0	0	0	102.4	0	0
	Nov	57.8	24.2	0	58.8	37.6	0
	Dec	17.6	63.2	0	17.2	65.4	0
2009	Jan	12.2	75	17.8	10.2	69.0	16.6
	Feb	2.8	37.8	46.2	3.6	39.8	51.6
	Mar	0	0	66.6	0	0	65.0

Table 3. The mean number of 1st instar *Hypoderma* larvae detected in various organs of yaks in Yushu and Haiyan counties of Qinghai between August 2008 and February 2009

Year	Month	Esophagus	Rumen	Small intestine	Colon	Diaphragm	Spinal canal
2008	Aug	8.8	7.6	4.6	3.2	0.5	0.9
	Sep	17.1	16.3	5.0	0.8		0.3
	Oct	56.2	53.0				
	Nov	31.5	26.8				
2009	Jan	9.9	7.5				
	Feb	7.3	3.9				

counties. First instar larvae appeared in the yak bodies in mid-August 2008 and were continually found until February 2009. The number of 1st instar larvae detected per yak was highest in October 2008 and then decreased gradually. The migration pattern of 1st instar larvae in yak bodies was quite similar in the two counties, and thus, the data are combined

and shown in Table 3. The majority of larvae were found in the subserosa of the esophagus and rumen. However, although restricted in August and September, parts of the larvae were also found in the subserosa of the small intestine and colon and in the diaphragm and spinal canal.

In November 2008, 2nd instar larvae first appeared on the

Table 4. The effect of administration of different doses of ivermectin on *Hypoderma* larval infection in yaks

County	Dose	No. treated	No. infected	Prevalence (%)	Mean No. lumps*
Yushu	500 µg/kg	1,100	0	0	0
	250 µg/kg	1,058	0	0	0
	125 µg/kg	1,350	2	0.2	1.5
	Control	1,064	992	93.2	12.9
Maqin	500 µg/kg	1,500	0	0	0
	250 µg/kg	1,258	0	0	0
	125 µg/kg	1,050	1	0.1	1.0
	Control	1,164	1,092	93.2	12.6
Haiyan	500 µg/kg	1,800	0	0	0
	250 µg/kg	1,558	0	0	0
	125 µg/kg	1,250	2	0.2	1.0
	Control	1,364	1,290	94.6	13.7

\*Mean number of *Hypoderma* lumps found per yak.

Table 5. Prevalence of *Hypoderma* larval infection and mean number of lumps found in different age groups of yaks after administration of ivermectin in four counties in Qinghai

County	Age group	No. treated	No. examined	No. infected	Prevalence (%)	Mean No. lumps*
Yushu	1-year-old	38,050	3,797	34	0.9	1.2
	2-year-old	37,542	3,916	35	0.9	1.1
	3-year-old	32,039	3,114	27	0.9	1.0
	Adult	25,873	2,561	22	0.9	1.0
Maqin	1-year-old	44,644	4,576	45	1.0	1.1
	2-year-old	40,021	4,083	35	0.9	1.4
	3-year-old	31,943	3,306	24	0.7	1.1
	Adult	24,434	2,592	14	0.5	1.1
Haiyan	1-year-old	42,316	4,483	37	0.8	1.1
	2-year-old	40,034	4,206	35	0.8	1.0
	3-year-old	30,618	3,158	19	0.6	1.0
	Adult	23,463	2,486	13	0.5	1.0
Guinan	1-year-old	51,722	5,282	43	0.8	1.2
	2-year-old	43,720	4,535	37	0.8	1.2
	3-year-old	31,934	3,266	24	0.7	1.4
	Adult	24,642	2,612	18	0.7	1.0

\*Mean number of *Hypoderma* lumps found per yak.

back of animal in both counties. The number of 2nd instar larvae detected per yak increased in December 2008, peaked in January 2009 and disappeared by March. All of the larvae were found in the subcutaneous tissue of the back (Table 2).

In January 2009, 3rd instar larvae appeared on the back of animals, and the number detected per yak peaked in March 2009. Since 3rd instar larvae were found from April to June 2008, it is suspected that they parasitize yaks between January and June (Table 2).

Third instar larvae could easily be distinguished from 2nd instar larvae by the body size and color. The 2nd instar larvae were 11–15 mm long and 3–6 mm wide, while 3rd instar larvae were 23–30 mm long and 12–15 mm wide. The body color of 2nd instar larvae was yellowish white, while that of 3rd instar larvae was yellowish to dark brown.

*An effective dose of ivermectin for Hypoderma larvae:* The effect of different doses of ivermectin administration on

*Hypoderma* larval infection in yaks is shown in Table 4. The trials conducted in all three counties (Yushu, Maqin and Haiyan) showed similar results. The prevalence in the control groups was very high (93.2–94.6%), and the mean number of lumps found was around 13. In contrast, no *Hypoderma* lumps were found in the groups treated with 500 and 250 µg/kg bw ivermectin, and hence, no infected animals were found. The prevalence in the groups treated with 125 µg/kg bw ivermectin was also extremely low (0.1–0.2%), and the mean number of lumps found was 1.0 to 1.5.

*A successful control of wable fly larval infection:* Cumulative data from the trial conducted over three years in four counties are shown in Table 5. There was no obvious difference in the annual data between 2009 and 2011 in the four counties. Some yaks remained infected in all counties and in all age groups, but the overall prevalence was <1.0% and the mean number of lumps found was below 1.5 at 5 to 7 months

post-administration of ivermectin. There was no obvious difference among the data obtained from different age groups or different counties.

## DISCUSSION

This survey was conducted in four counties located in the middle-east Qinghai to Qinghai lake area (Haiyan county and Guinan county) and southeast Qinghai area (Yushu county and Maqin county). These two areas are 1,000 km apart with average elevation of 3,000 and 4,000 m, respectively. They have different ecological environments with annual fluctuations in mean atmospheric pressure, temperature and precipitation that could affect the distribution of *Hypoderma*. However, we did not find any obvious differences in the prevalence of *Hypoderma* in yaks from these two areas. The results indicate that *Hypoderma* is distributed widely in the east Qinghai-Tibetan plateau.

It should be emphasized that the prevalence in young yaks was high (>94%) compared to adult yaks which was around 50%. The reason for this observation is not clear, but it could be partially explained by difference in the movement constraint used in young and adult yaks. Yak's milk is an important source of income for Tibetan nomadic people. Thus, young yaks are tethered to stakes so that they cannot move and drink milk freely from their mothers. In contrast, adult yaks were able to move freely and escape from adult warble fly attack by swishing their tails or running away. Therefore, it would be relatively difficult for warble flies to lay their eggs on adult yaks. Also, acquired resistance against *Hypoderma* infection may play a significant role in the infection in adult yaks [10, 16].

The seasonal development and migration patterns of *Hypoderma* larvae in yak bodies in the study areas were also quite similar. It has been reported that the 1st instar larvae of *H. lineatum* and *H. sinense* parasitized in the esophagus and those of *H. bovis* in the epidural fat of the spinal canal [2, 4, 15, 21]. However, as observed in our previous study [13], a large proportion of 1st instar larvae were found not only in the esophagus but also in the rumen. Moreover, the larvae were detected in other organs in late summer to early autumn (August and September). Unfortunately, species identification was not conducted in this study. Although morphological identification for the species of 1st instar larvae is impossible [15], DNA identification of species would help to understand the migration patterns of *H. bovis*, *H. lineatum* and *H. sinense* in yaks.

*Hypoderma* in the study areas showed the similar seasonal larval development, and therefore, ivermectin was administered at the same time (October) to yaks grazing the different sites. The timing of the ivermectin administration was chosen based on our previous research that showed that administration in October resulted in complete elimination of the larvae from yaks, but that administration in September or November resulted in incomplete elimination [13].

Ivermectin has been previously shown to be effective for deworming *Hypoderma* larvae [1, 7, 11, 14, 18, 20]. However, the continuous use or overuse of ivermectin could be

accompanied by side effects to the animals treated [5, 9], environmental pollution [3], induction of drug-resistant flies or impairment of human health as a result of consumption of meat containing the chemical [22]. Therefore, it is ideal to use the minimum effective dose of ivermectin with the minimum number of treatments per year [14]. This study showed that a single administration of full and half doses of ivermectin could completely eliminate the parasites from yaks. A single quarter dose administration was also quite effective and eliminated most of the larvae from yaks. Therefore, we determined that a single dose of 125  $\mu\text{g}/\text{kg}$  bw of ivermectin was an optimal dose for treatment.

As expected, a large-scale control trial conducted in this study was successful. Although complete elimination of *Hypoderma* larvae from treated yaks was not achieved and a small number of *Hypoderma* lumps remained in some yaks after treatment, the trial showed a high cost-performance ratio. We believe that the following factors may also have contributed to the successful control. Before implementation of this project, a training program for the Tibetan local nomads was conducted to teach the method and the appropriate time of year for administration of ivermectin to yaks. In addition, since restraining yaks is very difficult, administration of the drug by splashing onto the animals seemed to be a suitable method which is also less laborious, easier and safer than injection. In conclusion, the control method used in this study would be suitable for use in the Qinghai-Tibetan plateau to reduce economic losses due to *Hypoderma* infection and contribute to the improvement of animal health and subsequently human health.

We assumed multiple species of *Hypoderma* were detected in this study, because we had detected two species of *Hypoderma*, *H. bovis* and *H. lineatum*, in yaks at the same sites of this study in a previous study (unpublished). However, regrettably, we did not identify the species of *Hypoderma* detected in the present study. Because the difference in the species construction could affect the local epidemiology and larval migration pattern, species identification by morphological and DNA sequence characteristics should be incorporated into further studies.

**ACKNOWLEDGMENT.** This work was supported by the Strategic Japanese-Chinese Cooperative Program on "Research on health problems related to the environment" granted by the Japan Science and Technology Agency (JST) and the Ministry of Science and Technology of the People's Republic of China (MOST).

## REFERENCES

1. Alva-Valdes, R., Wallace, D. H., Holste, J. E., Egerton, J. R., Cox, J. L., Wooden, J. W. and Barrick, R. A. 1986. Efficacy of ivermectin in a topical formulation against induced gastrointestinal and pulmonary nematode infections, and naturally acquired grubs and lice in cattle. *Am. J. Vet. Res.* **47**: 2389–2392. [[Medline](#)]
2. Andrews, A. H. 1978. Warble fly: the life cycle, distribution, economic losses and control. *Vet. Rec.* **103**: 348–353. [[Medline](#)] [[CrossRef](#)]
3. Bloom, R. A. and Matheson, J. C. 1993. Environmental assess-

- ment of avermectins by the US Food and drug administration. *Vet. Parasitol.* **48**: 281–294. [Medline] [CrossRef]
4. Boulard, C. 2002. Durably controlling bovine hypodermosis. *Vet. Res.* **33**: 455–464. [Medline] [CrossRef]
  5. Button, C., Barton, R., Honey, P. and Rickford, P. 1988. Avermectin toxicity in calves and an evaluation of picrotoxin as an antidote. *Aust. Vet. J.* **65**: 157–158. [Medline] [CrossRef]
  6. Cai, J., Ma, Y., Yu, M., Wang, M., Li, C., Liu, S. and Wang, Y. 2006. *Hypoderma* larvae of Tibetan antelope infection status report. *Chin. J. Vet. Med.* **42**: 25–26 (in Chinese with English summary).
  7. De Chaneet, G. C., Casey, R., Dixon, F. F., Besier, R. B. and Mitchell, R. K. 1988. Effect of avermectin B1 and benzimidazole anthelmintics on worm egg output of treated cattle. *Aust. Vet. J.* **65**: 85–86. [Medline] [CrossRef]
  8. Gao, X., Cai, J., Xu, M., Yu, M., Cao, J., Li, C., Guo, M., Wang, Y., Zhu, S., Huang, Y. and He, G. 2008. Species identification of *H. sinense* infesting chiru by molecular characterization. *Chin. J. Vet. Sci.* **28**: 1037–1039 (in Chinese with English summary).
  9. Guzzo, C. A., Furtak, C. I., Porras, A. G., Chen, C., Tipping, R., Clineschmidt, C. M., Sciberras, D. G., Hsieh, J. Y. and Lasseter, K. C. 2002. Safety, tolerability, and pharmacokinetics of escalating high doses of ivermectin in healthy adult subjects. *J. Clin. Pharmacol.* **42**: 1122–1133. [Medline] [CrossRef]
  10. Hassan, M. U., Khan, M. N., Abubakar, M., Waheed, H. M., Iqbal, Z. and Hussain, M. 2010. Bovine hypodermosis—a global aspect. *Trop. Anim. Health Prod.* **42**: 1615–1625. [Medline] [CrossRef]
  11. Heinze-Mutz, E. M., Barth, D., Cramer, L. G., Gross, S. J. and Visser, M. 1993. Efficacy of abamectin against ectoparasites of cattle. *Vet. Rec.* **132**: 455–457. [Medline] [CrossRef]
  12. Li, W., Nasu, T., Ma, Y., Xiyan, Z. and Makimura, S. 2004. Scanning electron microscopic study of third-instar warbles in yak in China. *Vet. Parasitol.* **121**: 167–172. [Medline] [CrossRef]
  13. Li, W., Nasu, T., Ma, Y., Xiyan, Z., Rende, S. and Makimura, S. 2004. Migration of warble fly larvae in the yak and optimum timing of ivermectin treatment. *J. Vet. Med. Sci.* **66**: 891–892. [Medline] [CrossRef]
  14. Ma, M., Guan, G., Lu, B., Liu, A., Liu, Z., Chang, Z., Li, F., Chang, F., Luo, J., Lu, W., Zhang, Q., Yuan, G., Yin, H. and Boulard, C. 2003. Efficacy of different dosages of ivermectin injectable against the *Hypoderma* spp. in yaks. *Vet. Parasitol.* **117**: 147–151. [Medline] [CrossRef]
  15. Otranto, D., Paradies, P., Testini, G., Lia, R. P., Giangaspero, A., Traversa, D. and Colwell, D. D. 2006. First description of the endogenous life cycle of *Hypoderma sinense* affecting yaks and cattle in China. *Med. Vet. Entomol.* **20**: 325–328. [Medline] [CrossRef]
  16. Panadero, R., Dacal, V., López, C., Vázquez, L., Cienfuegos, S., Díaz, P., Morrondo, P. and Díez-Baños, P. 2009. Immunomodulatory effect of *Hypoderma lineatum* antigens: *in vitro* effect on bovine lymphocyte proliferation and cytokine production. *Parasite Immunol.* **31**: 72–77. [Medline] [CrossRef]
  17. Pang, C., Cai, J., Gao, X., Li, C., Cao, J., Xu, M., Liu, S., Huang, Y., Zhu, S., Boulard, C. and He, G. 2008. A fourth species of *Hypoderma* found from Tibetan antelope in China. *Acta Entomologica Sinica* **51**: 1099–1102 (in Chinese with English summary).
  18. Prichard, R. K. and Ranjan, S. 1993. Anthelmintics. *Vet. Parasitol.* **46**: 113–120. [Medline] [CrossRef]
  19. R Development Core Team. 2012. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna. ISBN 3-900051-07-0, Available from <http://www.R-project.org/>.
  20. Roncalli, R. A. and Usher, C. B. 1988. Efficacy of ivermectin against *Dermatobia hominis* in cattle. *Vet. Parasitol.* **28**: 343–346. [Medline] [CrossRef]
  21. Scholl, P. J. 1993. Biology and control of cattle grubs. *Annu. Rev. Entomol.* **38**: 53–70. [Medline] [CrossRef]
  22. Shoop, W. L., Haines, H. W., Michael, B. F. and Eary, C. H. 1993. Mutual resistance to avermectins and milbemycins: oral activity of ivermectin and moxidectin against ivermectin-resistant and susceptible nematodes. *Vet. Rec.* **133**: 445–447. [Medline] [CrossRef]
  23. Wang, C., He, H., Li, M., Lei, F., Root, J. J., Wu, Y. and Qin, J. 2009. Parasite species associated with wild plateau pika (*Ochotona curzoniae*) in southeastern Qinghai Province, China. *J. Wildl. Dis.* **45**: 288–294. [Medline] [CrossRef]
  24. Yin, H., Ma, M., Yuan, G., Huang, S., Liu, Z., Luo, J. and Guan, G. 2003. Hypodermosis in China. *J. Anim. Vet. Adv.* **2**: 179–183.