

UNCLASSIFIED

AD NUMBER
AD833372
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; MAR 1968. Other requests shall be referred to Department of the Army, Fort Detrick, Attn: Technical Releases Branch, Frederick, MD 21701.
AUTHORITY
Army Biological Defense Research Lab ltr dtd 29 Sep 1971

THIS PAGE IS UNCLASSIFIED

AD833372

AD

TECHNICAL MANUSCRIPT 429

INHIBITION OF STAPHYLOCOCCAL
ENTEROTOXIN B FORMATION
BY CELL WALL BLOCKING AGENTS
AND OTHER COMPOUNDS

Mischa E. Friedman

MARCH 1968

This document is intended for the use of the Department of the Army and is not to be distributed outside the Department of the Army. Reproduction of this document is prohibited without the express permission of the Department of the Army.

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland 21701

TECHNICAL MANUSCRIPT 429

INHIBITION OF STAPHYLOCOCCAL ENTEROTOXIN B FORMATION
BY CELL WALL BLOCKING AGENTS AND OTHER COMPOUNDS

Mischa E. Friedman

Medical Bacteriology Division
BIOLOGICAL SCIENCES LABORATORY

Project 1C522301A059

March 1968

ACKNOWLEDGMENT

The excellent technical assistance of Guy M. Dayhoff is acknowledged.

ABSTRACT

Enterotoxin B formation by Staphylococcus aureus S6 was inhibited by Tween 80, oleic acid, sodium deoxycholate, penicillin, D-cycloserine, or bacitracin. Toxin formation by strain 243 was sensitive to oleic acid, sodium deoxycholate, sodium lauryl sulfate, D-cycloserine, or bacitracin. The effect of D-cycloserine was reversed by D-alanine with strain 243 but not with strain S6. Neither penicillin nor bacitracin inhibited α -hemolysin or coagulase activity of strain S6; however, 0.118 μ moles of D-cycloserine per ml of liquid medium increased the α -hemolysin titer more than eightfold. Pigmentation of strain 243 was reduced by oleic acid, sodium deoxycholate, or methicillin and was completely inhibited by D-cycloserine or bacitracin. Glucose was required for the inhibition by spermine of valine-C¹⁴ incorporation into cellular protein of strain S6. It is suggested that these data indicate that the cell surface may contain sites important to the synthesis of enterotoxin B.

I. INTRODUCTION

The inhibition of the formation of staphylococcal enterotoxin B by various nutritional inhibitors was reported previously.¹ Although the nature of the inhibitory compounds suggested possible mechanisms involved, none have been elucidated. However, fluorescent antibody studies by Friedman and White² and more recently by Genigeorgis and Sadler³ strongly suggested that enterotoxin was cell surface - associated. This view was strengthened further by Hartman and Goodgal,⁴ who wrote that enterotoxin had the characteristics of a cell surface constituent of bacteria.

The present paper reports experiments designed to test the ability of Staphylococcus aureus to produce enterotoxin B in the presence of compounds known to affect bacterial cell walls. The results strengthen the hypothesis that staphylococcal enterotoxin is cell surface - associated.

II. MATERIALS AND METHODS

A. ORGANISMS AND CULTURE METHODS

S. aureus strains 86⁵ and 243⁶ were used. Stock cultures, maintained on Trypticase Soy agar (BBL) supplemented with 0.001% thiamine and nicotinic acid, were stored at 4 C and transferred every 2 weeks. Broth cultures were grown in PHP medium consisting of 2% Protein Hydrolysate Powder (Mead Johnson) and 0.001% thiamine and nicotinic acid, initial pH 7.7. Oleic acid, the penicillins, D-cycloserine, and bacitracin were prepared as sterile solutions and added aseptically to autoclaved medium. All other additions were autoclaved with the medium. Cells for the preparation of inocula were grown at 37 C for 6 to 7 hours on Trypticase Soy agar slants, removed with sterile deionized water, and adjusted to an optical density of 0.26 (655 m μ). This represented approximately 1.1×10^9 cells/ml. This suspension (0.2 ml) was inoculated into 25 ml of medium in a 250-ml Erlenmeyer flask so that the initial cell count was approximately 9×10^6 per ml. Incubation was for 16 hours at 37 C on a reciprocating shaker making 100 excursions per minute. At the end of the growth period, when the cultures were in the phase of decline, the contents of duplicate flasks were pooled and turbidity measurements at 655 m μ were made on aliquots diluted tenfold in fresh medium, with fresh medium as a blank. The cell density values presented do not represent the maximal growth attained by the cultures, but are given merely to indicate that sufficient growth did occur for the cells to form enterotoxin were it not for the presence of the inhibitor. A supernatant solution collected by centrifugation of the culture at 3,200 rpm for 15 minutes then was subjected to serological assay

for enterotoxin. Merthiolate (final concentration of 100 $\mu\text{g/ml}$) was added to maintain sterility. The packed cells of strain 243 were observed for loss of pigmentation.

B. ENTEROTOXIN ASSAY

Enterotoxin B content of the culture supernatant fluid was assayed quantitatively by a modification⁷ of the single agar diffusion method of Oudin,⁸ sensitive to a minimal concentration of approximately 5 μg toxin per ml. Semiquantitative assays were done using the Ouchterlony double diffusion technique.⁹ Precipitin bands representing the reaction of antibody with as little as 1 μg of antigen per ml could be observed following overnight incubation at 30 C if the plate was washed with 0.85% NaCl repeatedly for a minimum of 4 hours and then flooded with a saturated solution of HgCl_2 . Antitoxin was prepared as described by Silverman.¹⁰

C. α -HEMOLYSIN AND COAGULASE ASSAYS

Culture supernatant solutions without added merthiolate were used in these studies. Erythrocytes from heparinized normal rabbit blood were separated from the plasma by centrifugation at 3,000 rpm for 20 minutes, washed three times, and resuspended to 5% (v/v) in a diluent consisting of 0.85% NaCl, 0.03% $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, and 0.03% KCl. To 0.4 ml of serially diluted sample, 0.1 ml of 5% erythrocytes was added; this mixture then was incubated at 37 C for 1 hour followed by centrifugation at 1,500 rpm for 3 min. Complete hemolysis was designated as a 4+ reaction. For the coagulase test, 0.5 ml of undiluted sample plus 0.5 ml of heparinized plasma was incubated at 37 C and examined every 15 minutes for clotting.

D. ISOTOPIC EXPERIMENT

The incorporation of L-valine- C^{14} into the protein of whole cells of strain S6 was examined as described by Friedman and Bachrach¹¹ except that 1% PHP broth was used and the inoculum was more dense. Radioactivity was measured in a Nuclear Chicago windowless gas flow counter.

E. CHEMICALS

Penicillin G, crystalline sodium (Upjohn); sodium methicillin (Staphcillin, Bristol Laboratories); sodium ampicillin (Polycillin-N, Bristol Laboratories); bacitracin (Calbiochem); and D-cycloserine (Sigma) were used in these studies. Uniformly labeled L-valine- C^{14} (specific activity, 200 mc/mM) was purchased from Schwarz BioResearch.

III. RESULTS

A. EFFECT OF DETERGENT-LIKE SUBSTANCES

Tween 80 effectively inhibited enterotoxin formation by strain S6 but increased toxin yield by strain 243 (Table 1). The supernatant solution obtained following centrifugation of an S6 culture containing Tween 80 appeared turbid, whereas that from strain 243 did not, indicating that strain S6 but not 243 cleaved insoluble oleic acid from the Tween 80. Subsequently, when oleic acid was included in the growth medium, toxin formation by both strains was inhibited. In general, strain 243 was more sensitive to inhibition by oleic acid, sodium deoxycholate, or sodium lauryl sulfate than strain S6. In fact, toxin formation by strain S6 was increased in the presence of sodium lauryl sulfate.

B. EFFECT OF PENICILLINS

Penicillin inhibited toxin formation by strain S6 at the low concentration of 0.0056-0.0393 μ moles/ml (Table 2). Methicillin and ampicillin also were effective. In the case of strain 243 (penicillin- and ampicillin-resistant), 0.449 μ moles/ml of penicillin allowed full growth of the organism but inhibited toxin formation 12%. Growth of this strain was retarded by 0.0015 μ moles/ml of methicillin (penicillinase-resistant); however, the toxin yield was almost twice that of the control culture containing no antibiotic.

C. EFFECT OF D-CYCLOSERINE AND REVERSAL BY ALANINE

D-cycloserine, also active against bacterial cell wall synthesis, but at a different site than penicillin,^{1a} was tested. The toxin synthesizing systems of both strains were sensitive although the growth of S6 was decreased by lower concentrations of D-cycloserine (Table 3). It is to be noted that at 0.118 μ moles/ml of inhibitor, cell turbidity was equal to the control, but toxin was reduced 83%. However, it was consistently observed that with a slight increase in concentration, growth decreased but toxin yield did not. When inhibitor content was increased further, both growth and toxin yield decreased. The activity toward strain 243 was reversed to a great extent by D-alanine but not by the L- or DL-isomers; however, none of these compounds antagonized the effect on strain S6 (Table 4). The reason for this became apparent when it was shown that D-alanine itself was inhibitory to enterotoxin formation by strain S6 but not by strain 243.

TABLE 1. INHIBITION OF STAPHYLOCOCCAL ENTEROTOXIN B SYNTHESIS BY DETERGENT-LIKE SUBSTANCES

Inhibitor, μ moles/ml	<i>S. aureus</i> S6			<i>S. aureus</i> 243		
	OD ^a / (655 m μ)	Enterotoxin, μ g/ml	Inhibition, %	OD ^a / (655 m μ)	Enterotoxin, μ g/ml	Inhibition, %
None	0.29	29		0.31	42	
Tween 80 ^b / 8.0	0.36	9	69	0.24	112	0
16.0	0.40	10	66	0.24	132	0
24.0	0.38	9	69	0.21	110	0
Oleic acid 0.354	0.28	30	0	0.40	9	79
0.885	0.30	33	0	0.43	0	100
1.42	0.24	11	62	0.34	0	100
Sodium deoxycholate 0.048	0.33	13	55	0.29	15	64
0.145	0.21	18	38	0.31	7	83
0.386	0.26	15	48	0.24	3	93
Sodium lauryl sulfate 0.014	0.28	46	0	0.32	36	14
0.028	0.34	49	0	0.29	28	33
0.042	0.36	43	0	0.33	18	57

a. Optical density of a tenfold-diluted 16-hour culture.
b. Milligrams per milliliter.

TABLE 2. INHIBITION OF STAPHYLOCOCCAL ENTEROTOXIN B BY PENICILLIN G

Penicillin, μ moles/ml	<i>S. aureus</i> S6			<i>S. aureus</i> 243		
	OD ^a / (655 m μ)	Enterotoxin, μ g/ml	Inhibition, %	OD ^a / (655 m μ)	Enterotoxin, μ g/ml	Inhibition, %
0.0	0.30	39		0.31	49	
0.0056	0.26	34	13			
0.0112	0.25	21	46			
0.0168	0.21	14	64			
0.0224	0.19	7	82			
0.0281	0.19	4	90			
0.0335	0.15	3	92			
0.0393	0.24	0	100			
0.2245				0.32	43	12
0.4490				0.32	43	12

a. Optical density of a tenfold-diluted 16-hour culture.

TABLE 3. INHIBITION OF STAPHYLOCOCCAL ENTEROTOXIN B SYNTHESIS BY D-CYCLOSERINE

D-Cycloserine, μ moles/ml	<i>S. aureus</i> S6			<i>S. aureus</i> 243		
	OD ^a / (655 m μ)	Enterotoxin, μ g/ml	Inhibition, %	OD ^a / (655 m μ)	Enterotoxin, μ g/ml	Inhibition, %
0.0	0.29	29		0.27	49	
0.039	0.29	21	28	0.27	29	41
0.078	0.34	21	28	0.26	26	47
0.118	0.29	5	83	0.29	26	47
0.157	0.13	19	34	0.30	16	67
0.196	0.04	8	74	0.31	11	78
0.235	0.02	4	86	0.29	11	78
0.274	0.01	0	100	0.18	7	86

a. Optical density of a tenfold-diluted 16-hour culture.

TABLE 4. EFFECT OF ALANINE ON INHIBITORY ACTIVITY OF D-CYCLOSERINE TOWARD THE FORMATION OF STAPHYLOCOCCAL ENTEROTOXIN B

Additions to Medium, μ moles/ml	OD ^a / (655 m μ)	Enterotoxin, μ g/ml	Inhibition, %
<i>S. aureus</i> S6			
None	0.30	19	
+ D-cycloserine (0.118)	0.32	3	84
+ L-alanine (1.35)	0.30	4	79
+ L-alanine (2.02)	0.33	5	74
+ D-alanine (0.45)	0.29	5	74
+ D-alanine (0.90)	0.32	6	68
+ DL-alanine (0.45)	0.32	3	84
+ DL-alanine (0.90)	0.35	3	84
<i>S. aureus</i> 243			
None	0.29	54	
+ D-cycloserine (0.235)	0.29	12	78
+ L-alanine (1.57)	0.29	12	78
+ L-alanine (2.54)	0.33	14	74
+ D-alanine (2.02)	0.38	38	30
+ D-alanine (2.54)	0.30	37	31
+ DL-alanine (1.57)	0.38	10	81
+ DL-alanine (2.54)	0.38	14	74

a. Optical density of a tenfold-diluted 16-hour culture.

D. EFFECT OF BACITRACIN

Bacitracin, also active against cell wall formation,¹³ inhibited the synthesis of enterotoxin B (Table 5). The toxin forming system of strain S6 was more sensitive to bacitracin than that of 243; however, the growth of strain 243 was inhibited at a lower concentration of bacitracin than that of S6.

TABLE 5. INHIBITION OF STAPHYLOCOCCAL ENTEROTOXIN B SYNTHESIS BY BACITRACIN

Bacitracin, $\mu\text{moles/ml}^a$	<i>S. aureus</i> S6			<i>S. aureus</i> 243		
	OD ^b / (655 m μ)	Enterotoxin, $\mu\text{g/ml}$	Inhibition, %	OD ^b / (655 m μ)	Enterotoxin, $\mu\text{g/ml}$	Inhibition, %
0.0	0.29	32		0.28	54	
0.004	0.29	31	3	0.29	43	20
0.009	0.32	9	72	0.32	39	28
0.013	0.32	9	72	0.32	28	48
0.017	0.32	8	75	0.23	14	74
0.021	0.28	8	75	0.12	12	78
0.026	0.28	6	81	0.02	0	100
0.030	0.15	2	94	0.00	0	100

a. Assuming a molecular weight of 1411.

b. Optical density of a tenfold-diluted 16-hour culture.

E. INHIBITORY EFFECTS ON OTHER MACROMOLECULAR SUBSTANCES

Penicillin and bacitracin, at concentrations that did not inhibit growth, had little or no effect on the α -hemolysin or coagulase activity of cultures of strain S6. However, α -hemolysin titers of the supernatant solutions of cultures grown in the presence of 0.118 to 0.157 $\mu\text{moles/ml}$ of D-cycloserine were at least eightfold greater than those of control cultures. At this range of concentration of inhibitor, the yield of enterotoxin was consistently greater than that when the organism was grown in the presence of the next lower concentration of antibiotic. D-Cycloserine had no effect on coagulase activity. Bacitracin and D-cycloserine did not affect α -hemolysin or coagulase activity of strain 243. The control supernatant solution of strain 243 cultures contained more than eight times the α -hemolysin activity of the strain S6 control culture.

Pigmentation of strain 243 was not affected by growth in the presence of Tween 80, sodium lauryl sulfate, or penicillin. Slight loss occurred in the presence of greater than 1.24 μ moles/ml of oleic acid, 0.145 μ moles/ml of sodium deoxycholate, or 0.0015 μ moles/ml of methicillin. Pigmentation loss was complete in the presence of 0.021 μ moles/ml of bacitracin and approximately 0.235 μ moles/ml of D-cycloserine; however, the pigment was regained when the action of D-cycloserine was reversed by D-alanine. Subculture of nonpigmented cultures in the absence of inhibitor also resulted in the regain of pigmentation.

F. VALINE-C¹⁴ INCORPORATION AND THE EFFECT OF SPERMINE

Spermine, an inhibitor of enterotoxin formation,² decreased the incorporation of valine-C¹⁴ by log phase cells of *S. aureus* S6, but only if glucose was in the growth medium (Fig. 1). The inhibition was not complete, because uptake of radioactivity proceeded at a rate which, though slower than the control, was steady.

IV. DISCUSSION

The formation of staphylococcal enterotoxin B can be inhibited nutritionally by a number of compounds without affecting the growth of the organism.¹ The mechanisms of inhibition most likely vary; none has been elucidated nor has the site of synthesis been discovered through these studies. However, there is strong indication that the cell surface is very important to these phenomena.³ The studies reported here in which detergent-like compounds and agents blocking cell wall synthesis displayed the ability to inhibit enterotoxin formation serve to strengthen the hypothesis that enterotoxin is associated with the cell surface.

Because penicillin, D-cycloserine, and bacitracin affect cell wall synthesis at different sites, it is most likely that their activities toward enterotoxin were not specific but affected the cell surface in such a way as to preclude the formation of enterotoxin. Growth of *S. aureus* in the presence of penicillin resulted in structural changes of the cell wall characterized by thinning and disturbance of the deposition of new wall material.¹⁴ It is possible that such "lesions" would occur at sites important to enterotoxin elaboration resulting in complete loss of formation or premature sloughing off of toxic material and/or precursors. The presence of enterotoxin in cultures of low density and the increase in enterotoxin and α -hemolysin concentrations in cultures containing 0.118 to 0.157 μ moles/ml of D-cycloserine over that observed in the presence of lower amounts of the inhibitor suggest that structural changes may have




Figure 1. Effect of Spermine on Incorporation of Valine-C¹⁴ into Growing Log-Phase Cells of *Staphylococcus aureus* S6. (A) Medium containing no added glucose. (B) Medium prepared with 0.2% glucose. The pH of the cultures was adjusted to 7.8 with concomitant addition of spermine and L-valine-C¹⁴ (0.09 µg or 0.05 µg/ml). Organisms were grown in 20-ml cultures in 250-ml Erlenmeyer flasks with shaking in a water bath (37 C).

progressed to a point where "leaking" of antigenic material occurred. If this were so, then observations made here and elsewhere¹⁸ that enterotoxin is elaborated as a metabolic product that apparently does not accumulate in the whole cell point further to the cell surface as the site of formation.

Further support for the hypothesis that the cell surface is the major site of enterotoxin synthesis is derived from the results of the following experiments. It was demonstrated that a drop in the pH of a culture from 7.7 to 5.4 caused by the metabolism of glucose in the medium resulted in the reversal of the ability of spermine to inhibit enterotoxin formation.¹ Cells of S. aureus S6 grown in the presence of 0.2% glucose incorporated 21% of the spermine-C¹⁴ measured in the whole culture, but cells grown in the absence of glucose incorporated 12% of the label* and experiments reported here demonstrated that glucose was necessary for the inhibition by spermine of the incorporation of valine-C¹⁴ into cellular protein. The role of glucose in the activity of spermine was confirmed by the work of Razin and Rozansky,¹⁶ who showed that glucose was necessary in a buffer solution for the maintenance of S. aureus in an active metabolic state in order to demonstrate the bactericidal activity of spermine toward the cells. Also, Schlenk and Daenko¹⁷ found that the inhibitory effect of spermine toward ribonuclease activity on the cell surface of yeast did not occur in the presence of glucose, which was necessary for the assimilation of the polyamine into the cells. Thus, if spermine inhibits enterotoxin formation only in the absence of glucose, and little spermine enters the cell in the absence of glucose, we suggest that spermine acts at the cell surface to inhibit enterotoxin formation. In addition, the failure of penicillin, D-cycloserine, or bacitracin to block α -hemolysin or coagulase indicates that their activities were not toward bacterial protein in general, but more specifically toward those cellular products whose origin was at or near the cell wall and therefore this outer surface must contain the site(s) of enterotoxin synthesis.

* M.E. Friedman, unpublished data.

LITERATURE CITED

1. Friedman, M.E. 1966. Inhibition of staphylococcal enterotoxin B formation in broth cultures. *J. Bacteriol.* 92:277-278.
2. Friedman, M.E.; White, J.D. 1965. Immunofluorescent demonstration of cell-associated staphylococcal enterotoxin B. *J. Bacteriol.* 89:1155.
3. Genigeorgis, C.; Sadler, W.W. 1966. Immunofluorescent detection of staphylococcal enterotoxin B: I. Detection in culture media. *J. Food Sci.* 31:441-449.
4. Hartman, P.E.; Goodgal, S.H. 1959. Bacterial genetics. *Ann. Rev. Microbiol.* 13:465-504.
5. Surgalla, M.J.; Bergdoll, M.S.; Dack, G.M. 1953. Some observations on the assay of staphylococcal enterotoxin by the monkey-feeding test. *J. Lab. Clin. Med.* 41:782-788.
6. Casman, E.P.; Bergdoll, M.S.; Robinson, J. 1963. Designation of staphylococcal enterotoxins. *J. Bacteriol.* 85:715-716.
7. Schantz, E.J.; Roessler, W.G.; Wagman, J.; Spero, L.; Dunnery, D.A.; Bergdoll, M.S. 1965. Purification of staphylococcal enterotoxin B. *Biochemistry* 4:1011-1016.
8. Oudin, J. 1946. Méthode d'analyse immuno-chimique par précipitation spécifique en milieu gélatiné. *Compt. Rend.* 222:115-116.
9. Ouchterlony, Ö. 1948. In vitro method for testing the toxin-producing capacity of diphtheria bacteria. *Acta Pathol. Microbiol. Scand.* 25:166-191.
10. Silverman, S.J. 1963. Serological assay of culture filtrate for Staphylococcus enterotoxin. *J. Bacteriol.* 85:955-956.
11. Friedman, M.E.; Bachrach, U. 1966. Inhibition of protein synthesis by spermine in growing cells of Staphylococcus aureus. *J. Bacteriol.* 92:49-55.
12. Strominger, J.L.; Ito, E.; Threnn, R.H. 1960. Competitive inhibition of enzymatic reactions by oxamycin. *J. Amer. Chem. Soc.* 82: 998-999.
13. Abraham, E.P.; Newton, G.G.F. 1958. Structure and function of some sulphur-containing peptides, p. 205-223. In Ciba Found. Symp. Amino Acids Peptides Antimetab. Activ.

14. Murray, R.G.E.; Francombe, W.H.; Mayall, B.H. 1959. The effect of penicillin on the structure of staphylococcal cell walls. *Can. J. Microbiol.* 5:641-648.
15. Martin, W.J.; Marcus, S. 1964. Relation of pyrogenic and emetic properties of enterobacteriaceal endotoxin and of staphylococcal enterotoxin. *J. Bacteriol.* 87:1019-1026.
16. Rasin, S.; Rozansky, R. 1959. Mechanism of the antibacterial action of spermine. *Arch. Biochem. Biophys.* 81:36-54.
17. Schlenk, F.; Dainko, J.L. 1966. Effects of ribonuclease and spermine on yeast cells. *Arch. Biochem. Biophys.* 113:127-133.

Unclassified
Security Classification

17

DOCUMENT CONTROL DATA - R & D		
<i>(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)</i>		
1. ORIGINATING ACTIVITY (Corporate number) Department of the Army Fort Detrick, Frederick, Maryland, 21701		2A. REPORT SECURITY CLASSIFICATION Unclassified
3. REPORT TITLE INHIBITION OF STAPHYLOCOCCAL ENTEROTOXIN B FORMATION BY CELL WALL BLOCKING AGENTS AND OTHER COMPOUNDS		2B. GROUP
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5. AUTHOR(S) (First name, middle initial, last name) Mischa E. Friedman		
6. REPORT DATE March 1968	7A. TOTAL NO. OF PAGES 17	7B. NO. OF REFS 17
8A. CONTRACT OR GRANT NO. A. PROJECT NO. 1C522301A059	8B. ORIGINATOR'S REPORT NUMBER(S) Technical Manuscript 429	
C. 4.	8C. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
10. DISTRIBUTION STATEMENT Qualified requesters may obtain copies of this publication from DDC. Foreign announcement and dissemination of this publication by DDC is not authorized. Release or announcement to the public is not authorized.		
11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY Department of the Army Fort Detrick, Frederick, Maryland, 21701	
13. ABSTRACT Enterotoxin B formation by <u>Staphylococcus aureus</u> S6 was inhibited by Tween 80, oleic acid, sodium deoxycholate, penicillin, D-cycloserine, or bacitracin. Toxin formation by strain 243 was sensitive to oleic acid, sodium deoxycholate, sodium lauryl sulfate, D-cycloserine, or bacitracin. The effect of D-cycloserine was reversed by D-alanine with strain 243 but not with strain S6. Neither penicillin nor bacitracin inhibited α -hemolysin or coagulase activity of strain S6; however, 0.118 μ moles of D-cycloserine per ml of liquid medium increased the α -hemolysin titer more than eightfold. Pigmentation of strain 243 was reduced by oleic acid, sodium deoxycholate, or methicillin and was completely inhibited by D-cycloserine or bacitracin. Glucose was required for the inhibition by spermine of valine-C ¹⁴ incorporation into cellular protein of strain S6. It is suggested that these data indicate that the cell surface may contain sites important to the synthesis of enterotoxin B.		
14. Key Words * <u>Staphylococcus aureus</u> *Inhibition *Enterotoxin D-cycloserine Bacitracin Penicillin		

DD FORM 1473

REPLACES DD FORM 1473, 1 JAN 64, WHICH IS OBSOLETE FOR ARMY USE.

Unclassified
Security Classification