



Body image in social anxiety disorder, obsessive–compulsive disorder, and panic disorder



Idan M. Aderka^{a,b,*}, Cassidy A. Gutner^a, Amit Lazarov^c, Haggai Hermesh^{c,d}, Stefan G. Hofmann^a, Sofi Marom^{d,e}

^a Boston University, Boston, MA, USA

^b University of Haifa, Mount Carmel, Israel

^c Tel-Aviv University, Tel-Aviv, Israel

^d Geha Mental Health Center, Petach-Tikva, Israel

^e Rupin Academic Center, Netanya, Israel

ARTICLE INFO

Article history:

Received 10 October 2012

Received in revised form 3 September 2013

Accepted 4 September 2013

Keywords:

Body dysmorphic disorder

Body image

Obsessive–compulsive disorder

Social anxiety disorder

Panic disorder

ABSTRACT

Body dysmorphic disorder falls under the category of obsessive–compulsive and related disorders, yet research has suggested it may also be highly associated with social anxiety disorder. The current study examined body image variables among 68 outpatients with primary obsessive–compulsive disorder (OCD; $n=22$), social anxiety disorder (SAD; $n=25$), and panic disorder (PD; $n=21$). Participants filled out self-report measures of body image disturbance, attitudes toward one's appearance, and anxiety. Body image disturbance and attitudes toward appearance did not significantly differ between the groups. However, SAD symptoms predicted body image disturbance, Appearance Evaluation and Body Areas Satisfaction, and OCD symptoms predicted Appearance Orientation. These findings suggest that SAD and OCD may be associated with different facets of body image. Implications for the treatment of anxiety disorders and for future research are discussed.

© 2013 Elsevier Ltd. All rights reserved.

Introduction

Body dysmorphic disorder (BDD) falls under the category of obsessive–compulsive and related disorders and is characterized by an excessive preoccupation with a minor or imagined defect in appearance that causes clinically significant distress or impairment (American Psychiatric Association, 2013). Current prevalence rates of BDD are 1.7–1.8% in community samples (Buhlmann et al., 2010; Rief, Buhlmann, Wilhelm, Borkenhagen, & Brähler, 2006) and the disorder has been associated with higher rates of unemployment, lower income, and lower likelihood of living with a partner (Rief et al., 2006). Some individuals with BDD report suicidal ideation and are at increased risk for suicide attempts due to concerns about their physical appearance (Buhlmann et al., 2010; Rief et al., 2006).

Rates of comorbidity between BDD and obsessive–compulsive disorder (OCD) are as high as 30% (Gunstad & Phillips, 2003; Phillips, Menard, Fay, & Weisberg, 2005), and both disorders share similarities in demographics, onset and illness duration (Phillips, Pinto, Menard, Eisen, Mancebo, & Rasmussen, 2007). In addition, research has demonstrated that both disorders share familial and genetic

components (Bienvenu et al., 2000; Phillips, Gunderson, Mallya, McElroy, & Carter, 1998). OCD and BDD also share similar cognitive patterns characterized by recurrent, persistent and intrusive unwanted thoughts (Phillips, McElroy, Keck, Pope, & Hudson, 1993), and behavioral patterns such as compulsive checking (Phillips, Menard, Fay, & Weisberg, 2005; Phillips et al., 2010). Finally, BDD symptoms are common among individuals with OCD (Phillips et al., 2007). Thus, there is substantial evidence suggesting that the two disorders are related.

At the same time, research has identified some key differences between OCD and BDD. For example, individuals with BDD demonstrate less insight and more delusional beliefs compared to individuals with OCD (Eisen, Phillips, Coles, & Rasmussen, 2004; Phillips et al., 2007). BDD is also associated with higher levels of suicidal ideation, and higher levels of major depressive disorder and substance use disorder compared to OCD (Phillips et al., 2007). Moreover, individuals with BDD have been found to have significantly greater body image impairment compared to individuals with OCD (Hrabosky et al., 2009). Finally, in comparison to OCD, the content of beliefs in BDD seems to focus more on unacceptability of the self (Veale & Riley, 2001).

Recently, there has been growing evidence for the association between BDD and social anxiety disorder (SAD; see Fang & Hofmann, 2010 for a review). Research has shown that among individuals with SAD, 4.8–12% are diagnosed with BDD, and among

* Corresponding author at: Department of Psychology, Boston University, 648 Beacon Street, 6th Floor, Boston, MA, USA.

E-mail addresses: iaderka@bu.edu, iaderka@psy.haifa.ac.il (I.M. Aderka).

Table 1
Demographic and clinical measures.

	Panic disorder (n = 21)	Social anxiety disorder (n = 25)	Obsessive–compulsive disorder (n = 22)	Statistic	p
Age, <i>M</i> (<i>SD</i>)	38.67 (13.67) _a	30.24 (5.97) _b	40.05 (11.97) _a	$F_{(2, 65)} = 5.27$	<.01
Gender, n %				$\chi^2_{(2)} = 0.35$.84
Female	11 (52.38%)	11 (44.00%)	11 (50.00%)		
Male	10 (47.62%)	14 (56.00%)	11 (50.00%)		
Education (years), <i>M</i> (<i>SD</i>)	12.57 (2.18)	13.72 (2.01)	12.82 (1.87)	$F_{(2, 65)} = 2.10$.13
Marital status, n %				Fisher's exact test statistic = 19.33	<.001
Single	5 (23.81%)	21 (84.00%)	13 (59.09%)		
Married	14 (66.67%)	4 (16.00%)	7 (31.81%)		
Divorced	1 (4.76%)	0 (0.00%)	2 (9.10%)		
Widowed	1 (4.76%)	0 (0.00%)	0 (0.00%)		
LSAS, <i>M</i> (<i>SD</i>)	16.29 (13.12) _b	68.29 (28.49) _a	25.36 (20.70) _b	$F_{(2, 64)} = 36.46$	<.001
OCL, <i>M</i> (<i>SD</i>)	14.71 (9.12) _b	19.06 (14.12) _b	36.86 (12.01) _a	$F_{(2, 57)} = 21.22$	<.001
BIDQ, <i>M</i> (<i>SD</i>)	1.72 (0.50)	2.22 (0.82)	2.07 (1.17)	$F_{(2, 63)} = 1.86$.16
MBSRQ-AS, <i>M</i> (<i>SD</i>)					
Appearance Evaluation	3.48 (0.73)	3.40 (0.79)	3.65 (0.81)	$F_{(2, 64)} = 0.56$.58
Appearance Orientation	3.41 (0.60)	3.70 (0.55)	3.67 (0.59)	$F_{(2, 64)} = 1.63$.21
Body Areas Satisfaction	3.43 (0.63)	3.22 (0.61)	3.45 (0.81)	$F_{(2, 64)} = 0.77$.47

Note. Subscripts indicate significant differences: $a > b$. Post hoc tests were Dunnett T3 tests that do not assume variance equality across the groups. LSAS=Liebowitz Social Anxiety Scale, OCL=Obsessive Compulsive Inventory, BIDQ=Body Image Disturbance Questionnaire, MBSRQ-AS=Multidimensional Body-Self Relations Questionnaire—Appearance Scales.

individuals with BDD, 12–68.8% have SAD (Fang & Hofmann, 2010). Research on BDD and SAD has highlighted several commonalities including high social anxiety and social avoidance (Coles et al., 2006; Kelly, Walters, & Phillips, 2010; Pinto & Phillips, 2005), low extraversion (Naragon-Gainey, Watson, & Markon, 2009; Phillips & McElroy, 2000), and high levels of embarrassment and shame (Buhlmann & Wilhelm, 2004; Conroy, Menard, Fleming-Ives, Modha, Cerullo, & Phillips, 2008; Fuchs, 2002). Other shared characteristics include elevated rates of suicidal ideation (Cogle, Keough, Riccardi, & Sachs-Ericsson, 2009; Phillips & Menard, 2006; Phillips et al., 2007), negative interpretation bias for ambiguous social information (Amir, Foa, & Coles, 1998; Buhlmann, Wilhelm, McNally, Tuschen-Caffier, Baer, & Jenike, 2002; Heinrichs & Hofmann, 2001; Hofmann, 2007), and a typical age of onset in childhood to early adulthood (Phillips, Menard, Fay, & Weisberg, 2005; Wittchen & Fehm, 2003).

Most studies have examined the relationship between BDD and either OCD or SAD. To our knowledge, only a single study has examined BDD among individuals with both disorders. Lochner and Stein (2010) compared rates of BDD among individuals with OCD, SAD, and panic disorder (PD) and found no significant differences between the three groups. However, the rate of BDD in the SAD group (12.2%) was approximately twice the rate of BDD in the OCD group (6.5%) and the PD group (5.4%) despite being a non-significant difference. In the present study we sought to extend these findings by examining continuous levels of body image disturbance and attitudes toward one's appearance (as opposed to a dichotomous diagnosis of BDD) among individuals with SAD, OCD and PD. Body image disturbance and attitudes toward one's appearance are on a continuum and are related to BDD such that individuals with the disorder have higher levels of body image disturbance and more dysfunctional attitudes toward their appearance compared to individuals without the disorder. Using continuous rather than dichotomous variables can help assess non-clinical levels of body image variables that may be found in the anxiety disorders and can complicate their treatment. The examination of continuous body image variables in the anxiety disorders is also important as it can shed light on whether body image disturbance and attitudes toward one's appearance are specifically related to one or more of these anxiety disorders, and on the relative strength of the associations.

A group with PD was chosen as the clinical control group for several reasons. First, as mentioned above, the literature on BDD

ties the disorder, its symptoms and its etiology to OCD and SAD but not to other anxiety disorders such as PD. Second, a PD group can help determine whether body image variables are specifically related to OCD and SAD or rather to anxiety disorders in general. Third, a PD group can facilitate comparison of the results with the only previous study that explicitly examined the presence of BDD in individuals with primary OCD, SAD and PD (Lochner & Stein, 2010).

We expected that body image disturbance and dysfunctional attitudes toward one's appearance would be elevated in both the OCD group and the SAD group compared to the PD group, as both OCD and SAD have been previously found to be associated with BDD (e.g., Fang & Hofmann, 2010; Mataix-Cols, Pertusa, & Leckman, 2007). In addition, we also expected obsessive–compulsive and social anxiety symptoms to be significantly associated with body image variables across the three groups. We included individuals in the PD group because individuals with primary PD may experience non-clinical levels of OCD and SAD symptoms that may in turn be associated with body image variables. Finally, we explored whether body image variables would be more strongly related to obsessive–compulsive symptoms or social anxiety symptoms.

Method

Participants

Participants were 68 consecutive outpatients who sought treatment at a large public health center. Of the total sample, 21 individuals were diagnosed with primary panic disorder, 25 were diagnosed with primary social anxiety disorder, and 22 were diagnosed with primary obsessive–compulsive disorder. Participants' mean age was 36.01 ($SD = 11.97$), and 48.50% were female. Most participants were single (57.40%), 36.80% were married, 4.40% were divorced, and 1.50% were widowed. The average participant had a mean of 13.07 ($SD = 2.05$) years of education. Table 1 provides demographic and clinical measures for each of three groups separately. Participants were excluded from the present study if they: (a) received a primary diagnosis other than panic disorder, social anxiety disorder, or obsessive–compulsive disorder, (b) received a secondary diagnosis of panic disorder, social anxiety disorder, or obsessive–compulsive disorder, or (c) if they received a past or present diagnosis of psychotic episode or schizophrenia. Thus, participants in the OCD group did not have comorbid SAD or PD, participants in the SAD group did not have comorbid OCD or PD,

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات