

## Beetroot juice and exercise: pharmacodynamic and dose-response relationships

Lee J. Wylie,<sup>1</sup> James Kelly,<sup>1</sup> Stephen J. Bailey,<sup>1</sup> Jamie R. Blackwell,<sup>1</sup> Philip F. Skiba,<sup>1</sup> Paul G. Winyard,<sup>2</sup> Asker E. Jeukendrup,<sup>3</sup> Anni Vanhatalo,<sup>1</sup> and Andrew M. Jones<sup>1</sup>

<sup>1</sup>Sport and Health Sciences, College of Life and Environmental Sciences, University of Exeter, St. Luke's Campus, Exeter, United Kingdom; <sup>2</sup>University of Exeter Medical School, St. Luke's Campus, Exeter, United Kingdom; and <sup>3</sup>Gatorade Sports Science Institute, Barrington, Illinois

Submitted 25 March 2013; accepted in final form 25 April 2013

**Wylie LJ, Kelly J, Bailey SJ, Blackwell JR, Skiba PF, Winyard PG, Jeukendrup AE, Vanhatalo A, Jones AM.** Beetroot juice and exercise: pharmacodynamic and dose-response relationships. *J Appl Physiol* 115: 325–336, 2013. First published May 2, 2013; doi:10.1152/jappphysiol.00372.2013.—Dietary supplementation with beetroot juice (BR), containing approximately 5–8 mmol inorganic nitrate ( $\text{NO}_3^-$ ), increases plasma nitrite concentration ( $[\text{NO}_2^-]$ ), reduces blood pressure, and may positively influence the physiological responses to exercise. However, the dose-response relationship between the volume of BR ingested and the physiological effects invoked has not been investigated. In a balanced crossover design, 10 healthy men ingested 70, 140, or 280 ml concentrated BR (containing 4.2, 8.4, and 16.8 mmol  $\text{NO}_3^-$ , respectively) or no supplement to establish the effects of BR on resting plasma  $[\text{NO}_3^-]$  and  $[\text{NO}_2^-]$  over 24 h. Subsequently, on six separate occasions, 10 subjects completed moderate-intensity and severe-intensity cycle exercise tests, 2.5 h postingestion of 70, 140, and 280 ml BR or  $\text{NO}_3^-$ -depleted BR as placebo (PL). Following acute BR ingestion, plasma  $[\text{NO}_2^-]$  increased in a dose-dependent manner, with the peak changes occurring at approximately 2–3 h. Compared with PL, 70 ml BR did not alter the physiological responses to exercise. However, 140 and 280 ml BR reduced the steady-state oxygen ( $\text{O}_2$ ) uptake during moderate-intensity exercise by 1.7% ( $P = 0.06$ ) and 3.0% ( $P < 0.05$ ), whereas time-to-task failure was extended by 14% and 12% (both  $P < 0.05$ ), respectively, compared with PL. The results indicate that whereas plasma  $[\text{NO}_2^-]$  and the  $\text{O}_2$  cost of moderate-intensity exercise are altered dose dependently with  $\text{NO}_3^-$ -rich BR, there is no additional improvement in exercise tolerance after ingesting BR containing 16.8 compared with 8.4 mmol  $\text{NO}_3^-$ . These findings have important implications for the use of BR to enhance cardiovascular health and exercise performance in young adults.

nitrate; nitrite; nitric oxide; blood pressure; exercise economy;  $\text{O}_2$  uptake; exercise tolerance

NITRIC OXIDE (NO) IS A GASEOUS signaling molecule that modulates human physiological function via its role in, for example, the regulation of blood flow, neurotransmission, immune function, glucose and calcium homeostasis, muscle contractility, and mitochondrial respiration (9, 36).<sup>1</sup> NO is generated through the oxidation of the amino acid L-arginine in a reaction catalyzed by NO synthase (NOS), with nitrite ( $\text{NO}_2^-$ ) and nitrate ( $\text{NO}_3^-$ ) being oxidation products of NO (30). It is now appreciated that under appropriate physiological conditions, NO can also be produced via the reduction of  $\text{NO}_2^-$ , a process that may be particularly important in situations where oxygen

( $\text{O}_2$ ) availability is low, and/or NOS function is impaired (12). Interestingly, administration of dietary inorganic  $\text{NO}_3^-$  has been shown to increase plasma  $\text{NO}_2^-$  concentration ( $[\text{NO}_2^-]$ ) and to produce NO-like bioactivity (19, 23, 39). Up to 25% of ingested  $\text{NO}_3^-$  enters the enterosalivary circulation and is concentrated in the saliva, whereupon facultative, anaerobic bacteria in the oral cavity reduce the  $\text{NO}_3^-$  to  $\text{NO}_2^-$  (30). When swallowed into the acidic environment of the stomach, some of the  $\text{NO}_2^-$  is converted further into NO, whereas the remainder is absorbed to increase circulating plasma  $[\text{NO}_2^-]$ . This  $\text{NO}_2^-$  may be reduced further to NO and other reactive nitrogen intermediates, particularly in tissues that may be relatively hypoxic, such as contracting skeletal muscle (30).

We (3, 37) and others (19, 23, 39) have demonstrated that  $\text{NO}_3^-$  ingestion, either in the form of  $\text{NO}_3^-$  salts or via the consumption of high  $\text{NO}_3^-$  vegetable products, such as beetroot juice (BR), reduces resting blood pressure (BP) profoundly and consistently. Consequently, dietary  $\text{NO}_3^-$  supplementation has emerged as a potential nutritional agent for the prevention and treatment of hypertension and cardiovascular disease (30). Webb et al. (39) assessed the effects of acute BR consumption (~23 mmol  $\text{NO}_3^-$ ) on plasma  $[\text{NO}_2^-]$  and BP over 24 h. Plasma  $[\text{NO}_2^-]$  peaked 3 h postingestion, remained close to peak values until 5 h postingestion, and returned to baseline after 24 h (39). The systolic and diastolic BP and the mean arterial pressure (MAP) were reduced significantly, by ~10, ~8, and ~8 mmHg, respectively, at 2.5–3 h after BR intake. The same research group later reported a dose-dependent increase in plasma  $[\text{NO}_3^-]$  and  $[\text{NO}_2^-]$  and reduction in BP following ingestion of potassium  $\text{NO}_3^-$  ( $\text{KNO}_3$ ) (19). In this study, plasma  $[\text{NO}_2^-]$  rose by ~1.3-, approximately two-, and approximately fourfold following consumption of 4, 12, and 24 mmol  $\text{KNO}_3$ , respectively. The peak rise in plasma  $[\text{NO}_2^-]$  was accompanied by significant reductions in both systolic BP (of ~2, ~6, and ~9 mmHg, respectively) and diastolic BP (of ~4, ~4, and ~6 mmHg, respectively). However, since BR contains polyphenols and antioxidants, which can facilitate the synthesis of NO from  $\text{NO}_2^-$  in the stomach (30), it is unclear whether BP is similarly impacted when different doses of BR are ingested compared with equivalent doses of  $\text{NO}_3^-$  salts. Given the growing interest in dietary  $\text{NO}_3^-$  supplementation in the form of BR amongst athletes and the general population, it is important to determine the pharmacokinetic-pharmacodynamic relationship between different volumes of BR consumption and changes in plasma  $[\text{NO}_2^-]$  and BP to establish an optimal dose for beneficial effects.

Recent investigations suggest that dietary  $\text{NO}_3^-$  supplementation has the potential to influence human physiology beyond

<sup>1</sup>This article is the topic of an Invited Editorial by L. Burke (5a).

Address for reprint requests and other correspondence: A. M. Jones, College of Life and Environmental Sciences, Univ. of Exeter, St. Luke's Campus, Exeter EX1 2LU, UK (e-mail: a.m.jones@exeter.ac.uk).

the above hemodynamic effects (3, 26). Specifically, we (2, 3, 22) and others (6, 24–26) have demonstrated that 3–6 days of dietary  $\text{NO}_3^-$  supplementation reduces the  $\text{O}_2$  cost of moderate-intensity exercise and may enhance exercise tolerance in healthy, young adults. It appears that these effects are related to  $\text{NO}_2^-$  or NO-mediated enhancements of muscle contractile function (2, 17) and/or mitochondrial efficiency (24) and/or enhanced muscle blood flow, especially to *type II* fibers (14). Importantly, a reduction of the  $\text{O}_2$  cost of exercise (25, 37) and improved exercise performance (21) has also been reported as early as 2.5 h following a single dose of dietary  $\text{NO}_3^-$ , which is consistent with the time required for the peak plasma  $[\text{NO}_2^-]$  to be attained (39). However, since all exercise-performance studies completed to date with BR have administered approximately 5–8 mmol  $\text{NO}_3^-$ , it is unclear whether a dose-response relationship exists between acute  $\text{NO}_3^-$  intake and the physiological responses to exercise. The establishment of the dose-response relationship between  $\text{NO}_3^-$  intake and the physiological responses to exercise and the ascertainment of the optimal  $\text{NO}_3^-$  dose for enhancing exercise performance are important, given the increasing popularity of BR supplementation in both basic research and applied exercise settings.

Therefore, the purpose of the present study was twofold: firstly, to characterize the plasma  $[\text{NO}_3^-]$  and  $[\text{NO}_2^-]$  pharmacokinetics and the changes in BP after ingestion of three different quantities of  $\text{NO}_3^-$ -rich BR; and secondly, to investigate the dose-response relationship between BR/ $\text{NO}_3^-$  intake and the physiological responses to exercise. In two separate experiments, we administered a BR concentrate that enabled a substantial  $\text{NO}_3^-$  load to be ingested quickly and easily. We investigated: 1) the influence of acute  $\text{NO}_3^-$  doses of 4.2, 8.4, and 16.8 mmol consumed in 70, 140, and 280 ml concentrated BR on plasma  $[\text{NO}_3^-]$  and  $[\text{NO}_2^-]$  and BP over a 24-h period; and 2) the physiological responses to step transitions to moderate- and severe-intensity exercise, 2.5 h postingestion of the same  $\text{NO}_3^-$  doses. We hypothesized that the effects of dietary inorganic  $\text{NO}_3^-$  on plasma  $[\text{NO}_3^-]$  and  $[\text{NO}_2^-]$ , BP, the  $\text{O}_2$  cost of moderate-intensity exercise, and exercise tolerance (assessed as the time-to-task failure) during severe-intensity exercise would be dose dependent.

## METHODS

The study was conducted in two phases [*study 1* ( $S_1$ ), pharmacokinetics; and  $S_2$ , dose response], with the results generated in  $S_1$  used to inform the experimental design in  $S_2$ . There was distinct subject recruitment for each experiment. Ten healthy, recreationally active men volunteered for each experiment [mean  $\pm$  SD:  $S_1$ , age  $23 \pm 5$  yr, height  $1.79 \pm 0.07$  m, body mass (BM)  $79 \pm 9$  kg;  $S_2$ , age  $22 \pm 5$  yr, height  $1.77 \pm 0.05$  m, BM  $74 \pm 8$  kg]. None of the subjects in  $S_1$  and  $S_2$  was a tobacco smoker or user of dietary supplements. All subjects recruited for  $S_2$  were fully familiar with laboratory exercise-testing procedures, having participated previously in studies using cycle ergometry in our laboratory. The procedures used in  $S_1$  and  $S_2$  were granted full ethics approval by the Institutional Research Ethics Committee. All subjects gave their written, informed consent to participate after the experimental procedures, associated risks, and potential benefits of participation had been explained in detail.

All subjects in  $S_1$  and  $S_2$  were instructed to keep a food and physical-activity diary in the 24 h preceding their first laboratory visit and to replicate food consumption and physical activity in the 24 h preceding subsequent visits. The subjects were required to arrive at the laboratory in a rested and fully hydrated state, following an

overnight fast, and to avoid strenuous activity in the 24 h preceding each testing session. Subjects were instructed to refrain from caffeine and alcohol-containing drinks for 6 and 24 h before each laboratory visit, respectively, and to abstain from using antibacterial mouthwash and chewing gum throughout the study, because these are known to eradicate the oral bacteria that are necessary for the conversion of  $\text{NO}_3^-$  to  $\text{NO}_2^-$  (16).

## $S_1$ : Pharmacokinetics and Pharmacodynamics

**Procedures.** All subjects reported to the laboratory on four separate occasions over a period of 3 wk. Upon arrival to the laboratory, resting BP was measured, and a venous blood sample was obtained for the measurement of plasma  $[\text{NO}_2^-]$  and  $[\text{NO}_3^-]$ . Subjects then consumed an acute dose of 70, 140, or 280 ml  $\text{NO}_3^-$ -rich BR (organic BR containing  $\sim 4.2$ ,  $\sim 8.4$ , or  $\sim 16.8$  mmol  $\text{NO}_3^-$ , respectively; Beet It; James White Drinks, Ipswich, UK) or 140 ml water [control (CON)], in addition to a standardized breakfast (72 g porridge oats with 180 ml semiskimmed milk). BP was measured, and a venous blood sample was obtained, 1, 2, 4, 8, 12, and 24 h postingestion. For each 24-h period of data collection, subjects were provided with a standardized, low  $\text{NO}_3^-$  diet. The quantity and timing of food and drink intake were recorded on *visit 1* and replicated in subsequent visits. A washout period of at least 3 days separated the laboratory visits.

**Measurements.** The BP of the brachial artery was measured using an automated sphygmomanometer (Dinamap Pro; GE Medical Systems, Tampa, FL), with the subjects in a seated position. After arrival at the laboratory and following 10 min of rest in an isolated room, four measurements were recorded, and the mean of the final three measurements was used for data analysis.

Venous blood samples were drawn into lithium-heparin tubes (7.5 ml Monovette lithium heparin; Sarstedt, Leicester, UK). Samples were centrifuged at 4,000 rpm and  $4^\circ\text{C}$  for 7 min, within 1 min of collection. Plasma was subsequently extracted and immediately frozen at  $-80^\circ\text{C}$  for later analysis of  $[\text{NO}_2^-]$  and  $[\text{NO}_3^-]$ .

All glassware, utensils, and surfaces were rinsed with deionized water to remove residual  $[\text{NO}_2^-]$  and  $[\text{NO}_3^-]$  before blood analyses. The  $[\text{NO}_2^-]$  of the undiluted (nonproteinized) plasma was determined by its reduction to NO in the presence of glacial acetic acid and 4% (w/v) aqueous sodium iodide. The spectral emission of electronically excited nitrogen dioxide product, from the NO reaction with ozone, was detected by a thermoelectrically cooled, red-sensitive photomultiplier tube, housed in a Sievers gas-phase chemiluminescence NO analyzer (NOA; Sievers NOA 280i; Analytix, Durham, UK). The  $[\text{NO}_2^-]$  was determined by plotting signal (mV) area against a calibration plot of 100 nM–1  $\mu\text{M}$  sodium  $\text{NO}_2^-$ . Before determination of  $[\text{NO}_3^-]$ , samples were deproteinized using zinc sulfate ( $\text{ZnSO}_4$ )/sodium hydroxide (NaOH) precipitation. Aqueous  $\text{ZnSO}_4$  [400  $\mu\text{l}$  10% (w/v)] and 400  $\mu\text{l}$  0.5 M NaOH were added to 200  $\mu\text{l}$  of sample and vortexed for 30 s before being left to stand at room temperature for 15 min. Thereafter, samples were centrifuged at 4,000 rpm for 5 min, and the supernatant was removed for subsequent analysis. The  $[\text{NO}_3^-]$  of the deproteinized plasma sample was determined by its reduction to NO in the presence of 0.8% (w/v) vanadium trichloride in 1 M HCl. The production of NO was detected using the chemiluminescence NOA, as described above.

To determine more precisely the time-to-peak plasma  $[\text{NO}_2^-]$  following  $\text{NO}_3^-$  ingestion, a one-compartment model with first-order absorption and elimination kinetics was used, as described in the following equation

$$Y = (\exp(-Ke \times X)/(Ke/Ka)) - \exp(-Ke \times X)/(Ke/Ka - 1)$$

where Y represents fraction absorbed; X represents time; and, Ka and Ke represent the first-order absorption and elimination rate constants, respectively.

**Statistical analysis.** Two-way repeated-measures ANOVA was used to assess the difference across conditions (4.2, 8.4, and 16.8

mmol  $\text{NO}_3^-$  and CON) and across time (0, 1, 2, 4, 8, 12, and 24 h) for plasma  $[\text{NO}_2^-]$  and  $[\text{NO}_3^-]$  and BP. Significant main or interaction effects were analyzed further using simple contrasts. One-way repeated-measures ANOVA was used to assess the differences in time-to-peak plasma  $[\text{NO}_2^-]$ . Relationships between plasma  $[\text{NO}_2^-]$  and BP were analyzed using Pearson product moment correlation coefficients. Statistical significance was accepted at  $P < 0.05$ . Results are presented as mean  $\pm$  SD unless stated otherwise.

## S<sub>2</sub>: Dose Response

**Protocol.** Subjects were required to report to the laboratory on seven separate occasions, over a 4- to 5-wk period. During the first visit to the laboratory, subjects completed a ramp incremental exercise test for determination of peak  $\dot{V}\text{O}_2$  (peak) and gas-exchange threshold (GET). All tests were performed on an electronically braked cycle ergometer (Lode Excalibur Sport, Groningen, The Netherlands). Initially, each subject completed 3 min of “unloaded” baseline cycling; then, the work rate was increased by 30 W/min until the subject was unable to continue. The subjects cycled at a self-selected pedal rate (70–90 rpm), and this pedal rate along with the saddle and handlebar height and configuration were recorded and reproduced in subsequent tests. The breath-by-breath pulmonary gas-exchange data were collected continuously during the incremental tests and averaged over consecutive 10-s periods.  $\dot{V}\text{O}_{2\text{peak}}$  was taken as the highest 30-s mean value attained before the subject’s volitional exhaustion. The GET was determined as described previously (3, 37). The work rates that would require 80% of the GET (moderate-intensity exercise) and 75% of the difference between the power output at GET and  $\dot{V}\text{O}_{2\text{peak}}$  plus the power output at GET, i.e., severe-intensity exercise ( $\Delta$ ) were subsequently calculated.

On test days, subjects arrived at the laboratory at ~8 AM. A venous blood sample was drawn for measurement of plasma  $[\text{NO}_2^-]$  and  $\text{NO}_3^-$ . Subjects then ingested 70, 140, or 280 ml  $\text{NO}_3^-$ -rich BR (containing 4.2, 8.4, or 16.8 mmol  $\text{NO}_3^-$ , respectively; Beet It) or 70, 140, or 280 ml  $\text{NO}_3^-$ -depleted BR as a placebo (PL70, PL140, or PL280; containing ~0.04, ~0.08, or ~0.12 mmol  $\text{NO}_3^-$ ; Beet It). All BR and PL doses were administered using a randomized, double-blind crossover design. Subjects were asked to consume the beverage within a 5-min period and, after doing so, were served a standardized breakfast (72 g porridge with 180 ml semiskimmed milk). A washout period of at least 72 h separated each visit.

After ingestion of the beverage, subjects were given a period of 2.5 h, during which they were allowed to leave the laboratory but were asked to refrain from strenuous physical activity. Subjects were also asked to fast during this time, although water was permitted ad libitum. Following this 2.5-h period, a second venous blood sample was drawn for measurement of plasma  $[\text{NO}_2^-]$  and  $[\text{NO}_3^-]$ . Subjects then completed “step” exercise tests, from a 20-W baseline to moderate-intensity ( $93 \pm 11$  W) and severe-intensity ( $258 \pm 23$  W) work rates for the determination of pulmonary  $\dot{V}\text{O}_2$  dynamics. On each visit, subjects completed two, 5-min bouts of moderate-intensity exercise and one bout of severe-intensity exercise that was continued until task failure as a measure of exercise tolerance. All bouts of exercise on each day were separated by 5 min of passive rest. The time-to-task failure was recorded when the pedal rate fell by  $>10$  rpm below the self-selected pedal rate. In the severe-intensity bouts, the subjects were verbally encouraged to continue for as long as possible.

**Measurements.** During all exercise tests, pulmonary gas exchange and ventilation were measured breath by breath, with subjects wearing a nose clip and breathing through a low dead-space (90 ml), low-resistance ( $0.75 \text{ mmHg l}^{-1} \cdot \text{s}^{-1}$  at 15 l/s) mouthpiece and impeller turbine assembly (Jaeger Triple-V; Jaeger GmbH, Hoechberg, Germany). The inspired and expired gas volume and gas concentration signals were sampled continuously at 100 Hz—the latter using paramagnetic ( $\text{O}_2$ ) and infrared [carbon dioxide ( $\text{CO}_2$ )] analyzers (Oxycon Pro; Jaeger GmbH) via a capillary line connected to the mouthpiece.

These analyzers were calibrated before each test with gases of known concentration, and the turbine volume transducer was calibrated using a 3-liter syringe (Hans Rudolph, Kansas City, MO). The volume and concentration signals were time aligned by accounting for the delay in capillary gas transit and analyzer rise time relative to volume signal.  $\text{O}_2$  uptake,  $\text{CO}_2$  output, and minute ventilation were calculated using a standard formula and displayed breath by breath. Heart rate (HR) was measured using short-range radiotelemetry (model RS400; Polar Electro Oy, Kempele, Finland).

Capillary blood samples were collected from the fingertip into a capillary tube during the baseline, preceding each step transition in work rate; during the final 30 s of each moderate-intensity exercise bout; and following exhaustion in the severe-intensity exercise bout. These samples were analyzed immediately to determine blood lactate concentration ([lactate]; model YSI 1500; Yellow Springs Instrument, Yellow Springs, OH). Venous blood samples were treated and analyzed as described in S<sub>1</sub>.

The breath-by-breath data from each exercise test were linearly interpolated to provide second-by-second values, and the two identical, moderate-intensity repetitions performed on each visit were time aligned to the start of exercise and ensemble averaged. Baseline  $\dot{V}\text{O}_2$  ( $\dot{V}\text{O}_{2\text{baseline}}$ ), expired  $\text{CO}_2$  at baseline ( $\dot{V}\text{CO}_{2\text{baseline}}$ ), and respiratory exchange ratio (RER) at baseline were defined as the mean values measured over the final 90 s of baseline pedaling. The end-exercise  $\dot{V}\text{O}_2$ ,  $\dot{V}\text{CO}_2$ , and RER were defined as the mean values measured over the final 30 s of exercise. The amplitude of the  $\dot{V}\text{O}_2$  response was calculated by subtracting  $\dot{V}\text{O}_{2\text{baseline}}$  from  $\dot{V}\text{O}_2$  at the end of exercise. Subsequently, the functional gain of the entire response was calculated by dividing the  $\dot{V}\text{O}_2$  amplitude by the change ( $\Delta$ ) in work rate. The amplitude of the  $\dot{V}\text{O}_2$  slow component during the severe-intensity exercise bout was estimated by subtracting the mean  $\dot{V}\text{O}_2$  at 2 min from the mean  $\dot{V}\text{O}_2$  at 6 min.

**Statistical analysis.** Two-way repeated-measures ANOVA was used to assess the difference in pulmonary gas-exchange variables, blood [lactate], and HR across dose (70, 140, and 280 ml) and treatment (PL and BR). Differences in pre- and postplasma  $[\text{NO}_2^-]$  and  $[\text{NO}_3^-]$  were assessed separately in PL and BR, across dose and time (pre and post) using two-way repeated-measures ANOVAs. Significant main and interaction effects were analyzed further using simple contrasts. Statistical significance was accepted at  $P < 0.05$ . Results are presented as mean  $\pm$  SD unless stated otherwise.

## RESULTS

Ingestion of BR was tolerated well by all subjects in S<sub>1</sub> and S<sub>2</sub>. Subjects did, however, report beeturia (red urine) and red stools, consistent with previous studies (3, 39). The absolute  $\text{NO}_3^-$  doses used in S<sub>1</sub> and S<sub>2</sub> (4.2, 8.4, and 16.8 mmol) were equivalent to  $\sim 0.05 \pm 0.01$  (range: 0.05–0.07),  $\sim 0.11 \pm 0.01$  (range: 0.09–0.13), and  $\sim 0.22 \pm 0.03$  mmol (range: 0.19–0.26)  $\text{NO}_3^-/\text{kg BM}$ , respectively.

### S<sub>1</sub>: Pharmacokinetics and Pharmacodynamics

The effects of different volumes of BR (and therefore, different amounts of ingested  $\text{NO}_3^-$ ) on plasma  $[\text{NO}_3^-]$  and  $[\text{NO}_2^-]$  are presented in Fig. 1. There were significant main effects by dose and time and an interaction effect for both plasma  $[\text{NO}_3^-]$  (Fig. 1A; all  $P < 0.01$ ) and plasma  $[\text{NO}_2^-]$  (Fig. 1B; all  $P < 0.01$ ).

At resting baseline, before the ingestion of any beverage, plasma  $[\text{NO}_3^-]$  was not significantly different between doses (Fig. 1A; all  $P > 0.05$ ). ANOVA analyses revealed significant dose-dependent increases in plasma  $[\text{NO}_3^-]$  following BR supplementation ( $P < 0.05$ ). The peak elevation above baseline in plasma  $[\text{NO}_3^-]$  occurred 1 h postadministration of 4.2 ( $160 \pm$






Fig. 1. Plasma nitrate concentration ( $[\text{NO}_3^-]$ ; A) and nitrite concentration ( $[\text{NO}_2^-]$ ; B) following consumption of water (control; ●) and 4.2 (▲), 8.4 (■), and 16.8 (◆) mmol  $\text{NO}_3^-$  (group mean  $\pm$  SE). Plasma  $[\text{NO}_3^-]$  and  $[\text{NO}_2^-]$  rose significantly in a dose-dependent manner. See text for further details. \*Significant difference from presupplementation baseline ( $P < 0.05$ ); <sup>a</sup>significant difference from control ( $P < 0.05$ ); <sup>b</sup>significant difference from 4.2 mmol  $\text{NO}_3^-$  ( $P < 0.05$ ); <sup>c</sup>significant difference from 8.4 mmol  $\text{NO}_3^-$  ( $P < 0.05$ ).

43  $\mu\text{M}$ ) and 8.4 mmol  $\text{NO}_3^-$  ( $269 \pm 92 \mu\text{M}$ ) and 2 h postadministration of 16.8 mmol  $\text{NO}_3^-$  ( $581 \pm 209 \mu\text{M}$ ; Fig. 1A; all  $P < 0.05$ ). Plasma  $[\text{NO}_3^-]$  remained elevated above baseline and CON at all time points after administration of 4.2, 8.4, and 16.8 mmol  $\text{NO}_3^-$  ( $P < 0.05$ ).

At baseline, before ingestion of any beverage, plasma  $[\text{NO}_2^-]$  was not significantly different between doses (Fig. 1B;  $P > 0.05$ ). ANOVA analyses revealed significant dose-dependent increases in plasma  $[\text{NO}_2^-]$  following BR supplementation ( $P < 0.05$ ). The peak elevation above baseline in plasma  $[\text{NO}_2^-]$  occurred 2 h postadministration of 4.2 ( $220 \pm 104 \text{ nM}$ ) and 8.4 mmol  $\text{NO}_3^-$  ( $374 \pm 173 \text{ nM}$ ) and 4 h postadministration of 16.8 mmol  $\text{NO}_3^-$  ( $653 \pm 356 \text{ nM}$ ; Fig. 1B; all  $P < 0.05$ ). Kinetic analyses revealed that plasma  $[\text{NO}_2^-]$  peaked significantly later ( $198 \pm 64 \text{ min}$ ; range: 130–367 min) following ingestion of 16.8 mmol relative to both 8.4 mmol ( $146 \pm 38 \text{ min}$ ; range: 77  $\pm$  213 min;  $P < 0.05$ ) and 4.2 mmol BR ( $106 \pm 39 \text{ min}$ ; range: 63–192 min;  $P < 0.05$ ). Peak plasma  $[\text{NO}_2^-]$ , following ingestion of 8.4 mmol, tended to occur later compared with 4.2 mmol ( $P = 0.06$ ). Plasma  $[\text{NO}_2^-]$  remained elevated above baseline and CON at 1, 2, 4, and 8 h after administration of 4.2, 8.4, and 16.8 mmol  $\text{NO}_3^-$  (all  $P < 0.05$ ). At 12 h, plasma  $[\text{NO}_2^-]$  remained elevated above baseline and 4.2 mmol BR following ingestion of 8.4 and 16.8 mmol  $\text{NO}_3^-$  (all  $P < 0.05$ ). In addition, plasma  $[\text{NO}_2^-]$  remained elevated at

24 h following administration of 16.8 mmol  $\text{NO}_3^-$  compared with all other doses ( $P < 0.05$ ).

The effects of different volumes of BR (and therefore, different amounts of ingested  $\text{NO}_3^-$ ) on systolic and diastolic BP and MAP are presented in Fig. 2. The changes in systolic BP across all conditions are presented in Fig. 2A. There were significant main effects by dose and time and an interaction effect on systolic BP (all  $P < 0.05$ ). Systolic BP at baseline, before administration of any beverage, was lower ( $P < 0.05$ ) in the 16.8-mmol  $\text{NO}_3^-$  condition ( $118 \pm 5 \text{ mmHg}$ ) relative to CON ( $121 \pm 5 \text{ mmHg}$ ) but not relative to 4.2 ( $119 \pm 6 \text{ mmHg}$ ) and 8.4 mmol  $\text{NO}_3^-$  ( $120 \pm 6 \text{ mmHg}$ ). Compared with baseline, systolic BP was lowered significantly following ingestion of 4.2, 8.4, and 16.8 mmol  $\text{NO}_3^-$  (all  $P < 0.05$ ). The peak reduction in systolic BP occurred 4 h postadministration of 4.2 ( $5 \pm 5 \text{ mmHg}$ ), 8.4 ( $10 \pm 5 \text{ mmHg}$ ), and 16.8 mmol  $\text{NO}_3^-$  ( $9 \pm 4 \text{ mmHg}$ ), respectively, relative to baseline (all  $P < 0.05$ ). Systolic BP was reduced relative to baseline, CON, and 4.2 mmol  $\text{NO}_3^-$ , at 2, 4, and 8 h postadministration of 8.4 mmol and 16.8 mmol  $\text{NO}_3^-$  (all  $P < 0.05$ ). There were no differences in systolic BP between 8.4 and 16.8 mmol  $\text{NO}_3^-$  at any time point ( $P > 0.05$ ). At 24 h, systolic BP remained significantly lower (by  $5 \pm 5 \text{ mmHg}$ ) than baseline, following consumption of 16.8 mmol  $\text{NO}_3^-$  ( $P < 0.05$ ). In contrast, systolic BP was not significantly different than CON or baseline at 24 h postad-




Fig. 2. Change ( $\Delta$ ) relative to presupplementation baseline in systolic blood pressure (BP; A), diastolic BP (B), and mean arterial pressure (MAP; C) following consumption of water (control;  $\bullet$ ) and 4.2 ( $\blacktriangle$ ), 8.4 ( $\blacksquare$ ), and 16.8 ( $\blacklozenge$ ) mmol  $\text{NO}_3^-$  (group mean  $\pm$  SE). \*Significant difference from presupplementation baseline ( $P < 0.05$ ); <sup>a</sup>significant difference from control ( $P < 0.05$ ); <sup>b</sup>significant difference from 4.2 mmol  $\text{NO}_3^-$  ( $P < 0.05$ ).

ministration of 4.2 and 8.4 mmol  $\text{NO}_3^-$  ( $P > 0.05$ ). Overall, the mean systolic BP across 24 h, relative to CON, was lowered dose dependently by  $\sim 3$ ,  $\sim 4$ , and  $\sim 6$  mmHg after administration of 4.2, 8.4, and 16.8 mmol  $\text{NO}_3^-$ , respectively (all  $P < 0.05$ ). The change in systolic BP was correlated with the change in plasma  $[\text{NO}_3^-]$  ( $r = -0.27$ ;  $P < 0.05$ ) and the change in plasma  $[\text{NO}_2^-]$  ( $r = -0.37$ ;  $P < 0.05$ ). The peak reduction in systolic BP was not correlated with the baseline systolic BP.

The changes in diastolic BP following the ingestion of different doses of  $\text{NO}_3^-$ -rich BR are presented in Fig. 2B. There was a significant interaction effect (dose  $\times$  time) on diastolic BP ( $P < 0.05$ ). Diastolic BP at baseline was not significantly different among conditions (CON:  $67 \pm 5$ ; 4.2 mmol:  $68 \pm 4$ ; 8.4 mmol:  $68 \pm 6$ ; 16.8 mmol:  $67 \pm 6$  mmHg;  $P > 0.05$ ). Follow-up tests revealed that ingestion of 8.4 and 16.8 but not

4.2 mmol  $\text{NO}_3^-$  reduced diastolic BP significantly, relative to baseline and CON (all  $P < 0.05$ ). The peak reduction in diastolic BP from baseline occurred at 4 h postadministration of 8.4 mmol  $\text{NO}_3^-$  ( $3 \pm 3$  mmHg) and 2 h postadministration of 16.8 mmol  $\text{NO}_3^-$  ( $4 \pm 4$  mmHg; both  $P < 0.05$ ) relative to baseline (both  $P > 0.05$ ) and returned to near-baseline values by 24 h ( $P > 0.05$ ). There were no differences in diastolic BP between 8.4 and 16.8 mmol  $\text{NO}_3^-$  at any time point ( $P > 0.05$ ). The change in diastolic BP was correlated with the change in plasma  $[\text{NO}_3^-]$  ( $r = -0.35$ ;  $P < 0.05$ ) and the change in plasma  $[\text{NO}_2^-]$  ( $r = -0.39$ ;  $P < 0.05$ ). Moreover, the peak change in diastolic BP was correlated with the baseline diastolic BP ( $r = -0.49$ ;  $P < 0.05$ ).

The changes in MAP following the ingestion of different doses of  $\text{NO}_3^-$ -rich BR are presented in Fig. 2C. There were significant main effects by dose and time and an interaction effect on MAP (all  $P < 0.05$ ). At baseline, before the ingestion of any beverage, MAP was not significantly different among conditions (CON:  $85 \pm 4$ ; 4.2 mmol:  $85 \pm 4$ ; 8.4 mmol:  $85 \pm 5$ ; 16.8 mmol:  $84 \pm 5$  mmHg;  $P > 0.05$ ). MAP was significantly lower following ingestion of 4.2, 8.4, and 16.8 mmol  $\text{NO}_3^-$  relative to baseline and CON (all  $P < 0.05$ ). Following ingestion of 4.2 mmol  $\text{NO}_3^-$ , the peak reduction ( $2 \pm 2$  mmHg) in MAP occurred at 1 h, and MAP remained reduced by  $\sim 2$  mmHg at 2 h relative to baseline ( $P < 0.05$ ). In contrast, the peak reduction in MAP ( $5 \pm 3$  mmHg) occurred 4 h postadministration of 8.4 and 16.8 mmol  $\text{NO}_3^-$  relative to baseline ( $P < 0.05$ ). MAP was not different between 8.4 and 16.8 mmol  $\text{NO}_3^-$  at any time point ( $P > 0.05$ ). Overall, the mean MAP across 24 h, relative to CON, was reduced dose dependently by  $\sim 1$ ,  $\sim 2$ , and  $\sim 4$  mmHg after administration of 4.2, 8.4, and 16.8 mmol  $\text{NO}_3^-$ , respectively (all  $P < 0.05$ ). The change in MAP was correlated significantly with the change in plasma  $[\text{NO}_3^-]$  ( $r = -0.35$ ;  $P < 0.05$ ) and the change in plasma  $[\text{NO}_2^-]$  ( $r = -0.41$ ;  $P < 0.05$ ).

## S<sub>2</sub>: Dose Response

**Plasma  $[\text{NO}_3^-]$  and  $[\text{NO}_2^-]$ .** The group mean plasma  $[\text{NO}_3^-]$  and  $[\text{NO}_2^-]$  responses in the BR and PL conditions are illustrated in Fig. 3, A and B, respectively. Presupplementation plasma  $[\text{NO}_3^-]$  was not significantly different between conditions ( $P > 0.05$ ), and no significant change in plasma  $[\text{NO}_3^-]$  was observed following PL supplementation ( $P > 0.05$ ). ANOVA analyses revealed a significant dose-dependent increase in plasma  $[\text{NO}_3^-]$  at 2.5 h following BR supplementation ( $P < 0.05$ ). An elevation in plasma  $[\text{NO}_3^-]$  above baseline was apparent following 4.2 ( $130 \pm 17$   $\mu\text{M}$ ;  $P < 0.05$ ), 8.4 ( $282 \pm 54$   $\mu\text{M}$ ;  $P < 0.05$ ), and 16.8 mmol  $\text{NO}_3^-$  ( $580 \pm 89$   $\mu\text{M}$ ;  $P < 0.05$ ). Presupplementation plasma  $[\text{NO}_2^-]$  was not significantly different among conditions ( $P > 0.05$ ), and no significant change in plasma  $[\text{NO}_2^-]$  was observed following PL supplementation ( $P > 0.05$ ). ANOVA analyses revealed a significant dose-dependent increase in plasma  $[\text{NO}_2^-]$  at 2.5 h following BR supplementation ( $P < 0.05$ ). Following administration of 4.2, 8.4, and 16.8 mmol  $\text{NO}_3^-$ , plasma  $[\text{NO}_2^-]$  was elevated above baseline by  $150 \pm 73$  nM,  $291 \pm 145$  nM, and  $425 \pm 225$  nM, respectively (all  $P < 0.05$ ). Plasma  $[\text{NO}_2^-]$  was significantly greater after ingestion of 16.8 mmol compared with 4.2 mmol  $\text{NO}_3^-$  ( $P < 0.05$ ) and tended to be greater compared with 8.4 mmol  $\text{NO}_3^-$  ( $P = 0.06$ ). Plasma  $[\text{NO}_2^-]$  was




Fig. 3. Mean  $\pm$  SE plasma  $[\text{NO}_3^-]$  (A) and  $[\text{NO}_2^-]$  (B) preingestion (black bars) and 2.5-h postingestion (gray bars) of 70, 140, and 280 ml  $\text{NO}_3^-$ -rich beetroot juice (BR) ( $\text{NO}_3^-$ ) or  $\text{NO}_3^-$ -depleted BR [placebo (PL)]. See text for further details. \*Significant difference from baseline ( $P < 0.05$ ); <sup>a</sup>significant difference postconsumption of 70 ml  $\text{NO}_3^-$ -rich BR ( $P < 0.05$ ); <sup>b</sup>significant difference from postconsumption of 140 ml  $\text{NO}_3^-$ -rich BR ( $P < 0.05$ ).

significantly greater following ingestion of 8.4 mmol  $\text{NO}_3^-$  compared with 4.2 mmol  $\text{NO}_3^-$  ( $P < 0.05$ ).

**Moderate-intensity exercise.** The pulmonary gas exchange and ventilatory responses to moderate-intensity exercise across all doses and conditions are summarized in Table 1. The  $\dot{V}_{\text{O}_2}$  measured during the period of baseline cycling at 20 W was not affected by dose or condition ( $P > 0.05$ ). However, the absolute end-exercise  $\dot{V}_{\text{O}_2}$ , measured over the final 30 s of moderate-intensity exercise, was altered significantly by BR ingestion ( $P < 0.05$ ; Fig. 4A). Follow-up tests indicated that end-exercise  $\dot{V}_{\text{O}_2}$  was lowered significantly by  $\sim 3\%$  following administration of 16.8 mmol  $\text{NO}_3^-$  relative to the respective PL (PL280:  $1.65 \pm 0.19$  vs. BR280,  $1.60 \pm 0.23$  l/min;  $P < 0.05$ ). In addition, there was a trend toward a significant reduction ( $\sim 2\%$ ) in end-exercise  $\dot{V}_{\text{O}_2}$  following administration of 8.4 mmol  $\text{NO}_3^-$  relative to the respective PL (PL140:  $1.67 \pm 0.21$  vs. BR140,  $1.64 \pm 0.23$  l/min;  $P = 0.06$ ). The change in plasma  $[\text{NO}_2^-]$  from baseline to postingestion of 4.2, 8.4, and 16.8 mmol  $\text{NO}_3^-$  was correlated with the change in end-exercise  $\dot{V}_{\text{O}_2}$  ( $r = -0.47$ ;  $P < 0.05$ ). There was no significant difference in end-exercise  $\dot{V}_{\text{O}_2}$  following ingestion of 4.6 mmol  $\text{NO}_3^-$  (BR70) compared with PL70 ( $P > 0.05$ ).

The amplitude of the  $\dot{V}_{\text{O}_2}$  response (end-exercise  $-\dot{V}_{\text{O}_2\text{baseline}}$ ; Table 1) was affected by dose ( $P < 0.05$ ) and tended to be affected by condition ( $P = 0.07$ ). Follow-up tests revealed that there was a trend toward a significant reduction in the  $\dot{V}_{\text{O}_2}$  amplitude (by  $\sim 6\%$ ) after administration of 16.8 mmol  $\text{NO}_3^-$  compared with 8.4 mmol  $\text{NO}_3^-$  (BR140:  $0.70 \pm 0.16$  vs.

BR280:  $0.66 \pm 0.16$  l/min;  $P = 0.06$ ). The change in plasma  $[\text{NO}_2^-]$  from baseline to postingestion of 4.2, 8.4, and 16.8 mmol  $\text{NO}_3^-$  was correlated with the change in  $\dot{V}_{\text{O}_2}$  amplitude ( $r = -0.38$ ;  $P < 0.05$ ). There was no significant difference in  $\dot{V}_{\text{O}_2}$  amplitude between PL and BR at any dose ( $P > 0.05$ ).

The  $\dot{V}_{\text{CO}_2\text{baseline}}$ , measured over the last 90 s of 20 W pedaling, and the end-exercise  $\dot{V}_{\text{CO}_2}$ , measured over the last 30 s of exercise, were affected by dose ( $P < 0.05$  for both) but not condition ( $P > 0.05$  for both; Table 1). Follow-up tests revealed that  $\dot{V}_{\text{CO}_2\text{baseline}}$  was increased significantly, as the volume of supplement ingested increased ( $P < 0.05$ ), irrespective of the condition (i.e., PL or BR). Specifically,  $\dot{V}_{\text{CO}_2\text{baseline}}$  was increased by  $\sim 7\%$  and  $\sim 5\%$  following consumption of 280 ml of supplement relative to 70 and 140 ml, respectively ( $P < 0.05$  for both). There were no significant differences in  $\dot{V}_{\text{CO}_2}$  between the ingestion of 70 and 140 ml of supplement ( $P > 0.05$ ). Furthermore, post hoc analysis revealed that the end-exercise  $\dot{V}_{\text{CO}_2}$  was significantly higher following ingestion of both 140 and 280 ml of supplement relative to 70 ml ( $P < 0.01$  for both). There was, however, no significant difference in end-exercise  $\dot{V}_{\text{CO}_2}$  between ingestion of 140 and 280 ml of supplement ( $P > 0.05$ ).

Baseline and end-exercise RER were affected by dose ( $P < 0.05$  for both) but not condition ( $P > 0.05$ ). The follow-up tests indicated that RER increased as the volume of supplement ingested increased ( $P < 0.05$ ; Table 1). Specifically, RER at baseline was increased by  $\sim 5\%$  and  $\sim 4\%$ , following consumption of 280 ml of supplement relative to 70 and 140 ml, respectively ( $P < 0.05$  for both). Although there was no significant interaction effect or main effect by condition, baseline RER tended to be higher (by  $\sim 3\%$ ) following administration of 16.8 mmol  $\text{NO}_3^-$  compared with the respective PL ( $P = 0.08$ ). End-exercise RER was increased significantly by  $\sim 4\%$  and  $\sim 3\%$ , following consumption of 280 ml compared with 70 and 140 ml of supplement, respectively ( $P < 0.05$  for both). In addition, the ingestion of 140 ml increased end-exercise RER compared with ingestion of 70 ml of supplement ( $P < 0.05$ ). The baseline, end-exercise, and change in blood [lactate] and HR were not altered significantly by dose or condition (Table 2;  $P > 0.05$ ).

**Severe-intensity exercise.** The pulmonary gas exchange and ventilatory responses to severe-intensity exercise across all doses and conditions are summarized in Table 1. In contrast to the effects observed for moderate-intensity exercise, the  $\dot{V}_{\text{O}_2}$  and  $\dot{V}_{\text{CO}_2}$  measured at baseline and at task failure were not altered by dose or treatment (all  $P > 0.05$ ). Moreover, neither the dose nor the treatment altered the  $\dot{V}_{\text{O}_2}$  slow component amplitude ( $P > 0.05$  for both). There was a trend toward significant main effects by dose ( $P = 0.09$ ) and treatment ( $P = 0.08$ ) but no interaction effect on RER at baseline ( $P > 0.05$ ). Follow-up tests revealed that there was a trend toward significant increases in RER at baseline by  $\sim 4\%$  and  $\sim 3\%$  following consumption of 280 ml of supplement compared with the consumption of 70 ( $P = 0.06$ ) or 140 ml ( $P = 0.08$ ) of supplement, respectively. RER, at task failure, was not altered by dose or treatment ( $P > 0.05$ ). The baseline, end-exercise, and change in blood [lactate] and HR were not altered significantly by dose or condition (Table 2;  $P > 0.05$ ).

There was a significant main effect by condition ( $P < 0.05$ ) but not dose ( $P > 0.05$ ) on time-to-task failure (Table 1 and

Table 1. Pulmonary gas-exchange variables during moderate- and severe-intensity exercise following supplementation with 3 different volumes of beetroot juice and placebo

	70 ml		140 ml		280 ml	
	Placebo	Nitrate, 4.2 mmol	Placebo	Nitrate, 8.4 mmol	Placebo	Nitrate, 16.8 mmol
Moderate-intensity exercise						
$\dot{V}O_2$						
Baseline, l/min	0.94 ± 0.10	0.93 ± 0.09	0.92 ± 0.12	0.94 ± 0.13	0.95 ± 0.12	0.94 ± 0.08
End-exercise, l/min	1.64 ± 0.21	1.61 ± 0.21	1.67 ± 0.21 <sup>a</sup>	1.64 ± 0.23	1.65 ± 0.19	1.60 ± 0.18 <sup>b</sup>
Primary amplitude, l/min	0.70 ± 0.16	0.68 ± 0.16	0.74 ± 0.16	0.70 ± 0.16	0.70 ± 0.14	0.66 ± 0.16
Primary gain, ml · min <sup>-1</sup> · W <sup>-1</sup>	9.5 ± 1.0	9.2 ± 1.1	10.1 ± 0.9	9.5 ± 0.9	9.6 ± 0.6	9.0 ± 1.1 <sup>c</sup>
$\dot{V}CO_2$						
Baseline, l/min	0.82 ± 0.07	0.81 ± 0.05	0.82 ± 0.09	0.83 ± 0.12	0.86 ± 0.07 <sup>a</sup>	0.89 ± 0.07 <sup>a,d</sup>
End-exercise, l/min	1.48 ± 0.17	1.45 ± 0.17	1.51 ± 0.17	1.50 ± 0.17	1.52 ± 0.14 <sup>a</sup>	1.52 ± 0.17 <sup>d</sup>
$\dot{V}_E$						
Baseline, l/min	23 ± 3	22 ± 2	23 ± 3	23 ± 4	24 ± 3	23 ± 2
End-exercise, l/min	37 ± 5	36 ± 5	37 ± 5	37 ± 5	38 ± 5	37 ± 4
RER						
Baseline	0.88 ± 0.05	0.88 ± 0.04	0.89 ± 0.04	0.89 ± 0.04	0.91 ± 0.05	0.94 ± 0.04 <sup>c,d</sup>
End-exercise	0.91 ± 0.04	0.90 ± 0.04	0.91 ± 0.03	0.92 ± 0.05	0.93 ± 0.04	0.95 ± 0.04 <sup>c,d</sup>
Severe-intensity exercise						
$\dot{V}O_2$						
Baseline, l/min	1.00 ± 0.10	0.99 ± 0.11	0.99 ± 0.13	0.99 ± 0.11	0.99 ± 0.11	0.97 ± 0.11
End-exercise, l/min	3.89 ± 0.40	3.97 ± 0.34	3.96 ± 0.38	3.99 ± 0.40	3.98 ± 0.35	3.94 ± 0.28
Overall gain, ml · min <sup>-1</sup> · W <sup>-1</sup>	12.1 ± 0.8	12.5 ± 1.0	12.5 ± 0.9	12.6 ± 1.2	12.6 ± 0.9	12.5 ± 0.8
Slow-phase amplitude, 6–2 min; l/min	0.66 ± 0.14	0.65 ± 0.15	0.67 ± 0.17	0.62 ± 0.17	0.75 ± 0.09	0.69 ± 0.11
$\dot{V}CO_2$						
Baseline, l/min	0.91 ± 0.06	0.90 ± 0.08	0.89 ± 0.08	0.91 ± 0.09	0.92 ± 0.08	0.93 ± 0.11
End-exercise, l/min	4.16 ± 0.36	4.17 ± 0.27	4.21 ± 0.38	4.18 ± 0.32	4.20 ± 0.25	4.20 ± 0.31
RER						
Baseline	0.91 ± 0.05	0.91 ± 0.06	0.91 ± 0.06	0.92 ± 0.06	0.94 ± 0.05	0.96 ± 0.05
End-exercise	1.07 ± 0.06	1.05 ± 0.05	1.06 ± 0.05	1.05 ± 0.05	1.06 ± 0.05	1.07 ± 0.05
Time-to-task failure(s)	470 ± 81	508 ± 102	498 ± 113	570 ± 153 <sup>e</sup>	493 ± 114	552 ± 117 <sup>b</sup>

Values are means ± SD.  $\dot{V}O_2$ , oxygen uptake;  $\dot{V}CO_2$ , expired carbon dioxide;  $\dot{V}_E$ , ventilation; RER, respiratory exchange ratio. <sup>a</sup>Significantly different from placebo (PL)70 ( $P < 0.05$ ); <sup>b</sup>significantly different from PL280 ( $P < 0.05$ ); <sup>c</sup>Significantly different from beetroot juice (BR)140 ( $P < 0.05$ ); <sup>d</sup>significantly different from BR70 ( $P < 0.05$ ); <sup>e</sup>significantly different from PL140 ( $P < 0.05$ ).

Fig. 4B). Follow-up tests revealed that consumption of 8.4 mmol  $NO_3^-$  (BR140) and 16.8 mmol  $NO_3^-$  (BR280) resulted in a significant increase in time-to-task failure by  $71 \pm 77$  s and  $59 \pm 61$  s, respectively, relative to PL140 and PL280 ( $P < 0.05$ ; Fig. 4B). There was no difference in time-to-task failure between BR70 and PL70 ( $P > 0.05$ ). The change in plasma  $[NO_2^-]$  from baseline to postingestion of 4.2, 8.4, and 16.8 mmol  $NO_3^-$  was correlated significantly with the change in time-to-task failure ( $r = 0.55$ ;  $P < 0.05$ ). There was no significant difference in time-to-task failure among 4.2, 8.4, and 16.8 mmol BR (all  $P > 0.05$ ) or among PL70, PL140, and PL280 ( $P > 0.05$ ).

In terms of positive changes in time-to-task failure, there were three “nonresponders” in the 4.2-mmol condition, two in the 8.4-mmol condition, and one in the 16.8-mmol condition. Individual subjects who did not respond at lower doses did respond at higher doses. The increase in plasma  $[NO_2^-]$  from baseline to pre-exercise for the nonresponders was similar to the other subjects who did respond. For example, the three nonresponders at the lowest  $NO_3^-$  dose had an increase in plasma  $[NO_2^-]$  of 140, 208, and 161 nM compared with a group mean increase of 150 nM. In addition, the nonresponders did not have high baseline values of plasma  $[NO_2^-]$  (70–121 nM) compared with the group mean.

## DISCUSSION

This study is the first to characterize the pharmacokinetic-pharmacodynamic effects of  $NO_3^-$ -rich BR ingestion and to

investigate the dose-response relationship between BR ingestion and the physiological responses to exercise. Specifically, we studied how acute ingestion of three different BR volumes (and thus three different  $NO_3^-$  doses) impacted on plasma  $[NO_3^-]$  and  $[NO_2^-]$ , resting BP, the pulmonary gas-exchange responses to moderate- and severe-intensity exercise, and exercise tolerance. Our principal findings were that plasma  $[NO_3^-]$  and  $[NO_2^-]$  increased dose dependently up to 16.8 mmol  $NO_3^-$  with there being a dose-dependent peak reduction in BP up to 8.4 mmol  $NO_3^-$ . A  $NO_3^-$  dose of 16.8 mmol was required to elicit a significant reduction in the  $O_2$  cost of moderate-intensity cycle exercise, although there was a trend ( $P = 0.06$ ) for a reduction with 8.4 mmol. A significant improvement in time-to-task failure during severe-intensity exercise was evident after ingestion of 8.4 mmol  $NO_3^-$ , with no further benefits observed following the ingestion of 16.8 mmol  $NO_3^-$ .

### *S<sub>1</sub>: BR Pharmacokinetics and Pharmacodynamics—Effects on Plasma $[NO_3^-]$ , $[NO_2^-]$ , and BP*

The results of *S<sub>1</sub>* demonstrated that concentrated BR consumption causes dose-dependent increases in plasma  $[NO_3^-]$  and  $[NO_2^-]$ . Plasma  $[NO_3^-]$  increased by approximately five- and eightfold, 1 h after the ingestion of 4.2 and 8.4 mmol  $NO_3^-$ , and by ~18-fold, 2 h after the ingestion of 16.8 mmol  $NO_3^-$ . In contrast, the increase in plasma  $[NO_2^-]$  occurred later, peaking at approximately 2–2.5 h postadministration of 4.2 and 8.4 mmol  $NO_3^-$  and ~3 h postadministration of 16.8 mmol  $NO_3^-$ .






Fig. 4. Mean  $\pm$  SE steady-state oxygen consumption ( $\dot{V}O_2$ ) during moderate-intensity exercise (A) and time-to-task failure during severe-intensity exercise (B), following consumption of 70, 140, and 280 ml  $NO_3^-$ -rich BR (gray bars) or  $NO_3^-$ -depleted BR (PL; black bars). End-exercise  $\dot{V}O_2$  during moderate-intensity exercise was reduced significantly following the ingestion of 280 ml BR. Time-to-task failure during severe-intensity exercise was extended after consumption of 140 ml BR with no further increase following 280 ml BR. \*Significant difference from PL ( $P < 0.05$ ).

As expected, the rise in plasma  $[NO_2^-]$  was smaller compared with plasma  $[NO_3^-]$ , with peak increases of  $\sim 2.5$ -fold, approximately fourfold, and approximately eightfold, respectively. The delayed peak increases in plasma  $[NO_2^-]$  compared with plasma  $[NO_3^-]$  reflect the importance of the enterosalivary circulation and subsequent reduction of  $NO_3^-$  to  $NO_2^-$  by

lingual bacteria (16, 39). These pharmacokinetic responses to BR supplementation are consistent with those reported previously following acute ingestion of  $KNO_3$  (19). Together, these data suggest that the pharmacokinetics of plasma  $[NO_3^-]$  and  $[NO_2^-]$  are dose dependent when  $NO_3^-$  is administered, either as  $NO_3^-$  salt or in the form of a natural vegetable supplement.

Ingestion of concentrated BR dose dependently lowered systolic BP and MAP up to an intake of 8.4 mmol  $NO_3^-$ . More specifically, acute ingestion of 4.2, 8.4, and 16.8 mmol inorganic  $NO_3^-$ , administered in the form of BR, resulted in peak reductions of systolic BP of  $\sim 5$ ,  $\sim 10$ , and  $\sim 9$  mmHg and peak reductions of MAP of  $\sim 2$ ,  $\sim 5$ , and  $\sim 5$  mmHg, respectively. Moreover, BR ingestion resulted in a similar “threshold” effect on diastolic BP, with peak reductions of  $\sim 3$  and  $\sim 4$  mmHg following administration of 8.4 and 16.8 mmol  $NO_3^-$ ; however, ingestion of 4.2 mmol  $NO_3^-$  did not reduce diastolic BP significantly. These reductions in BP are similar to those reported by Kapil et al. (19) following acute administration of  $KNO_3$ , except that Kapil et al. (19) reported a dose-dependent reduction in BP up to 24 mmol  $KNO_3$ . The reason for this discrepancy between studies is unclear. Interestingly, compared with Kapil et al. (19), who reported 6 mmHg and 9 mmHg reductions in systolic BP following the consumption of 12 mmol and 24 mmol  $KNO_3$ , respectively, we observed larger reductions in BP following the consumption of BR (e.g., a peak reduction of 10 mmHg in systolic BP with 8.4 mmol  $NO_3^-$  contained in 140 ml BR). It is possible that this apparent greater potency of BR compared with  $NO_3^-$  salt in reducing BP is related to the polyphenols and other antioxidants present in BR, which may facilitate a more efficient conversion of  $NO_3^-$  to  $NO_2^-$  (30). Interestingly, although the peak reduction in BP was not significantly different between 8.4 and 16.8 mmol  $NO_3^-$ , the mean reduction in BP over 24 h was dose dependent, with MAP, for example, reduced by 1, 2, and 4 mmHg following administration of 4.2, 8.4, and 16.8 mmol  $NO_3^-$ , respectively.

The results of the present study suggest that BR (and presumably other  $NO_3^-$ -rich vegetable) consumption can provide a natural approach to maintaining or improving BP and

Table 2. Heart rate and blood lactate responses to moderate- and severe-intensity exercise following supplementation with 3 different volumes of beetroot juice and placebo

	70 ml		140 ml		280 ml	
	Placebo	Nitrate, 4.2 mmol	Placebo	Nitrate, 8.4 mmol	Placebo	Nitrate, 16.8 mmol
Moderate-intensity exercise						
Heart rate, beats/min						
Baseline	89 $\pm$ 9	89 $\pm$ 8	88 $\pm$ 8	88 $\pm$ 8	89 $\pm$ 8	89 $\pm$ 6
End-exercise	116 $\pm$ 11	116 $\pm$ 12	115 $\pm$ 10	116 $\pm$ 8	115 $\pm$ 9	115 $\pm$ 10
Blood [lactate], mM						
Baseline	1.1 $\pm$ 0.3	1.1 $\pm$ 0.5	1.1 $\pm$ 0.4	1.0 $\pm$ 0.4	1.0 $\pm$ 0.3	1.1 $\pm$ 0.3
End-exercise	1.2 $\pm$ 0.2	1.2 $\pm$ 0.5	1.2 $\pm$ 0.5	1.1 $\pm$ 0.4	1.1 $\pm$ 0.4	1.2 $\pm$ 0.5
$\Delta$	0.1 $\pm$ 0.2	0.1 $\pm$ 0.4	0.1 $\pm$ 0.3	0.1 $\pm$ 0.2	0.1 $\pm$ 0.3	0.1 $\pm$ 0.2
Severe-intensity exercise						
Heart rate, beats/min						
Baseline	99 $\pm$ 9	100 $\pm$ 8	99 $\pm$ 9	100 $\pm$ 10	100 $\pm$ 10	99 $\pm$ 8
End-exercise	186 $\pm$ 11	186 $\pm$ 12	185 $\pm$ 12	187 $\pm$ 10	186 $\pm$ 11	185 $\pm$ 10
Blood [lactate], mM						
Baseline	0.9 $\pm$ 0.4	0.9 $\pm$ 0.4	0.9 $\pm$ 0.4	0.9 $\pm$ 0.6	1.0 $\pm$ 0.3	0.9 $\pm$ 0.2
Task failure	9.7 $\pm$ 1.4	9.4 $\pm$ 1.6	9.4 $\pm$ 1.6	9.6 $\pm$ 1.8	9.5 $\pm$ 1.5	9.5 $\pm$ 1.1
$\Delta$	8.7 $\pm$ 1.2	8.5 $\pm$ 1.7	8.5 $\pm$ 1.7	8.7 $\pm$ 1.3	8.5 $\pm$ 1.4	8.6 $\pm$ 1.2

Values are means  $\pm$  SD. [lactate], lactate concentration;  $\Delta$ , change.



vascular health in young adults. The reductions in BP, evident in the present study, are noteworthy. For example, it has been suggested that lowering systolic BP by 10 mmHg may reduce the risk of ischemic heart disease by ~25% and the risk of stroke by ~35% (27–29, 31). The beneficial hemodynamic effects of  $\text{NO}_3^-$  supplementation are thought to be due to the reduction of  $\text{NO}_3^-$  to  $\text{NO}_2^-$  and then to NO within the blood vessel (13), resulting in arterial dilatation and a reduced peripheral resistance (39). However, it is possible that  $\text{NO}_2^-$  itself may also exert a direct effect on the vascular system, independent of NO formation (1). There are several advantages to using inorganic rather than organic  $\text{NO}_3^-$  for the prevention or treatment of hypertension (33). These include a slow and controlled increase in plasma  $[\text{NO}_2^-]$  following inorganic  $\text{NO}_3^-$  intake (due to  $\text{NO}_3^-$  uptake into the enterosalivary circulation) compared with the more abrupt changes in plasma  $[\text{NO}_2^-]$  (perhaps to toxic levels) and BP, which can occur with organic  $\text{NO}_3^-$  administration (33). Moreover, unlike the chronic administration of organic  $\text{NO}_3^-$ , inorganic  $\text{NO}_3^-$  does not appear to lead to the development of tolerance (37) and endothelial dysfunction (33).

### *S<sub>2</sub>: Dose Response*

The results of *S<sub>2</sub>* confirm that concentrated BR consumption causes a dose-dependent increase in plasma  $[\text{NO}_3^-]$  by 334%, 778%, and 1,556% and plasma  $[\text{NO}_2^-]$  by 121%, 218%, and 338%, 2.5 h postingestion of 4.2, 8.4, and 16.8 mmol  $\text{NO}_3^-$ , respectively. The magnitude of the increase in plasma  $[\text{NO}_2^-]$  following consumption of 8.4 and 16.8 mmol  $\text{NO}_3^-$  in the present study was much larger than the approximate 15–150% rise in plasma  $[\text{NO}_2^-]$ , reported previously, following acute (approximately 4–6 mmol) (5, 21, 25, 37) and chronic (approximately 5–6 mmol/day) (2, 3, 22, 26, 37) dietary  $\text{NO}_3^-$  supplementation. This finding is likely a consequence of the relatively higher  $\text{NO}_3^-$  doses (8.4 and 16.8 mmol  $\text{NO}_3^-$ ) administered in the present study. Interestingly, the group mean plasma  $[\text{NO}_3^-]$  and  $[\text{NO}_2^-]$  reported in *S<sub>2</sub>* are somewhat lower than those reported at 2–4 h postingestion of BR in *S<sub>1</sub>*. Given that there was distinct subject recruitment for *S<sub>1</sub>* and *S<sub>2</sub>*, it is likely that this discrepancy is due to individual variations in the pharmacokinetic response to BR consumption. For example, when the individual plasma  $[\text{NO}_2^-]$  responses to the ingestion of 16.8 mmol  $\text{NO}_3^-$  in *S<sub>1</sub>* are considered, peak concentrations ranged from 493 to 1,523 nM, and the time-to-peak concentration ranged from 130 to 367 min. The cause of this wide interindividual variability in the response of plasma  $[\text{NO}_2^-]$  to  $\text{NO}_3^-$  ingestion is unclear, although it may depend, in part, on salivary flow rate; also, it is known that the reduction of  $\text{NO}_3^-$  to  $\text{NO}_2^-$  is highly dependent on the activity of oral bacteria (16, 39). Another consideration is that the absolute  $\text{NO}_3^-$  doses administered in the present study (4.2, 8.4, and 16.8 mmol in 1, 2, and 4 BR shots, respectively) resulted in somewhat different  $\text{NO}_3^-$  doses when expressed relative to BM (0.05–0.07, 0.09–0.13, and 0.19–0.25 mmol  $\text{NO}_3^-$ /kg BM, respectively).

### *Dose Response: Moderate-Intensity Exercise*

This is the first study to assess the acute dose-dependent physiological responses to exercise following dietary  $\text{NO}_3^-$  supplementation in humans. We assessed the acute response to three different doses of BR at 2.5 h postingestion, based on the

significant dose-dependent elevation in plasma  $[\text{NO}_2^-]$  observed at 2–3 h postingestion in *S<sub>1</sub>* (Fig. 1B). The steady-state  $\dot{V}\text{O}_2$  measured over the final 30 s of moderate-intensity cycle exercise was unaffected by 4.2 mmol  $\text{NO}_3^-$ , tended to be lower (~30 ml/min) following administration of 8.4 mmol  $\text{NO}_3^-$ , and was reduced significantly (by ~50 ml/min) following administration of 16.8 mmol  $\text{NO}_3^-$ .

The reduction in steady-state  $\dot{V}\text{O}_2$  (~3%), observed following acute ingestion of 16.8 mmol  $\text{NO}_3^-$  (~0.23 mmol/kg BM), is similar to that reported 2.5 h postingestion of 5.2 mmol  $\text{NO}_3^-$  (~0.07 mmol/kg BM) in the form of nonconcentrated BR (37) but is smaller than the 6% reduction reported 1 h postingestion of 0.033 mmol/kg BM sodium nitrate (25). In contrast to acute ingestion, longer-term BR supplementation (3–6 days at approximately 5–7 mmol  $\text{NO}_3^-$ /day) resulted in an approximate 5–7% reduction in steady-state  $\dot{V}\text{O}_2$  during moderate-intensity cycling (3, 26) and running (22).

Previous studies have indicated that the lowering of submaximal exercise  $\dot{V}\text{O}_2$ , following dietary  $\text{NO}_3^-$  supplementation, may result from improved mitochondrial efficiency (25) and/or a reduction in the ATP cost of muscle force production (4). Alterations in protein expression have been proposed as the mechanistic basis for these effects (17, 24); however, it is unlikely that these alterations occur quickly enough to explain the effects observed so soon (1–2.5 h) after  $\text{NO}_3^-$  ingestion (25, 37). Alternatively, NO may acutely and reversibly impact protein function through post-translational protein modifications. For instance, S-nitrosation of adenine nucleotide translocase or other mitochondrial or calcium-handling proteins (35) may contribute to the acute reduction in  $\text{O}_2$  cost of exercise following BR ingestion. The mechanistic basis for the acute changes in the  $\text{O}_2$  cost of exercise following BR ingestion warrants further investigation.

An interesting observation was the dose-dependent increase in baseline and end-exercise  $\dot{V}\text{CO}_2$ , irrespective of condition (i.e., PL or BR). This small but significant rise in  $\dot{V}\text{CO}_2$  led to a dose-dependent increase in RER that was more pronounced during baseline cycling compared with the exercising steady-state. An elevation in RER is indicative of a shift in substrate use toward a relatively greater reliance on carbohydrate and is likely due to the sugar content of the concentrated BR and PL beverages (~16 g/70 ml).

### *Dose Response: Severe-Intensity Exercise*

A novel finding of the present study was that 8.4 and 16.8 mmol  $\text{NO}_3^-$ , but not 4.2 mmol  $\text{NO}_3^-$ , administered acutely in the form of concentrated BR, significantly improved the time-to-task failure by 14% and 12%, respectively, during severe-intensity exercise. These findings are similar to the 14–16% improvement in exercise tolerance reported previously following 5–6 days of BR supplementation at a lower dose (5–6 mmol  $\text{NO}_3^-$ ) (3, 22). Although the mechanism(s) responsible for the ergogenic potential of  $\text{NO}_3^-$  supplementation remain uncertain, they are believed to be mediated via a biochemical reduction of ingested  $\text{NO}_3^-$  to biologically active  $\text{NO}_2^-$  and NO (4).

NO has been linked to the efficiency of aerobic respiration (9) and the regulation of muscle contraction (35). Indeed, both more efficient mitochondrial oxidative phosphorylation, via a reduced proton leak across the inner mitochondrial membrane

(24) and a reduced ATP and phosphocreatine cost of muscle force production (2, 15), has been reported following dietary  $\text{NO}_3^-$  supplementation. In addition, recent evidence suggests that BR supplementation results in a marked increase in muscle blood flow during exercise in rats, with the blood flow preferentially distributed to muscle groups that principally contain *type II* fibers, which are recruited during severe-intensity exercise (14). Furthermore,  $\text{NO}_3^-$  supplementation has been shown to increase muscle force production in mice via modulation of intracellular calcium ion ( $\text{Ca}^{2+}$ ) handling in fast-twitch fibers (17). It is possible that these mechanisms operate simultaneously and/or synergistically, resulting in enhanced exercise tolerance. It is, however, important to note that the studies that demonstrated effects of  $\text{NO}_3^-$  supplementation on muscle metabolic and vascular control mechanisms (2, 14, 17, 24) used chronic, rather than acute,  $\text{NO}_3^-$  supplementation protocols. On the other hand, Cosby et al. (10) reported acutely increased blood flow to exercising forearm muscle following infusion of  $\text{NO}_2^-$  into the brachial artery. It is possible that the improved time-to-task failure that we observed with 8.4 and 16.8 mmol  $\text{NO}_3^-$  was related to improved blood flow to muscle or to a NO-mediated enhancement of local matching of  $\text{O}_2$  delivery to metabolic rate. This would be consistent with reports that BR supplementation results in a preferential distribution of blood flow to *type II* fibers (14) and improves oxidative function in hypoxic muscle (38). The lack of a further improvement in time-to-task failure with 16.8 mmol compared to 8.4 mmol  $\text{NO}_3^-$  mirrors the lack of an additional effect of consuming the higher  $\text{NO}_3^-$  dose on the peak reduction in BP that we observed in  $S_1$ , suggesting that the acute effects of BR ingestion on exercise tolerance may be related, at least in part, to effects on the vasculature. Further studies are needed to establish which mechanisms may be responsible for the ergogenic potential of  $\text{NO}_3^-$ , at least at high doses, as early as 2.5 h after ingestion of BR.

The results of the present study indicate a dose-dependent effect of BR supplementation on exercise tolerance up to 8.4 mmol, with no further benefit (indeed a small reduction in exercise tolerance compared with 8.4 mmol) following ingestion of 16.8 mmol  $\text{NO}_3^-$ . A possible explanation for this threshold might be a NO-dependent reduction in skeletal muscle force via modulation of excitation-contraction coupling. It has been reported that the opening of the  $\text{Ca}^{2+}$  release channels of the sarcoplasmic reticulum (SR) is inhibited by NO (32, 35) and highly related to NO availability (35). In addition,  $\text{Ca}^{2+}$  transport (35), SR  $\text{Ca}^{2+}$ -ATPase activity (18), and cytochrome c-oxidase inhibition (9) may be influenced by NO and contribute to a dose-dependent modulation of excitation-contraction coupling. Therefore, whereas an increase in NO bioavailability may result in a more efficient mitochondrial function (24) and changes to *type II* fiber contractility (17) and blood flow (14), it is possible that these positive effects may be offset by impairments of mitochondrial or contractile function at higher NO levels that might promote nitrate stress. These suggestions are naturally speculative and await further investigation.

The improvements in time-to-task failure during severe-intensity exercise, following ingestion of 8.4 and 16.8 mmol  $\text{NO}_3^-$  in the present study, were evident without any significant changes in the  $\dot{V}_{\text{O}_2}$  response to exercise. Neither the amplitude of the  $\dot{V}_{\text{O}_2}$  slow component nor the end-exercise  $\dot{V}_{\text{O}_2}$  was influenced by acute ingestion of up to 16.8 mmol  $\text{NO}_3^-$ . This

finding is consistent with some (20) but not all previous reports (3, 22). For example, Bailey et al. (3) reported that 3 days of BR supplementation reduced the  $\dot{V}_{\text{O}_2}$  slow-component amplitude by 23% and improved exercise tolerance by ~16%. In contrast, Kelly et al. (20) reported that 3 days of BR supplementation improved exercise tolerance at three different severe intensities by 12–17%, without any accompanying changes in the  $\dot{V}_{\text{O}_2}$  response. We found no difference in end-exercise  $\dot{V}_{\text{O}_2}$  between BR and PL at any dose. In the severe exercise-intensity domain, the  $\dot{V}_{\text{O}_2}$  at the point of volitional exhaustion would be expected to equal the maximum  $\dot{V}_{\text{O}_2}$  ( $\dot{V}_{\text{O}_2 \text{max}}$ ) (11). Our results are therefore consistent with some (3, 37) but not all (5, 25) previous studies that indicate that  $\text{NO}_3^-$  supplementation does not reduce  $\dot{V}_{\text{O}_2 \text{max}}$ . Interestingly, there was a disconnect between the effects of BR on steady-state  $\dot{V}_{\text{O}_2}$  during moderate-intensity exercise (where the greatest reduction occurred at the highest dose of  $\text{NO}_3^-$ ) and the effects of BR on exercise tolerance (where the increased time-to-task failure was similar with 8.4 and 16.8 mmol  $\text{NO}_3^-$ ). Collectively, these results appear to indicate that the effects of BR on severe-intensity exercise performance may be independent from the effects of BR on the  $\text{O}_2$  cost of submaximal exercise.

It should be noted that while an approximate 12–14% extension of time-to-task failure during severe-intensity, constant work-rate exercise, following acute BR ingestion, may appear impressive, this is likely to translate into no more than a 1–2% reduction in the time to complete a given distance, for example, during a short endurance time-trial (TT) event (34). This is similar to the magnitude of improvement in performance reported previously for 4 km and 16.1 km TT after acute BR ingestion (21) and for 10 km TT following 6 days of BR supplementation (6). A 1% improvement in performance is highly meaningful in elite sport. For example, it could improve 1,500-m running performance by ~2 s or 3,000-m running performance by approximately 4–5 s in international standard athletes. It remains unclear, however, whether elite athletes may confer a performance benefit from  $\text{NO}_3^-$  supplementation. Several studies now indicate that at least when  $\text{NO}_3^-$  is ingested acutely, TT performance is not enhanced in highly trained endurance athletes (7, 8, 40). This may be related to factors such as greater NOS activity, better muscle oxygenation and mitochondrial efficiency, and a lower fraction of *type II* fibers in the muscles of highly endurance trained compared with moderately trained subjects (40). It is possible that the dose-response relationship between  $\text{NO}_3^-$  ingestion and changes in exercise performance are different in elite compared with sub-elite subjects, such that larger  $\text{NO}_3^-$  doses and/or longer supplementation periods may be required to elicit improved exercise performance. The significant correlation between the change in plasma  $[\text{NO}_2^-]$  and the change in time-to-task failure indicates that the dietary  $\text{NO}_3^-$  intervention must be sufficient to increase plasma  $[\text{NO}_2^-]$  if performance is to be improved. In this regard, an important consideration may be the timing of supplementation relative to the start of exercise. The present study shows that on average, plasma  $[\text{NO}_2^-]$  takes longer to peak when larger doses of  $\text{NO}_3^-$  are imbibed. However, there are appreciable interindividual differences in the speed with which ingested  $\text{NO}_3^-$  is reduced to  $\text{NO}_2^-$ , which may preclude any more specific advice other than to consume  $\text{NO}_3^-$  some 2–3 h before the start of exercise.



It has been suggested previously that there may be “responders” and nonresponders to dietary  $\text{NO}_3^-$  supplementation (40), and there was evidence of this in the present study. Interestingly, the number of nonresponders (in terms of exercise capacity) decreased as the dose ingested increased. For example, there were three nonresponders in the 4.2-mmol condition, two in the 8.4-mmol condition, and one in the 16.8-mmol condition. Two of the subjects who did not respond at the lowest dose did respond to the larger doses, and one subject who did not respond following administration of 4.2 or 8.4 mmol did respond to the 16.8-mmol dose. This suggests that some individuals will require a larger acute dose than others to elicit any positive effects on exercise capacity from dietary  $\text{NO}_3^-$  ingestion. Unlike in our previous study (40), the increase in plasma  $[\text{NO}_2^-]$  from baseline to pre-exercise for the nonresponders was not smaller than that measured in other subjects who did respond, and the nonresponders did not have particularly high baseline plasma  $[\text{NO}_2^-]$ . In a recent study, we found that the subjects who demonstrated improvement in high-intensity, intermittent exercise performance following dietary  $\text{NO}_3^-$  supplementation were those whose plasma  $[\text{NO}_2^-]$  fell significantly during exercise (41). We did not measure plasma  $[\text{NO}_2^-]$  postexercise in the present study. The explanation for the existence of responders and nonresponders to dietary  $\text{NO}_3^-$  supplementation is presently obscure.

In conclusion, dietary supplementation with  $\text{NO}_3^-$ -rich BR dose dependently increased plasma  $[\text{NO}_3^-]$  and  $[\text{NO}_2^-]$  up to 16.8 mmol  $\text{NO}_3^-$  and caused peak reductions in systolic BP and MAP dose dependently, up to 8.4 mmol  $\text{NO}_3^-$ . These results suggest that the consumption of high  $\text{NO}_3^-$  foodstuffs may be an effective strategy for maintaining and perhaps enhancing vascular health in young adults. The present study also demonstrated that the  $\text{O}_2$  cost of moderate-intensity exercise is reduced dose dependently, up to 16.8 mmol  $\text{NO}_3^-$ . Supplementation with 4.2 mmol  $\text{NO}_3^-$  did not enhance time-to-task failure relative to PL; however, supplementation with 8.4 mmol  $\text{NO}_3^-$  significantly improved time-to-task failure relative to PL, with no further improvement evident following supplementation with 16.8 mmol  $\text{NO}_3^-$ . Although the mechanistic bases for the reduction in the  $\text{O}_2$  cost of submaximal exercise and enhancements in exercise tolerance following acute dietary BR remain unclear, these results provide important, practical information that may underpin the potential use of BR/ $\text{NO}_3^-$  supplementation for improving cardiovascular health in the general population and for enhancing exercise performance in athletes.

#### ACKNOWLEDGMENTS

The authors thank Beet It for providing the beverages used in this study, gratis.

#### GRANTS

This study was funded in part by a research grant of GSSI, a division of PepsiCo, Inc. The views expressed in this manuscript are those of the authors and do not necessarily reflect the position or policy of PepsiCo, Inc.

#### DISCLOSURES

The views expressed in this article are those of the authors and do not necessarily reflect the position or policy of PepsiCo.

#### AUTHOR CONTRIBUTIONS

Author contributions: L.J.W., S.J.B., P.G.W., A.E.J., A.V., and A.M.J. conception and design of research; L.J.W., J.K., and J.R.B. performed exper-

iments; L.J.W., J.K., P.F.S., and A.V. analyzed data; L.J.W., S.J.B., P.F.S., A.V., and A.M.J. interpreted results of experiments; L.J.W. and P.F.S. prepared figures; L.J.W. and A.M.J. drafted manuscript; L.J.W., J.K., S.J.B., P.G.W., A.E.J., A.V., and A.M.J. edited and revised manuscript; L.J.W., J.K., S.J.B., J.R.B., P.F.S., P.G.W., A.E.J., A.V., and A.M.J. approved final version of manuscript.

#### REFERENCES

1. Alzawahra WF, Talukder MA, Liu X, Samouilov A, Zwiuer JL. Heme proteins mediate the conversion of nitrite to nitric oxide in the vascular wall. *Am J Physiol Heart Circ Physiol* 295: H499–H508, 2008.
2. Bailey SJ, Fulford J, Vanhatalo A, Winyard PG, Blackwell JR, DiMenna FJ, Wilkerson DP, Benjamin N, Jones AM. Dietary nitrate supplementation enhances muscle contractile efficiency during knee-extensor exercise in humans. *J Appl Physiol* 109: 135–148, 2010.
3. Bailey SJ, Winyard P, Vanhatalo A, Blackwell JR, DiMenna FJ, Wilkerson DP, Tarr J, Benjamin N, Jones AM. Dietary nitrate supplementation reduces the  $\text{O}_2$  cost of low-intensity exercise and enhances tolerance to high-intensity exercise in humans. *J Appl Physiol* 107: 1144–1155, 2009.
4. Benjamin N, O'Driscoll F, Dougall H, Duncan C, Smith L, Golden M, McKenzie H. Stomach NO synthesis. *Nature* 368: 502–503, 1994.
5. Bescos R, Rodriguez FA, Iglesias X, Ferrer MD, Iborra E, Pons A. Acute administration of inorganic nitrate reduces  $\dot{\text{V}}\text{O}_{2\text{peak}}$  in endurance athletes. *Med Sci Sports Exerc* 43: 1979–1986, 2011.
- 5a. Burke LM. To beet or not to beet. *J Appl Physiol*. doi:10.1152/jappphysiol.00612.2013.
6. Cermak NM, Gibala MJ, van Loon LJ. Nitrate supplementation's improvement of 10-km time-trial performance in trained cyclists. *Int J Sport Nutr Exerc Metab* 22: 64–71, 2012.
7. Cermak NM, Res P, Stinkens R, Lundberg JO, Gibala MJ, van Loon LJ. No improvement in endurance performance following a single dose of beetroot juice. *Int J Sport Nutr Exerc Metab* 22: 470–478, 2012.
8. Christensen PM, Nyberg M, Bangsbo J. Influence of nitrate supplementation on  $\dot{\text{V}}\text{O}_2$  kinetics and endurance of elite cyclists. *Scand J Med Sci Sports* 23: e21–e31, 2013.
9. Clerc P, Rigoulet M, Leverve X, Fontaine E. Nitric oxide increases oxidative phosphorylation efficiency. *J Bioenerg Biomembr* 39: 158–166, 2007.
10. Cosby K, Partovi KS, Crawford JH, Patel RP, Reiter CD, Martyr S, Yang BK, Wacławski MA, Zalos G, Xu X, Huang KT, Shields H, Kim-Shapiro DB, Schechter AN, Cannon III RO, Gladwin MT. Nitrite reduction to nitric oxide by deoxyhemoglobin vasodilates the human circulation. *Nat Med* 9: 1498–1505, 2003.
11. Day JR, Rossiter HB, Coats EM, Skasick A, Whipp BJ. The maximally attainable  $\dot{\text{V}}\text{O}_2$  during exercise in humans: the peak vs. maximum issue. *J Appl Physiol* 95: 1901–1907, 2003.
12. Dejam A, Hunter CJ, Schechter AN, Gladwin MT. Emerging role of nitrite in human biology. *Blood Cells Mol Dis* 32: 423–429, 2004.
13. Feelisch M, Fernandez BO, Bryan NS, Garcia-Saura MF, Bauer S, Whitlock DR, Ford PC, Janero DR, Rodriguez J, Ashrafian H. Tissue processing of nitrite in hypoxia: an intricate interplay of nitric oxide-generating and -scavenging systems. *J Biol Chem* 283: 33927–33934, 2008.
14. Ferguson SK, Hirai DM, Copp SW, Holdsworth CT, Allen JD, Jones AM, Musch TI, Poole DC. Impact of dietary nitrate supplementation via beetroot juice on exercising muscle vascular control in rats. *J Physiol* 591: 547–557, 2013.
15. Fulford J, Winyard PG, Vanhatalo A, Bailey SJ, Blackwell JR, Jones AM. Influence of dietary nitrate supplementation on human skeletal muscle metabolism and force production during maximum voluntary contractions. *Pflugers Arch* 465: 517–528, 2013.
16. Govoni M, Jansson EA, Weitzberg E, Lundberg JO. The increase in plasma nitrite after a dietary nitrate load is markedly attenuated by an antibacterial mouthwash. *Nitric Oxide* 19: 333–337, 2008.
17. Hernández A, Schiffer TA, Ivarsson N, Cheng AJ, Bruton JD, Lundberg JO, Weitzberg E, Westerblad H. Dietary nitrate increases tetanic  $[\text{Ca}^{2+}]_i$  and contractile force in mouse fast-twitch muscle. *J Physiol* 590: 3575–3583, 2012.
18. Ishii T, Sunami O, Saitoh N, Nishio H, Takeuchi T, Hata F. Inhibition of skeletal muscle sarcoplasmic reticulum  $[\text{Ca}^{2+}]$ -ATPase by nitric oxide. *FEBS Lett* 440: 218–222, 1998.
19. Kapil V, Milsom AB, Okorie M, Maleki-Toyserkani S, Akram F, Rehman F, Arghandawi S, Pearl V, Benjamin N, Loukogeorgakis S,



- Macallister R, Hobbs AJ, Webb AJ, Ahluwalia A. Inorganic nitrate supplementation lowers blood pressure in humans: role for nitrite-derived NO. *Hypertension* 56: 274–281, 2010.
20. Kelly J, Vanhatalo A, Blackwell JR, Wilkerson DP, Wylie LJ, Jones AM. Effects of nitrate on the power-duration relationship for severe-intensity exercise. *Med Sci Sports Exerc*. In press.
21. Lansley KE, Winyard PG, Bailey SJ, Vanhatalo A, Wilkerson DP, Blackwell JR, Gilchrist M, Benjamin N, Jones AM. Acute dietary nitrate supplementation improves cycling time trial performance. *Med Sci Sports Exerc* 43: 1125–1131, 2011.
22. Lansley KE, Winyard PG, Fulford J, Vanhatalo A, Bailey SJ, Blackwell JR, DiMenna FJ, Gilchrist M, Benjamin N, Jones AM. Dietary nitrate supplementation reduces the O<sub>2</sub> cost of walking and running: a placebo-controlled study. *J Appl Physiol* 110: 591–600, 2011.
23. Larsen FJ, Ekblom B, Sahlin K, Lundberg JO, Weitzberg E. Effects of dietary nitrate on blood pressure in healthy volunteers. *N Engl J Med* 355: 2792–2793, 2006.
24. Larsen FJ, Schiffer TA, Borniquel S, Sahlin K, Ekblom B, Lundberg JO, Weitzberg E. Dietary inorganic nitrate improves mitochondrial efficiency in humans. *Cell Metab* 13: 149–159, 2011.
25. Larsen FJ, Weitzberg E, Lundberg JO, Ekblom B. Dietary nitrate reduces maximal oxygen consumption while maintaining work performance in maximal exercise. *Free Radic Biol Med* 48: 342–347, 2010.
26. Larsen FJ, Weitzberg E, Lundberg JO, Ekblom B. Effects of dietary nitrate on oxygen cost during exercise. *Acta Physiol (Oxf)* 191: 59–66, 2007.
27. Law MR, Wald NJ, Morris JK, Jordan RE. Value of low dose combination treatment with blood pressure lowering drugs: analysis of 354 randomised trials. *BMJ* 326: 1427–1434, 2003.
28. Lawes CM, Rodgers A, Bennett DA, Parag V, Suh I, Ueshima H, MacMahon S, Asia Pacific Cohort Studies Collaboration. Blood pressure and cardiovascular disease in the Asia Pacific region. *J Hyperten* 21: 707–717, 2003.
29. Lewington S, Clarke R, Qizibash N, Peto R, Collins R. Prospective Studies Collaboration. Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet* 360: 1903–1913, 2002.
30. Lundberg JO, Carlström M, Larsen FJ, Weitzberg E. Roles of dietary inorganic nitrate in cardiovascular health and disease. *Cardiovasc Res* 89: 525–532, 2011.
31. MacMahon S, Peto R, Cutler J, Collins R, Sorlie P, Neaton J, Abbott R, Godwin J, Dyer A, Stamler J. Blood pressure, stroke and coronary heart disease: part 1, prolonged differences in blood pressure: prospective observational studies corrected for the regression dilution bias. *Lancet* 335: 765–774, 1990.
32. Mészáros L, Minarovic I, Zahradnikova A. Inhibition of the skeletal muscle ryanodine receptor calcium release channel by nitric oxide. *FEBS Lett* 380: 49–52, 1996.
33. Omar SA, Artime E, Webb AJ. A comparison of organic and inorganic nitrates/nitrites. *Nitric Oxide* 26: 229–240, 2012.
34. Paton CD, Hopkins WG. Variation in performance of elite cyclists from race to race. *Eur J Sports Sci* 6: 25–31, 2006.
35. Reid MB. Role of nitric oxide in skeletal muscle: synthesis, distribution and functional importance. *Acta Physiol Scand* 162: 401–409, 1998.
36. Stamler JS, Meissner G. Physiology of nitric oxide in skeletal muscle. *Physiol Rev* 81: 209–237, 2001.
37. Vanhatalo A, Bailey SJ, Blackwell JR, DiMenna FJ, Pavey TG, Wilkerson DP, Benjamin N, Winyard PG, Jones AM. Acute and chronic effects of dietary nitrate supplementation on blood pressure and the physiological responses to moderate-intensity and incremental exercise. *Am J Physiol Regul Integr Comp Physiol* 299: R1121–R1131, 2010.
38. Vanhatalo A, Fulford J, Bailey SJ, Blackwell JR, Winyard PG, Jones AM. Dietary nitrate reduces muscle metabolic perturbation and improves exercise tolerance in hypoxia. *J Physiol* 589: 5517–5528, 2011.
39. Webb AJ, Patel N, Loukogeorgakis S, Okorie M, Aboud Z, Misra S, Rashid R, Miall P, Deanfield J, Benjamin N, MacAllister R, Hobbs AJ, Ahluwalia A. Acute blood pressure lowering, vasoprotective, and anti-platelet properties of dietary nitrate via bioconversion to nitrite. *Hypertension* 51: 784–790, 2008.
40. Wilkerson DP, Hayward GM, Bailey SJ, Vanhatalo A, Blackwell JR, Jones AM. Influence of acute dietary nitrate supplementation on 50 mile time trial performance in well-trained cyclists. *Eur J Appl Physiol* 112: 4127–4134, 2012.
41. Wylie LJ, Mohr M, Krstrup P, Jackman SR, Ermudis G, Kelly J, Black MI, Bailey SJ, Vanhatalo A, Jones AM. Dietary nitrate supplementation improves team sport-specific intense intermittent exercise performance. *Eur J Appl Physiol* 113: 1673–1684, 2013.