

Protein Kinase C-Dependent Dephosphorylation of Tyrosine Hydroxylase Requires the B56 δ Heterotrimeric Form of Protein Phosphatase 2A

Jung-Hyuck Ahn^{1*}, Yong Kim², Hee-Sun Kim³, Paul Greengard², Angus C. Nairn^{2,4*}

1 Department of Biochemistry, Ewha Womans University School of Medicine, Seoul, Korea, **2** Laboratory of Molecular and Cellular Neuroscience, The Rockefeller University, New York, New York, United States of America, **3** Department of Molecular Medicine and Tissue Injury Defense Research Center, Ewha Womans University, School of Medicine, Seoul, Korea, **4** Department of Psychiatry, Yale University School of Medicine, New Haven, Connecticut, United States of America

Abstract

Tyrosine hydroxylase, which plays a critical role in regulation of dopamine synthesis, is known to be controlled by phosphorylation at several critical sites. One of these sites, Ser40, is phosphorylated by a number of protein kinases, including protein kinase A. The major protein phosphatase that dephosphorylates Ser40 is protein phosphatase-2A (PP2A). A recent study has also linked protein kinase C to the dephosphorylation of Ser40 [1], but the mechanism is unclear. PP2A isoforms are comprised of catalytic, scaffold, and regulatory subunits, the regulatory B subunits being able to influence cellular localization and substrate selection. In the current study, we find that protein kinase C is able to phosphorylate a key regulatory site in the B56 δ subunit leading to activation of PP2A. In turn, activation of the B56 δ -containing heterotrimeric form of PP2A is responsible for enhanced dephosphorylation of Ser40 of tyrosine hydroxylase in response to stimulation of PKC. In support of this mechanism, down-regulation of B56 δ expression in N27 cells using RNAi was found to increase dopamine synthesis. Together these studies reveal molecular details of how protein kinase C is linked to reduced tyrosine hydroxylase activity via control of PP2A, and also add to the complexity of protein kinase/protein phosphatase interactions.

Citation: Ahn J-H, Kim Y, Kim H-S, Greengard P, Nairn AC (2011) Protein Kinase C-Dependent Dephosphorylation of Tyrosine Hydroxylase Requires the B56 δ Heterotrimeric Form of Protein Phosphatase 2A. PLoS ONE 6(10): e26292. doi:10.1371/journal.pone.0026292

Editor: Maria Gasset, Consejo Superior de Investigaciones Cientificas, Spain

Received: May 19, 2011; **Accepted:** September 24, 2011; **Published:** October 21, 2011

Copyright: © 2011 Ahn et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by grants from the National Institutes of Health (Grant DA10044, MH74866, and MH090963 to PG and ACN) and National Research Foundation of Korea (NRF) grant funded by Korea government (MEST) (2010-0029354). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: ahnj@ewha.ac.kr (JHA); angus.nairn@yale.edu (ACN)

Introduction

Tyrosine hydroxylase (TH) is the rate-limiting enzyme involved in the synthesis of catecholamines such as dopamine [2]. The activity of TH is controlled by phosphorylation of multiple sites, including Ser8, Ser19, Ser31 and Ser40, that is catalyzed by several different protein kinases, including protein kinase A (PKA), protein kinase C (PKC), calcium/calmodulin-dependent protein kinase II (CaMKII), and the MAP kinase, ERK [3–5]. In particular, phosphorylation of Ser40 by PKA has been found to play a critical role in activation of TH, and has been the subject of extensive study in dopaminergic neurons and other types of cell. Elucidation of the mechanisms involved in control of TH is essential for understanding its role in the normal function of dopaminergic neurons as well as in neurodegenerative diseases such as Parkinson's disease, where dopamine synthesis is impaired.

The dephosphorylation of TH has also been the subject of a number of studies. The major protein phosphatase that dephosphorylates Ser40 of TH is believed to be protein phosphatase 2A (PP2A) based on *in vitro* studies as well as in studies in intact cell systems using PP2A inhibitors [6–8]. A recent study has linked the δ isoform of PKC to enhanced PP2A activity and reduced TH activity through dephosphorylation of Ser40 [1]. However, the detailed mechanism is not known. PP2A is ubiquitously expressed in eukaryotic cells where it exists as a heterotrimeric enzyme

composed of a 36 kDa catalytic C subunit, a 64 kDa scaffolding A subunit, and multiple regulatory B subunits that are thought to influence enzyme activity, substrate specificity and subcellular localization [9–14]. We have recently found that the B56 δ subunit is phosphorylated by PKA at Ser566 leading to activation of PP2A, and enhanced dephosphorylation of certain sites in DARPP-32 [15–16], a key mediator of dopamine action in striatal medium spiny neurons [17]. Other recent studies suggest that the regulation of the B56 δ -containing heterotrimeric form of PP2A by PKA is not limited to medium spiny neurons and that therefore the control of protein dephosphorylation by cAMP/PKA/B56 δ /PP2A may be a more widespread phenomenon [18–20]. In the current study, we have found that PKC phosphorylates the B56 δ subunit at Ser566. Moreover, we find that the regulation of B56 δ by PKC plays an important role in activation of PP2A, which in turn is responsible for enhanced dephosphorylation of Ser40 and inactivation of TH.

Materials and Methods

Chemicals and antibodies

Rottlerin and phorbol-12-myristate-13-acetate (PMA) were obtained from Calbiochem (La Jolla, CA) Mouse tyrosine hydroxylase, phospho-Ser40 and phospho-Ser31 antibodies, were obtained from Chemicon (Temecula, CA). Anti-FLAG antibody,

benzoase, and heparin Type I pre-packed columns were obtained from Sigma-Aldrich (St. Louis, MO). Antibody to B56 δ and to the various phosphorylation sites in B56 δ were prepared as described [15]. Deuterated 2-(3,4-dihydroxyphenyl)ethyl-1,1-d₂-amine HCl (deuterated dopamine HCl) was obtained from CDN isotope INC (Quebec, Canada).

Cell culture

Neuro-2a (N2a) cells were purchased from ATCC (Manassas, VA), and cultured in 50% Opti-MEM and 50% DMEM containing 10% fetal bovine serum, 50 U penicillin, and 50 μ g/ml streptomycin. N27 cells were grown in RPMI 1640 medium with 2 mM L-glutamine, 10% fetal bovine serum, 50 U penicillin, and 50 μ g/ml streptomycin.

Transfection

N2a cells were cultured to 60–70% confluence in 50% Opti-MEM and 50% DMEM containing 10% fetal bovine serum without penicillin-streptomycin. Expression plasmids were transfected into N2a cells in six-well plate using Fugene 6 reagent (Roche). Media were replaced with fresh media containing 50 U penicillin and 50 μ g/ml streptomycin 12 h post-transfection. Forty-eight hours after transfection, cells were treated with DMSO vehicle or the indicated reagents for the indicated times. Cells were then lysed in 200 μ l of a buffer containing 50 mM Tris pH 8.0, 150 mM NaCl, 1% Triton X-100, 0.5% sodium deoxycholate, 0.1% SDS, protease inhibitor cocktail (Roche), and phosphatase inhibitor cocktail (Calbiochem), followed by brief sonication and centrifugation at 10,000 \times g. Supernatants were used for immunoblotting as described below.

Immunoblotting

Cell lysates were separated by 4–20% SDS-PAGE with pre-made Tris-glycine gels (Invitrogen) and immunoblotted onto polyvinylidene difluoride (PVDF) membranes (Immobilon-P; Millipore, Bedford, Massachusetts). The membranes were blocked for 2 h at room temperature in Superblock reagent (Pierce) and phosphatase inhibitor cocktail (1 ml in 100 ml of blocking reagent). Membranes were probed with the indicated antibodies. Antibody binding was detected using horseradish peroxidase-linked IgG (1:10000; Pierce) and the ECL immunoblotting detection system (GE Healthcare Life Science). For each experiment, values for levels of phospho-proteins were corrected for loading of total protein and then data was calculated relative to the value for the control in each experiment. Normalized data from multiple experiments were expressed as means as described in the Figure legends.

Immunoprecipitation and in vitro PKC phosphorylation assays

FLAG-B56 δ wt or FLAG-B56 δ S566A mutant were expressed in N2a cells and active heterotrimeric complexes were immunoprecipitated from cell lysates using anti-FLAG antibody as described [15,21]. PKC was purified from rat brain as described [22]. Immunoprecipitated PP2A was mixed with purified PKC (100 ng) without (control) or with PKC activators (0.1 mg/ml phosphatidylserine, 0.02 mg/ml diacylglycerol, 0.2 mM CaCl₂) and Mg ATP for 10 min in kinase reaction buffer (20 mM Tris-HCl, pH 7.4, 10 mM MgCl₂, 150 mM NaCl).

Viral production and purification of adeno-associated virus-RNAi (AAV-RNAi)

B56 δ oligonucleotides for RNAi production, GATCCCCC-CAGTGTGTGCTCCTCTTT CTTCCTGTCA AAAGAGGA-

CACAGCACTGGTTTTTTTGAAT and CTAGATTCCAAA-AAACCAGTGTGTGCTCCTCTTTTGACAGGAAG AAAG-AGGACACAGCACTGG GGG, were hybridized in buffer [100 mM Tris-HCl (pH 8.0), 10 mM MgCl₂, 150 mM NaCl], and then ligated into pAAV-H1 precut with Bgl II and Xba I. HEK293 cells (ATCC, Manassas, VA) were grown in ten 150 \times 25-mm dishes and co-transfected with pAAV-H1/B56 δ RNAi or pAAV-H1 (control AAV-RNAi) and helper plasmid using calcium phosphate. Cells were trypsinized, pelleted and resuspended in buffer [0.15 M NaCl and 50 mM Tris-HCl (pH 8.0)] 72-hour post-transfection. Virus was purified using a published protocol [23] with some modifications. Briefly, after three freeze-thaw cycles (liquid nitrogen to 37°C) to lyse the cells, benzoase was added (50 U/ml, final) and the cell lysate was incubated at 37°C for 30 min. The lysate was added to a centrifuge tube containing a 15%, 25%, 40% and 60% iodixanol step gradient. Samples were centrifuged at 350,000 g for 60 min at 18°C, the 40% fraction was collected, then added to an affinity column containing heparin type 1 (Sigma), washed with 0.1 M NaCl and eluted with 1 M NaCl. Elution buffer was exchanged with 25 volumes of PBS using an Amicon Ultra 100K NMWL centrifugal filter. The virus was titered using an AAV ELISA kit (Progen) and kept at –80°C until time of use. AAV (1 \times 10¹² pfu) was added to each well of a 6-well plate (cells were typically 70% confluent) for 48 hours. AAV without any RNAi oligonucleotide was used as a control.

Dopamine measurement

N27 cells expressing either control RNAi or B56 δ RNAi were lysed and sonicated in 50 mM ammonium bicarbonate buffer (pH 8.0) after 72 h. Dopamine was measured using LC-MS/MS. Dopamine-1,1,2,2-d₄ hydrochloride (CDN isotopes, Quebec, Canada) was used as a standard. The dopamine isotope was spiked into the samples and analyzed by nanoHPLC-MS with a QSTAR Q_qTOF mass spectrometer (Applied Biosystems).

Results

The PKC activator PMA increases phosphorylation of the B56 δ subunit of PP2A at Ser566

To initially address the ability of PKC to regulate the phosphorylation of the B56 δ subunit, we expressed FLAG-B56 δ in N2a cells, then treated cells with vehicle or the phorbol ester PMA under a variety of conditions (Figure 1). The phosphorylation level at four sites in B56 δ (Ser53, Ser68, Ser81, Ser566) was analyzed with their respective phospho-specific antibody. PMA treatment increased the phosphorylation level of Ser566 ~1.5-fold compared to vehicle-treated cells (Figure 1a). Exposure of N2a cells to different concentrations of PMA resulted in a dose responsive increase in Ser566 phosphorylation up to 10 nM (Figure 1b). In response to treatment with PMA, phosphorylation of B56 δ was rapid and maximal after 30 min (Figure 1c). In the presence of 5 nM PMA, the phosphorylation at Ser566 increased ~3.5-fold after 30 min compared to the vehicle-treated control.

PKC δ is involved in PMA-dependent phosphorylation of B56 δ

We next investigated the type of PKC isozyme involved in B56 δ phosphorylation. FLAG-B56 δ was expressed in N2a cells, and the effect of the PKC δ inhibitor rottlerin was examined (Figure 2). Cells were pre-incubated in the absence or presence of rottlerin for 30 min, treated with DMSO vehicle or PMA. PMA-dependent phosphorylation at Ser566 was inhibited by rottlerin. Rottlerin alone did not have any significant effect on Ser566 phosphorylation. Go6976, a PKC α specific inhibitor, did not affect Ser566




Figure 1. The PKC activator PMA increases phosphorylation of B56 δ at Ser566. (A) N2a cells expressing FLAG-B56 δ subunit were treated with DMSO (vehicle) or PMA (5 nM) for 5 min. Cells were lysed and proteins were analyzed by SDS-PAGE and immunoblotting with phospho-specific antibodies to the indicated sites in B56 δ . Total B56 δ was analyzed with anti-FLAG antibody. Upper panels show immunoblots. The bar graph shows quantification of immunoblot data normalized in each experiment to vehicle for each site as means \pm s.e.m. (n=3). *, P<0.001 compared with vehicle-treated control by student's t-test. (B) N2a cells expressing FLAG-B56 δ were treated with different concentrations of PMA (0, 1, 10, 100, 1000 nM) for 5 min. Phosphorylation of Ser566 and total B56 δ was assayed as in (A). The bar graph shows quantification of data normalized in each experiment to the zero PMA condition as means \pm s.e.m. (n=3). *, P<0.05, **, P<0.01 compared with vehicle-treated control by student's t-test. (C) N2a cells expressing FLAG-B56 δ were treated with 5 nM PMA for the indicated times. Phosphorylation of Ser566 and total B56 δ was assayed as in (A). The bar graph shows quantification of data normalized in each experiment to the zero time condition as means \pm s.e.m. (n=3). *, P<0.01, **, P<0.001 compared with vehicle treated control by student's t-test. doi:10.1371/journal.pone.0026292.g001

phosphorylation in a statistically significant manner (data not shown) suggesting that PKC δ is likely a major PKC isozyme involved in B56 δ phosphorylation in N2a cells.

PKC activates the B56 δ heterotrimeric form of PP2A *in vitro* via phosphorylation at Ser566

Our previous studies have shown that exogenous expression of the B56 δ subunit results in formation of a heterotrimeric complex that can be isolated from cell lysates (Ahn *et al.* 2007a). Wild-type B56 δ (FLAG-B56 δ wt) or a mutant in which Ser566 was replaced by alanine (FLAG-B56 δ S566A) were expressed in N2a cells, and

PP2A heterotrimers were immunoprecipitated with anti-FLAG antibody. The immuno-purified PP2A preparations were then incubated *in vitro* with PKC and MgATP in the absence or presence of a PKC activator mixture which contained phosphatidylserine, diacylglycerol, and Ca²⁺ (Figure 3). Using immunoblotting, for the PP2A preparation that contained wild-type B56 α subunit, a low level of phosphorylation at Ser566 was observed in the presence of PKC, and this was increased with addition of the PKC activator mixture (Figure 3A). However, as expected, only background phospho-Ser566 signal was detected in the PP2A preparation that contained B56 δ with the S566A mutation.




Figure 2. Rottlerin inhibits PMA-dependent phosphorylation of Ser566 in B56δ. N2a cells expressing FLAG-B56δ were pretreated without or with rottlerin (5 μM) for 30 min as indicated and then treated with DMSO (Control) or PMA (10 nM) for an additional 10 min. Phosphorylation of Ser566 and total B56δ was assayed as above. The bar graph shows quantification of immunoblot data (not shown) normalized in each experiment to control as means ± s.e.m. (n=3). *, P<0.001, **, P<0.01 compared with vehicle-treated vector control by one-way ANOVA with Newman-Keuls multiple comparison test. doi:10.1371/journal.pone.0026292.g002

In parallel assays, PP2A activity was measured. PKC incubation resulted in activation of the PP2A preparation containing wild-type B56δ as measured using a synthetic phospho-peptide as substrate. However, there was no effect of PKC on PP2A activity associated with the B56δ S566A mutant. We also carried out assays where ATP was replaced by [³²P] ATP. Phosphorylation of the A, B and C subunits was detected in the absence of the PKC activator mixture, while addition of the activator mixture led to increased phosphorylation of the B56δ subunit, with a smaller increase also being observed for the C subunit, and no increase being observed for the A subunit (see supplementary Figure S1). Mutation of Ser566 prevented the increase in phosphorylation of the B56δ mutant protein, seen in the presence of the PKC activator mixture. However, mutation of Ser566 had no effect on phosphorylation of either the A or C subunits. Taken together with the phosphatase activity assays, these results indicate that phosphorylation of Ser566 in the B56δ subunit by PKC is solely responsible for the activation of PP2A activity observed in the in vitro assays.

PP2A containing the B56δ subunit is activated by PMA, and selectively dephosphorylates Ser40 of tyrosine hydroxylase

Among several sites such as Ser8, Ser19, Ser31, and Ser40, phosphorylation of Ser40 by PKA has been found previously to be associated with increased activity of TH [5]. Interestingly, a previous study found that PKCδ was associated with activation of PP2A and dephosphorylation of Ser40 [1], but the precise mechanism was not elucidated. To address the role of PKC-mediated activation of B56δ in PP2A-mediated dephosphorylation of TH, we analyzed the phosphorylation status of Ser40 of TH in rat mesencephalic neuronal N27 cells that express significant endogenous levels of the enzyme and which produce dopamine following dibutyryl cAMP-induced differentiation [24]. Notably, N27 cells expressed higher levels of B56δ than N2a cells, and the effect of PMA on B56δ S566 phosphorylation was robust in N27 cells (data not shown).




Figure 3. PKC activates PP2A/B56δ in vitro. FLAG-B56δ wt or FLAG-B56δ S566A mutant were expressed in N2a cells and active heterotrimeric complexes were immunoprecipitated from cell lysates using anti-FLAG antibody. Immunoprecipitated PP2A was mixed with purified PKC without (control) or with PKC activators (0.1 mg/ml phosphatidylserine, 0.02 mg/ml diacylglycerol, 0.2 mM CaCl₂) and Mg ATP for 10 min. (A). Phosphorylation at Ser566 of B56δ was assessed by immunoblotting using anti-phospho-Ser566 antibody. As a loading control, total B56δ subunit and co-immunoprecipitated PP2A C subunit were assessed by immunoblotting. (B). Immobilized PP2A was suspended in phosphatase assay buffer then PP2A activity was measured using a phospho-peptide substrate (Promega, 5 min assay). Phosphatase activity is shown as means ± s.e.m. (n=3). *, P<0.001, compared with no PKC activator control by student's t-test. doi:10.1371/journal.pone.0026292.g003

N27 cells expressing either vector control, FLAG-B56δ wt, or FLAG-B56δ S566A, were treated with PMA, then the phosphorylation level of Ser40 of tyrosine hydroxylase was analyzed by immunoblotting (Figure 4). In control cells (vector alone), upon PMA stimulation the phosphorylation on Ser40 was decreased. Expression of FLAG-B56δ wt resulted in a slightly larger effect of PMA but this was not statistically significant. In contrast, in cells expressing FLAG-B56δ S566A, there was an increased basal level of phosphorylation of Ser40 which was unaffected by PMA. Phosphorylation of Ser31, which was used as a control, was unaffected by any condition examined.

These results suggest that the FLAG-B56δ S566A mutant may be acting in a dominant-negative manner to antagonize the endogenous action of B56δ. We further examined the PKC-dependent regulation of PP2A/B56δ using RNAi to down-regulate endogenous B56δ expression in N27 cells (Figure 5). B56δ-specific RNAi, expressed via adeno-associated virus (AAV) in N27 cells, resulted in reduced expression of endogenous B56δ protein by the B56δ-specific RNAi (less than 20% compared to control RNAi). Control RNAi expression had no effect compared to non-infected cells (not shown). Control or B56δ knock-down




Figure 4. Phosphorylation of B56 δ at Ser566 mediates PMA-dependent dephosphorylation of Ser40 of tyrosine hydroxylase. N27 cells expressing either vector, FLAG-B56 δ wt or FLAG-B56 δ S566A mutant were treated with either DMSO (-) or PMA (10 nM) for 30 min. Cells were lysed and proteins were analyzed by SDS-PAGE and immunoblotting with phospho-specific antibodies to Ser40 and Ser31 in tyrosine hydroxylase (TH), and a total tyrosine hydroxylase antibody (upper panels). The bar graph shows quantification of the immunoblot data normalized in each experiment to control as means \pm s.e.m. (n=3). *, $P<0.001$, ** $P<0.01$, *** $P<0.05$ compared with vehicle-treated vector control by one-way ANOVA with Newman-Keuls multiple comparison test.

doi:10.1371/journal.pone.0026292.g004

N27 cells were treated with DMSO vehicle, PMA or PMA together with rottlerin. PMA treatment resulted in dephosphorylation of phospho-Ser40 in control RNAi-infected cells, and this was inhibited by rottlerin. However, there was no effect of PMA treatment in cells in which B56 δ expression was knocked down. Notably, the basal level of phospho-Ser40 was increased following knockdown of B56 δ . Phosphorylation of Ser31 was unaffected under any of the conditions used.

PP2A containing the B56 δ subunit regulates dopamine synthesis in N27 cells

Previous studies have indicated that PKC δ -deficient N27 cells showed an increase in dopamine synthesis through enhanced TH activity [1]. We further examined the physiological role of TH regulation by PKC-mediated control of the B56 δ subunit by measuring dopamine synthesis in N27 cells (Figure 6). Expression of B56 δ was knocked down in N27 cells using AAV-mediated expression of B56 δ RNAi. After 72 h, cells were harvested, and dopamine production was analyzed using LC/MS. N27 cells with reduced B56 δ expression showed significantly increased dopamine levels compared to cells expressing a control RNAi plasmid. Dopamine levels were determined to be 1.59 ± 0.16 $\mu\text{g}/\text{mg}$ of




Figure 5. PMA-dependent dephosphorylation of Ser40 in tyrosine hydroxylase is inhibited by down-regulation of B56 δ . N27 cells were infected with either AAV-control RNAi or AAV-B56 δ RNAi for 48 h. N27 cells were then pretreated with either vehicle or rottlerin (5 μM) for 30 min, then treated with PMA (10 nM) for 10 min as indicated. Phosphorylation of Ser40 and Ser31 of tyrosine hydroxylase was measured as above. B56 δ levels were measured by immunoblotting. The bar graph shows quantification of immunoblot data normalized in each experiment to control as means \pm s.e.m. (n=3). *, $P<0.001$ compared with vehicle-treated control by one-way ANOVA with Newman-Keuls multiple comparison test.

doi:10.1371/journal.pone.0026292.g005

protein in control RNAi cells compared to 3.69 ± 0.21 $\mu\text{g}/\text{mg}$ protein in B56 δ RNAi expressing cells.

Discussion

The results obtained in this study indicate that Ser566 in the B56 δ subunit of PP2A is phosphorylated in response to activation of PKC in intact cells. Previously we have found that




Figure 6. Dopamine synthesis in N27 cells is increased following down-regulation of B56 δ . N27 cells were infected with AAV-control RNAi or AAV-B56 δ RNAi for 72 h. Dopamine was measured in cell lysates by LC/MS with tritium labeled DA as a standard. Data are shown as means \pm s.e.m. (n=3). *, $P<0.01$ compared with control RNAi infected cells by student's t-test.

doi:10.1371/journal.pone.0026292.g006

phosphorylation of this residue by PKA results in activation of phosphatase activity of the heterotrimeric PP2A complex containing B56 δ (Ahn *et al.* 2007a). Consistent with our previous results, phosphorylation of Ser566 of B56 δ was accompanied by activation of PP2A. The results obtained from studies with over-expression of wild-type and mutant B56 δ , as well as of knockdown of B56 δ expression, provide strong support for the conclusion that phosphorylation of B56 δ s causally involved in mediating the effect of PKC activation on dephosphorylation of Ser40 in TH. Our results are consistent with a role for the PKC δ isoform in regulation of the B56 δ /PP2A heterotrimer based on the use of the relatively specific inhibitor, rottlerin. Various studies have found that several different isoforms of PKC are able to interact with PP2A, although nothing appears to be known about the identity of the B subunit in these complexes [25–28]. We therefore cannot rule out that PKC isoforms other than PKC δ might be able to regulate PP2A activity via phosphorylation of B56 δ .

In previous studies, Zhang *et al* have shown that PKC could enhance PP2A activity resulting in dephosphorylation of Ser40 of TH [1]. Based on *in vitro* experiments, their study suggested that PKC could phosphorylate the C subunit of PP2A, apparently leading to increased phosphatase activity. However, the *in vitro* evidence for this was not strong, and the mechanism involved in intact cells was not studied. Our results indicate that B56 δ phosphorylation at Ser566 is responsible for all of the effects we observed of PKC activation on TH Ser40 dephosphorylation.

A number of studies have implicated PKC in the regulation of TH in several different cellular systems [5]. However, the precise actions of PKC have been difficult to assess. *In vitro*, PKC is able to phosphorylate TH at Ser40 and stimulate its activity [29]. However, another *in vitro* study found that while PKC could phosphorylate TH at Ser40, there was no effect on activity [30]. The exact reason for the differences in these studies is not clear but may be related to the stoichiometry of phosphorylation that was achieved in the two studies. Several studies using intact cell preparations have indicated that activation of PKC leads to phosphorylation and activation of TH [31–33]. Some of these studies indicate that Ser40 is regulated by PKC while others suggest the likelihood that Ser31 may be the site that is targeted. Our results as well as those of Zhang *et al* [1] show a consistent ability of PKC activation to result in dephosphorylation of TH at Ser40, with no obvious effect at Ser31, in a number of different types of cell types in culture as well as *in vivo*. Presumably, the differences observed in the different studies reflect the use of various types of cells and the methods used to stimulate PKC. Based on our results, it is also possible that the cell types used in these previous studies expressed variable levels of B56 δ , which has been found to have restricted cell and tissue expression [34]. In the absence of B56 δ , PP2A would not be activated and therefore PKC activation might lead alternatively to phosphorylation and activation of TH.

Our recent studies have found that PP2A is subject to novel forms of regulation by second messenger-mediated signaling mechanisms. We have found that cAMP/PKA can be coupled to protein dephosphorylation via the ability of PKA to phosphorylate B56 δ and activate PP2A [15]. In other studies, we have found that Ca²⁺ can activate the PR72-containing heterotrimer of PP2A [21]. The present study extends these results and shows that pathways traditionally thought to increase phosphorylation can also be coupled to pathways that result in decreased phosphorylation of selected substrates. The B subunits of PP2A likely influence the specificity of PP2A towards certain substrates through direct protein:protein interactions [10,35]. The crystal structures of a PP2A heterotrimer containing the core

domain of the B56 γ isoform has provided important insight into holoenzyme assembly, and the role of the B56 subunit in substrate recognition [35–38]. The central core of the B56 subunit contains a number of HEAT-like repeats that make multiple contacts with the C subunit, suggesting mechanisms whereby the specificity of substrate interactions with the active site of PP2A can be influenced. A previous study found that the B56 β isoform of PP2A was able to confer specificity for PP2A towards Ser40 of TH [8]. A recent follow-up study found that a specific Glu residue (Glu153) in B56 β is an important molecular determinant for PP2A specificity towards TH [39]. Notably, a residue equivalent to Glu153 is found in other B56 isoforms including B56 δ . The precise mechanism whereby phosphorylation of Ser566 of B56 δ is able to increase PP2A activity is not known. However, phosphorylation may be able to increase PP2A affinity for TH or other selected substrates through additional protein:protein interactions, perhaps by influencing the interaction of Glu153 with specific substrates like TH.

Phosphorylation of Ser40 of TH by PKA is known to play a critical role in the regulation of enzyme activity [5]. Based on our previous studies of regulation of B56 δ , one might speculate that activation of PKA could also lead to dephosphorylation of TH at Ser40. While we did not investigate this in detail, our preliminary studies suggest that stimulation of PKA results in increased Ser40 phosphorylation in the cell lines used in this study. We interpret these results and studies of TH in other cell systems as indicating that the balance between PKA-mediated phosphorylation and PKA/B56 δ /PP2A-mediated dephosphorylation favors the phosphorylation of Ser40 by PKA. A similar situation is found for phosphorylation of Thr34 of DARPP-32 by PKA [15].

Our initial studies of B56 δ and regulation of PP2A by cAMP/PKA-dependent signaling focused on DARPP-32, a protein that plays an important role in dopaminergic signaling in striatal neurons [16,17]. Other more recent studies have suggested that this mechanism is more widespread and we have implicated its role in regulation of PP2A in a number of different cellular signaling processes [18–20]. By comparison, it seems likely that the ability of PKC to regulate PP2A via phosphorylation of B56 δ may also play a role in the dephosphorylation of substrates other than TH in neuronal and non-neuronal cell types. In one study of B56 δ , regulation of PP2A activity has been found to be involved in a complex signaling process in cardiac muscle involving protein:protein interactions between PP2A, PKA and the muscle A kinase anchoring protein, and the phosphodiesterase PDE4D3 [19]. This results in both feed-forward and feed-back mechanisms that are presumably designed to fine-tune cAMP signaling in muscle. A number of studies have found that PP2A interacts with various isoforms of PKC, and that in some cases PP2A is able to dephosphorylate and inactivate certain PKC isoforms including PKC δ [40]. This suggests by analogy with the example of B56 δ -dependent regulation of PDE4D3 in cardiac muscle, that PKC is able to phosphorylate and activate PP2A, but that PP2A can inactivate PKC through a feed-back mechanism leading to transient control of this signaling system.

Supporting Information

Figure S1 PKC-mediated phosphorylation of PP2A with B56 δ subunit. N2a cells were transfected either with FLAG-B56 δ S566A mutant or FLAG-B56 δ wt then the lysates were immunoprecipitated using anti-FLAG antibody. Immunoprecipitated PP2A holoenzymes were eluted by adding FLAG peptide (100 μ g/ml, Sigma-Aldrich, St. Louis, MO) for 10 min. Eluted PP2A with B56 δ subunit complex was incubated with purified

PKC and [γ - 32 P] ATP without or with PKC activator for 5 min. Immune-complexes were separated with 4–20% SDS-PAGE, and phosphorylation of PP2A subunits measured using autoradiography. After autoradiography, bands corresponding to each subunit (A, B, and C) were excised and 32 P incorporation measured using a scintillation counter (cpm). The bar graph represents the cpm as means \pm s.e.m. ($n = 3$). *, $P < 0.001$, ** $P < 0.01$, *** $P < 0.05$ compared with PKC-only treated control by student's t-test. (EPS)

References

- Zhang D, Kanthasamy A, Yang Y, Anantharam V, Kanthasamy A (2007) Protein kinase C delta negatively regulates tyrosine hydroxylase activity and dopamine synthesis by enhancing protein phosphatase-2A activity in dopaminergic neurons. *J Neurosci* 27: 5349–5362.
- Kumer SC, Vrana KE (1996) Intricate regulation of tyrosine hydroxylase activity and gene expression. *J Neurochem* 67: 443–462.
- Zigmond RE, Schwarzschild MA, Rittenhouse AR (1989) Acute regulation of tyrosine hydroxylase by nerve activity and by neurotransmitters via phosphorylation. *Annu Rev Neurosci* 12: 415–461.
- Haycock JW (1990) Phosphorylation of tyrosine hydroxylase in situ at serine 8, 19, 31, and 40. *J Biol Chem* 265: 11682–11691.
- Dunkley PR, Bobrovskaya L, Graham ME, von Nagy-Felsobuki EI, et al. (2004) Tyrosine hydroxylase phosphorylation: regulation and consequences. *J Neurochem* 91: 1025–1043.
- Haavik J, Schelling DL, Campbell DG, Andersson KK, Flatmark T, et al. (1989) Identification of protein phosphatase 2A as the major tyrosine hydroxylase phosphatase in adrenal medulla and corpus striatum: evidence from the effects of okadaic acid. *FEBS Lett* 251: 36–42.
- Leal RB, Sim AT, Goncalves CA, Dunkley PR (2002) Tyrosine hydroxylase dephosphorylation by protein phosphatase 2A in bovine adrenal chromaffin cells. *Neurochem Res* 27: 207–213.
- Saraf A, Virshup DM, Strack S (2007) Differential expression of the B β regulatory subunit of protein phosphatase 2A modulates tyrosine hydroxylase phosphorylation and catecholamine synthesis. *J Biol Chem* 282: 573–580.
- Virshup DM (2000) Protein phosphatase 2A: a panoply of enzymes. *Curr Opin Cell Biol* 12: 180–185.
- Janssens V, Goris J (2001) Protein phosphatase 2A: a highly regulated family of serine/threonine phosphatases implicated in cell growth and signalling. *Biochem J* 353: 417–439.
- Price NE, Mumby MC (1999) Brain protein serine/threonine phosphatases. *Curr Opin Neurobiol* 9: 336–342.
- Price NE, Mumby MC (2000) Effects of regulatory subunits on the kinetics of protein phosphatase 2A. *Biochemistry* 39: 11312–11318.
- Van Kanegan MJ, Adams DG, Wadzinski BE, Strack S (2005) Distinct protein phosphatase 2A heterotrimers modulate growth factor signaling to extracellular signal-regulated kinases and Akt. *J Biol Chem* 280: 36029–36036.
- Letourmeux C, Rocher G, Porteu F (2006) B56-containing PP2A dephosphorylate ERK and their activity is controlled by the early gene IEX-1 and ERK. *EMBO J* 25: 727–738.
- Ahn JH, McAvoy T, Rakhilin SV, Nishi A, Greengard P, et al. (2007) Protein kinase A activates protein phosphatase 2A by phosphorylation of the B56{delta} subunit. *Proc Natl Acad Sci U S A* 104: 2979–2984.
- Stipanovich A, Valjent E, Matamales M, Nishi A, Ahn JH, et al. (2008) A phosphatase cascade by which rewarding stimuli control nucleosomal response. *Nature* 453: 879–884.
- Greengard P, Allen PB, Nairn AC (1999) Beyond the dopamine receptor: the DARPP-32/protein phosphatase-1 cascade. *Neuron* 23: 435–447.
- Flynn MP, Maizels ET, Karlsson AB, McAvoy T, Ahn JH, et al. (2008) Luteinizing hormone receptor activation in ovarian granulosa cells promotes protein kinase A-dependent dephosphorylation of microtubule-associated protein 2D. *Mol Endocrinol* 22: 1695–1710.
- Dodge-Kafka KL, Bauman A, Mayer N, Henson E, Heredia L, et al. (2010) cAMP-stimulated protein phosphatase 2A activity associated with muscle A kinase-anchoring protein (mAKAP) signaling complexes inhibits the phosphorylation and activity of the cAMP-specific phosphodiesterase PDE4D3. *J Biol Chem* 285: 11078–11086.
- Ceglia I, Kim Y, Nairn AC, Greengard P (2010) Signaling pathways controlling the phosphorylation state of WAVE1, a regulator of actin polymerization. *J Neurochem* 114: 182–190.

Acknowledgments

We thank Dr. Michael Kaplitt for the pAAV-H1 vector used for RNAi expression, and Dr. Curt Freed for the N27 cell line.

Author Contributions

Conceived and designed the experiments: JA PG ACN. Performed the experiments: JA YK HK. Analyzed the data: JA ACN. Contributed reagents/materials/analysis tools: ACN PG. Wrote the paper: JA ACN.

- Ahn JH, Sung JY, McAvoy T, Nishi A, Janssens V, et al. (2007) The B β /PR72 subunit mediates Ca $^{2+}$ -dependent dephosphorylation of DARPP-32 by protein phosphatase 2A. *Proc Natl Acad Sci U S A* 104: 9876–9881.
- Woodgett JR, Hunter T (1987) Isolation and characterization of two distinct forms of protein kinase C. *J Biol Chem* 262: 4836–4843.
- Zolotukhin S, Byrne BJ, Mason E, Zolotukhin I, Potter M, et al. (1999) Recombinant adeno-associated virus purification using novel methods improves infectious titer and yield. *Gene Ther* 6: 973–985.
- Zhou W, Hurlbert MS, Schaack J, Prasad KN, Freed CR (2000) Overexpression of human alpha-synuclein causes dopamine neuron death in rat primary culture and immortalized mesencephalon-derived cells. *Brain Res* 866: 33–43.
- Boudreau RT, Garduno R, Lin TJ (2002) Protein phosphatase 2A and protein kinase C α are physically associated and are involved in Pseudomonas aeruginosa-induced interleukin 6 production by mast cells. *J Biol Chem* 277: 5322–5329.
- Nunbhakdi-Craig V, Schuechner S, Sontag JM, Montgomery L, Pallas DC, et al. (2007) Expression of protein phosphatase 2A mutants and silencing of the regulatory B alpha subunit induce a selective loss of acetylated and detyrosinated microtubules. *J Neurochem* 101: 959–971.
- Huang W, Batra S, Atkins BA, Mishra V, Mehta KD (2005) Increases in intracellular calcium dephosphorylate histone H3 at serine 10 in human hepatoma cells: potential role of protein phosphatase 2A-protein kinase CbetaII complex. *J Cell Physiol* 205: 37–46.
- Lee IH, Lim HJ, Yoon S, Seong JK, Bae DS, et al. (2008) Ahnak protein activates protein kinase C (PKC) through dissociation of the PKC-protein phosphatase 2A complex. *J Biol Chem* 283: 6312–6320.
- Albert KA, Helmer-Matyjek E, Nairn AC, Muller TH, Haycock JW, et al. (1984) Calcium/phospholipid-dependent protein kinase (protein kinase C) phosphorylates and activates tyrosine hydroxylase. *Proc Natl Acad Sci U S A* 81: 7713–7717.
- Funakoshi H, Okuno S, Fujisawa H (1991) Different effects on activity caused by phosphorylation of tyrosine hydroxylase at serine 40 by three multifunctional protein kinases. *J Biol Chem* 266: 15614–15620.
- Tachikawa E, Tank AW, Weiner DH, Mosimann WF, Yanagihara N, et al. (1987) Tyrosine hydroxylase is activated and phosphorylated on different sites in rat pheochromocytoma PC12 cells treated with phorbol ester and forskolin. *J Neurochem* 48: 1366–1376.
- Cahill AL, Horwitz J, Perlman RL (1989) Phosphorylation of tyrosine hydroxylase in protein kinase C-deficient PC12 cells. *Neuroscience* 30: 811–818.
- Bobrovskaya L, Cheah TB, Bunn SJ, Dunkley PR (1998) Tyrosine hydroxylase in bovine adrenal chromaffin cells: angiotensin II-stimulated activity and phosphorylation of Ser19, Ser31, and Ser40. *J Neurochem* 70: 2565–2573.
- Nagase T, Murakami T, Nozaki H, Inoue R, Nishito Y, et al. (1997) Tissue and subcellular distributions, and characterization of rat brain protein phosphatase 2A containing a 72-kDa delta/B β subunit. *J Biochem (Tokyo)* 122: 178–187.
- Mumby M (2007) The 3D structure of protein phosphatase 2A: new insights into a ubiquitous regulator of cell signaling. *ACS Chem Biol* 2: 99–103.
- Xu Y, Xing Y, Chen Y, Chao Y, Lin Z, et al. (2006) Structure of the protein phosphatase 2A holoenzyme. *Cell* 127: 1239–1251.
- Cho US, Xu W (2006) Crystal structure of a protein phosphatase 2A heterotrimeric holoenzyme. *Nature* 445: 53–57.
- Shi Y (2009) Serine/threonine phosphatases: mechanism through structure. *Cell* 139: 468–484.
- Saraf A, Oberg EA, Strack S (2010) Molecular determinants for PP2A substrate specificity: charged residues mediate dephosphorylation of tyrosine hydroxylase by the PP2A/B β regulatory subunit. *Biochemistry* 49: 986–995.
- Srivastava J, Goris J, Dilworth SM, Parker PJ (2002) Dephosphorylation of PKCdelta by protein phosphatase 2Ac and its inhibition by nucleotides. *FEBS Lett* 516: 265–269.