

The present and future of veterinary vaccines for Japanese encephalitis in Korea

Clin Exp Vaccine Res 2015;4:130-136
<http://dx.doi.org/10.7774/cevr.2015.4.2.130>
 pISSN 2287-3651 • eISSN 2287-366X

**Jin-Ju Nah, Dong-Kun Yang,
Ha-Hyun Kim, Jae-Young Song**
 Animal and Plant Quarantine Agency, MAFRA,
 Anyang, Korea

Received: March 26, 2015
 Revised: May 3, 2015
 Accepted: May 20, 2015

Corresponding author: Jin-Ju Nah, VMD, MSc
 Animal and Plant Quarantine Agency, 175 Anyang-
 ro, Anyang 430-757, Korea
 Tel: +82-31-467-1791, Fax: +82-31-467-1797
 E-mail: nahjj75@korea.kr

No potential conflict of interest relevant to this article was reported.

Japanese encephalitis (JE) is a mosquito-borne zoonotic disease that affects approximately 50,000 people annually in Asia, causing 10,000 deaths. Considering the role of pigs as the virus-amplifying host and the economic loss in the swine industry, JE is an important disease for both public and animal health. A nationwide JE virus (JEV) vaccination program has been conducted annually for more than 30 years to prevent severe reproductive disorders in the Korean sow population. Remarkable progress in molecular biology has made it possible to analyze the genome of the vaccine strain at the nucleotide and amino acid levels. However, the scientific record of the current JEV veterinary vaccine has not been reported. Therefore, this article outlines the current JEV vaccine strain used in animals and discusses future directions for developing new veterinary JEV vaccines.

Keywords: Japanese encephalitis, Japanese encephalitis viruses, Vaccines, Attenuated, Animals

Introduction

Recorded outbreaks of Japanese encephalitis (JE) in Asia date back to the late 19th century. JE is a major cause of viral encephalitis and is endemic in several regions of Asia and the Pacific, causing an estimated 50,000 infections and 10,000 deaths annually [1,2]. JE virus (JEV) is maintained in a natural cycle between *Culex* mosquitoes and water birds, and is amplified in pigs. Accidental infection occurs in dead-end hosts, including humans and horses. The most important mosquito vector in Korea is *Culex tritaeniorhynchus* [3,4]. In regions where JE is endemic, vaccination against JEV is the most effective measure to control infection in humans and domestic pigs [5].

JEV can infect humans and a several other animals. However, most animals show no overt signs of infection, except for humans and horses. In horses, most infections are inapparent, and the mortality rate of viral encephalitis is less than 5%. Pigs are affected only during pregnancy; JEV infection manifests as abortion, mummified fetuses, and stillborn or weak piglets, depending on the stage of pregnancy [6]. JEV infection is an important economic disease in the domestic swine industry.

Japanese Encephalitis Virus

JEV, a member of the genus *Flavivirus* in the family *Flaviviridae*, clusters in the JE sero-




Fig. 1. Genome structure of Japanese encephalitis virus. C, capsid; prM, pre-membrane and membrane; E, envelope; NS, non-structural.




Fig. 2. Annual Japanese encephalitis (JE) cases in humans and swine in Korea. Human cases from KCDC; Swine cases from QIA, KAHIS. KCDC, Korea Centers for Disease Control and Prevention; QIA, Animal and Plant Quarantine Agency.

complex group. JEV has an 11-kb encapsidated RNA genome composed of a single large open reading frame that encodes three structural and seven non-structural viral proteins (Fig. 1) [7,8]. The outer membrane of the virion consists of 90 envelope (E) protein homo-dimers, which play roles in the attachment to cellular receptors and membrane fusion [8]. Although the E protein was reported to be the major target for protection, animal experiments have shown that protection from disease is also mediated by antibodies against NS1, which is secreted from infected cells [9,10].

The Current Status of JE in Korea

From 1984 to 2009, few cases (a mean of 2.1) of JE in humans were reported annually due to the national vaccination policy for Korean infants [11]. In 2010 and 2012, however, there were human outbreaks involving 26 cases with 7 deaths and 20 cases, respectively [12]. There has been no official notification of an outbreak in the Korean pig population since 2007 (Fig. 2). Swine JE is the second notifiable disease in the Act on the Prevention of Contagious Animal Disease [13]. A JE vaccination program for the entire Korean sow population has

been performed annually to prevent reproductive disorders. In addition, regular serosurveillance has been conducted in the pig population to evaluate herd immunity in sows and monitor JEV in fattening pigs.

Epidemiology of JE: Genotype Shift Phenomenon

JEV can be classified into five genotypes based on similarities in the E gene (Fig. 3A) [14]. The replacement of JEV genotype 3 (G3) by genotype 1 (G1) was first reported in 1994 in Japan [15]. Similar to many other Southeast Asian countries, G1 viruses were introduced into Korea in 1993 and subsequently completely replaced G3 viruses as the dominant circulating JEV genotype (Fig. 3B) [3,16]. In addition, a JE G5 virus was first reported from a *Culex* mosquito in 2011; it is closely related to the Muar strain found in Malaysia in 1952 [17]. The genotypes differ by 10%-20% and 2%-6% at the nucleotide and amino acid levels, respectively. It has been suggested that JEV comprises a single serotype due to its relatively limited diversity at the amino acid level [18]. A mouse experiment demonstrated that vaccine strains induced significant, but lower, cross-neutralization against heterologous viruses [8,19]. However, Fan et al. [5] reported recently that live attenuated G3 swine vaccine (at222 strain) was partially protective against G1 viruses with very low cross-protective antibody titers. It is widely accepted that the current JEV vaccines derived from G3 strains are also able to protect against the now predominant G1 viruses. However, it is controversial whether the vaccine strain (G3) can induce a protective immune response against the now dominant JEV G1 strain.

The Current JE Vaccine for Animals

History of sequential attenuation of the JE vaccine strain

The JEV Anyang strain was isolated from the spleen of a newborn piglet in 1969 [20], and was sequentially attenuated in chicken embryo fibroblast primary cells (CEF) over 300 passages (Fig. 4). During the attenuation process, chorioallantoic

embryos or placentas, while the parent virus was recovered from two embryos. These results indicate that the vaccine strain induced protective immunity, but completely lost its pathogenic features and the capacity for mosquito transmission via pigs (Table 2) [23].

Immunity studies of the vaccine strain in animals

In a preliminary test of piglets, the vaccine strain induced significant neutralizing antibody titers (> 1:100; data not shown). In comparison, in a field test of sows, the first inoculation of the vaccine strain induced only low neutralizing antibody titers, close to the minimum protective titer of 1:10. The second inoculation increased the neutralizing antibody titer significantly. Based on a field test, the standard program for vaccinating sows against JE was established as two injections at a 3-week interval (Fig. 5A) [24]. In horses, the vaccine strain induced significant neutralizing antibodies with a single injection (Fig. 5B) [25].

Table 1. Pathogenicity of the Japanese encephalitis vaccine strain in mice

Mouse	LD ₅₀		
	Suckling mouse S.C.	3-week-old I.P.	4-week-old I.C.
Anyang parent (4 × 10 ⁷ pfu/mL)	0.01 pfu	80 pfu	0.01 pfu
Anyang285 ^{a)} (1 × 10 ⁷ pfu/mL)	-	-	9 × 10 ³ pfu

These results were originated from Kwon et al. [23]. S.C., subcutaneous; I.P., intraperitoneal; I.C., intracerebral route; pfu, plaque forming unit.

^{a)}The virus passage number in chicken primary embryo fibroblast cells.

Molecular Characterization of the Vaccine Strain

When the genome sequence of the vaccine strain was compared with those of other JEV strains, it was grouped with G3 viruses. At the nucleotide level, it showed high homology with other G3 viruses but differed clearly from G1 viruses [26]. At the start of the 3'-untranslated region, the genome of G3 viruses is characterized by specific nucleotide insertions right after the stop codon [16]. At the deduced amino acid level, there are several differences between G3 and G1 viruses scattered throughout the genome, including nonstructural proteins. There are also several differences from all other JEV strains, which were thought to be related to the viral attenuation or cell adaptation that occurred during the many passages in CEF cells. In the prM and E protein genes, which are

Table 2. Pathogenicity of the Japanese encephalitis vaccine strain in pigs

	Pathogenicity (pfu/mL)	
	Parent	Anyang300
Piglets (pfu/mL)		
Titer	4 × 10 ⁶	3 × 10 ⁵
Viremia	Up to 5DPI	-
NT titers ^{a)} (mean)	2,560	400
Sow (pfu/mL)		
Titer	4 × 10 ⁷	1 × 10 ⁷
Virus recovery from embryo	2/8	0/20
From placenta	0/1	0/3

Inoculation route: intramuscular. These results were originated from Kwon et al. [24].

^{a)}Neutralizing antibody titers.




Fig. 5. (A, B) Field test of the Japanese encephalitis vaccine strain. These results were originated from Kwon et al. [24,25]. SN titer, serum neutralizing antibody titer; WPI, weeks after inoculation.




Fig. 6. Molecular characterization of the Japanese encephalitis (JE) vaccine strain in animals. (A) Comparison of 3' UTR region between the vaccine strain and other JE viruses (JEVs). (B) Difference between the vaccine strain and other JEVs at the amino acid level. prM, pre-membrane and membrane; E, envelope; NS, non-structural; CEF, chicken primary embryo fibroblast cells. ^{a)}Amino acid position of the each viral proteins.

important for protective immunity, there are four amino acid differences compared to the G1 virus group, especially two amino acid differences related to the induction of neutralizing antibodies in the E domain III region. These results indicate that the vaccine strain clearly has the molecular character of JEV G3, with several amino acid differences from the JEV G1 strains that now prevail in Korea (Fig. 6).

Development of New JE Vaccines for Animals

Inactivated JE vaccine

A live attenuated JE vaccine (Anyang300 strain, G3) has been used in veterinary medicine in Korea for more than 30 years. However, the low immunogenicity of the current vaccine in pigs and the recent genotype shift phenomenon (G3 to G1) in

Asia have raised questions regarding the protective immunity generated with the current vaccine strain. New JEV G1 vaccines with enhanced immunogenicity are required for pigs. A new inactivated G1 JEV, which originated from a virus obtained from the blood of an infected piglet in 1999 (KV1899 strain), is now being developed for pigs, and its protective ability and safety are being evaluated in various animal species. The new vaccine induced significantly high neutralizing antibody levels in pigs and will be commercialized for pigs soon.

Live attenuated JE vaccine

To induce significantly high protective immunity in animals and considering the economic efficiency of the vaccine, a new live attenuated JE vaccine should be developed. To replace the current G3 vaccine, a recent G1 isolate, KV1899, was se-

quentially attenuated in African monkey kidney cells (Vero cells) until the virus completely lost its pathogenicity in various animals. The development and evaluation of this new live attenuated G1 virus are now in progress.

Other new JE vaccine candidates

For human vaccines, many new JE vaccine candidates have been reported, including recombinant virus-like particles [27], DNA vaccines containing the E and NS1 genes [28], and chimeric flavivirus vaccines [29,30]. Unfortunately, few researchers are attempting to develop JE veterinary vaccines using advanced technologies. We anticipate that novel, effective veterinary vaccines will be developed using reverse genetics systems in the near future.

Conclusion

In the JE endemic region, vaccination against JEV is the most effective measure for controlling JEV infection in humans and domestic pigs. This article reviewed the history of JEV vaccine development and the molecular characterization of the vaccine strain in animals. The low immunogenicity of the current vaccine in pigs and the recent genotype shift phenomenon (G3 to G1) in Korea have increased the demand for new G1 vaccines. New inactivated and live attenuated G1 vaccines have been developed to induce more effective immunity in animals. Furthermore, novel, effective vaccines should be developed for animals using advanced technologies so that we can deal promptly with newly emerging JE genotype viruses.

ORCID

Jin-Ju Nah <http://orcid.org/0000-0002-5021-0941>

Dong-Kun Yang <http://orcid.org/0000-0001-5765-3043>

Ha-Hyun Kim <http://orcid.org/0000-0001-6473-0035>

Jae-Young Song <http://orcid.org/0000-0002-9554-9252>

References

- Pan XL, Liu H, Wang HY, et al. Emergence of genotype I of Japanese encephalitis virus as the dominant genotype in Asia. *J Virol* 2011;85:9847-53.
- Mackenzie JS, Williams DT, Smith DW. Japanese encephalitis virus: the geographic distribution, incidence, and spread of a virus with a propensity to emerge in new areas. In: Tabor E, editor. *Emerging viruses in human populations*. Amsterdam: Elsevier B.V.; 2007. p.201-68.
- Yun SM, Cho JE, Ju YR, et al. Molecular epidemiology of Japanese encephalitis virus circulating in South Korea, 1983-2005. *Virology* 2010;7:127.
- Seo HJ, Kim HC, Klein TA, et al. Molecular detection and genotyping of Japanese encephalitis virus in mosquitoes during a 2010 outbreak in the Republic of Korea. *PLoS One* 2013;8:e55165.
- Fan YC, Chen JM, Chen YY, Lin JW, Chiou SS. Reduced neutralizing antibody titer against genotype I virus in swine immunized with a live-attenuated genotype III Japanese encephalitis virus vaccine. *Vet Microbiol* 2013;163:248-56.
- World Organisation for Animal Health (OIE). *Terrestrial manuals, chapter 2.1.7. Japanese encephalitis* [Internet]. Paris: OIE; [cited 2015 Mar 1]. Available from: http://www.oie.int/fileadmin/Home/eng/Health_standards/tahm/2.01.07_JEV.pdf.
- Heinz FX, Stiasny K. Flaviviruses and their antigenic structure. *J Clin Virol* 2012;55:289-95.
- Beasley DW, Lewthwaite P, Solomon T. Current use and development of vaccines for Japanese encephalitis. *Expert Opin Biol Ther* 2008;8:95-106.
- Heinz FX, Stiasny K. Flaviviruses and flavivirus vaccines. *Vaccine* 2012;30:4301-6.
- Li Y, Counor D, Lu P, Duong V, Yu Y, Deubel V. Protective immunity to Japanese encephalitis virus associated with anti-NS1 antibodies in a mouse model. *Virology* 2012;9:135.
- Sohn YM. Japanese encephalitis immunization in South Korea: past, present, and future. *Emerg Infect Dis* 2000;6:17-24.
- Jung Y, Cha G. Activity of Japanese encephalitis virus in the Republic of Korea, 2012. *Public Health Wkly Rep* 2013;6:325-8.
- Ministry of Agriculture and Forestry (MAF), Animal, Plant and Fisheries Quarantine and Inspection Agency (QIA). *Act of The Prevention of Contagious Animal Diseases in Korea*. Anyang: Ministry of Agriculture and Forestry (MAF), Animal, Plant and Fisheries Quarantine and Inspection Agency (QIA); 2013.
- Solomon T, Ni H, Beasley DW, Ekkelenkamp M, Cardoso MJ, Barrett AD. Origin and evolution of Japanese encephalitis virus in southeast Asia. *J Virol* 2003;77:3091-8.
- Morita K. Molecular epidemiology of Japanese encephalitis in East Asia. *Vaccine* 2009;27:7131-2.
- Yang DK, Kim BH, Kweon CH, Kwon JH, Lim SI, Han HR.

- Molecular characterization of full-length genome of Japanese encephalitis virus (KV1899) isolated from pigs in Korea. *J Vet Sci* 2004;5:197-205.
17. Takhampunya R, Kim HC, Tippayachai B, et al. Emergence of Japanese encephalitis virus genotype V in the Republic of Korea. *Virology* 2011;8:449.
 18. Tsarev SA, Sanders ML, Vaughn DW, Innis BL. Phylogenetic analysis suggests only one serotype of Japanese encephalitis virus. *Vaccine* 2000;18 Suppl 2:36-43.
 19. Kurane I, Takasaki T. Immunogenicity and protective efficacy of the current inactivated Japanese encephalitis vaccine against different Japanese encephalitis virus strains. *Vaccine* 2000;18 Suppl 2:33-5.
 20. Kwon HJ, Lee CK, Kang BJ, Lim YM. Studies on Japanese encephalitis live vaccine. I. Isolation of Japanese encephalitis virus (Anyang strain) from a new-born piglet. *Res Rep Off Rural Dev* 1974;16:1-5.
 21. Kwon HJ, Kang BJ, Lim YM, Lee CK. Studies on Japanese encephalitis live vaccine. II. Development of an attenuated strain of virus (Anyang strain). *Res Rep Off Rural Dev* 1975;17:95-100.
 22. Kwon HJ, Kang BJ, Lim YM, Lee CK. Studies on Japanese encephalitis live vaccine. V. Multiplicity of Japanese encephalitis virus (Anyang strain) in chick embryo fibroblast and duck embryo fibroblast cell cultures. *Res Rep Off Rural Dev* 1978;20:11-6.
 23. Kwon HJ, Kang BJ, Lim YM, et al. Studies on Japanese encephalitis live vaccine. III. Pathogenicity of tissue culture attenuated strain of virus (Anyang Strain). *Res Rep Off Rural Dev* 1976;18:21-8.
 24. Kwon HJ, Kang BJ, Lim YM, Lee CK, Jeon YS. Studies on Japanese encephalitis live vaccine. VI. Field application of Japanese encephalitis live virus (Anyang attenuated strain) vaccine. *Res Rep Off Rural Dev* 1978;20:17-28.
 25. Kwon HJ, Kang BJ, Lim YM, Lee CK, Jeon YS. Studies on Japanese encephalitis live vaccine. VII. Pathogenicity and immunogenicity of horses with Anyang strain of attenuated virus. *Res Rep Off Rural Dev* 1978;20:29-34.
 26. Lee JA, Yang DK, Kim HH, et al. Evaluation of Japanese encephalitis virus vaccine strains currently used in pigs by molecular characterization. *Korean J Vet Serv* 2012;35:169-74.
 27. Yamaji H, Segawa M, Nakamura M, Katsuda T, Kuwahara M, Konishi E. Production of Japanese encephalitis virus-like particles using the baculovirus-insect cell system. *J Biosci Bioeng* 2012;114:657-62.
 28. Gao N, Chen W, Zheng Q, et al. Co-expression of Japanese encephalitis virus prM-E-NS1 antigen with granulocyte-macrophage colony-stimulating factor enhances humoral and anti-virus immunity after DNA vaccination. *Immunol Lett* 2010;129:23-31.
 29. Ishikawa T, Widman DG, Bourne N, Konishi E, Mason PW. Construction and evaluation of a chimeric pseudoinfectious virus vaccine to prevent Japanese encephalitis. *Vaccine* 2008;26:2772-81.
 30. Ishikawa T, Wang G, Widman DG, et al. Enhancing the utility of a prM/E-expressing chimeric vaccine for Japanese encephalitis by addition of the JEV NS1 gene. *Vaccine* 2011;29:7444-55.