

This article was downloaded by: [Loughborough University]

On: 17 January 2012, At: 03:20

Publisher: Routledge

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Sports Sciences

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/rjsp20>

Energy availability in athletes

Anne B. Loucks^a, Bente Kiens^b & Hattie H. Wright^c

^a Department of Biological Sciences, Ohio University, Athens, Ohio, USA

^b The Molecular Physiology Group, Department of Exercise and Sport Sciences, University of Copenhagen, Copenhagen, Denmark

^c Center of Excellence for Nutrition, Faculty of Health Sciences, North-West University, Potchefstroom, South Africa

Available online: 28 Jul 2011

To cite this article: Anne B. Loucks, Bente Kiens & Hattie H. Wright (2011): Energy availability in athletes, Journal of Sports Sciences, 29:sup1, S7-S15

To link to this article: <http://dx.doi.org/10.1080/02640414.2011.588958>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Energy availability in athletes

ANNE B. LOUCKS¹, BENTE KIENS², & HATTIE H. WRIGHT³

¹Department of Biological Sciences, Ohio University, Athens, Ohio, USA, ²The Molecular Physiology Group, Department of Exercise and Sport Sciences, University of Copenhagen, Copenhagen, Denmark, and ³Center of Excellence for Nutrition, Faculty of Health Sciences, North-West University, Potchefstroom, South Africa

(Accepted 13 May 2011)

Abstract

This review updates and complements the review of energy balance and body composition in the Proceedings of the 2003 IOC Consensus Conference on Sports Nutrition. It argues that the concept of energy availability is more useful than the concept of energy balance for managing the diets of athletes. It then summarizes recent reports of the existence, aetiologies, and clinical consequences of low energy availability in athletes. This is followed by a review of recent research on the failure of appetite to increase *ad libitum* energy intake in compensation for exercise energy expenditure. The review closes by summarizing the implications of this research for managing the diets of athletes.

Keywords: Energy availability, energy balance, diet, exercise, appetite

Introduction

In the 2003 IOC Consensus Conference on Sports Nutrition, evidence was presented that many athletes, most often female athletes, were deficient in energy, and especially energy in the form of carbohydrates, resulting in impaired health and performance (Loucks, 2004). It was emphasized, however, that energy balance is not the objective of athletic training whenever athletes seek to modify their body size and composition to achieve performance objectives. They then need to carefully manage their diet and exercise regimens to avoid compromising their health.

Distinctions between energy availability and energy balance

In the field of bioenergetics, the concept of energy availability recognizes that dietary energy is expended in several fundamental physiological processes, including cellular maintenance, thermoregulation, growth, reproduction, immunity, and locomotion (Wade & Jones, 2004). Energy expended in one of these processes is not available for others. Therefore, bioenergeticists investigate the effects of a particular metabolic demand on physiological systems in terms

of energy availability. They define energy availability as dietary energy intake minus the energy expended in the particular metabolic demand of interest. In experiments investigating effects of cold exposure, for example, energy availability would be defined, quantified, and controlled as dietary energy intake minus the energy cost of thermogenesis.

Exercise training increases, and in endurance sports may double or even quadruple, the amount of energy expended in locomotion. In exercise physiology, therefore, energy availability is defined as dietary energy intake minus the energy expended in *exercise* ($EA = EI - EEE$). As the amount of dietary energy remaining after exercise training for all other metabolic processes, energy availability is an *input to* the body's physiological systems.

In the field of dietetics, the concept of energy balance has been the usual basis of research and practice. Defined as dietary energy intake minus *total* energy expenditure ($EB = EI - TEE$), energy balance is the amount of dietary energy added to or lost from the body's energy stores after the body's physiological systems have done all their work for the day. Thus energy balance is an *output from* those systems. For healthy young adults, $EB = 0 \text{ kcal} \cdot \text{day}^{-1}$ when $EA = 45 \text{ kcal} \cdot \text{kg FFM}^{-1} \cdot \text{day}^{-1}$ (where FFM = fat-free mass).




Figure 1. Negative energy balance rising at a rate of ≈ 90 kcal · day⁻¹ as metabolic processes were suppressed while the energy intake (2770 kcal · day⁻¹), exercise energy expenditure (840 kcal · day⁻¹), and energy availability (2770 - 840 = 1930 kcal · day⁻¹) of eight lean, untrained men remained constant. EI (+) = energy intake, TEE (□) = total energy expenditure, EEE (○) = exercise energy expenditure, EA (●) = energy availability, EB (■) = energy balance. Original figure based on data in Stubbs et al. (2004).

The contrast between energy availability and energy balance is illustrated in Figure 1, which shows data collected while eight lean, untrained men lived in a room calorimeter for a week (Stubbs et al., 2004). During that week, their energy intake (2770 kcal · day⁻¹), exercise energy expenditure (840 kcal · day⁻¹), and energy availability (2770 - 840 = 1930 kcal · day⁻¹ ≈ 30 kcal · kg FFM⁻¹ · day⁻¹) were constant. Meanwhile, the magnitude of their negative energy balance (2770 - 4500 = -1730 kcal · day⁻¹ on Day 1) decreased towards zero at a rate of ~ 90 kcal · day⁻¹ as various physiological processes slowed down. At this rate, they would have recovered EB = 0 kcal · day⁻¹ (a pathological state of energy balance achieved by suppressing physiological systems) in 3 weeks, while remaining in severely low energy availability.

Undergraduate nutrition textbooks assert that energy requirements can be determined by measuring energy expenditure, but measures of energy expenditure contain no information about whether physiological systems are functioning in a healthy manner. Because physiological processes are suppressed by severely low energy availability, measurements of total or resting energy expenditure will underestimate a chronically undernourished athlete's energy requirements.

Therefore, because energy balance is an output from, rather than an input to, physiological systems, because it does not contain reliable information about energy requirements, and because it is not even the objective of athletic training, energy balance is not a useful concept for managing an athlete's diet.

Energy deficiency in athletes: Existence, aetiologies, and consequences

At the 2003 IOC Consensus Conference, the existence of widespread energy deficiency in athletes was still questioned. Since then, the IOC Medical Commission has published two position stands (Sangenis et al., 2005, 2006) and the American College of Sports Medicine (ACSM) has published a revised position stand (Nattiv, Loucks, Manore, Sundgot-Borgen, & Warren, 2007) on the "female athlete triad". In addition, a coaches' handbook on Managing the Female Athlete Triad developed by the co-chairs of the athlete interest group of the Academy of Eating Disorders has been published by the US National Collegiate Athletics Association (NCAA) (Sherman & Thompson, 2005). All four publications attribute the functional hypothalamic menstrual disorders and low bone mineral density found in many female athletes to energy deficiency, but the ACSM position stand differs from the other three in that it excludes disordered eating and eating disorders as necessary components of the triad. The ACSM emphasizes that athletes who expend large amounts of energy in prolonged exercise training can become energy deficient without eating disorders, disordered eating or even dietary restriction.

The ACSM identified three distinct origins of energy deficiency in athletes. The first is obsessive eating disorders with their attendant clinical mental illnesses. The second is intentional and rational but mismanaged efforts to reduce body size and fatness to qualify for and succeed in athletic competitions. This mismanagement may or may not include disordered eating behaviours such as fasting, diet pills, laxatives, diuretics, enemas, and vomiting that are entrenched parts of the culture and lore of some sports. The third is the inadvertent failure to increase energy intake to compensate for the energy expended in exercise. The percentages of cases of the female athlete triad originating from these three sources are unknown, but ACSM emphasized that any epidemiological study requiring the presence of an eating disorder or disordered eating for diagnosing cases of the triad (e.g. Schtscherbyna, Soares, de Oliveira, & Ribeiro, 2009) will underestimate its prevalence.

Sports vary greatly in the relative importance of various factors for competitive success. As they strive to achieve sport-specific mixes of these factors, athletes engage in different diet and exercise behaviours that impact energy availability. In endurance sports, prolonged exercise training greatly reduces energy availability, unless energy intake is increased to replace the energy expended in exercise. In sports where less energy is expended in training, dietary restriction may be a prominent part of the strategy for reducing energy availability to modify body size and composition.

Female athletes may also under-eat for reasons unrelated to sport. Around the world about twice as many young women as men *at every decile of body mass index* perceive themselves to be overweight (Wardle, Haase, & Steptoe, 2006). The disproportionate numbers actively trying to lose weight are even higher, and this disproportion *increases* as body mass index declines, so that almost nine times as many lean women as lean men are actively trying to lose weight! Indeed, more young female athletes report improvement of appearance than improvement of performance as a reason for dieting (Martinsen, Bratland-Sanda, Eriksson, & Sundgot-Borgen, 2010). Thus issues unrelated to sport may need to be addressed to persuade female athletes to eat appropriately.

The controversy about whether female athletes can increase glycogen stores as much as male athletes is instructive in this regard. An experiment in which participants consumed diets containing high and low percentages of carbohydrates found that women could not do so (Tarnopolsky, Atkinson, Phillips, & MacDougall, 1995). Subsequently, it was noted that the total energy intake (per kilogram of body weight) of the women in that study had been so low that the amount of carbohydrate they consumed on the high percent carbohydrate diet was no greater than the amount consumed by the men on the low percent carbohydrate diet. Later research showed that women could, indeed, load glycogen like men when they ate as much as men (per kilogram of body weight) (James et al., 2001; Tarnopolsky et al., 2001).

In the 2003 IOC Consensus Conference, the disruption of reproductive function at energy availabilities $< 30 \text{ kcal} \cdot \text{kg FFM}^{-1} \cdot \text{day}^{-1}$ was discussed in some detail and the low bone mineral density (BMD) found in amenorrhoeic athletes was represented as being mediated by oestrogen deficiency (Loucks, 2004). Since then, oestrogen-independent mechanisms by which low energy availability can reduce BMD have also been identified (Ihle & Loucks, 2004). As energy availability declines, the rate of bone protein synthesis declines along with insulin, which enhances amino acid uptake, in a linear dose-response manner. By contrast, the rate of bone mineralization declines abruptly as energy availability declines below $30 \text{ kcal} \cdot \text{kg FFM}^{-1} \cdot \text{day}^{-1}$, as do concentrations of insulin-like growth factor-1 and tri-iodothyronine. These effects occurred within 5 days of the onset of energy deficiency, and without a reduction in oestrogen concentration.

In older adults, fracture risk doubles for each reduction of one standard deviation below mean peak young adult BMD. In adolescents, fracture risk can rise even as BMD increases. Because BMD

normally doubles during the decade of adolescence, a child entering adolescence with a high BMD relative to others of the same age can accrue bone mineral so slowly that adulthood is entered with a relatively low BMD. Because low BMD is an aetiological factor in stress fractures, anything that impairs bone mineral accrual during adolescence is undesirable. Unfortunately, this is exactly what was found in a study of 183 interscholastic competitive female athletes, of whom 93 were endurance runners and 90 were non-runners (Barrack, Rauh, & Nichols, 2010). The BMD *z*-scores were similar in runners and non-runners aged 13–15 years, but were significantly lower in runners than non-runners at 16–18 years of age.

Also questioned at the 2003 IOC Consensus Conference was whether energy deficiency and its clinical consequences were a problem among elite athletes. Since then a study of 50 British national or higher standard middle- and long-distance runners found BMD to be lower in amenorrhoeic runners and higher in eumenorrhoeic runners compared with European reference data (Gibson, Mitchell, Harries, & Reeve, 2004). The duration of eumenorrhoea was positively associated with spine BMD, and the rate of bone mineralization was reduced in the amenorrhoeic runners. Alone, the Eating Attitudes Test (EAT) is not clinically diagnostic for eating disorders, but in this study scores on the EAT classified one of 24 amenorrhoeic runners and none of nine oligomenorrhoeic runners as having an eating disorder, while eight amenorrhoeic runners and three oligomenorrhoeic runners were classified as practising disordered eating behaviours. This left 63% of the cases of amenorrhoea and 67% of the cases of oligomenorrhoea unaccounted for by the EAT test (Figure 2). Similarly, a low body mass index ($< 18.5 \text{ kg} \cdot \text{m}^{-2}$) failed to account for 67% of the cases of amenorrhoea and 67% of the cases of oligomenorrhoea (Figure 3). Another study diagnosed low BMD (*z*-score less than -1) in the lumbar spine of 34% and osteoporosis (*z*-score less than -2) in the radius of 33% of 44 elite British female endurance runners (Pollock et al., 2010). Reductions in BMD over time were associated with training volume. These findings led the authors to recommend that all female endurance athletes undergo dual-energy X-ray absorptiometry screening.

Ovarian function depends critically upon the frequency with which the pituitary gland secretes luteinizing hormone (LH) into the bloodstream, and LH pulsatility in exercising women depends on energy availability, rather than energy intake or energy expenditure separately (Loucks, Verdun, & Heath, 1998). Furthermore, exercise has no suppressive effect on LH pulsatility beyond the impact of its energy cost on energy availability (Loucks et al.,

1998). Reproductive function (Loucks & Thuma, 2003) and bone formation (Ihle & Loucks, 2004) are impaired abruptly and promptly below a threshold of energy availability ($< 30 \text{ kcal} \cdot \text{kg FFM}^{-1} \cdot \text{day}^{-1}$), which corresponds closely to resting metabolic rate. Figure 4 shows sleeping metabolic rate (SMR) measured by indirect calorimetry in young adult men ($n=20$) and women ($n=17$) (Westerterp, 2003). Sleeping metabolic rate is slightly less than resting metabolic rate by the small energy expendi-




Figure 2. Logistic relationship between the order of menstrual dysfunction (right ordinate scale: 0 (+)=eumenorrhoea; 1 (x)=oligomenorrhoea; 2 (•)=amenorrhoea) and total EAT score ($P=0.014$ for model). Left ordinate scale is a scale of probability. The right ordinate scale shows the proportions of participants in each category. DE = classified as disordered eating. ED = classified as eating disorder. Figure modified from Gibson et al. (2004).




Figure 3. Logistic relationship between the order of menstrual dysfunction (right ordinate scale: 0 (+)=eumenorrhoea; 1 (x)=oligomenorrhoea; 2 (•)=amenorrhoea) and body mass index (BMI) ($P < 0.001$ for model). Left ordinate scale is a scale of probability. The right ordinate scale shows the proportions of participants in each category. BMI < 18.5 = underweight. Figure modified from Gibson et al. (2004).

ture associated with being awake. In Figure 4, the solid regression line relating sleeping metabolic rate to fat-free mass ($\text{SMR} [\text{MJ} \cdot \text{day}^{-1}] = 2.27 + 0.091 \times \text{FFM} [\text{kg}]$) has a significant y -intercept. The dashed line through the data and the origin has a slope of $30 \text{ kcal} \cdot \text{kg FFM}^{-1} \cdot \text{day}^{-1}$. As Figure 4 shows, energy availabilities $< 30 \text{ kcal} \cdot \text{kg FFM}^{-1} \cdot \text{day}^{-1}$ provide less energy than is required for physiological systems in young adults to function at rest.

Observational and experimental data indicate that low energy availability also suppresses Type 1 immunity. The immune system mounts different defences against two types of pathogens. Type 1 defences are mounted against intracellular pathogens like viruses, while Type 2 defences are mounted against extracellular pathogens like bacteria. Endurance athletes frequently suffer upper respiratory tract infections (URTI) caused by viruses. A survey of all members of Swedish teams participating in the Olympic Games of 2002 and 2004 found that those participating in disciplines emphasizing leanness made more frequent attempts to lose weight, trained longer, and reported almost twice as many illnesses, primarily URTI, during the preceding 3 months (Hagmar, Hirschberg, Berglund, & Berglund, 2008). The results of a recent experiment challenge the hypothesis that Type 1 immunity in athletes might be suppressed by exercise itself (Lancaster et al., 2005). Participants expended 2200 kcal of energy by exercising for 2½ h at 65% of maximal oxygen uptake ($\dot{V}O_{2\text{max}}$). Replacing just 23% of this energy with carbohydrate reduced the suppression of Type 1 defences by an average of 65%. Thus, ingesting sufficient energy and nutrients is vital for supporting




Figure 4. Sleeping metabolic rate plotted as a function of fat-free mass (FFM). • = females; ○ = males. The solid regression line has a significant non-zero intercept. The dashed line is $30 \text{ kcal} \cdot \text{kg FFM}^{-1} \cdot \text{day}^{-1}$. Figure modified from Westerterp (2003).

immune function, and even more so for immune-compromised individuals, such as those infected with HIV (Fenton & Silverman, 2008), whose resting energy expenditure is elevated (Mangili, Murman, Zampini, & Wanke, 2006). Therefore, HIV-infected athletes should take special care to reach their energy and nutrient needs.

Seventeen years have now passed since the first IOC Consensus Conference on Nutrition. Yet studies of energy intake and total energy expenditure continue to report that elite American figure skaters (Ziegler, Nelson, Barratt-Fornell, Fiveash, & Drewnowski, 2001), elite Kenyan runners (Fudge et al., 2006), and high-performance Canadian athletes in several sports (Lun, Erdman, & Reimer, 2009) train in substantial negative energy balance. Depending on the level of energy availability, such negative energy balance may either impair or benefit health and performance. In the current review period, two studies of elite athletes reported energy intake and exercise energy expenditure so that their average energy availability could be estimated. In the week before a race, the high percent carbohydrate diet of male Kenyan runners provided an energy availability of $34 \text{ kcal} \cdot \text{kg FFM}^{-1} \cdot \text{day}^{-1}$ (Onywera, Kiplamai, Boit, & Pitsiladis, 2004). This energy availability may or may not have been appropriate depending on their athletic objectives at the time. If their performance in that particular race was less important than losing weight to improve performance in a later race, it was fine. However, no athletic objective would appear to justify professional male cyclists training for the Tour de France 6 months later at an energy availability of only $8 \text{ kcal} \cdot \text{kg FFM}^{-1} \cdot \text{day}^{-1}$ (Vogt et al., 2005). Such observations indicate that the diet and exercise regimens of elite athletes range widely and are sometimes dangerously energy deficient.

Effects of prolonged exercise on hunger and energy intake

Eating disorders may be intractable, and weight and fat loss programmes may be challenging to manage effectively and safely, but these two origins of low energy availability in athletes are at least familiar to sports dietitians. The third origin, the suppression of appetite by prolonged exercise, appears to be less familiar. Neglect of appetite as an important factor in sports nutrition is indicated by the appearance of the word “appetite” only once (and then only in a discussion of fluid losses at high altitude) in the recently revised position stand on nutrition and athletic performance jointly adopted by the American Dietetic Association, the Dietitians of Canada, and the American College of Sports Medicine (Rodriguez, DiMarco, & Langley, 2009).

Some of the then available evidence that appetite is not a reliable indicator of energy needs in athletes was reviewed at the 2003 IOC Consensus Conference (Loucks, 2004). Shortly afterwards, the suppressive effect of prolonged exercise on *ad libitum* energy intake was clearly demonstrated by the experiment in which eight lean, untrained men expended $\sim 840 \text{ kcal} \cdot \text{day}^{-1}$ by cycle ergometry as they lived in a room calorimeter for 7 days (Stubbs et al., 2004). During that week, their *ad libitum* energy intake was similar to another week in the room calorimeter when they did not exercise. This inadvertent failure to increase energy intake in compensation for exercise energy expenditure reduced their *ad libitum* energy availability by $\sim 10 \text{ kcal} \cdot \text{kg FFM}^{-1} \cdot \text{day}^{-1}$.

The participants in that experiment actually performed the experiment four times, with and without exercise while consuming equally palatable 62% and 37% carbohydrate diets in a 2×2 crossover design. Compared with weeks when the participants ate the 37% carbohydrate diet, their *ad libitum* energy intake declined by $\sim 1000 \text{ kcal} \cdot \text{day}^{-1}$ on the 62% carbohydrate diet, reducing their *ad libitum* energy availability by $\sim 16 \text{ kcal} \cdot \text{kg FFM}^{-1} \cdot \text{day}^{-1}$. Moreover, the suppressive effects of prolonged exercise and the high percent carbohydrate diet were additive. *Ad libitum* energy availability declined from $\sim 47 \text{ kcal} \cdot \text{kg FFM}^{-1} \cdot \text{day}^{-1}$ when the participants were sedentary on the 37% carbohydrate diet to $\sim 21 \text{ kcal} \cdot \text{kg FFM}^{-1} \cdot \text{day}^{-1}$ when they exercised on the 62% carbohydrate diet (Figure 5).

These findings in lean, untrained men exercising in a laboratory confirmed a previous report of a high

	Intake (kcal · day ⁻¹)		Balance (kcal · day ⁻¹)		Availability (kcal · kg ⁻¹ FFM · day ⁻¹)	
	840	0	840	0	840	0
37% Dietary CHO	3350	3060	-910	+170	38	47
62%	2230	2110	-1960	-480	21	32
	840 0		840 0		840 0	
	Exercise energy expenditure (kcal · day ⁻¹)					

Figure 5. The *ad libitum* energy intake, energy balance, and energy availability of eight lean men living in a laboratory for 7 days during an experiment contrasting two diets (50% fat, 32% carbohydrate [CHO]; and 25% fat, 67% CHO) and two levels of exercise (840 and 0 kcal · day⁻¹) (data from Stubbs et al., 2004). Energy availability was estimated assuming 16% body fat. Appetite failed to match energy intake to activity-induced energy expenditure on either diet. Appetite also failed to drive energy intake on a low fat, high carbohydrate diet to match energy expenditure at either activity level. These effects were additive. Reproduced from Loucks (2007) with permission from Adis, a Wolters Kluwer business (© Adis Data Information BV 2007. All rights reserved).

2010), as well as on male and female endurance trained runners who ran for 90 min at $\sim 60\%$ of $\dot{V}O_{2\max}$ before a 10-km time-trial as fast as possible on a treadmill after 2 days of carbohydrate loading and a standardized 70% carbohydrate breakfast (Russel, Willis, Ravussin, & Larson-Meyer, 2009).

Other experiments have investigated individual differences in the effects of exercise training on *ad libitum* energy intake over a period of 12 weeks (King, Hopkins, Caudwell, Stubbs, & Blundell, 2008; King et al., 2009). These experiments on overweight and obese men and women confirmed the exercise-induced suppression of *ad libitum* energy intake mediated by suppressed hunger and increased satiety that had been found in shorter term experiments, but they also revealed a high degree of individual variability in weight and fat loss. When participants were retrospectively classified as compensators or non-compensators based on actual weight loss compared with the weight loss expected from exercise energy expenditure, *ad libitum* energy intake was found to have increased in the compensators and decreased in the non-compensators (King et al., 2008). Exercise induced a similar increase in satiety in both groups, which was even greater after 12 weeks of training than before, but compensators became progressively hungrier during the experiment (King et al., 2009).

Thus, findings in both trained and untrained male and female participants consistently demonstrate that a single bout of diverse forms of exercise acutely suppresses *ad libitum* energy intake and that exercise training chronically maintains the resulting energy deficiency for many weeks. This effect has been interpreted to be at least partially responsible for exercise-induced anorexia (Russel et al., 2009). Whether some female athletes become hungrier and increase *ad libitum* energy intake as their training progresses, and thereby avoid developing functional hypothalamic amenorrhoea (like overweight participants who do not lose weight on an exercise training programme), has yet to be investigated.

Implications for managing the diets of athletes

The studies of *ad libitum* energy intake cited above demonstrate that appetite is an unreliable indicator of energy requirements for athletes engaged in prolonged exercise training, just as thirst is an unreliable indicator of water requirements during a marathon race. Marathon runners are advised not to wait until they are thirsty before they begin drinking during a race. Similarly, athletes who engage in prolonged exercise training should be advised to eat by discipline, that is, to eat specific amounts of particular foods at planned times, instead of waiting

until they are hungry and then eating only until they are satisfied.

The recommendation for athletes in endurance sports to consume a diet containing a high percentage of carbohydrates should also be reconsidered. The original evidentiary basis for this recommendation was the finding that high carbohydrate intake for a few days before a high-intensity endurance event increased glycogen storage and improved performance (Costill, 1988). It may be reasonable to expect endurance athletes to override their appetites to consume a high volume of a high percent carbohydrate diet for glycogen loading for a day or two as a pre-race tactic, but it may not be realistic to expect them to override the appetite-suppressive effects of both a high percent carbohydrate diet and prolonged exercise as a lifestyle. Meanwhile, research is needed to determine whether the suppression of *ad libitum* energy intake by a high percent carbohydrate diet can be ameliorated by increasing the proportion of refined carbohydrate in the diet (Mann et al., 2007).

The American Academy of Pediatrics (AAP, 2005), the IOC Medical Commission (Sangenis et al., 2005), and ACSM (Nattiv et al., 2007) have all recommended that national and international governing bodies of sports and athletic organizations put policies and procedures in place to eliminate potentially harmful weight loss practices of female athletes. Procedures and policies were not specified, because best practices may be sport-specific. These recommendations followed the establishment of such policies and procedures by the governing bodies of US men's collegiate wrestling in the late 1990s (Oppliger, Utter, Scott, Dick, & Klossner, 2006) and men's international ski jumping in 2004 (FIS, 2004). The procedures specified by NCAA Wrestling Rule 3 are expensive, intrusive for the athlete, labour-intensive for athletic trainers, and bureaucratic with local and national databases, but they seem to have been effective in reducing unhealthy weight loss behaviours and promoting competitive equity (Oppliger et al., 2006). Based on this success, a call has gone out for the International Judo Federation (IJF) to implement regulations to improve weight management behaviours among judo competitors, and for these regulations to be adopted by all National and Regional Federations (Artoli et al., 2010a).

By contrast, the International Ski Federation (FIS) employed a very different strategy for preventing excessive weight loss practices in ski jumpers (FIS, 2004). Instead of policing athlete behaviour, the FIS removed the motivation for ski jumpers to pursue excessive weight loss objectives. Ski jumpers have their height and weight measured at the top of the hill immediately before their jump, whereupon they are simply issued skis that are shorter or longer in

proportion to their body mass index. Shorter skis reduce aerodynamic lift to compensate for the advantage a lighter jumper would otherwise gain, so that a jumper's success depends on their skill rather than their body weight. Adoption of this new rule reduced the percentage of underweight ski jumpers in the next World Cup competition from 23% to 8% (Muller, Groschl, Muller, & Sudi, 2006). Replacing body mass index with mass index (MI = body mass divided by the square of sitting height) has been proposed to further improve the rule (Muller, 2009).

Periodization of training may require a periodization of energy availability to achieve training objectives. Experimental evidence indicates that athletes should follow diet and exercise regimens that provide energy availabilities of 30–45 kcal · kg FFM⁻¹ · day⁻¹ while training to reduce body size or fatness. However, if athletes in other sports are like judo competitors, personal counselling of athletes by sports dietitians may not be the most effective way to moderate athlete behaviour to prevent excessive energy deficiency. Among seven different types of advisors for weight management behaviour, judo competitors ranked dietitians second to last, ahead only of physicians (Artioli et al., 2010b). The most influential advisor was the coach. In that context, a more effective way to modify athlete behaviour may be for sports dietitians and nutritionists to educate coaches, using workshops and handbooks that are specifically targeted at them, about the importance of energy availability and practical techniques for managing it.

References

- AAP (2005). American Academy of Pediatrics Committee on Sports Medicine and Fitness: Promotion of healthy weight-control practices in young athletes. *Pediatrics*, *116*, 1557–1564.
- Artioli, G., Franchini, E., Nicastro, H., Sterkowicz, S., Solis, M. Y., & Lancha, A. H., Jr. (2010a). The need of a weight management control program in judo: A proposal based on the successful case of wrestling. *Journal of the International Society of Sports Nutrition*, *7*, 15.
- Artioli, G. G., Gualano, B., Franchini, E., Scagliusi, F. B., Takesian, M., Fuchs, M., et al. (2010b). Prevalence, magnitude, and methods of rapid weight loss among judo competitors. *Medicine and Science in Sports and Exercise*, *42*, 436–442.
- Ballard, T. P., Melby, C. L., Camus, H., Cianciulli, M., Pitts, J., Schmidt, S., et al. (2009). Effect of resistance exercise, with or without carbohydrate supplementation, on plasma ghrelin concentrations and postexercise hunger and food intake. *Metabolism*, *58*, 1191–1199.
- Barrack, M. T., Rauh, M. J., & Nichols, J. F. (2010). Cross-sectional evidence of suppressed bone mineral accrual among female adolescent runners. *Journal of Bone and Mineral Research*, *25*, 1850–1857.
- Batterham, R. L., Le Roux, C. W., Cohen, M. A., Park, A. J., Ellis, S. M., Patterson, M., et al. (2003). Pancreatic polypeptide reduces appetite and food intake in humans. *Journal of Clinical Endocrinology and Metabolism*, *88*, 3989–3992.
- Bergouignan, A., Momken, I., Schoeller, D. A., Normand, S., Zahariev, A., Lescure, B., et al. (2010). Regulation of energy balance during long-term physical inactivity induced by bed rest with and without exercise training. *Journal of Clinical Endocrinology and Metabolism*, *95*, 1045–1053.
- Borer, K. T. (2010). Nonhomeostatic control of human appetite and physical activity in regulation of energy balance. *Exercise and Sport Sciences Reviews*, *38*, 114–121.
- Costill, D. L. (1988). Carbohydrates for exercise: Dietary demands for optimal performance. *International Journal of Sports Medicine*, *9*, 1–18.
- Fenton, M., & Silverman, E. (2008). Medical nutrition therapy for human immunodeficiency virus (HIV) infection and acquired immunodeficiency syndrome (AIDS). In L. Mahan & S. Escott-Stump (Eds.), *Krause's food, nutrition and diet therapy* (pp. 1008–1016). Philadelphia, PA: W. B. Saunders.
- FIS (2004). International Ski Federation World Congress, Miami, FL, USA.
- Fudge, B. W., Westerterp, K. R., Kiplamai, F. K., Onyvera, V. O., Boit, M. K., Kayser, B., et al. (2006). Evidence of negative energy balance using doubly labelled water in elite Kenyan endurance runners prior to competition. *British Journal of Nutrition*, *95*, 59–66.
- Gibson, J. H., Mitchell, A., Harries, M. G., & Reeve, J. (2004). Nutritional and exercise-related determinants of bone density in elite female runners. *Osteoporosis International*, *15*, 611–618.
- Hagmar, M., Hirschberg, A. L., Berglund, L., & Berglund, B. (2008). Special attention to the weight-control strategies employed by Olympic athletes striving for leanness is required. *Clinical Journal of Sports Medicine*, *18*, 5–9.
- Horvath, P. J., Eagen, C. K., Fisher, N. M., Leddy, J. J., & Pendergast, D. R. (2000a). The effects of varying dietary fat on performance and metabolism in trained male and female runners. *Journal of the American College of Nutrition*, *19*, 52–60.
- Horvath, P. J., Eagen, C. K., Ryer-Calvin, S. D., & Pendergast, D. R. (2000b). The effects of varying dietary fat on the nutrient intake in male and female runners. *Journal of the American College of Nutrition*, *19*, 42–51.
- Ihle, R., & Loucks, A. B. (2004). Dose–response relationships between energy availability and bone turnover in young exercising women. *Journal of Bone and Mineral Research*, *19*, 1231–1240.
- James, A. P., Lorraine, M., Cullen, D., Goodman, C., Dawson, B., Palmer, T. N., et al. (2001). Muscle glycogen supercompensation: Absence of a gender-related difference. *European Journal of Applied Physiology*, *85*, 533–538.
- King, J. A., Miyashita, M., Wasse, L. K., & Stensel, D. J. (2010). Influence of prolonged treadmill running on appetite, energy intake and circulating concentrations of acylated ghrelin. *Appetite*, *54*, 492–498.
- King, N. A., Caudwell, P. P., Hopkins, M., Stubbs, J. R., Naslund, E., & Blundell, J. E. (2009). Dual-process action of exercise on appetite control: Increase in orexigenic drive but improvement in meal-induced satiety. *American Journal of Clinical Nutrition*, *90*, 921–927.
- King, N. A., Hopkins, M., Caudwell, P., Stubbs, J. R., & Blundell, J. E. (2008). Individual variability following 12 weeks of supervised exercise: Identification and characterization of compensation for exercise-induced weight loss. *International Journal of Obesity (London)*, *32*, 177–184.
- Lancaster, G. I., Khan, Q., Drysdale, P. T., Wallace, F., Jeukendrup, A. E., Drayson, M. T., et al. (2005). Effect of prolonged exercise and carbohydrate ingestion on type 1 and type 2 T lymphocyte distribution and intracellular cytokine production in humans. *Journal of Applied Physiology*, *98*, 565–571.
- Loucks, A. B. (2004). Energy balance and body composition in sports and exercise. *Journal of Sports Sciences*, *22*, 1–14.

- Loucks, A. B. (2007). Low energy availability in the marathon and other endurance sports. *Sports Medicine*, 37, 348–352.
- Loucks, A. B., & Thuma, J. R. (2003). Luteinizing hormone pulsatility is disrupted at a threshold of energy availability in regularly menstruating women. *Journal of Clinical Endocrinology and Metabolism*, 88, 297–311.
- Loucks, A. B., Verdun, M., & Heath, E. M. (1998). Low energy availability, not stress of exercise, alters LH pulsatility in exercising women. *Journal of Applied Physiology*, 84, 37–46.
- Lun, V., Erdman, K. A., & Reimer, R. A. (2009). Evaluation of nutritional intake in Canadian high-performance athletes. *Clinical Journal of Sports Medicine*, 19, 405–411.
- Mangili, A., Murman, D. H., Zampini, A. M., & Wanke, C. A. (2006). Nutrition and HIV infection: Review of weight loss and wasting in the era of highly active antiretroviral therapy from the Nutrition for Healthy Living cohort. *Clinical Infectious Diseases*, 42, 836–842.
- Mann, J., Cummings, J. H., Englyst, H. N., Key, T., Liu, S., Riccardi, G., et al. (2007). FAO/WHO scientific update on carbohydrates in human nutrition: Conclusions. *European Journal of Clinical Nutrition*, 61 (suppl. 1), S132–S137.
- Martins, C., Morgan, L. M., Bloom, S. R., & Robertson, M. D. (2007). Effects of exercise on gut peptides, energy intake and appetite. *Journal of Endocrinology*, 193, 251–258.
- Martinsen, M., Bratland-Sanda, S., Eriksson, A. K., & Sundgot-Borgen, J. (2010). Dieting to win or to be thin? A study of dieting and disordered eating among adolescent elite athletes and non-athlete controls. *British Journal of Sports Medicine*, 44, 70–76.
- Muller, W. (2009). Determinants of ski-jump performance and implications for health, safety and fairness. *Sports Medicine*, 39, 85–106.
- Muller, W., Groschl, W., Muller, R., & Sudi, K. (2006). Underweight in ski jumping: The solution of the problem. *International Journal of Sports Medicine*, 27, 926–934.
- Nattiv, A., Loucks, A. B., Manore, M. M., Sundgot-Borgen, J., & Warren, M. P. (2007). American College of Sports Medicine Position Stand: The female athlete triad. *Medicine and Science in Sports and Exercise*, 39, 1867–1882.
- Onywera, V. O., Kiplamai, F. K., Boit, M. K., & Pitsiladis, Y. P. (2004). Food and macronutrient intake of elite Kenyan distance runners. *International Journal of Sport Nutrition and Exercise Metabolism*, 14, 709–719.
- Oppliger, R. A., Utter, A. C., Scott, J. R., Dick, R. W., & Klossner, D. (2006). NCAA rule change improves weight loss among national championship wrestlers. *Medicine and Science in Sports and Exercise*, 38, 963–970.
- Pollock, N., Grogan, C., Perry, M., Pedlar, C., Cooke, K., Morrissey, D., et al. (2010). Bone-mineral density and other features of the female athlete triad in elite endurance runners: A longitudinal and cross-sectional observational study. *International Journal of Sport Nutrition and Exercise Metabolism*, 20, 418–426.
- Rodriguez, N. R., DiMarco, N. M., & Langley, S. (2009). Position of the American Dietetic Association, Dietitians of Canada, and the American College of Sports Medicine: Nutrition and athletic performance. *Journal of the American Dietetic Association*, 109, 509–527.
- Russel, R., Willis, K. S., Ravussin, E., & Larson-Meyer, E. D. (2009). Effects of endurance running and dietary fat on circulating ghrelin and peptide YY. *Journal of Sports Science and Medicine*, 8, 574–583.
- Sangenis, P., Drinkwater, B. L., Loucks, A. B., Sherman, R. T., Sundgot-Borgen, J., & Thompson, R. A. (2005). International Olympic Committee Medical Commission Working Group on Women in Sport: Position Stand on the Female Athlete Triad. Retrieved from: http://multimedia.olympic.org/pdf/en_report_917.pdf
- Sangenis, P., Drinkwater, B. L., Mountjoy, M., Constantini, N., Schamasch, P., Elwani, R., et al. (2006). IOC Consensus Statement: The Female Athlete Triad. *Medicine and Science in Tennis*, 11, 20.
- Schtscherbyna, A., Soares, E. A., de Oliveira, F. P., & Ribeiro, B. G. (2009). Female athlete triad in elite swimmers of the city of Rio de Janeiro, Brazil. *Nutrition*, 25, 634–639.
- Sherman, R., & Thompson, R. (2005). *Managing the female athlete triad: NCAA coaches handbook*. Indianapolis, IN: National Collegiate Athletic Association.
- Staten, M. A. (1991). The effect of exercise on food intake in men and women. *American Journal of Clinical Nutrition*, 53, 27–31.
- Stubbs, R. J., Hughes, D. A., Johnstone, A. M., Whybrow, S., Horgan, G. W., King, N., et al. (2004). Rate and extent of compensatory changes in energy intake and expenditure in response to altered exercise and diet composition in humans. *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology*, 286, R350–R358.
- Stubbs, R. J., Sepp, A., Hughes, D. A., Johnstone, A. M., Horgan, G. W., King, N., et al. (2002a). The effect of graded levels of exercise on energy intake and balance in free-living men, consuming their normal diet. *European Journal of Clinical Nutrition*, 56, 129–140.
- Stubbs, R. J., Sepp, A., Hughes, D. A., Johnstone, A. M., King, N., Horgan, G. W., et al. (2002b). The effect of graded levels of exercise on energy intake and balance in free-living women. *International Journal of Obesity and Related Metabolic Disorders*, 26, 866–869.
- Tarnopolsky, M. A., Atkinson, S. A., Phillips, S. M., & MacDougall, J. D. (1995). Carbohydrate loading and metabolism during exercise in men and women. *Journal of Applied Physiology*, 78, 1360–1368.
- Tarnopolsky, M. A., Zawada, C., Richmond, L. B., Carter, S., Shearer, J., Graham, T., et al. (2001). Gender differences in carbohydrate loading are related to energy intake. *Journal of Applied Physiology*, 91, 225–230.
- Ueda, S. Y., Yoshikawa, T., Katsura, Y., Usui, T., & Fujimoto, S. (2009). Comparable effects of moderate intensity exercise on changes in anorectic gut hormone levels and energy intake to high intensity exercise. *Journal of Endocrinology*, 203, 357–364.
- Vogt, S., Heinrich, L., Schumacher, Y. O., Grosshauser, M., Blum, A., Konig, D., et al. (2005). Energy intake and energy expenditure of elite cyclists during pre-season training. *International Journal of Sports Medicine*, 26, 701–706.
- Wade, G. N., & Jones, J. E. (2004). Neuroendocrinology of nutritional infertility. *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology*, 287, R1277–R1296.
- Wardle, J., Haase, A. M., & Steptoe, A. (2006). Body image and weight control in young adults: International comparisons in university students from 22 countries. *International Journal of Obesity (London)*, 30, 644–651.
- Westertorp, K. (2003). Energy metabolism and body composition: General principles. *European Respiratory Monograph*, 24, 1–10.
- Wren, A. M., Seal, L. J., Cohen, M. A., Brynes, A. E., Frost, G. S., Murphy, K. G., et al. (2001). Ghrelin enhances appetite and increases food intake in humans. *Journal of Clinical Endocrinology and Metabolism*, 86, 5992–5995.
- Ziegler, P., Nelson, J. A., Barratt-Fornell, A., Fiveash, L., & Drewnowski, A. (2001). Energy and macronutrient intakes of elite figure skaters. *Journal of the American Dietetic Association*, 101, 319–325.