

A Novel Cytosolic Isoform of Mitochondrial Trans-2-Enoyl-CoA Reductase Enhances Peroxisome Proliferator-Activated Receptor α Activity

Dong-Gyu Kim¹, Jae Cheal Yoo¹, Eunju Kim², Young-Sun Lee¹, Oleg V. Yarishkin², Da Yong Lee², Kun Ho Lee³, Seong-Geun Hong¹, Eun Mi Hwang², Jae-Yong Park^{1,2}

¹Department of Physiology, Institute of Health Science, Medical Research Center for Neural Dysfunction, Gyeongsang National University School of Medicine, Jinju; ²Center for Functional Connectomics, Korea Institute of Science and Technology, Seoul; ³Department of Marine Life Science, Chosun University College of Natural Sciences, Gwangju, Korea

Background: Mitochondrial trans-2-enoyl-CoA reductase (MECR) is involved in mitochondrial synthesis of fatty acids and is highly expressed in mitochondria. MECR is also known as nuclear receptor binding factor-1, which was originally reported with yeast two-hybrid screening as a binding protein of the nuclear hormone receptor peroxisome proliferator-activated receptor α (PPAR α). However, MECR and PPAR α are localized at different compartment, mitochondria, and the nucleus, respectively. Therefore, the presence of a cytosolic or nuclear isoform of MECR is necessary for functional interaction between MECR and PPAR α .

Methods: To identify the expression pattern of MECR and the cytosolic form of MECR (cMECR), we performed reverse transcription polymerase chain reaction (RT-PCR) with various tissue samples from Sprague-Dawley rats. To confirm the interaction between cMECR and PPAR α , we performed several binding assays such as yeast two-hybrid, coimmunoprecipitation, and bimolecular fluorescence complementation. To observe subcellular localization of these proteins, immunocytochemistry was performed. A luciferase assay was used to measure PPAR α activity.

Results: We provide evidence of an alternatively spliced variant of the rat MECR gene that yields cMECR. The cMECR lacks the N-terminal 76 amino acids of MECR and shows uniform distribution in the cytoplasm and nucleus of HeLa cells. cMECR directly bound PPAR α in the nucleus and increased PPAR α -dependent luciferase activity in HeLa cells.

Conclusion: We found the cytosolic form of MECR (cMECR) was expressed in the cytosolic and/or nuclear region, directly binds with PPAR α , and enhances PPAR α activity.

Keywords: Trans-2-enoyl-CoA reductase (NADPH); Cytosolic mitochondrial trans-2-enoyl-CoA reductase; PPAR alpha; Alternative splicing; Mitochondrial targeting sequences

INTRODUCTION

Mammalian mitochondria perform nicotinamide adenine di-

nucleotide phosphate (NADPH)-dependent *de novo* fatty acid synthesis (FAS) [1]. This mitochondrial FAS pathway resembles the well-understood bacterial FAS II pathway [2]. Mito-

Received: 13 August 2013, Accepted: 11 October 2013

Corresponding author: Jae-Yong Park

Department of Physiology, Gyeongsang National University School of Medicine, 15 Jinju-daero 816beon-gil, Jinju 660-751, Korea

Tel: +82-55-772-8043, Fax: +82-55-772-8049, E-mail: jaeyong@gnu.ac.kr

Copyright © 2014 Korean Endocrine Society

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

chondrial trans-2-enoyl-CoA reductase (MECR) is a component of the mitochondrial FAS II pathway and catalyzes the fatty acid elongation cycle in the last step, which is the NADPH-dependent reduction of the enoyl-acyl carrier protein substrate [3-5]. MECR is primarily localized in mitochondria [6-8], and the N-terminal amino acids of this protein are important for its mitochondrial localization [9]. MECR is also known as nuclear receptor binding factor-1 (NRBF-1) because, using yeast two-hybrid screening, it was identified as a binding protein of the nuclear hormone receptor peroxisome proliferator-activated receptor α (PPAR α) [10]. However, the biological meaning of the interaction between MECR and PPAR α has not been determined.

PPAR α is a ligand-activated transcription factor that is one of three different PPAR subtypes: PPAR α , PPAR β/δ , and PPAR γ . The PPARs play important roles in nutrient homeostasis [11-13] and are localized in the nucleus. Although MECR was previously reported as a binding protein of PPAR α [10], interaction between MECR and PPAR α seems not to occur in mammalian cells due to their different subcellular localizations, mitochondria and the nucleus, respectively. Therefore, the presence of a cytosolic or nuclear isoform of MECR is necessary for functional interaction between MECR and PPAR α in the nuclei of cells.

Here, we analyzed the expression pattern of MECR in several rat tissues and found a novel splice variant of MECR in which an additional exon was inserted between exon 1 and exon 2. The protein generated from this splicing variant has an N-terminal region that does not contain the mitochondrial targeting signal peptide and thus is not localized in mitochondria. Moreover, this MECR variant bound PPAR α in the nucleus and enhanced PPAR α transcriptional activity. Based on these results, we propose that this novel variant of MECR, cytosolic MECR (cMECR), plays a role in intracellular signal pathways as an interacting partner of PPAR α .

METHODS

RNA extraction and reverse transcription polymerase chain reaction

Total RNA was extracted from tissues of male Sprague-Dawley rats (14 weeks old) using TRIzol reagent (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's protocol. Total RNA (1 μ g) from each sample was reverse transcribed using random primers (50 pmol), SuperScript III Reverse Transcriptase (Invitrogen), and dNTPs (1 mM) at 42°C for 1 hour. The forward primer (5'-ATGTTGGTCAGCCGGC-

GACT-3') and reverse primer (5'-TCACATAGTGAGAATCTGCT-3') were designed to amplify full-length MECR. Additional primers were used for specific detection of MECR and cMECR cDNA: forward primer (5'-GTGCTGGAAGCGGCATGTTG-3') and reverse primer (5'-TGAGCTCCAGGTTCTTCAGT-3'). cMECR and PPAR α fragments were amplified under the following cycle conditions: denaturation at 94°C for 30 seconds, annealing at 55°C for 30 seconds, and extension at 72°C for 30 seconds. This cycle was repeated 35 times. Glyceraldehyde 3-phosphate dehydrogenase fragments were also amplified under the same conditions except that 25 cycles were run. Polymerase chain reaction (PCR) products were analyzed with 2.0% agarose gel electrophoresis, purified, and ligated into the pGEM-T Easy vector (Promega, Madison, WI, USA). The recombinant plasmids were sequenced.

Construction of plasmids and antibodies

The rat full-length MECR (GenBank accession no. AB015724), cytosolic variant of MECR, and PPAR α (GenBank accession no. NM_001001928) were cloned using the Gateway Cloning System (Invitrogen) as previously described [14]. After amplification of the three genes, PCR products were cloned into the pDONR207 vector and then subcloned into self-constructed destination vectors such as pDS_XB-HA, pDs_XB-Flag, pDS_XB-enhanced green fluorescent protein (EGFP), and pDS_XB-mCherry expression vectors. Anti-Flag (F6531) antibody was purchased from Sigma (St. Louis, MO, USA). Anti-EGFP (B-2) and antihemagglutinin (anti-HA) (F-7) antibodies were obtained from Santa Cruz Biotechnology (Santa Cruz, CA, USA).

Cell culture and transfection

HeLa cells were maintained in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum (Invitrogen) and 1% penicillin-streptomycin in a humid atmosphere containing 5% CO₂ at 37°C. When needed, cells were seeded onto cover slips for imaging analysis or 60-mm dishes for preparation of lysates for Western blot analysis. Transfection of expression vectors was performed with lipofectamine 2000 reagent (Invitrogen) according to the manufacturer's protocol. Transfected cells were cultured for an additional 24 hours in growth medium and then used for further analysis.

Coimmunoprecipitation and Western blot analysis

Transfected cells were lysed with radioimmunoprecipitation assay buffer (50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 5 mM ethylenediaminetetraacetic acid, 1 mM phenylmethanesulfo-

nyl fluoride, and 1% NP-40) containing a protease inhibitor cocktail (Sigma). Whole-cell lysates were incubated on ice for 30 minutes and then cleared at 13,000 rpm for 20 minutes at 4°C. The immune complexes were incubated for 1 hour at 4°C with gentle rotation with 20 μ L protein A/G PLUS-agarose beads (Santa Cruz Biotechnology) that had been prewashed and suspended in 100 mL cold lysis buffer. The samples were washed three times in 1 mL cold lysis buffer and eluted in 20 μ L sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) sample buffer. The proteins were separated with 10% SDS-PAGE and blotted onto polyvinylidene fluoride membranes. The blots were blocked with 5% skim milk in Tris-buffered saline with Tween 20 (20 mM Tris-buffered saline and 0.05% Tween 20, pH 7.5) at room temperature for 20 minutes and incubated overnight at 4°C with anti-GFP antibody (1:1,000), anti-HA antibody (1:1,000), or anti-FLAG antibody (1:1,000). Blots were then washed and incubated with horseradish peroxidase-conjugated secondary antibody (1:3,000), followed by washing and detection of immunoreactivity with enhanced chemiluminescence (Amersham, Piscataway, NJ, USA).

Imaging analysis

Subcellular distribution of MECR, cMECR, and PPAR α were confirmed according to the EGFP and mCherry fluorescence detection method with a confocal microscope (Olympus Fluoview FV1000, Olympus, Tokyo, Japan). For imaging analysis, cells were plated onto glass cover slips. After transfection, the cells were grown for 24 hours and treated with the mitochondrion-selective fluorescent dye mito-tracker red CMXRos (Molecular Probes Europe BV, Leiden, Netherlands) according to the manufacturer's instructions.

Yeast two-hybrid assay

cMECR was cloned into pGADT7 encoding the activation domain, and PPAR α was ligated into pGBKT7 encoding the GAL4 DNA binding domain. To evaluate the protein-protein interaction between cMECR and PPAR α , both pGAD-cMECR and pGBK-PPAR α were cotransformed into the yeast strain AH109. This strain is unable to synthesize histidine. However, interaction between cMECR and PPAR α enables the yeast to make the His3 enzyme, thereby permitting histidine biosynthesis and growth on His minimal medium.

Bimolecular fluorescence complementation assays

C- and N-terminal venus plasmids were purchased from Add-

gene and modified to contain cMECR or PPAR α with sub-cloning. Cells were transfected with the lipofectamine 2000 reagent (Invitrogen) with venus plasmids containing cMECR and PPAR α . After 24 hours, cells were imaged with confocal microscopy as described above.


Luciferase assay

HeLa cells were plated at a density of 1.0 to 1.5 \times 10⁵ cells per well onto six-well dishes. Twenty-four hours later, cells were cotransfected with 1 μ g of PPAR responsive element (PPRE)-3x-TK-Luc, the firefly luciferase reporter gene containing PPRE; 0.2 μ g of the PPAR α expression vector; and either the MECR expression vector or cMECR expression vector. Transfections were accomplished with lipofectamine 2,000 reagent according to the manufacturer's recommendations. The PPRE-3x-TK-Luc plasmid includes three PPREs upstream of an inducible thymidine kinase promoter controlling transcription of the firefly luciferase gene. Cells were harvested 24 hours after the start of exposure to vehicle or inducing agent. Luciferase assays were performed using the Luciferase Assay System (Promega) according to the manufacturer's protocol.

RESULTS

Identification of a novel alternative transcript variant of MECR

During reverse transcription (RT)-PCR analysis of MECR expression in the rat brain, two different MECR transcripts were detected. One was the expected MECR mRNA (210 bp), which corresponds to the known MECR mRNA (GenBank accession no. AB015724). The other was a longer 306-bp PCR fragment (Fig. 1C). Because PCR was carried out with specific primers corresponding to exons 1 and 2 of rat MECR cDNA, alternative splicing was proposed to account for the presence of this fragment. Fig. 1A shows the 10 exons of rat MECR cDNA. The novel MECR variant has an additional 96 nucleotides between exon 1 and exon 2. Basic local alignment search tool analysis of the aligned sequences of rat genomic DNA (chromosome 5 genomic contig NW_047784) and cDNA of this MECR variant showed that the flanking nucleotides of convergent points between exon 1 and exon 2 followed the GT/AG rule of alternative splicing (Supplementary Fig. 1A). This addition of 96 nucleotides generated a frame-shifted MECR, and the N-terminal 76 amino acids were deleted by use of another start codon in exon 3 (Fig. 1B). Because this splice variant lacked the N-terminal mitochondrial targeting sequences, we



Supplementary Fig. 1. (A) Nucleotide sequence showing exon/intron junctions for the generation of cytosolic form of mitochondrial trans-2-enoyl-CoA reductase (cMECR): all exon-intron boundaries conform to consensus splice-donor and splice-acceptor sites (GT-AG). (B) Amino acid alignment of rat cMECR with human MECR isoform 2: the amino acid sequences of both proteins were aligned using the Uniprot alignment program. Sequence identity is 87.2%. Asterisks mean same amino acid.

named it cMECR.


Next, to compare expression of cMECR and MECR, RT-PCR analyses were done using RNA samples from eight different tissues with a specific primer set as shown in Fig. 1B. As shown in Fig. 1C, cMECR (306 bp) and MECR (210 bp) were ubiquitously expressed in most tissues examined, although the expression level of cMECR was lower than that of MECR. Furthermore, we found amino acid sequences of human MECR isoform 2 (GenBank accession no. NP001019903.1) were well matched with that of rat cMECR (Supplementary Fig. 1B). These data raise the possibility that cMECR can be produced in other species.

cMECR localized in the cytoplasm and nucleus of HeLa cells

To visualize the intracellular distribution of MECR in HEK cells, we generated two constructs containing EGFP-tagged full-length rat MECR. As shown in Supplementary Fig. 2, when EGFP was tagged to the C-terminal end of MECR (MECR-EGFP), this protein was distinctly expressed in mito-

chondria, as previously reported [3]. However, the N-terminal EGFP-tagged MECR (EGFP-MECR) was diffusely expressed in the cytoplasm of HEK cells. N-terminal tagging of the protein with EGFP might have disrupted the proper mitochondrial localization of MECR, perhaps because the N-terminal mitochondrial targeting sequence of MECR was masked by the EGFP. These results suggest that N-terminal mitochondrial targeting sequences are critical for mitochondrial localization of MECR proteins.

Because cMECR lacks the N-terminal mitochondrial targeting sequences, the intracellular distribution of this protein was unclear. The intracellular distribution of MECR and cMECR was thus compared by tagging both cDNAs with EGFP at their C-terminal ends and then transiently expressing them in HeLa cells (Fig. 2). MECR-EGFP showed a well-defined punctate pattern that clearly colocalized with mito tracker, a marker of mitochondria (Fig. 2B). However, cMECR-EGFP was expressed in cytosolic and nuclear regions and did not colocalize with Mito tracker (Fig. 2C).



Supplementary Fig. 2. (A) Schematic drawing of the effect of enhanced green fluorescent protein (EGFP) tags on the localization of mitochondrial trans-2-enoyl-CoA reductase (MECR). (B) N-terminal EGFP-tagged MECR is mostly primarily located in cytosolic regions. C-terminal EGFP-tagged MECR localizes to mitochondrial regions. HeLa cells were transiently transfected with EGFP-MECR or MECR-EGFP. Scale bar=20 μ m.




Fig. 2. Subcellular localization of mitochondrial trans-2-enoyl-CoA reductase (MECR) and cytosolic form of MECC (cMECR) in HeLa cells. (A) The two constructs used in this study: MECR and cMECR were tagged at the C-terminus with enhanced green fluorescent protein (EGFP; green oval). The black box in MECR indicates the mitochondrial targeting domain. HeLa cells were transiently transfected with (B) MECR-EGFP or (C) cMECR-EGFP and then stained with Mito Tracker, a mitochondrial marker (red). Scale bar=20 μ m.

cMECR directly interacts with PPAR α

MECR, also known as NRBF-1, was previously identified with yeast two-hybrid screening as a PPAR α -interacting protein [10], although explaining how mitochondrial-targeted MECR interacted with the nuclear hormone receptor PPAR α was difficult. In contrast to MECR, cMECR was localized in the cytoplasm and nucleus of HeLa cells, and its amino acid sequence is exactly the same as that of MECR, except for the N-terminal 76 amino acids that compose the mitochondrial targeting sequence. Thus, cMECR, rather than mitochondrial MECR, could interact with PPAR α in the nucleus. Indeed, the yeast two-hybrid assay showed a direct interaction between cMECR and PPAR α (Fig. 3A). Immunoprecipitation and colocalization analyses also showed that cMECR interacted with PPAR α in HeLa cells (Fig. 3B). Furthermore, cMECR-EGFP colocalized with mCherry-tagged PPAR α in the nucleus of HeLa cells (Fig. 3C). In contrast, wild-type MECR was not colocalized with PPAR α in HeLa cells (Supplementary Fig. 3). Taken together, these data strongly indicate that cMECR, but not MECR, is a bona fide binding partner of PPAR α .

cMECR binds with PPAR α in the nucleus and potentiates PPAR α activity

To more clearly investigate whether the interaction between cMECR and PPAR α can occur in the nucleus of HeLa cells, we performed a bimolecular fluorescence complementation (BiFC) experiment. The BiFC assay provides a direct approach for the visualization of molecular interactions in living





Fig. 3. Cytosolic form of mitochondrial trans-2-enoyl-CoA reductase (cMECR) directly interacts with peroxisome proliferator-activated receptor α (PPAR α) *in vitro*. (A) Yeast two-hybrid assay between cMECR and PPAR α : on a Trp-/Leu-/His- plate, colonies indicate an interaction between the two genes. (B) Coimmunoprecipitation assays with cMECR and PPAR α : cMECR-FLAG was cotransfected with hemagglutinin (HA)-PPAR α in HeLa cells. The lysates were immunoprecipitated with anti-FLAG antibody and then immunoblotted with anti-HA antibody. (C) Colocalization of cMECR and PPAR α : HeLa cells were transiently transfected with cMECR-enhanced green fluorescent protein (EGFP; green) and mCherry-PPAR α (red). Scale bar = 20 μ m.



Supplementary Fig. 3. Localization of mitochondrial trans-2-enoyl-CoA reductase (MECR) and peroxisome proliferator-activated receptor α (PPAR α): wild-type MECR is localized in cytosolic regions, whereas PPAR α is localized in nuclear regions. HeLa cells were transiently transfected with MECR-enhanced green fluorescent protein (EGFP; green) and mCherry-PPAR α (red). Scale bar = 20 μ m.

cells [15,16]. We constructed the N-terminal part of venus plasmids tagged to cMECR (cMECR-VN) and the C-terminal portion of venus plasmids tagged to PPAR α (VC-PPAR α) (Fig. 4A). Consistent with the immunoprecipitation and colo-

calization analyses, the BiFC experiment also indicated protein-protein interaction between cMECR and PPAR α (Fig. 4B). Only when both constructs were cotransfected was the BiFC signal seen in the nucleus of HeLa cells. Individual con-




Fig. 4. Cytosolic form of mitochondrial trans-2-enoyl-CoA reductase (cMECR) potentiates peroxisome proliferator-activated receptor α (PPAR α) activity. (A) Schematic diagram to explain the use of the venus fusion proteins for bimolecular fluorescence complementation (BiFC). (B) BiFC of cMECR with PPAR α : plasmids containing cMECR-N terminal fragment of venus (VN) and C terminal fragment of venus (VC)-PPAR α were cotransfected into HeLa cells. These cells were stained with 4',6-diamidino-2-phenylindole (DAPI). Venus fluorescence is pseudocolored green. Scale bar=20 mm. (C) Luciferase assay of PPAR α : PPAR α activity was measured in HeLa cells expressing the PPAR-Luciferase reporter plasmid, HA-PPAR α , and cMECR-FLAG. All values are mean \pm s.e.m. *P* values were obtained with Student's *t* test. ^a*P*<0.01.

structs (cMECR-VN or VC-PPAR α) did not produce fluorescence alone (Fig. 4B). These results imply that cMECR interacts with PPAR α in the nucleus of HeLa cells.

cMECR interacts with PPAR α , a well-known nuclear hormone receptor. Thus, to examine whether cMECR can modulate PPAR α activity, we transfected cMECR into HeLa cells together with PPAR α and a luciferase reporter plasmid containing a PPAR-binding site (Fig. 4C). Compared with cells transfected with PPAR α only, PPAR-dependent luciferase expression levels were significantly increased in the presence of cMECR. These results indicate that cMECR can act as a co-activator of PPAR α in the nucleus of HeLa cells.

DISCUSSION

MECR has been reported to be a mitochondrial protein [7] that catalyzes NADPH-dependent reduction of trans-2-enoyl thioesters to the corresponding saturated acyl thioesters [5]. However, NRBF-1 (also known as MECR) was originally identified as a binding protein of the nuclear receptor PPAR α [10]. In that paper, the authors did not perform any assessment of endogenous binding in cells or report the biological meaning of this protein-protein interaction, although NRBF-1 binds with PPAR α in the yeast two-hybrid system.

Here, we report a cMECR that is generated by alternative splicing. cMECR is an N-terminal-76-amino-acid truncated

isoform of MECR. Because mitochondrial targeting signal sequences are contained between amino acids 1 and 53 of MECR [8], cMECR lacks this mitochondrial targeting signal sequence and was preferentially localizes to cytosolic and nuclear regions and was not present in mitochondria (Fig. 2C). In addition, our data show that N-terminal EGFP-tagged MECR also lost its proper intracellular localization (Supplementary Fig. 2). Therefore, the N-terminal region of MECR is critical for mitochondrial localization of the MECR protein.

Different intracellular distribution of cMECR strongly implies that its biological functions differ from the functions of MECR. In addition, cMECR mRNA was ubiquitously expressed in eight tested tissues (Fig. 1C), implying that cMECR could perform common essential functions in the various tissues. Although further studies should be performed to elucidate the function(s) of cMECR, we present its first function in this study. Various nuclear receptors activate transcription of their target genes through interactions with coactivators or corepressors [17,18]. Like other coactivators, cMECR directly bound PPAR α in the nucleus and potentiated PPAR α activity (Fig. 4).

Generally, PPAR α is found in tissues where fatty acid metabolism is important and regulates genes involved in lipid and lipoprotein metabolism as well as glucose homeostasis [19-21]. Because cMECR potentiates PPAR α activity in HeLa cells, cMECR could be involved in these biological phenomena. An earlier study used the yeast two-hybrid assay and demonstrated that MECR (NRBF-1) also interacts with various nuclear hormone receptors such as TR β , RAR α , RXR α , and HNF-4 [10]. Because our data strongly support the idea that cMECR is a bona fide binding partner of PPAR α and its amino acid sequence is the same as that of MECR except that it lacks the N-terminal 76 amino acids, cMECR might also bind with other nuclear hormone receptors. This possibility should be tested in future studies.

In summary, our results suggest that cMECR, a novel alternatively spliced variant of MECR, directly interacts with PPAR α in the nucleus and could be a positive regulator in the physiological regulation of PPAR α .

CONFLICTS OF INTEREST

No potential conflict of interest relevant to this article was reported.

ACKNOWLEDGMENTS

This research was supported by the Basic Research Promotion Fund (NRF-2008-313-C00758).

REFERENCES

1. Chen Z, Leskinen H, Liimatta E, Sormunen RT, Miinalainen IJ, Hassinen IE, Hiltunen JK. Myocardial overexpression of Mecn, a gene of mitochondrial FAS II leads to cardiac dysfunction in mouse. *PLoS One* 2009;4:e5589.
2. White SW, Zheng J, Zhang YM, Rock. The structural biology of type II fatty acid biosynthesis. *Annu Rev Biochem* 2005;74:791-831.
3. Torkko JM, Koivuranta KT, Miinalainen IJ, Yagi AI, Schmitz W, Kastaniotis AJ, Airene TT, Gurvitz A, Hiltunen KJ. Candida tropicalis Etr1p and Saccharomyces cerevisiae Ybr026p (Mrf1'p), 2-enoyl thioester reductases essential for mitochondrial respiratory competence. *Mol Cell Biol* 2001;21:6243-53.
4. Chen ZJ, Pudas R, Sharma S, Smart OS, Juffer AH, Hiltunen JK, Wierenga RK, Haapalainen AM. Structural enzymological studies of 2-enoyl thioester reductase of the human mitochondrial FAS II pathway: new insights into its substrate recognition properties. *J Mol Biol* 2008;379:830-44.
5. Hoffmeister M, Piotrowski M, Nowitzki U, Martin W. Mitochondrial trans-2-enoyl-CoA reductase of wax ester fermentation from Euglena gracilis defines a new family of enzymes involved in lipid synthesis. *J Biol Chem* 2005;280:4329-38.
6. Pfanner N. Protein sorting: recognizing mitochondrial presequences. *Curr Biol* 2000;10:R412-5.
7. Emanuelsson O, Nielsen H, Brunak S, von Heijne G. Predicting subcellular localization of proteins based on their N-terminal amino acid sequence. *J Mol Biol* 2000;300:1005-16.
8. Claros MG, Vincens P. Computational method to predict mitochondrially imported proteins and their targeting sequences. *Eur J Biochem* 1996;241:779-86.
9. Chen JQ, Delannoy M, Cooke C, Yager JD. Mitochondrial localization of ERalpha and ERbeta in human MCF7 cells. *Am J Physiol Endocrinol Metab* 2004;286:E1011-22.
10. Masuda N, Yasumo H, Furusawa T, Tsukamoto T, Sadano H, Osumi T. Nuclear receptor binding factor-1 (NRBF-1), a protein interacting with a wide spectrum of nuclear hor-

- mone receptors. *Gene* 1998;221:225-33.
11. Desvergne B, Wahli W. Peroxisome proliferator-activated receptors: nuclear control of metabolism. *Endocr Rev* 1999; 20:649-88.
 12. Kersten S, Desvergne B, Wahli W. Roles of PPARs in health and disease. *Nature* 2000;405:421-4.
 13. Evans RM, Barish GD, Wang YX. PPARs and the complex journey to obesity. *Nat Med* 2004;10:355-61.
 14. Park JY, Hwang EM, Yarishkin O, Seo JH, Kim E, Yoo J, Yi GS, Kim DG, Park N, Ha CM, La JH, Kang D, Han J, Oh U, Hong SG. TRPM4b channel suppresses store-operated Ca²⁺ entry by a novel protein-protein interaction with the TRPC3 channel. *Biochem Biophys Res Commun* 2008; 368:677-83.
 15. Hu CD, Chinenov Y, Kerppola TK. Visualization of interactions among bZIP and Rel family proteins in living cells using bimolecular fluorescence complementation. *Mol Cell* 2002;9:789-98.
 16. Shyu YJ, Liu H, Deng X, Hu CD. Identification of new fluorescent protein fragments for bimolecular fluorescence complementation analysis under physiological conditions. *Biotechniques* 2006;40:61-6.
 17. Kallenberger BC, Love JD, Chatterjee VK, Schwabe JW. A dynamic mechanism of nuclear receptor activation and its perturbation in a human disease. *Nat Struct Biol* 2003; 10:136-40.
 18. McKenna NJ, Lanz RB, O'Malley BW. Nuclear receptor coregulators: cellular and molecular biology. *Endocr Rev* 1999;20:321-44.
 19. Plutzky J. The PPAR-RXR transcriptional complex in the vasculature: energy in the balance. *Circ Res* 2011;108: 1002-16.
 20. Ziouzenkova O, Plutzky J. Lipolytic PPAR activation: new insights into the intersection of triglycerides and inflammation? *Curr Opin Clin Nutr Metab Care* 2004;7:369-75.
 21. Han SH, Quon MJ, Koh KK. Beneficial vascular and metabolic effects of peroxisome proliferator-activated receptor-alpha activators. *Hypertension* 2005;46:1086-92.