

# Diabetes causes marked changes in function and metabolism of rat neutrophils

T C Alba-Loureiro, S M Hirabara, J R Mendonça, R Curi  
and T C Pithon-Curi<sup>1</sup>

Department of Physiology and Biophysics, Institute of Biomedical Sciences, University of São Paulo, Av. Prof. Lineu Prestes 1524, 05508-900 São Paulo SP, Brazil

<sup>1</sup>Institute of Health Sciences, University of São Judas Tadeu, Taquari, 546-Moooca, 03166-000 São Paulo SP, Brazil

(Requests for offprints should be addressed to T C Alba-Loureiro; Email: tatica@icb.usp.br)

## Abstract

Several studies have shown impairment of neutrophil function, a disorder that contributes to the high incidence of infections in diabetes. Since glucose and glutamine play a key role in neutrophil function, we investigated their metabolism in neutrophils obtained from the peritoneal cavity of streptozotocin-induced diabetic rats. The activities of hexokinase, glucose-6-phosphate dehydrogenase (G6PDH), phosphofructokinase (PFK), citrate synthase, phosphate-dependent glutaminase, NAD<sup>+</sup>-linked and NADP<sup>+</sup>-linked isocitrate dehydrogenase were assayed. Glucose, glutamine, lactate, glutamate and aspartate, and the decarboxylation of [U-<sup>14</sup>C], [1-<sup>14</sup>C] and [6-<sup>14</sup>C]glucose; [U-<sup>14</sup>C]palmitic acid; and [U-<sup>14</sup>C]glutamine were measured in 1-h incubated neutrophils. Phagocytosis capacity and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) production were also determined. All measurements were carried out in neutrophils from control, diabetic and

insulin-treated (2–4IU/day) diabetic rats. Phagocytosis and phorbol myristate acetate (PMA)-stimulated H<sub>2</sub>O<sub>2</sub> production were decreased in neutrophils from diabetic rats. The activities of G6PDH and glutaminase were decreased, whereas that of PFK was raised by the diabetic state. The activities of the remaining enzymes were not changed. Diabetes decreased the decarboxylation of [1-<sup>14</sup>C]glucose and [U-<sup>14</sup>C]glutamine; however, [6-<sup>14</sup>C]glucose and [U-<sup>14</sup>C]palmitic acid decarboxylation was increased. These observations indicate that changes in metabolism may play an important role in the impaired neutrophil function observed in diabetes. The treatment with insulin abolished the changes induced by the diabetic state even with no marked change in glycemia. Therefore, insulin may have a direct effect on neutrophil metabolism and function.

*Journal of Endocrinology* (2006) **188**, 295–303

## Introduction

Patients with diabetes mellitus have increased susceptibility to and severity of infections. Several studies have shown alterations in neutrophil function, an effect that contributes to the high incidence of infections in diabetic patients. The observed changes in neutrophils include impairment of the following: adhesion to endothelium and migration to the site of inflammation (Pereira *et al.* 1987), chemotaxis (Mowat & Baum 1971), bactericidal activity (Tan *et al.* 1974), phagocytosis (Nolan *et al.* 1978) and production of reactive oxygen species (ROS) (Sagone *et al.* 1983).

Neutrophil functions require energy (Mowat & Baum 1971, McMurray *et al.* 1990, Walrand *et al.* 2004), which is produced mainly by the metabolization of glucose to lactate (Beck & Valentine 1952, Borregaard & Herlin 1982). Only 2–3% of glucose is oxidized through the Krebs cycle in neutrophils (Beck 1958, Wood *et al.* 1963).

These cells also utilize glutamine at high rates, which is mainly converted to glutamate, aspartate, lactate and CO<sub>2</sub> (Pithon-Curi *et al.* 1997).

High levels of glucose and ketone bodies seem to influence neutrophil function through production of polyols (Wilson *et al.* 1986). Decreased rates of glycolysis and glycogen synthesis were observed in leukocytes of diabetic patients. These changes are abolished by *in vivo* insulin administration (Esmann 1983). Walrand *et al.* (2004) postulated that insulin may normalize neutrophil functions, not only by re-establishing the control of the intermediary metabolism, but also through a direct effect of the hormone on the cells. These authors postulated that changes in neutrophil metabolism might play a key role in the impaired function of these cells in diabetes.

In the present study, the metabolism of glucose and glutamine in neutrophils obtained from streptozotocin (STZ)-induced diabetic rats was investigated. Key enzyme activities of glycolysis (hexokinase and

phosphofructokinase), the pentose-phosphate pathway (glucose-6-phosphate dehydrogenase), the Krebs cycle (citrate synthase) and glutaminolysis (phosphate-dependent glutaminase) were determined. In addition, the activities of NAD-linked and NADP-linked isocitrate dehydrogenase were also determined. Decarboxylation of [ $U-^{14}C$ ], [ $1-^{14}C$ ] and [ $6-^{14}C$ ]glucose; [ $U-^{14}C$ ]palmitic acid; and [ $U-^{14}C$ ]glutamine was measured in incubated neutrophils. Glucose and glutamine utilization, and production of lactate, glutamate and aspartate by incubated neutrophils were determined. Phagocytosis and production of hydrogen peroxide were also evaluated. The measurements were carried out in neutrophils from control, diabetic and insulin-treated diabetic rats.

## Materials and Methods

### Animals

Male Wistar rats weighing  $200 \pm 20$  g (about 2 months of age) were obtained from the Department of Physiology and Biophysics, Institute of Biomedical Sciences, University of São Paulo. The rats were maintained at  $23 \pm 2$  °C under a cycle of 12-h light:12-h darkness, and were allowed free access to food and water. The animal ethical committee of the Institute of Biomedical Sciences approved the experimental procedure of this study.

### Induction of diabetes

The experimental type 1 diabetes was induced by intravenous injection of 65 mg/kg STZ dissolved in citrate buffer (pH 4.2). Control rats were injected with buffer only. At 48 h after STZ injection, the diabetic state was confirmed by blood glucose levels above 200 mg/dl estimated with the aid of a glucose meter (Roche). Blood samples were obtained from the cut tip of the animal's tail. One group of diabetic rats was treated with neutral protamine Hagedorn (NPH) insulin. The daily dose was 2–4 IU by unique s.c. injection at 0900–1000 h for 2 weeks. The NPH insulin treatment was not sufficient to restore the glucose blood levels to control values.

### Experimental procedure

After 2 weeks of diabetes induction, fed rats were killed by decapitation without anesthesia at 1100–1200 h. Neutrophils were obtained by intraperitoneal lavage with 30 ml PBS, 4 h after the intraperitoneal injection of 10 ml 1% (w/v) glycogen solution (Sigma type II, from oyster) in PBS. The cell suspension was centrifuged at 4 °C (500 g for 10 min). The number of viable cells (>95% neutrophils) was determined in a Neubauer chamber under an

optical microscope by Trypan blue exclusion (Pires de Melo *et al.* 1998, Pithon-Curi *et al.* 2002).

### Phagocytosis

Neutrophils ( $1 \times 10^6$ ) were incubated for 40 min at 37 °C in 1 ml RPMI 1640 medium with opsonized particles of zymosan. The particles ( $1 \times 10^7$ ) were opsonized by incubation in the presence of control rat serum for 30 min at 37 °C. The cells that were capable of phagocytizing three or more particles were determined in a Neubauer chamber under an optical microscope by the violet crystal coloring method. The percentage of phagocytosis was expressed by the number of cells that had three or more particles of zymosan per total number of cells counted.

### Hydrogen peroxide production

Hydrogen peroxide ( $H_2O_2$ ) production was measured by the method of Pick and Mizel (1981), which is based on horseradish peroxidase-dependent conversion of phenol red by  $H_2O_2$  to a colored compound. Briefly, the cells were incubated in the presence of 5 mM glucose and a solution of phenol red (0.5%) and horseradish peroxidase (5 mg/ml) (222 units/mg) at 37 °C for 1 h. The production of  $H_2O_2$  was measured at rest and after stimulation with phorbol myristate acetate (PMA) (20 nM). The reaction was terminated by addition of 10  $\mu$ l 1 M NaOH solution, and the amount of product formed was measured by spectrophotometry at 620 nm.

### Assay of the enzyme activities

The activities of hexokinase (E.C. 2.7.1.1), glucose-6-phosphate dehydrogenase (G6PDH) (E.C. 1.1.1.49), phosphofructokinase (PFK) (E.C. 2.7.1.11), citrate synthase (E.C. 4.1.3.7), phosphate-dependent glutaminase (E.C. 3.5.1.2), NAD<sup>+</sup>-linked isocitrate dehydrogenase (E.C. 1.1.1.41) and NADP<sup>+</sup>-linked isocitrate dehydrogenase (E.C. 1.1.1.42) were determined as previously described (Mansour 1963, Crabtree & Newsholme 1972, Curthoys & Lowry 1973, Bergmeyer & Bernt 1974, Sugden & Newsholme 1975, Alp *et al.* 1976). Enzyme activities were expressed as nmol of substrate utilized/min per mg protein. A similar procedure was used in our previous studies (Guimarães *et al.* 1993, Costa-Rosa *et al.* 1996, Otton *et al.* 2002).

### Cell incubation for measurement of consumption and production of metabolites

Neutrophils ( $1.0 \times 10^6$ ) were incubated for 1 h at 37 °C in PBS with glucose (5 mM) or glutamine (2 mM). After incubation, the cells were disrupted by the addition of 0.2 ml 25% (w/v) trichloroacetic acid solution. Protein was removed by centrifugation, and the supernatant fluid

was neutralized with a Tris (hydroxymethyl-amino-methane)/KOH (0.5–2.0 M) solution for the measurement of the metabolites.

#### Assays of the metabolites

Samples of the incubation medium were used for measurements of glucose (Berham & Trinder 1972), glutamine (Windmueller & Spaeth 1974), lactate (Engel & Jones 1978), glutamate (Bernt & Bergmeyer 1974) and aspartate (Bergmeyer *et al.* 1974).

#### Spectrophotometric conditions of the enzyme assays and metabolite measurements

Activities of hexokinase, G6PDH, citrate synthase and PFK were assayed at 25 °C and activity of glutaminase at 37 °C. Citrate synthase activity was assayed by following the rate of change in absorbance at 412 nm and the remaining enzymes at 340 nm. The final volume of the assay mixtures in all cases was 1.0 ml. The production of NADH or NADPH was monitored in a Pharmacia Biotech spectrophotometer (model: Ultraspec 3000).

#### Incubation procedure for determination of metabolite decarboxylation

Neutrophils ( $1 \times 10^7$ ) were incubated for 60 min at 37 °C in 1 ml PBS with the following labeled metabolites: [ $U$ - $^{14}C$ ], [ $1$ - $^{14}C$ ] or [ $6$ - $^{14}C$ ]glucose (5.0 mM and 0.2  $\mu$ Ci/ml); [ $U$ - $^{14}C$ ]palmitic acid (0.1 mM and 0.2  $\mu$ Ci/ml); or [ $U$ - $^{14}C$ ]glutamine (2.0 mM and 0.2  $\mu$ Ci/ml). Palmitic acid was previously dissolved in ethanol (20 mM) for addition to incubation medium. The incubation was stopped by adding 0.2 ml chloridric acid.  $^{14}CO_2$  produced from these metabolites was collected as previously described (Leighton *et al.* 1985) in a special apparatus containing phenylethylamine and methanol solution (1:1 v/v), and the radioactivity was measured in a Beckman-LS 5000TD scintillator (Beckman Instruments, Fullerton, CA, USA). A similar procedure was used in our previous studies (Otton *et al.* 2002, Hirabara *et al.* 2003). Another procedure was used to estimate the palmitic acid decarboxylation. The decrease of [ $U$ - $^{14}C$ ]palmitic acid from the medium and cell preparation after 1-h incubation was determined.

#### Protein determination

The total protein content of neutrophils was measured by the method of Bradford (1976), using BSA as standard.

#### Statistical analysis

All results are expressed as means  $\pm$  S.E.M. Student's *t*-test or ANOVA followed by the Tukey–Kramer test was used to

**Table 1** Body weight gain and blood glucose levels of the control, diabetic and insulin-treated diabetic groups

Groups	<i>n</i>	Body weight gain (g)	Blood glucose levels (mg/dL)
Control	15	52 $\pm$ 5 <sup>a</sup>	81 $\pm$ 4 <sup>a</sup>
Diabetic	10	25 $\pm$ 4 <sup>b</sup>	365 $\pm$ 25
Insulin-treated diabetic	11	44 $\pm$ 2	301 $\pm$ 13

Rats were rendered diabetic by streptozotocin injection (65 mg/kg, i.v.) and the measurements were carried out 2 weeks afterward. NPH insulin (2–4 IU/day) was subcutaneously administered during the experiment period. Values are presented as means  $\pm$  S.E.M. *n* indicates the number of rats used in each group. <sup>a</sup>*P*<0.001 as compared to diabetic and insulin-treated rats; <sup>b</sup>*P*<0.05 as compared to insulin-treated rats.

assess the significance of differences between groups. Data were considered as statistically significant at *P*  $\leq$  0.05.

## Results


Relative to controls, animals rendered diabetic by STZ injection exhibited a significant reduction in body weight gain during the experimental period (Table 1). Blood glucose levels were significantly elevated in comparison to controls (by 3.5-fold). The treatment of diabetic rats with NPH insulin (2–4 IU/day) did not restore the blood glucose levels to the values of the control group (Table 1).

Neutrophil counts were performed 4 h after the intraperitoneal injection of 10 ml 1% (w/v) glycogen solution. In all groups, a similar number of cells had migrated into the peritoneal cavity. The mean values ( $\times 10^7$ ;  $\pm$  S.E.M.) were as follows: 23.3  $\pm$  2.7 (control, *n*=11), 21.1  $\pm$  2.6 (diabetic, *n*=6) and 23.4  $\pm$  2.5 (insulin treated, *n*=7).

Phagocytosis and PMA-stimulated hydrogen peroxide production by neutrophils from the diabetic group were decreased (by 17% and 31% respectively) as compared with controls. The daily treatment with insulin restored phagocytosis and H<sub>2</sub>O<sub>2</sub> production to the values found in neutrophils from control rats (Fig. 1A and B).

G6PDH activity was decreased (by 59%) in neutrophils from diabetic rats as compared with controls. The diabetic state also decreased phosphate-dependent glutaminase activity (by 22%) and increased PFK activity (by 32%) (Fig. 2). The treatment with insulin abolished the effect of diabetes on G6PDH, glutaminase and PFK activities. Hexokinase, citrate synthase and NAD<sup>+</sup>-linked and NADP<sup>+</sup>-linked isocitrate dehydrogenase activities were not changed by diabetes (Fig. 2).

Glucose and glutamine utilization by incubated neutrophils was not markedly affected by the diabetic state. Lactate production was decreased (by 24%) in neutrophils from diabetic rats as compared with controls, but the production of glutamate and aspartate was not



**Figure 1** Phagocytosis capacity (A) and hydrogen peroxide production (B) by neutrophils. The values are presented as means  $\pm$  S.E.M. of 10 determinations from at least five animals in each group. <sup>a</sup> $P < 0.05$  as compared to control, <sup>b</sup> $P < 0.01$  as compared to insulin-treated diabetic rats and <sup>c</sup> $P < 0.05$  as compared to diabetic rats.

modified. The treatment of diabetic rats with insulin restored the production of lactate by neutrophils to the values of the control group (Table 2).


Diabetes decreased the decarboxylation of [1-<sup>14</sup>C]glucose (by 31%) and [U-<sup>14</sup>C]glutamine (by 22%) by 1-h incubated neutrophils. On the other hand, [6-<sup>14</sup>C]glucose (by 55%) and [U-<sup>14</sup>C]palmitic acid (by 83%) decarboxylation was higher in neutrophils obtained from diabetic rats. The treatment with insulin abolished the changes induced by diabetes (Fig. 3). Similar results for palmitic acid decarboxylation (data not shown) were obtained by measuring the decrease of [U-<sup>14</sup>C]palmitic acid in the medium and cell preparation; there was an increase of 61% ( $P < 0.01$ ) in neutrophils obtained from diabetic rats, and the treatment with insulin abolished the changes induced by the diabetic state. The decarboxylation of [U-<sup>14</sup>C]glucose was not affected by the diabetic state; glucose decarboxylation by neutrophils of the control and diabetic groups was  $14.9 \pm 0.6$  and  $14.0 \pm 1.1$  nmol/h per  $10^7$  cells (mean  $\pm$  S.E.M. of two determinations from six animals in each group) respectively.

## Discussion

Neutrophils constitute the first line of defense against bacterial and fungal infections. Phagocytosis stimulates the respiratory burst of neutrophils as a result of activation of nicotinamide adenine dinucleotide phosphate (NADPH) oxidase (Bellavite 1988). The superoxide anion ( $O_2^-$ ) is the first metabolite generated through reduction of molecular oxygen by NADPH oxidase (Tauber *et al.* 1985, Bellavite 1988). The  $O_2^-$  formed dismutates spontaneously or via superoxide dismutase to  $H_2O_2$ , which is converted to hypochlorous acid by myeloperoxidase released from neutrophil granules (Tauber *et al.* 1985).

Conflicting results have been reported about the production of  $H_2O_2$  by neutrophils in diabetes. In unstimulated neutrophils from diabetic patients, no significant effect on  $H_2O_2$  production was found (Noritake *et al.* 1992, Inoue *et al.* 1996), while Zozulinska *et al.* (1996) found it to be increased. After incubation with PMA, neutrophils from diabetic patients have shown low (Noritake *et al.* 1992, Inoue *et al.* 1996) or unchanged (Zozulinska *et al.* 1996) production of  $H_2O_2$  as compared with cells from healthy subjects. Neutrophils from diabetic patients with nephropathy have shown increased production of  $H_2O_2$  as compared with those without kidney complications (Watanabe 1992). In our study, neutrophils from diabetic rats showed low production of  $H_2O_2$  after PMA stimulation was abolished by treatment with insulin. There was no other study on  $H_2O_2$  production by rat neutrophils in diabetes. An increase in  $H_2O_2$  production by nonactivated neutrophils in the insulin-treated group was observed. Some studies have shown that insulin activates a plasma membrane enzyme system that presents properties of a Nox (NADPH oxidase enzymes). The enzyme system catalyzes the reduction of oxygen to superoxide ( $2 O_2^- + NADP^{++}H^+$ ), which is spontaneously or by action of superoxide dismutase converted to  $H_2O_2$  (Mukherjee *et al.* 1978, Rhee *et al.* 2003, Goldstein *et al.* 2005).

Impairment of phagocytosis and decreased release of lysosomal enzymes in neutrophils from diabetic patients have been observed by many researchers (Bybee & Rogers 1964, Bagdade *et al.* 1972, Sagone *et al.* 1983, Wilson & Reeves 1986). Studies with diabetic rats and mice also showed a decreased neutrophil phagocytosis capacity (Canturk *et al.* 1998, Panneerselvam & Govindasamy 2003). The lowering of blood glucose levels by insulin treatment has been reported to have significant correlation with the improvement of phagocytosis capacity by neutrophils (Jakelic *et al.* 1995, Delamaire *et al.* 1997).



**Figure 2** Enzyme activities of neutrophils. The results are expressed as nmol/min per mg protein, and the values are presented as means  $\pm$  S.E.M. of 10 determinations from five animals in each group. <sup>a</sup> $P < 0.01$  as compared to control, <sup>b</sup> $P < 0.05$  as compared to diabetic rats, <sup>c</sup> $P < 0.05$  as compared to control and <sup>d</sup> $P < 0.01$  as compared to diabetic rats.

In our study, the decreased capacity of neutrophils from diabetic rats to phagocytize zimosan particles was reversed by insulin treatment even under high blood glucose levels.

Previous studies on the effect of diabetes on glucose metabolism in neutrophils are controversial. Esmann (1972)

and Munroe & Shipp (1965) did not observe differences in glucose utilization by neutrophils from healthy and diabetic patients. Nevertheless, decreased utilization of glucose by neutrophils from diabetic patients was found by others (Martin *et al.* 1954, Esmann 1983).

**Table 2** Consumption and production of metabolites by 1-h incubated neutrophils from the control, diabetic and insulin-treated diabetic groups

Groups	n	Glucose consumption	Lactate production	Glutamine consumption	Glutamate	Production of aspartate	Lactate
Control	6	855 ± 170	945 ± 65	1110 ± 95	390 ± 5	515 ± 50	215 ± 5
Diabetic	5	755 ± 150	700 ± 65 <sup>a</sup>	1015 ± 70	380 ± 10	555 ± 65	230 ± 40
Insulin-treated diabetic	6	815 ± 150	935 ± 35	1115 ± 85	390 ± 5	510 ± 60	230 ± 35


Consumption of glucose and glutamine and production of lactate, glutamate and aspartate were determined in 1 h incubated neutrophils. The cells were obtained from the peritoneal cavity of the rats, and the results are expressed as nmol/h per mg protein. The values are presented as means ± S.E.M. n represents the number of rats per group. <sup>a</sup>*P*>0.05 as compared to control and insulin-treated diabetic rats.

Munroe & Shipp (1965) did not observe alteration, whereas Martin *et al.* (1954) and Esmann (1983) found decreased production of lactate by neutrophils from diabetic patients. In the present study, there was no alteration in glucose utilization by neutrophils from diabetic rats, whereas lactate production was decreased. These findings support the proposition that the diabetic state reduces glycolysis activity in neutrophils.

Neutrophils from diabetic rats had no significant change in glucose oxidation, citrate synthase, and NAD<sup>+</sup>-linked and NADP<sup>+</sup>-linked isocitrate dehydrogenase activities, suggesting that the flux of substrates through the Krebs cycle was not altered. Decreased citrate synthase

activity was observed in lymphocytes from diabetic rats (Otton *et al.* 2002), whereas no significant change was found in macrophages (Costa-Rosa *et al.* 1996). There was no study on glucose oxidation, or NAD<sup>+</sup>-linked and NADP<sup>+</sup>-linked isocitrate dehydrogenase activities by rat neutrophils in diabetes.

The pentose-phosphate pathway oxidizes glucose-6-phosphate to intermediates of the glycolytic pathway, generating NADPH and ribose-5-phosphate for fatty acid and nucleotide synthesis respectively (Casazza & Veech 1986). NADPH is important for NADPH oxidase activity and for glutathione reductase to recycle oxidized glutathione in neutrophils (Pithon-Curi *et al.* 1998, 1999).



**Figure 3** Decarboxylation of labeled metabolites by 1 h incubated neutrophils. The values are presented as means ± S.E.M. of eight determinations from four animals in each group. <sup>a</sup>*P*<0.01 as compared to control, <sup>b</sup>*P*<0.05 as compared to diabetic rats and <sup>c</sup>*P*<0.05 as compared to control and insulin treated diabetic rats.

Similarly, glutamine is also probably very important in neutrophils to provide glutamate for glutathione synthesis (Newsholme *et al.* 2003).

To examine the flux of glucose through the pentose-phosphate pathway of neutrophils from diabetic rats, G6PDH activity and decarboxylation of [1-<sup>14</sup>C] and [6-<sup>14</sup>C]glucose were determined. The difference between <sup>14</sup>CO<sub>2</sub> production from [1-<sup>14</sup>C] and [6-<sup>14</sup>C]glucose estimates the flux of glucose through the pentose-phosphate pathway (Larrabee 1989). The difference for neutrophils (10<sup>7</sup>; ± s.e.m.) from the diabetic group was 2.35 ± 0.2 nmol/h compared with 3.79 ± 0.15 nmol/h for the control group (decrease of 38%; *P* < 0.001). Insulin treatment abolished the difference between the decarboxylation of [1-<sup>14</sup>C] and [6-<sup>14</sup>C]glucose. The activity of G6PDH was decreased in neutrophils from diabetic rats, as was also found in macrophages and lymphocytes (Costa-Rosa *et al.* 1996, Otton *et al.* 2002). These findings support the proposition that the activity of the pentose-phosphate pathway is decreased in leukocytes from diabetic rats. Leukocytes with deficiency of G6PDH activity present impaired phagocytosis, bactericidal capacity and superoxide production (Gray *et al.* 1973, Roos *et al.* 1999). Moreover, decreased flux of glucose through the pentose-phosphate pathway is expected to reduce the production of NADPH and ribose-5-phosphate. Therefore, reduced pentose-phosphate pathway activity may be related to impaired neutrophil function in the diabetic state. Although the production of NADP-linked isocitrate dehydrogenase activity was not affected by the diabetic state, this may not be sufficient to compensate for the reduced flux of substrates in the pentose-phosphate pathway.

Although decreased lactate production was observed in neutrophils from diabetic rats, the PFK maximal activity was increased. Similar results were observed in mesenteric lymph nodes and thymus lymphocytes from diabetic rats (Moreno-Aurioles *et al.* 1996, Otton *et al.* 2002). The activity of PFK is stimulated by fructose 2,6-bisphosphate (Wegener & Krause 2002) and inhibited by ATP at low fructose 6-phosphate content, but not at high fructose 6-phosphate concentration (Mansour 1963). In this study, we did not observe changes in the glucose consumption and oxidation in neutrophils from STZ-induced diabetic rats, but lactate production and pentose-phosphate pathway activity were markedly reduced. Therefore, it is expected that the content of intermediates of glycolysis, such as fructose 6-phosphate and fructose 2,6-bisphosphate, are elevated in this condition. This may partially explain the increase in the PFK activity. In accordance with this proposition, Moreno-Aurioles *et al.* (1996) found increased fructose 2,6-bisphosphate content and PFK activity in neutrophils from STZ-induced diabetic rat.

The production of glutamate and aspartate was not altered in neutrophils from diabetic rats incubated in the presence of 2 mM glutamine. Nevertheless, glutamine

oxidation and glutaminase activity were significantly decreased in neutrophils from diabetic rats. Glutamine plays an important role in protein (as amino-acid source), lipid (through NAD(P)H production) and nucleotide synthesis (through purine and pyrimidine production), and in NADPH oxidase activity (Newsholme *et al.* 2003, Curi *et al.* 2005). Glutamine raises the *in vitro* bacterial killing activity and the rate of ROS production by neutrophils (Ogle *et al.* 1994, Pithon-Curi *et al.* 1998, 2002). Pithon-Curi *et al.* (2003) showed that glutamine has a protective effect on neutrophil apoptosis. Therefore, decreased glutamine utilization may contribute to the impaired function by increasing the occurrence of apoptosis in neutrophils from diabetic rats. This issue remains to be investigated.

Diabetes leads to an increase in plasma levels of free fatty acids and triacylglycerols (Boden 1999, Kelley & Mandarino 2000). We have previously demonstrated increased fatty acid oxidation in lymphocytes from diabetic rats (Otton *et al.* 2002). Herein, we observed that palmitic acid decarboxylation is also elevated in neutrophils from diabetic rats.

In summary, neutrophils from diabetic rats present impaired metabolism of glucose and glutamine. On the other hand, palmitic acid oxidation is increased, and this may compensate for the reduction in glucose and glutamine utilization for ATP production. These metabolic changes may be related to impaired functions of these cells, such as phagocytosis and hydrogen peroxide production. The changes in metabolism and function of neutrophils from diabetic rats are fully abolished by insulin treatment, even with no marked changes in glycemia. These findings show that insulin may have a direct effect on neutrophil metabolism and function.

## Acknowledgements

The authors are indebted to the constant assistance of E P Portioli and G de Souza. This research is supported by FAPESP, CNPq and CAPES. The authors declare that there is no conflict of interest that would prejudice the impartiality of this scientific work.

## References

- Alp AR, Newsholme EA & Zammit VA 1976 Activities of citrate synthase and NAD<sup>+</sup>-linked and NAD<sup>+</sup>-linked isocitrate dehydrogenase in muscle from vertebrates and invertebrates. *Biochemistry Journal* **154** 689–700.
- Bagdade JD, Nielson KL & Bulger RJ 1972 Reversible abnormalities in phagocytic function in poorly controlled diabetic patients. *American Journal of Medical Science* **263** 451–456.
- Barham D & Trinder P 1972 An improved colour reagent for the determination of blood glucose by the oxidase system. *Analyst* **97** 142–145.

- Beck WS 1958 Occurrence and control of the phosphogluconate oxidation pathway in normal and leukemic leukocytes. *Journal of Biological Chemistry* **232** 271–283.
- Beck WS & Valentine WN 1952 The aerobic metabolism of leukocytes in health and leukemia. I. Glycolysis and respiration. *Cancer Research* **12** 818–822.
- Bellavite P 1988 The superoxide-forming enzymatic system of phagocytes. *Free Radical Biology and Medicine* **4** 225–261.
- Bergmeyer HU & Bernt E 1974 D-Glucose determination with hexokinase and glucose-6-phosphate dehydrogenase. In *Methods of Enzymatic Analysis*, pp 1196–1201. Ed HU Bergmeyer. London: Academic Press.
- Bergmeyer HU, Bernt E, Mollering H & Pfeleiderer ER 1974 L-Aspartate and L-asparagine. In *Methods of Enzymatic Analysis*, pp 1696–1700. Ed HU Bergmeyer. London: Academic Press.
- Bernt E & Bergmeyer HU 1974 L-Glutamate UV-assay with glutamate dehydrogenase and NAD<sup>+</sup>. In *Methods of Enzymatic Analysis*, pp 1704–1704. Ed HU Bergmeyer. London: Academic Press.
- Boden G 1999 Free fatty acids, insulin resistance, and type 2 diabetes mellitus. *Proceedings of the Association of American Physicians* **111** 241–248.
- Borregaard N & Herlin T 1982 Energy metabolism of human neutrophils during phagocytosis. *Journal of Clinical Investigation* **70** 550–557.
- Bradford M 1976 A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* **72** 248–254.
- Bybee JD & Rogers DE 1964 The phagocytic activity of polymorphonuclear leukocytes obtained from patients with diabetes mellitus. *Journal of Laboratory and Clinical Medicine* **64** 1–13.
- Canturk Z, Canturk NZ & Onen F 1998 Effects of rhG-CSF on neutrophil functions and survival in sepsis induced diabetic rats. *Endocrine Research* **24** 141–157.
- Casazza JP & Veech RL 1986 The measurement of xylulose 5-phosphate, ribulose 5-phosphate, and combined sedoheptulose 7-phosphate and ribose 5-phosphate in liver tissue. *Analytical Biochemistry* **159** 243–248.
- Costa-Rosa LF, Safi DA, Cury Y & Curi R 1996 The effect of insulin on macrophage metabolism and function. *Cell Biochemistry and Function* **14** 33–42.
- Crabtree B & Newsholme EA 1972 The activities of phosphorylase, hexokinase, phosphofruktokinase, lactate dehydrogenase and the glycerol-3-phosphate dehydrogenase in muscle from vertebrates and invertebrates. *Biochemistry Journal* **126** 49–58.
- Curi R, Newsholme P, Pithon-Curi TC, Pires-de-Melo M, Garcia C, Homem-de-Bittencourt PI Jr & Guimarães ARP 1999 Metabolic fate of glutamine in lymphocytes, macrophages and neutrophils. *Brazilian Journal of Medical Biology and Research* **32** 15–21.
- Curi R, Lagranha CJ, Doi SQ, Sellitti F, Procopio J, Pithon-Curi TC, Corless M & Newsholme P 2005 Molecular mechanisms of glutamine action. *Journal of Cell Physiology* **204** 392–401.
- Curthoys NP & Lowry OH 1973 The distribution of glutaminase isoenzymes in the various structures of the nephron in normal, acidotic and alkalotic rat kidney. *Journal of Biological Chemistry* **248** 162–168.
- Delamare M, Maugendre D, Moreno M, Le Goff MC, Allanic H & Genetet B 1997 Impaired leukocyte function in diabetic patients. *Diabetic Medicine* **14** 29–34.
- Engel PC & Jones JB 1978 Causes and elimination of erratic blanks in enzymatic metabolite assays involving the use of NAD<sup>+</sup> in alkaline hydrazine buffers: improved conditions for the assay of L-glutamate, L-lactate and other metabolites. *Analytical Biochemistry* **88** 475–484.
- Esmann V 1972 The diabetic leukocyte. *Enzyme* **13** 32–55.
- Esmann V 1983 The polymorphonuclear leukocyte in diabetes mellitus. *Journal of Clinical Chemistry and Clinical Biochemistry* **21** 561–567.
- Goldstein BJ, Kalyankar M & Wu X 2005 Insulin action is facilitated by insulin-stimulated reactive oxygen species with multiple potential signaling targets. *Diabetes* **54** 311–321.
- Gray GR, Klebanoff SJ, Stamatoyannopoulos G, Austin T, Naiman SC, Yoshida A, Kliman MR & Robinson GCF 1973 Neutrophil dysfunction, chronic granulomatous disease, and non-spherocytic haemolytic anaemia caused by complete deficiency of glucose-6-phosphate dehydrogenase. *Lancet* **2** 530–534.
- Guimarães AR, Costa-Rosa LF, Safi DA & Curi R 1993 Effect of a polyunsaturated fatty acid-rich diet on macrophage and lymphocyte metabolism of diabetic rats. *Brazilian Journal of Medical Biology and Research* **26** 813–818.
- Hirabara SM, Carvalho CR, Mendonça JR, Harber EP, Fernandes LC & Curi R 2003 Palmitate acutely raises glycogen synthesis in rat soleus muscle by a mechanism that requires its metabolization (Randle cycle). *FEBS Letters* **541**(1–3) 109–114.
- Inoue S, Lan Y, Muran J & Tsuji M 1996 Reduced hydrogen peroxide production in neutrophils from patients with diabetes. *Diabetes Research and Clinical Practices* **33** 119–127.
- Jakelic J, Kokic S, Hozo I, Maras J & Fabijanic D 1995 Nonspecific immunity in diabetes: hyperglycemia decreases phagocytic activity of leukocytes in diabetic patients. *Medical Archives* **49** 9–12.
- Kelley DE & Mandarino LJ 2000 Fuel selection in human skeletal muscle in insulin resistance: a reexamination. *Diabetes* **49** 677–683.
- Larrabee MG 1989 The pentose cycle (hexose monophosphate shunt). Rigorous evaluation of limits to the flux from glucose using <sup>14</sup>CO<sub>2</sub> data, with applications to peripheral ganglia of chicken embryos. *Journal of Biological Chemistry* **264** 15875–15879.
- Leighton B, Challiss RA, Lozeman FJ & Newsholme EA 1985 The effect of prostaglandins E1, E2 and F2 alpha and indomethacin on the sensitivity of glycolysis and glycogen synthesis to insulin in stripped soleus muscles of the rat. *Biochemistry Journal* **227** 337–340.
- Mansour TE 1963 Studies on heart phosphofruktokinase: purification, inhibition and activation. *Journal of Biological Chemistry* **238** 2285–2292.
- Martin SP, Chaudhuri SN, Green R & McKinney GR 1954 The effect of adrenal steroids on aerobic lactic acid formation in human leukocytes. *Journal of Clinical Investigation* **33** 358–360.
- McMurray RW, RW Bradsher, RW Steele & NS Pilkington 1990 Effect of prolonged modified fasting in obese persons on in vitro markers of immunity: lymphocyte function and serum effects on normal neutrophils. *American Journal of Medical Science* **299** 379–385.
- Moreno-Aurioles VR, Montano R, Conde M, Bustos R & Sobrino F 1996 Streptozotocin-induced diabetes increases fructose 2,6-biphosphate levels and glucose metabolism in thymus lymphocytes. *Life Sciences* **58** 477–484.
- Mowat AG & Baum J 1971 Chemotaxis of polymorphonuclear leukocytes from patients with diabetes mellitus. *New England Journal of Medicine* **284** 621–627.
- Mukherjee SP, Lane RH & Lynn WS 1978 Endogenous hydrogen peroxide and peroxidative metabolism in adipocytes in response to insulin and sulphhydryl reagents. *Biochemical Pharmacology* **27** 2589–2594.
- Munroe JF & Shipp JC 1965 Glucose metabolism in leucocytes from patients with diabetes mellitus, with and without hypercholesteremia. *Diabetes* **14** 584–590.
- Newsholme P, Lima MM, Procopio J, Pithon-Curi TC, Doi SQ, Bazotte RB & Curi R 2003 Glutamine and glutamate as vital metabolites. *Brazilian Journal of Medical Biology and Research* **36** 153–163.
- Nolan CN, Beaty HN & Bagdade JD 1978 Further characterization of the impaired bactericidal function of granulocytes in patients with poorly controlled diabetes. *Diabetes* **27** 889–894.
- Noritake M, Katsura Y, Shinomiya N, Kanatani M, Uwabe Y, Nagata N & Tsuru S 1992 Intracellular hydrogen peroxide production by peripheral phagocytes from diabetic patients. Dissociation between polymorphonuclear leukocytes and monocytes. *Clinical and Experimental Immunology* **88** 269–274.

- Ogle CK, Ogle JD, Mao JX, Simon J, Noel JG, Li BG & Alexander JW 1994 Effect of glutamine on phagocytosis and bacterial killing by normal and pediatric burn patient neutrophils. *Journal of Parenteral and Enteral Nutrition* **18** 128–133.
- Otton R, Mendonça JR & Curi R 2002 Diabetes causes marked changes in lymphocyte metabolism. *Journal of Endocrinology* **174** 55–61.
- Panneerselvam S & Govindasamy S 2003 Sodium molybdate improves the phagocytic function in alloxan-induced diabetic rats. *Chemical and Biological Interactions* **145** 159–163.
- Pereira MAA, Sannomiya P & Garcia-Leme J 1987 Inhibition of leukocyte chemotaxis by factor in alloxan-induced diabetic rat plasma. *Diabetes* **36** 1307–1314.
- Pick E & Mizel D 1981 Rapid microassays for the measurement of superoxide and hydrogen peroxide production by macrophages in culture using an automatic enzyme immunoassay reader. *Journal of Immunological Methods* **46** 211–226.
- Pires de Melo M, Curi TC, Miyasaka CK, Palanch AC & Curi R 1998 Effect of indole acetic acid on oxygen metabolism in cultured rat neutrophil. *General Pharmacology* **31** 573–578.
- Pithon-Curi TC, Melo MP, Azeveda RB, Zorn TMT & Curi R 1997 Glutamine utilization by rat neutrophils: presence of phosphate-dependent glutaminase. *American Physiology Society* **273** C1124–C1129.
- Pithon-Curi TC, Melo MP, Palanch AC, Miyasaka CK & Curi R 1998 Percentage of phagocytosis, production of O<sub>2</sub>, H<sub>2</sub>O<sub>2</sub>, and NO, and antioxidant enzyme activities of rat neutrophils in culture. *Cell Biochemistry and Function* **16** 43–49.
- Pithon-Curi TC, Levada AC, Lopes LR, Doi SQ & Curi R 2002 Evidence that glutamine is involved in neutrophil function. *Cell Biochemistry and Function* **20** 81–86.
- Pithon-Curi TC, Schumacher RI, Freitas JJS, Lagranha CJ, Newsholme P, Palanch AC, Doi SQ & Curi R 2003 Glutamine delays spontaneous apoptosis in neutrophils. *American Journal of Physiology* **284** C1355–1361.
- Rhee SG, Chang TS, Bae YS, Lee SR & Kang SW 2003 Cellular regulation by hydrogen peroxide. *Journal of the American Society of Nephrology* **14** S211–S215.
- Roos D, van Zwieten R, Wijnen JT, Gomez-Gallego F, de Boer M, Stevens D, Pronk-Admiraal CJ, de Rijk T, van Noorden CJ, Weening RS *et al.* 1999 Molecular basis and enzymatic properties of glucose 6-phosphate dehydrogenase volendam, leading to chronic nonspherocytic anemia, granulocyte dysfunction, and increased susceptibility to infections. *Blood* **94** 2955–2962.
- Sagone AL JR, Greenwald J, Kraut EH, Bianchine J & Singh D 1983 Glucose: a role as a free radical scavenger in biological systems. *Journal of Laboratory and Clinical Medicine* **101** 97–104.
- Sugden PH & Newsholme EA 1975 Activities of citrate synthase, NAD<sup>+</sup>-linked and NADP<sup>+</sup>-linked isocitrate dehydrogenase, glutamate dehydrogenase, aspartate aminotransferase and alanine aminotransferase in nervous tissues from vertebrates and invertebrates. *Biochemistry Journal* **150** 105–111.
- Tan JS, Anderson JL, Watanakunakorn C & Phair JP 1974 Neutrophil dysfunction in diabetes mellitus. *Clinical Experiments* **85** 26–33.
- Tauber AI, Wright J, Higson FK, Edelman SA & Waxman DJ 1985 Purification and characterization of the human neutrophil NADH-cytochrome b5 reductase. *Blood* **66** 673–678.
- Walrand S, Guillet C, Boirie Y & Vasson M-P 2004 In vivo evidences that insulin regulates human polymorphonuclear neutrophil functions. *Journal of Leukocyte Biology* **76** 1104–1110.
- Watanabe A 1992 The production of hydrogen peroxide by neutrophilic polymorphonuclear leukocytes in patients with non-insulin dependent diabetes mellitus *Nippon Jinzo Gakkai Shi* **34** 1219–1225.
- Wegener G & Krause U 2002 Different modes of activating phosphofructokinase, a key regulatory enzyme of glycolysis, in working vertebrate muscle. *Biochemistry Society Transactions* **30** 264–270.
- Wilson RM & Reeves WG 1986 Neutrophil phagocytosis and killing in insulin-dependent diabetes. *Clinical and Experimental Immunology* **63** 478–484.
- Wilson RM, Clarke P, Barkes H, Heller SR & Tattersall RB 1986 Starting insulin treatment as an outpatient. Report of 100 consecutive patients followed up for at least one year. *Journal of the American Medical Association* **256** 877–880.
- Windmueller HG & Spaeth AE 1974 Uptake and metabolism of plasma glutamine by the small intestine. *Journal of Biological Chemistry* **249** 5070–5079.
- Wood HG, Katz J & Landau BR 1963 Estimation of pathways of carbohydrate metabolism. *Biochemistry Journal* **338** 809–847.
- Zozulinska DA, Wierusz-Wysocka B, Wysocki H, Majchrzak AE & Wykretowicz A 1996 The influence of insulin-dependent diabetes mellitus (IDDM) duration on superoxide anion and hydrogen peroxide production by polymorphonuclear neutrophils. *Diabetes Research and Clinical Practices* **33** 139–144.

Received 17 October 2005

Accepted 10 November 2005